## PROTOCOLE OPERATOIRE / OPERATIVE REPORT

Site: Hopital Charles-LeMoyne Date: 2024-09-30 06:14

Anesthésiste / Anesthetist: Dr. Omar Fahmy Chirurgien / Surgeon: Dr. Marie-Claire Dubois

Assistant(s): Dr. resident Marc Gagnon

Diagnostic préopératoire / Pre-operative diagnosis:

APPENDICITIS WITH INTUSSUSCEPTION.

Diagnostic postopératoire / Post-operative diagnosis:

APPENDICITIS WITH PELVIC ABSCESS.

Opération / Operation:

LAPAROSCOPIC APPENDECTOMY WITH OMENTAL WRAPPING.

Tissu envoyé en pathologie / Tissue sent to pathology: Appendicolith

Anesthésie / Anesthesia: Total intravenous anesthesia

## Historique et constatations opératoires / History and operative findings:

A 12-year-old female who presented with persistent vomiting and abdominal pain. Initially evaluated 4 days prior and diagnosed with IBD. Now has normal WBC, elevated CRP, no fever. Imaging: MRI demonstrating RLQ inflammation. Recent travel history may be relevant.

## Procédure(s) opératoire(s) / Operative procedure(s):

Patient in supine position. General endotracheal anesthesia administered. Time-out was performed and abdomen prepped in sterile fashion. Transverse infraumbilical incision is performed and access gained via blunt dissection. Single-incision laparoscopic port is used. No iatrogenic injuries occurred during trocar placement. Upon entering the abdominal cavity, minimal adhesions noted. The appendix was phlegmonous with mild inflammation. A moderate amount of purulent material was present in the pelvis. No abnormal adhesions found. We proceed with careful dissection of the appendiceal attachments. We dissect the mesentery of the appendix and use the electrocautery to coagulate the artery. The appendix is ligated at its base with two absorbable sutures and then amputated. Specimen placed in EndoCatch bag for removal. Thorough irrigation of the abdominal cavity is performed, removing all purulent material. The umbilical fascia is reapproximated with interrupted PDS 2-0 sutures. Skin incisions are closed with non-absorbable Prolene 4-0.

The patient will be continued on IV antibiotics and advanced to diet as tolerated. The patient tolerated the procedure well with minimal blood loss.

Case ID: CASE-ICMTNY-11052