

# STAT 105 Exam I

## Reference Sheet

### Numeric Summaries

mean	$\bar{x} = \frac{1}{n} \sum_{i=1}^n x_i$
population variance	$\sigma^2 = \frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2$
population standard deviation	$\sigma = \sqrt{\frac{1}{n} \sum_{i=1}^n (x_i - \bar{x})^2}$
sample variance	$s^2 = \frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2$
sample standard deviation	$s = \sqrt{\frac{1}{n-1} \sum_{i=1}^n (x_i - \bar{x})^2}$

**Quantile Function**  $Q(p)$  For a dataset consisting of  $n$  values that are ordered so that  $x_1 \leq x_2 \leq \dots \leq x_n$  and value  $p$  where  $0 \leq p \leq 1$ , let  $i = \lfloor n \cdot p + 0.5 \rfloor$ . Then the quantile function at  $p$  is:

$$Q(p) = x_i + (n \cdot p + 0.5 - i)(x_{i+1} - x_i)$$

### Linear Relationships

Form	$y \approx \beta_0 + \beta_1 x$
Fitted linear relationship	$\hat{y} = b_0 + b_1 x$
Least squares estimates	$b_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2}$ $b_1 = \frac{\sum_{i=1}^n x_i y_i - n \bar{x} \bar{y}}{\sum_{i=1}^n x_i^2 - n \bar{x}^2}$ $b_0 = \bar{y} - b_1 \bar{x}$
Residuals	$e_i = y_i - \hat{y}_i$
sample correlation coefficient	$r = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum_{i=1}^n (x_i - \bar{x})^2 \sum_{i=1}^n (y_i - \bar{y})^2}}$ $r = \frac{\sum_{i=1}^n x_i y_i - n \bar{x} \bar{y}}{\sqrt{(\sum_{i=1}^n x_i^2 - n \bar{x}^2)(\sum_{i=1}^n y_i^2 - n \bar{y}^2)}}$
coefficient of determination	$R^2 = (r)^2$ $\frac{\sum_{i=1}^n (y_i - \bar{y})^2 - \sum_{i=1}^n (y_i - \hat{y}_i)^2}{\sum_{i=1}^n (y_i - \bar{y})^2}$

### Factorial Analysis (Two Factors)

Assuming

- Factor A with levels  $1, 2, \dots, I$ ,
- Factor B with levels  $1, 2, \dots, J$ ,

- $n$  is the total number of observations,
- $n_{ij}$  is the total number of observations with Factor A at level  $i$  and Factor B at level  $j$ ,
- $n_{i\cdot}$  is the total number of observations with Factor A at level  $i$ ,
- $n_{\cdot j}$  is the total number of observations with Factor B at level  $j$ .
- $y_{ijk}$  is the  $k$ th observation where Factor A is at level  $i$  and Factor B is at level  $j$ .

$$y_{\cdot\cdot} = \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K y_{ijk} \quad \bar{y}_{\cdot\cdot} = \frac{1}{n} \sum_{i=1}^I \sum_{j=1}^J \sum_{k=1}^K y_{ijk}$$

$$\bar{y}_{i\cdot} = \frac{1}{n_{i\cdot}} \sum_{j=1}^J \sum_{k=1}^K y_{ijk} \quad \bar{y}_{\cdot j} = \frac{1}{n_{\cdot j}} \sum_{i=1}^I \sum_{k=1}^K y_{ijk}$$

$$\text{Main effect of Factor A at level } i \quad a_i = \bar{y}_{i\cdot} - \bar{y}_{\cdot\cdot}$$

$$\text{Main effect of Factor B at level } j \quad b_j = \bar{y}_{\cdot j} - \bar{y}_{\cdot\cdot}$$

$$\text{Fitted Value} \quad \hat{y}_{ij} = a_i + b_j + \bar{y}_{\cdot\cdot}$$

### Discrete Random Variables

Probability function	$P[X = x] = f_X(x)$
Cumulative probability function	$P[X \leq x] = F_X(x)$
Expected Value	$\mu = E(X) = \sum_x x f_X(x)$
Variance	$\sigma^2 = Var(X) = \sum_x (x - \mu)^2 f_X(x)$
Standard Deviation	$\sigma = \sqrt{Var(X)}$

### Joint Distributions and Related Distributions

Joint Probability Function	$P[X = x, Y = y] = f(x, y)$
Marginal Probability Function	$P[X = x] = f_X(x) = \sum_{\text{all } y} f(x, y)$ $P[Y = y] = f_Y(y) = \sum_{\text{all } x} f(x, y)$
Conditional Probability Function	$P[X = x Y = y] = \frac{f(x, y)}{f_Y(y)}$ $P[Y = y X = x] = \frac{f(x, y)}{f_X(x)}$

### Geometric Random Variables

$X$  is the trial count upon which the first successful outcome is observed performing independent trials with probability of success  $p$ .

Possible Values	$x = 1, 2, 3, \dots$
Probability function	$P[X = x] = f_X(x) = p(1 - p)^{x-1}$
Expected Value	$\mu = E(X) = \frac{1}{p}$
Variance	$\sigma^2 = Var(X) = \frac{1-p}{p^2}$

### Binomial Random Variables

$X$  is the number of successful outcomes observed in  $n$  independent trials with probability of success  $p$ .

Possible Values	$x = 0, 1, 2, \dots, n$
Probability function	$P[X = x] = f_X(x) = \frac{n!}{(n-x)!x!} p^x (1-p)^{n-x}$
Expected Value	$\mu = E(X) = np$
Variance	$\sigma^2 = Var(X) = np(1-p)$

### Continuous Random Variables

Probability density function	$P[a \leq X \leq b] = \int_a^b f_X(x) dx$
Cumulative probability function	$P[X \leq x] = F_X(x) = \int_{-\infty}^x f_X(t) dt$
Expected Value	$\mu = E(X) = \int_{-\infty}^{\infty} x f_X(x) dx$
Variance	$\sigma^2 = Var(X) = \int_{-\infty}^{\infty} (x - \mu)^2 f_X(x) dx$
Standard Deviation	$\sigma = \sqrt{Var(X)}$

### Normal Random Variables

Let  $X$  be a normal random variable with mean  $\mu$  and variance  $\sigma^2$ .

Probability density function	$f_X(x) = \frac{1}{\sqrt{2\pi\sigma^2}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$
Expected Value	$E(X) = \mu$
Variance	$Var(X) = \sigma^2$

### Standard Normal Random Variables ( $Z$ )

A normal random variable with mean 0 and variance  $\sigma^2$ .  
If  $X$  is normal( $\mu, \sigma^2$ ) then  $P[a \leq X \leq b] = P\left[\frac{a-\mu}{\sigma} \leq Z \leq \frac{b-\mu}{\sigma}\right]$

Probability density function  $f_X(x) = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}x^2}$

### Functions of random variables

For  $X_1, X_2, \dots, X_n$  independent random variables and  $a_0, a_1, a_2, \dots, a_n$  constants if  $W = a_0 + a_1X_1 + \dots + a_nX_n$ :

- $E(W) = a_0 + a_1E(X_1) + a_2E(X_2) + \dots + a_nE(X_n)$
- $Var(W) = a_1^2Var(X_1) + a_2^2Var(X_2) + \dots + a_n^2Var(X_n)$

### Confidence Intervals and Hypothesis Tests

#### Confidence Intervals $n \geq 25$

$(1 - \alpha) \cdot 100\%$ Confidence interval for population mean	$\bar{x} \pm z_{1-\alpha/2} \sqrt{\frac{\sigma^2}{n}}$
$(1 - \alpha) \cdot 100\%$ Confidence lower bound	$\bar{x} - z_{1-\alpha} \sqrt{\frac{\sigma^2}{n}}$
$(1 - \alpha) \cdot 100\%$ Confidence upper bound	$\bar{x} + z_{1-\alpha} \sqrt{\frac{\sigma^2}{n}}$

#### Confidence Intervals $n < 25$

$(1 - \alpha) \cdot 100\%$ Confidence interval for population mean	$\bar{x} \pm t_{1-\alpha/2, n-1} \sqrt{\frac{\sigma^2}{n}}$
$(1 - \alpha) \cdot 100\%$ Confidence lower bound	$\bar{x} - t_{1-\alpha, n-1} \sqrt{\frac{\sigma^2}{n}}$
$(1 - \alpha) \cdot 100\%$ Confidence upper bound	$\bar{x} + t_{1-\alpha, n-1} \sqrt{\frac{\sigma^2}{n}}$

### Test statistics in hypothesis tests for population mean

$n \geq 25$	$\frac{\bar{x} - \mu}{\sqrt{\sigma^2/n}} \sim N(0, 1)$
$n < 25$	$\frac{\bar{x} - \mu}{\sqrt{\sigma^2/n}} \sim t$ with $\nu = n - 1$ degrees of freedom