

Recall:

of experiments probability

- Binomial distribution of

$X \sim \text{binomial}(n, p)$

of
success.

$X :=$ The number of successes out of "n" Bernoulli

trials.

$$x = 0, 1, 2, \dots, n$$

- each trial is independent of the other trials
- The probability of success, p , is the same over all n trials.
- NO closed form CDF. (e.g. $P(X \leq 4) = P(X=0 \text{ or } X=1 \text{ or } X=2 \text{ or } X=3 \text{ or } X=4)$)
- $E(X) = n \cdot p$
- $\text{Var}(X) = n p(1-p)$
- $SD(X) = \sqrt{\text{Var}(X)} = \sqrt{np(1-p)}$

- Geometric distribution ; $X \sim \text{Geom}(p)$

$X :=$ The number of trials until observing
the first success. $X = 1, 2, 3, \dots$

- Each trial is independent of others.

- The prob. of success, p , is the same for all trials.

$$(\text{e.g. } P(X \leq 20) = 1 - (1-p)^{20})$$

← - closed form CDF : $F_X(x) = 1 - (1-p)^x$

$$- E[X] = \frac{1-p}{p}$$

$$- \text{Var}[X] = \frac{1-p}{p^2}$$

$$- \text{SD}[X] = \sqrt{\text{Var}[X]} = \sqrt{\frac{1-p}{p^2}}$$

Mean

and

Variance

of Geometric Distribution

Common Distributions

Background

Bernoulli

Binomial

Geometric

The Geometric Distribution

$$X \sim \text{Geom.}(p)$$

Expected value:

$$E(X) = \frac{1}{p}$$

Variance:

$$\text{Var}(X) = \frac{1-p}{p^2}$$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Example

NiCad batteries: An experimental program was successful in reducing the percentage of manufactured NiCad cells with internal shorts to around 1%. Let T be the test number at which the first short is discovered. Then, $T \sim \text{Geom}(p)$.

Calculate

$$T \sim \text{Geom}(0.01), f(x) = P(X=x) = p(1-p)^{x-1} \quad x=1, 2, 3, \dots$$

- $P(\underline{\text{1st}} \text{ or } \underline{\text{2nd}} \text{ cell tested has the 1st short})$

$$P(T=1 \text{ or } T=2) = P(T=1) + P(T=2)$$

$$= P(1) + P(2) = p(1-p)^{1-1} + p(1-p)^{2-1} = (0.01)(0.99)^0 + (0.01)(0.99)^1 = 0.02$$

- $P(\text{at least 50 cells tested without finding a short})$

$$\underline{P(T > 50)} = 1 - P(T \leq 50)$$

$$< 1 - F_x(50) = 1 - \left[1 - (1 - 0.01)^{50} \right]$$

$$= (1 - 0.01)^{50} = (0.99)^{50} = 0.61$$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Example

NiCad batteries:

$$T \sim \text{Geom}(p=0.01)$$

Calculate the expected test number at which the first short is discovered and the variance in test numbers at which the first short is discovered.

$$E(T) = \frac{1}{p} = \frac{1}{0.01} = 100$$

(on average, you need to test 100 batteries until you observe the first short)

$$\text{Var}(T) = \frac{1-p}{p^2} = \frac{1-0.01}{(0.01)^2} = \frac{0.99}{(0.01)^2} = 9900$$

$$\text{SD}(T) = \sqrt{\text{Var}(T)} = \sqrt{9900}$$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Example

A shipment of 200 widgets arrives from a new widget distributor. The distributor has claimed that the widgets there is only a 10% defective rate on the widgets. Let X be the random variable associated with the number of trials until finding the first defective widgets.

- What is the probability distribution associated with this random variable X ? Precisely specify the parameter(s).

$$X \sim \text{Geom}(P=0.1)$$

- How many widgets would you expect to test before finding the first defective widget?

$$E X = \frac{1}{P} = \frac{1}{0.1} = 10$$

So, we expect to test 10 widgets (on average) until finding the first defective one.

Common Distributions

Background

Bernoulli

Binomial

Geometric

Example

You find your first defective widget while testing the third widget.

- What is the probability that the first defective widget would be found on the third test if there are only 10% defective widgets from in the shipment?

$$\begin{aligned} P(X = 3) &= p(1 - p)^{x-1} \\ &= 0.1(1 - 0.1)^{3-1} \\ &= 0.1(0.9)^2 = 0.081 \end{aligned}$$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Example

- What is the probability that the first defective widget would be found **by** the third test if there are only **10%** defective widgets from in the shipment?

$$P(\underbrace{x \leq 3}) = F_X(3) = 1 - (1 - p)^3$$
$$= 1 - (1 - .1)^3$$

$$= 1 - (0.9)^3 = \boxed{0.271}$$

$$\text{in Geom. : } F_x(x) = 1 - (1 - p)^x$$

$$\text{in Geom: Pmf : } f_x(x) := P_x(x) = p(1-p)^{x-1}, \quad x=1, 2, 3, \dots$$

The Poisson Distribution

Common Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

The Poisson Distribution

Origin: A rare occurrence is watched for over a specified interval of time or space.

It's often important to keep track of the total number of occurrences of some relatively rare phenomenon.

Definition

Consider a variable

X : the count of occurrences of a phenomenon across a specified interval of time or space

or

X: the number of times the rare occurrence is observed

Common
Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

The Poisson Distribution

probability function:

(λ)

The ~~Poisson (λ)~~ distribution is a discrete probability distribution with pmf

$$f(x) = \begin{cases} \frac{e^{-\lambda} \lambda^x}{x!} & x = 0, 1, \dots \\ 0 & \text{o. w.} \end{cases}$$

For $\lambda > 0$

\nwarrow parameter.
 $\text{Poisson}(\lambda)$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

The Poisson Distribution

These occurrences must:

- be independent
- be sequential in time (no two occurrences at once)
- occur at the same constant rate λ

λ the *rate parameter*, is the expected number of occurrences in **the specified interval of time or space** (i.e $E(X) = \lambda$)

Common Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

The Poisson Distribution

Examples that could follow a Poisson\$(\lambda)\$ distribution :

Y is the number of shark attacks off the coast of CA next **year**, $\lambda = 100$ attacks per **year**

Z is the number of shark attacks off the coast of CA **next month**, $\lambda = 100 / 12$ attacks per month

N is the **number** of α -particles emitted from a small bar of **polonium** registered by a counter in a minute, $\lambda = 459.21$ particles per **minute**

J is the number of particles **per hour**,
 $\lambda = 459.21 * 60 = 27,552.6$ particles per **hour**.

Common Distributions

Background


Bernoulli

Binomial


Geometric

Poisson

The Poisson Distribution



Right skewed with peak near λ



Common Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

The Poisson Distribution

(λ)

For X a Poisson(λ) random variable,

$$\mu = EX = \sum_{x=0}^{\infty} x \frac{e^{-\lambda} \lambda^x}{x!} = \lambda$$

$$\sigma^2 = \text{Var } X = \sum_{x=0}^{\infty} (x - \lambda)^2 \frac{e^{-\lambda} \lambda^x}{x!} = \lambda$$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

Example

Arrivals at the library

Some students' data indicate that between 12:00 and 12:10pm on Monday through Wednesday, an average of around 125 students entered Parks Library at ISU. Consider modeling

M: the number of students entering the ISU library between 12:00 and 12:01pm next Tuesday

Model $M \sim \text{Poisson}(\lambda)$. What would a reasonable choice of λ be?

$$\lambda = \frac{125}{10}$$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

Example

Arrivals at the library

Under this model, the probability that between 10 and 15 students arrive at the library between 12:00 and 12:01 PM is:

$$M \sim \text{Poisson}(\lambda = \frac{12.5}{10})$$

$$P_M(m) = \frac{e^{-\lambda} \lambda^m}{m!}, m=0, 1, 2, \dots$$

$$P(10 \leq M \leq 15) = P(10) + P(11) + \dots + P(15)$$

$$= \frac{e^{-12.5} (12.5)^{10}}{10!} + \frac{e^{-12.5} (12.5)^{11}}{11!} + \dots + \frac{e^{-12.5} (12.5)^{15}}{15!}$$

Common
Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

$$= 6.6$$

Shark attacks

No Closed Form Po
CDF in Poisson!

Let X be the number of unprovoked shark attacks that will occur off the coast of Florida next year. Model

$$X \sim \text{Poisson}(\lambda).$$

From the shark data at

<http://www.flmnh.ufl.edu/fish/sharks/statistics/FLactivity.html>,
246 unprovoked shark attacks occurred from 2000 to 2009.

What would a reasonable choice of λ be?

246 attacks
per 10 years.

The rate (λ) for next year (only)

one year)

$$\lambda = \frac{246}{10} = 24.6$$

Common Distributions

Background

Bernoulli

Binomial

Geometric

Poisson

Shark attacks

Under this model, calculate the following:

- $P(\text{no attacks next year})$

$$P(X=0) = P(0) = \frac{e^{-24.6} (24.6)^0}{0!} = e^{-24.6} = 2.07 \times 10^{-11}$$

- $P(\text{at least 5 attacks})$

$$P(X \geq 5) = 1 - P(X \leq 4) = 1 - P(X \leq 4)$$

$$= 1 - [P(X=0) + P(X=1) + P(X=2) + P(X=3) + P(X=4)]$$

$$= 1 - [P(X=0) + P(X=1) + \dots + P(X=4)]$$

- $P(\text{more than 10 attacks})$

$$P(X \geq 10) = 1 - P(X \leq 9)$$

$$1 - \left[e^{-24.6} \frac{(24.6)^0}{0!} + \frac{e^{-24.6}}{1!} \frac{(24.6)^1}{1!} + \dots \right]$$

$$+ \frac{e^{-24.6}}{4!} \frac{(24.6)^4}{4!} \right]$$

$$= 1 - e^{-24.6} (36266812)$$

$$= 0.999249$$
