

Banach Spaces of Analytic Functions

Ashish Kujur

Last Updated: January 30, 2023

Contents

§1 Analytic and Harmonic Functions	2
§1.1 The Cauchy and Poisson Kernels	2
§1.2 Boundary Values	2
§1.2.1 Weak* convergence of measures	2
§1.2.2 Convergence in norm	2
§1.2.3 Weak* convergence of bounded functions	2
§1.2.4 The entire picture!	3
§1.3 Fatou's Theorem	3
§1.4 Hardy Spaces – H^p spaces	5
§1.4.1 Series Representation of Harmonic Functions	6
§2 The space H^1	7
§2.1 Brief Recap!	7
§2.2 The Helson-Lowdenslager Approach	7
§2.3 Szegő's Theorem	11

§1 Analytic and Harmonic Functions

§1.1 The Cauchy and Poisson Kernels

Proposition §1.1.1. *Let $u : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ be a harmonic function. Then we have that*

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{it}) P_r(e^{i(\theta-t)}) dt \quad (\S 1.1.1)$$

§1.2 Boundary Values

§1.2.1 Weak* convergence of measures

Theorem §1.2.1. *Let $\{\varphi_i\}_i$ be an approximate identity on \mathbb{T} and let $\mu \in \mathcal{M}(\mathbb{T})$. Then for all i , $\varphi_i * \mu \in L^1(\mathbb{T})$ with*

$$\|\varphi_i * \mu\|_1 \leq C_\varphi \|\mu\|$$

and

$$\|\mu\| \leq \sup_i \|\varphi_i * \mu\|_1.$$

Moreover, the measures $d\mu_i = (\varphi_i * \mu)(e^{it}) dt/2\pi$ converge to $d\mu(e^{it})$ in the weak* topology, i.e.

$$\lim_i \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) (\varphi_i * \mu)(e^{it}) dt = \int_{\mathbb{T}} \varphi(e^{it}) d\mu(e^{it})$$

for all $f \in \mathcal{C}(\mathbb{T})$.

§1.2.2 Convergence in norm

Theorem §1.2.2. *Let $\{\varphi_i\}_i$ be an approximate identity on \mathbb{T} and let $f \in L^p(\mathbb{T})$ with $p \in [1, \infty)$. Then for all i , $\varphi_i * f \in L^p(\mathbb{T})$ with*

$$\|\varphi_i * f\|_p \leq C_\varphi \|f\|_p$$

and

$$\lim_i \|\varphi_i * f - f\|_p = 0.$$

§1.2.3 Weak* convergence of bounded functions

Theorem §1.2.3. *Let $\{\varphi_i\}_i$ be an approximate identity on \mathbb{T} and let $f \in L^\infty(\mathbb{T})$. Then for all i , $\varphi_i * \mu \in \mathcal{C}(\mathbb{T})$ with*

$$\|\varphi_i * \mu\|_\infty \leq C_\varphi \|\mu\|_\infty$$

and

$$\|f\|_{+\infty} \leq \sup_i \|\varphi_i * f\|_\infty.$$

Moreover, $\varphi_i * f$ converge to f in the weak* topology, i.e.

$$\lim_i \int_{-\pi}^{\pi} g(e^{it}) (\varphi_i * f)(e^{it}) dt = \int_{\mathbb{T}} g(e^{it}) f(e^{it}) dt$$

for all $g \in L^1(\mathbb{T})$.

§1.2.4 The entire picture!

Definition §1.2.4 (Poisson integral of some function or measure). Let $\tilde{f} : \mathbb{D} \rightarrow \mathbb{C}$ be a harmonic function. Then \tilde{f} is said to be the *Poisson integral* of the function $f : \mathbb{T} \rightarrow \mathbb{C}$ if

$$\tilde{f}(re^{i\theta}) = \frac{1}{2\pi} \int_{\mathbb{T}} f(e^{it}) P_r(e^{i(\theta-t)}) dt$$

In such a case, we will denote the function \tilde{f} by $P[f]$. Similarly, f is said to be the *Poisson integral* of a complex measure μ on T if

$$\tilde{f}(re^{i\theta}) = \frac{1}{2\pi} \int_{\mathbb{T}} P_r(e^{i(\theta-t)}) d\mu(e^{it})$$

In such a case, we will denote the function \tilde{f} by $P[\mu]$.

Theorem §1.2.5 (Ultimate Convergence). Let $f : \mathbb{D} \rightarrow \mathbb{C}$ be a harmonic function. Define for each $r \in [0, 1)$, the function $f_r : \mathbb{T} \rightarrow \mathbb{C}$ by

$$f_r(e^{i\theta}) = f(re^{i\theta})$$

The following statements holds:

1. If $1 < p \leq \infty$ then $f = P[g]$ for some $g \in L^p[\mathbb{T}]$ iff for each $r > 0$, $\|f_r\|_p < +\infty$.
2. If $p=1$ then $f = P[g]$ for some $g \in L^1[\mathbb{T}]$ iff f_r converge in the L^1 norm.
3. $f = P[\mu]$ for some $\mu \in \mathcal{M}(\mathbb{T})$ iff for each $r > 0$, $\|f_r\|_1 < +\infty$

§1.3 Fatou's Theorem

Theorem §1.3.1. Let μ be a complex measure on the unit circle \mathbb{T} , and let $f : \mathbb{D} \rightarrow \mathbb{C}$ be the harmonic function defined by

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_{\mathbb{T}} P_r(e^{i(\theta-t)}) d\mu(e^{it})$$

Let $e^{i\theta_0}$ be any point where μ is differentiable with respect to the normalised Lebesgue measure. Then

$$\lim_{r \rightarrow 1} f(re^{i\theta_0}) = \left(\frac{d\mu}{d\theta} \right) (e^{i\theta_0}) = \mu' (e^{i\theta_0})$$

In fact, $f(re^{i\theta}) \rightarrow \mu' (e^{i\theta_0})$ as $re^{i\theta}$ approaches $e^{i\theta_0}$ along any path in the open disc within the region of the form $|\theta - \theta_0| \leq c(1 - r)$ for some $c > 0$.

Corollary §1.3.2. *Let μ be a complex measure on \mathbb{T} . Then $P[\mu]$ has nontangential limits equal everywhere to the Radon Nikodym derivative of μ with respect to the normalised Lebesgue measure.*

Corollary §1.3.3. *Let $f : \mathbb{T} \rightarrow \mathbb{C}$ be L^1 . Then $P[f]$ has nontangential limits at almost everywhere and these limits equal to f almost everywhere.*

Corollary §1.3.4. *Let $f : \mathbb{D} \rightarrow \mathbb{C}$ be a harmonic function and $1 \leq p < \infty$. Suppose that for all $0 \leq r < 1$, we have that*

$$\|f_r\|_p < +\infty$$

Then for almost every θ the radial limits

$$\tilde{f}(e^{i\theta}) = \lim_{r \rightarrow 1} f(re^{i\theta})$$

exist and define a function \tilde{f} in $L^p(\mathbb{T})$. The following also holds:

1. *If $p > 1$ then $f = P[\tilde{f}]$.*
2. *If $p = 1$ then $f = P[\mu]$ for some complex measure μ whose absolutely continuous part is $f d\theta$.*
3. *If f is bounded then the boundary values exist almost everywhere and define a bounded measurable function \tilde{f} on \mathbb{T} such that $f = P[\tilde{f}]$.*

Proof. Suppose that for each $r \in [0, 1)$, we have $\|f_r\|_p < +\infty$. We need to prove that for almost every θ , $\lim_{r \rightarrow 1} f(re^{i\theta})$ exists. Then by Theorem §1.2.5, we have that $f = P[g]$ for some $g \in L^p(\mathbb{T})$. Since $L^p(\mathbb{T}) \subset L^1(\mathbb{T})$, we can use the previous corollary. By the previous corollary, we have that $P[g]$ has nontangential limits almost everywhere, we have that

$$\tilde{f}(e^{i\theta}) = \lim_{r \rightarrow 1} f(re^{i\theta}) = \lim_{r \rightarrow 1} P[g](re^{i\theta}) \quad (\S 1.3.1)$$

exists almost everywhere.

Now we proceed to prove part (1). Also by Theorem §1.2.5, we have that $f = P[g]$ for some $g \in L^p(\mathbb{T})$. Hence, we have that by Equation §1.3.1 that $\tilde{f}(e^{i\theta}) = \lim_{r \rightarrow 1} P[g](re^{i\theta})$ holds at almost every θ .

Also, by the previous corollary, $\lim_{r \rightarrow 1} P[g](re^{i\theta}) = g(e^{i\theta})$ for almost every θ . Hence, we have that $\tilde{f} = g$.

□

Corollary §1.3.5. *Let $f : \mathbb{D} \rightarrow \mathbb{R}_{\geq 0}$ be a harmonic function. Then f has nontangential limits at almost every point of \mathbb{T} . (Why demand nonnegative?)*

Let $h(\mathbb{D})$ denote the set of all harmonic functions on \mathbb{D} . Let $p \in [1, \infty]$. Define

$$h^p(\mathbb{D}) = \{f \in h(\mathbb{D}) \mid \{f_r\}_{0 \leq r < 1} \text{ is uniformly bounded in } L^p \text{ norm} \}$$

We define a norm on $h^p(\mathbb{D})$ by

$$\|f\|_{h^p(\mathbb{D})} = \sup_{0 \leq r < 1} \|f_r\|_{L^p(\mathbb{D})} = \begin{cases} \sup_{0 \leq r < 1} \left(\frac{1}{2\pi} \int_{-\pi}^{\pi} |f(re^{i\theta})|^p d\theta \right)^{\frac{1}{p}} & \text{if } p \in [1, \infty) \\ \sup_{0 \leq r < 1} \|f(re^{i\theta})\|_{L^\infty(\mathbb{D})} & \text{if } p = \infty \end{cases}$$

It is easy to see why $\|f\| < +\infty$ for any $f \in h^p(D)$. So we now proceed to show that $h^p(D)$ is a Banach space with this norm. First we show that it is indeed a normed linear space.

Clearly, $h(\mathbb{D})$ is a vector space. To show that $h^p(\mathbb{D})$ is a vector space, it suffices to check that $h^p(\mathbb{D})$ is a subspace.

Let $f, g \in h^p(\mathbb{D})$ and let $\alpha \in \mathbb{C}$. Then for any $r \in [0, 1)$, we have that

$$\begin{aligned} \|(f + \alpha g)_r\|_p &= \|f_r + \alpha g_r\| \\ &= \|f_r\|_p + \alpha \|g_r\|_p \end{aligned}$$

Take note of the use of Holder's inequality. After this is done, since $\{f_r\}_{r \in [0, 1)}$ and $\{g_r\}_{r \in [0, 1)}$ is uniformly bounded, we have that $\{f + \alpha g\}_{r \in [0, 1)}$ is uniformly bounded in L^p norm.

Now, we need to show that it is a normed linear space but this follows almost immediately.

To show that it is a Banach space, we show that

Theorem §1.3.6. *Let $p \in [1, \infty]$. If $u \in L^p(\mathbb{T})$ then $f = P * u \in h^p(\mathbb{D})$ and $\|f\|_p = \|u\|_p$. If $\mu \in \mathcal{M}(\mathbb{T})$ then $f = P * \mu \in h^1(\mathbb{D})$ and $\|f\|_1 = \|\mu\|$.*

Proof. We consider the case $p \in [1, \infty)$. The other cases can be dealt similarly. Consider the map

$$u \mapsto U$$

where $U(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} P_r(e^{i(\theta-t)}) u(e^{it}) dt$. By Theorem §1.2.2, we have that $\|U\| = \|u\|_p < +\infty$. Hence $U \in h^p(\mathbb{D})$.

Linearity is obvious. We need to check injectivity and surjectivity.

To check injectivity, let $u \in L^p(\mathbb{T})$ and suppose that $T(u) = P[u] = 0$. Now $\lim_{r \rightarrow 1} P[u](re^{i\theta}) = u$ for almost θ by Corollary §1.3.3 and hence $u = 0$ almost everywhere.

Surjectivity is clear from Theorem §1.2.5. □

§1.4 Hardy Spaces – H^p spaces

Let us denote the set of all analytic functions on \mathbb{D} by $H(\mathbb{D})$. Hence, $H(\mathbb{D}) \subset h(\mathbb{D})$. For $p \in (0, \infty]$, we consider the *Hardy classes* of analytic functions on the unit disc

$$H^p(\mathbb{D}) = \left\{ F \in H(\mathbb{D}) \mid \|F\|_p < \infty \right\}$$

Clearly,

$$H^p(\mathbb{D}) \subset h^p(\mathbb{D})$$

We will see that $H^p(\mathbb{D})$, $1 \leq p \leq +\infty$, is also a Banach spaces isomorphic to a closed subspace of $L^p(\mathbb{T})$ denoted by $H^p(\mathbb{T})$.

To prove that $H^p(\mathbb{D})$ is a closed subspace of $h^p(\mathbb{D})$, we are going to identify $H^p(\mathbb{D})$ with the closed subspace

$$\left\{ u \in L^p(\mathbb{T}) : \int_{-\pi}^{\pi} u(e^{it}) e^{ikt} dt = 0 \text{ for all } k \in \mathbb{N} \right\}$$

Let $\{u_n\}$ be a sequence of functions in the above subspace; suppose that $\{u_n\}$ converge to $u \in L^p(\mathbb{T})$. Now, let $k \in \mathbb{N}$ be arbitrary. Since $\{u_n\}$ converge to u in p -norm, we have that $\{u_n\}$ converge to u in 1-norm. Hence we have the following:

$$\left| \int_{-\pi}^{\pi} u_n(e^{it}) e^{ikt} dt - \int_{-\pi}^{\pi} u(e^{it}) e^{ikt} dt \right| \leq \int_{-\pi}^{\pi} |u_n(e^{it}) - u(e^{it})| dt$$

From the above inequality, it is evident that u is in the subspace mentioned above!

§1.4.1 Series Representation of Harmonic Functions

Theorem §1.4.1. *Let U be a harmonic on the disc $D_R = \{|z| < R\}$. Then, for each $n \in \mathbb{Z}$, the quantity*

$$a_n = \frac{\rho^{-|n|}}{2\pi} \int_{-\pi}^{\pi} U(\rho e^{it}) e^{-int} dt \quad (0 < \rho < R) \quad (\S 1.4.1)$$

is independent of ρ and we have

$$U(re^{i\theta}) = \sum_{n=-\infty}^{\infty} a_n r^{|n|} e^{in\theta} \quad (re^{i\theta} \in \mathbb{D}) \quad (\S 1.4.2)$$

The function

$$V(re^{i\theta}) = \sum_{n=-\infty}^{\infty} -i \operatorname{sgn}(n) a_n r^{|n|} e^{in\theta} \quad (re^{i\theta} \in \mathbb{D}) \quad (\S 1.4.3)$$

is the unique harmonic conjugate of U such that $V(0) = 0$. The series in §1.4.2 and §1.4.3 are absolutely and uniformly convergent on compact subsets of D_R

§2 The space H^1

§2.1 Brief Recap!

Theorem §2.1.1. *Let $u : \overline{\mathbb{D}} \rightarrow \mathbb{C}$ be a harmonic function. Then we have that*

$$u(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} u(e^{it}) P_r(e^{i(\theta-t)})$$

§2.2 The Helson-Lowdenslager Approach

Let $\mathcal{C}(\overline{\mathbb{D}})$ be the set of all continuous functions on $\overline{\mathbb{D}}$ and let $H(\mathbb{D})$ be the set of all holomorphic functions on the open disc \mathbb{D} . We define $\mathcal{A} = \mathcal{C}(\overline{\mathbb{D}}) \cap H(\mathbb{D})$.

We show that \mathcal{A} is a uniformly closed algebra of $\mathcal{C}(\overline{\mathbb{D}})$. Let $\{f_n\}$ be a sequence in \mathcal{A} converging uniformly to $f \in \mathcal{C}(\overline{\mathbb{D}})$.

We recall Morera's Theorem for analytic functions at this point:

Theorem §2.2.1 (Morera). *A continuous, complex valued function $f : D \rightarrow \mathbb{C}$ that satisfies $\oint_{\gamma} f(z) dz = 0$ for any closed piecewise C^1 path γ in D must be holomorphic on D .*

We use this theorem to prove what we want to prove. Now, let C be any closed curve in \mathbb{D} . Then for any $n \in \mathbb{N}$,

$$\oint_C f_n(z) dz = 0$$

So,

$$\oint_C f(z) dz = \oint_C \lim_{n \rightarrow \infty} f_n(z) dz = \lim_{n \rightarrow \infty} \oint_C f_n(z) dz = 0$$

Since C was arbitrary, f must be holomorphic. This shows that \mathcal{A} is uniformly closed. The fact that it is an algebra is easy to check ✓.

Now, note that since \mathbb{D} is a compact metric space, we have that $\mathcal{C}(\mathbb{D})$ is a complete metric space with supremum metric. Since the supremum metric can also be induced by a norm, namely the supremum norm, we have that $\mathcal{C}(\mathbb{D})$ is a Banach space with the supremum norm.

Thus, this is what we have proved so far:

Theorem §2.2.2. *The disc algebra $\mathcal{A} = \mathcal{C}(\overline{\mathbb{D}}) \cap H(\mathbb{D})$ is a Banach space under the sup norm*

$$\|f\|_{\infty} = \sup_{|z| \leq 1} |f(z)|$$

We make a couple of observations at this point:

1. Each $f \in \mathcal{A}$ is the Poisson integral of its boundary values:

$$f(re^{i\theta}) = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(e^{it}) P_r(e^{i(\theta-t)}) dt$$

2. It follows from the Maximum Modulus Theorem that

$$\|f\|_{\infty} = \sup |f(e^{it})|$$

Theorem §2.2.3 (Correspondence of \mathcal{A} with a closed subspace of $\mathcal{C}(\mathbb{T})$). *Consider the subspace*

$$\tilde{\mathcal{A}} = \left\{ f \in \mathcal{C}(\mathbb{T}) : \int_{-\pi}^{\pi} f(e^{it}) e^{in\theta} dt = 0 \text{ for } n = 1, 2, \dots \right\}$$

of $\mathcal{C}(\mathbb{T})$. Then there is an isomorphism of \mathcal{A} with $\tilde{\mathcal{A}}$.

Proof. First, we show that $\tilde{\mathcal{A}}$ is a closed subspace of $\mathcal{C}(\mathbb{T})$. Let $\{f_n\}$ be a sequence of functions in $\tilde{\mathcal{A}}$ converging to $f \in \mathcal{C}(\mathbb{T})$. Consider the following:

$$\begin{aligned} \left| \int_{-\pi}^{\pi} f(e^{it}) e^{ikt} dt \right| &= \left| \int_{-\pi}^{\pi} f(e^{it}) e^{ikt} dt - \int_{-\pi}^{\pi} f_n(e^{it}) e^{ikt} dt \right| \\ &= \int_{-\pi}^{\pi} |f(e^{it}) - f_n(e^{it})| dt \\ &\leq 2\pi \|f_n - f\|_{\infty} \rightarrow 0 \text{ as } n \rightarrow \infty \end{aligned}$$

This shows that $\tilde{\mathcal{A}}$ is closed under $\mathcal{C}(\mathbb{T})$ with supremum norm.

Now consider the linear map $T : \mathcal{A} \rightarrow \tilde{\mathcal{A}}$ given by

$$f \mapsto f|_{\mathbb{T}}$$

For the sake of convenience, we will write $f|_{\mathbb{T}}$ as $f_{\mathbb{T}}$. We first need to show this map is well defined! That is, we need to show that

$$\int_{-\pi}^{\pi} f_{\mathbb{T}}(e^{it}) e^{ikt} dt = 0$$

for all $k \in \mathbb{N}$ but this immediately follows from Cauchy's theorem.

Note that injectivity is clear from Theorem §2.1.1. To show surjectivity, let $f \in \tilde{\mathcal{A}}$. We need to show that there is a function $u \in \mathcal{A}$ such that $u_{\mathbb{T}} = f$. Consider the function

$$u(re^{i\theta}) = \begin{cases} (P * f)(re^{i\theta}) & \text{if } 0 \leq r < 1 \\ f(e^{i\theta}) & \text{if } r = 1 \end{cases}$$

This is the Dirichlet problem on the unit disc! So, u is continuous on $\overline{\mathbb{D}}$. It remains to show that u is analytic on \mathbb{D} . But note that for $r \in [0, 1)$,

$$\begin{aligned} u(re^{i\theta}) &= \sum_{n=-\infty}^{\infty} r^{|n|} \hat{f}(n) e^{int} \\ &= \sum_{n=0}^{\infty} r^{|n|} \hat{f}(n) e^{int} \end{aligned}$$

This completes the proof of the theorem! □

In view of the previous theorem, we will simply write $\tilde{\mathcal{A}}$ as \mathcal{A} .

Definition §2.2.4. A trigonometric polynomial p on the circle \mathbb{T} is of the form

$$p(e^{it}) = \sum_{k=0}^n a_k e^{ik\theta}$$

Proposition §2.2.5. *The set of the trigonometric polynomials is a dense subset of \mathcal{A} .*

Proof. It is clear that any trigonometric polynomial on the circle is a member of the disc algebra. Now, if $f : \mathbb{T} \rightarrow \mathbb{C}$ is in \mathcal{A} , then its negative Fourier coefficients are zero! Since, the Cesaro sum of f

$$s_n(x) = \sum_{k=-n}^{k=n} \hat{f}(n) e^{ikx} = \sum_{k=0}^n \hat{f}(n) e^{ikx}$$

converge to f uniformly and is a sequence of trigonometric polynomial, we are done! □

The following result is used in the proof of the next theorem, so, we prove it here:

Theorem §2.2.6. *The real parts of functions in \mathcal{A} are uniformly dense in $\mathcal{C}(\mathbb{T}, \mathbb{R})$. In other words, if μ is finite signed Borel measure on \mathbb{T} such that $\int f d\mu = 0$ for every $f \in \mathcal{A}$ then μ is the zero measure.*

Proof. We first show that any trigonometric polynomial of the form

$$p(e^{it}) = \sum_{k=-n}^n c_k e^{ikt} \tag{§2.2.1}$$

where $c_{-k} = \overline{c_k}$ for each $k \in \{1, \dots, n\}$ is a real part of a function $f \in \mathcal{A}$. Note that $p(e^{it})$ in Equation §2.2.1 is the real part of the function:

$$f(e^{it}) = c_0 + 2c_1 e^{it} + 2c_2 e^{2it} + \dots + 2c_n e^{int}$$

Now, we claim that every function $f \in C(\mathbb{T}, \mathbb{R})$ is a uniform limit of a trigonometric polynomial of the form §2.2.1. We will be done if we show that the negative Fourier coefficients of

real valued function is the conjugate of the its positive counterpart, that is, for each $n \in \mathbb{Z}_{\geq 0}$, we have that $\hat{f}(-n) = \overline{\hat{f}(n)}$. To show this, take any $n \in \mathbb{Z}_{\geq 0}$ and then observe that

$$\begin{aligned}\hat{f}(-n) &= \int_{\mathbb{T}} f(e^{it}) e^{int} \frac{dt}{2\pi} \\ &= \overline{\int_{\mathbb{T}} f(e^{it}) e^{-int} \frac{dt}{2\pi}} \\ &= \overline{\hat{f}(n)}\end{aligned}$$

This shows that the Cesaro means of a real valued function is a trigonometric polynomial of the form §2.2.1 and since the Cesaro means converges to f uniformly. Thus, the closure of the real parts of \mathcal{A} is indeed $\mathcal{C}(\mathbb{T}, \mathbb{R})$.

Now, let μ be a finite signed measure on \mathbb{T} such that $\int f d\mu = 0$ for every $f \in \mathcal{A}$. We show that μ is the zero measure. Notice that if $f \in \mathcal{A}$ then

$$0 = \int_{\mathbb{T}} f d\mu = \int_{\mathbb{T}} \Re(f) d\mu + i \int_{\mathbb{T}} \Im(f) d\mu$$

Hence, it follows that

$$\int_{\mathbb{T}} \Re(f) d\mu = 0$$

for every $f \in \mathcal{A}$. Now, if $g \in \mathcal{C}(\mathbb{T}, \mathbb{R})$ then by the first part of this theorem, there is a sequence $\{f_n\} \in \mathcal{A}$ such that $\Re(f_n)$ converges to g uniformly. By the Dominated Convergence Theorem (which holds, thanks to Jordan Decomposition Theorem), we have that $\int_{\mathbb{T}} g d\mu = 0$.

Now to prove that every $\mu = 0$, it suffices to show that $\hat{\mu}(n) = 0$ for every $n \in \mathbb{Z}^1$. Now, notice that for any $n \in \mathbb{Z}$, we have

$$\begin{aligned}\hat{\mu}(n) &= \int_{\mathbb{T}} e^{-int} d\mu(e^{it}) \\ &= \int_{\mathbb{T}} (\cos(nt) - i \sin(nt)) d\mu(e^{it}) \\ &= \int_{\mathbb{T}} \cos(nt) d\mu(e^{it}) - i \int_{\mathbb{T}} \sin(nt) d\mu(e^{it}) \\ &= 0\end{aligned}$$

This completes the proof! □

Corollary §2.2.7. *Let μ be a finite signed measure on \mathbb{T} such that $\int f d\mu = 0$ for every $f \in \mathcal{A}$ which vanishes at the origin then μ is a constant multiple of Lebesgue measure.*

Proof. We first prove the following claim: If $f \in \mathcal{A}$ then $\int_{\mathbb{T}} f d\mu = \frac{f(0)}{2\pi}$. Since the negative Fourier coefficients are zero and f is continuous, we have that the Cesaro means converge

¹See Page 41, Corollary 2.3 of Mashreghi.

uniformly to f , that is,

$$\sum_{k=0}^{\infty} \hat{f}(n) e^{int} \rightarrow f \text{ uniformly}$$

Thus,

$$\begin{aligned} \int_{\mathbb{T}} f(e^{it}) \frac{dt}{2\pi} &= \frac{1}{2\pi} \int_{\mathbb{T}} \left(\sum_{k=0}^{\infty} \hat{f}(n) e^{int} \right) dt \\ &= \frac{1}{2\pi} \sum_{k=0}^{\infty} \int_{\mathbb{T}} \left(\hat{f}(n) e^{int} \right) dt \\ &= \frac{f(0)}{2\pi} \end{aligned}$$

Now, we proceed to the proof. We define a measure $d\nu = d\mu - \frac{1}{2\pi}\mu(\mathbb{T}) dt$. Now, we have that

$$\begin{aligned} \int_{\mathbb{T}} f(e^{it}) d\nu(e^{it}) &= \int_{\mathbb{T}} [f - f(0)](e^{it}) d\mu(e^{it}) + f(0) \int_{\mathbb{T}} d\nu(e^{it}) \\ &= 0 \end{aligned}$$

Hence, we have that $d\mu = \frac{1}{2\pi}\mu(\mathbb{T}) dt$. □

Theorem §2.2.8 (F and M. Riesz).

§2.3 Szegő's Theorem