

# Shift Invariant Subspaces

...more specifically 1-unilateral shift.

by Ashish Kujur

on September 06, 2023

# Invariant Subspaces

## » Invariant Subspaces

Let  $V$  be any vector space over  $\mathbb{F}$  where  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{F} = \mathbb{C}$ .

### Definition (Invariant Subspaces)

Let  $T : V \rightarrow V$  be a linear operator. A subspace  $W$  of  $V$  is said to be an invariant subspace for  $T$  if  $T(W) \subset W$ .

## » Invariant Subspaces

Let  $V$  be any vector space over  $\mathbb{F}$  where  $\mathbb{F} = \mathbb{R}$  or  $\mathbb{F} = \mathbb{C}$ .

### Definition (Invariant Subspaces)

Let  $T: V \rightarrow V$  be a linear operator. A subspace  $W$  of  $V$  is said to be invariant subspace for  $T$  if  $T(W) \subset W$ .

Given any operator  $T: V \rightarrow V$ , the following are always invariant under  $T$ :

1.  $\{0\}$ ,
2.  $V$ ,
3.  $\ker T$ ,
4.  $\operatorname{im} T$ .

## » Questions that one may ask about invariant subspaces...

Let's separate out the finite and infinite dimensional cases.

## » Questions that one may ask about invariant subspaces...

Let's separate out the finite and infinite dimensional cases.

1. One may ask whether every linear operator on the finite dimensional vector space  $V$  over  $\mathbb{R}$  or  $\mathbb{C}$  admits a invariant subspace different from  $\{0\}$  and the whole space  $V$ .

## » Questions that one may ask about invariant subspaces...

Let's separate out the finite and infinite dimensional cases.

1. One may ask whether every linear operator on the finite dimensional vector space  $V$  over  $\mathbb{R}$  or  $\mathbb{C}$  admits a invariant subspace different from  $\{0\}$  and the whole space  $V$ . Answer: Yes! It always admits a one or two dimensional invariant subspace (See [Axl15]).

## » Questions that one may ask about invariant subspaces...

Let's separate out the finite and infinite dimensional cases.

1. One may ask whether every linear operator on the finite dimensional vector space  $V$  over  $\mathbb{R}$  or  $\mathbb{C}$  admits a invariant subspace different from  $\{0\}$  and the whole space  $V$ . Answer: Yes! It always admits a one or two dimensional invariant subspace (See [Axl15]).
2. The same question also has an affirmative response in infinite dimensions (...but this is an algebraic question.)



» A question about invariant subspaces  
that involves analysis...

## » A question about invariant subspaces that involves analysis...

### Question

*Let  $H$  be a separable complex Hilbert space of dimension  $> 1$ . Does every bounded linear operator on  $T: H \rightarrow H$  admit a nontrivial closed  $T$ -invariant subspace, that is, a closed linear subspace  $W$  of  $H$ , which is different from  $\{0\}$  and  $H$  such that  $T(W) \subset W$ ?*

## » A question about invariant subspaces that involves analysis...

### Question

*Let  $H$  be a separable complex Hilbert space of dimension  $> 1$ . Does every bounded linear operator on  $T: H \rightarrow H$  admit a nontrivial closed  $T$ -invariant subspace, that is, a closed linear subspace  $W$  of  $H$ , which is different from  $\{0\}$  and  $H$  such that  $T(W) \subset W$ ?*

This problem is still open.

## » A question about invariant subspaces that involves analysis...

### Question

*Let  $H$  be a separable complex Hilbert space of dimension  $> 1$ . Does every bounded linear operator on  $T: H \rightarrow H$  admit a nontrivial closed  $T$ -invariant subspace, that is, a closed linear subspace  $W$  of  $H$ , which is different from  $\{0\}$  and  $H$  such that  $T(W) \subset W$ ?*

This problem is still open. For the Banach space case, counterexamples were provided by Per Enflo (1976, 1987) and Charles Read on  $\ell^1$  (1984, 1985).



# Unilateral Shifts

» Right Shift on  $\ell^2(\mathbb{N}_0)$ 

Let  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

$\ell^2(\mathbb{N}_0) := \{(a_0, a_1, a_2, \dots) \mid \forall i \in \mathbb{N}_0, a_i \in \mathbb{C}; \sum_{i=0}^{\infty} |a_i|^2 < \infty\}$ .

$\ell^2(\mathbb{N}_0)$  is an Hilbert space with the following inner product:

$$\langle (a_n)_{n \in \mathbb{N}_0}, (b_n)_{n \in \mathbb{N}_0} \rangle = \sum_{n=0}^{\infty} a_n \bar{b}_n.$$

## » Right Shift on $\ell^2(\mathbb{N}_0)$

Let  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

$\ell^2(\mathbb{N}_0) := \{(a_0, a_1, a_2, \dots) \mid \forall i \in \mathbb{N}_0, a_i \in \mathbb{C}; \sum_{i=0}^{\infty} |a_i|^2 < \infty\}$ .

$\ell^2(\mathbb{N}_0)$  is an Hilbert space with the following inner product:

$$\langle (a_n)_{n \in \mathbb{N}_0}, (b_n)_{n \in \mathbb{N}_0} \rangle = \sum_{n=0}^{\infty} a_n \bar{b}_n.$$

Consider the map  $\mathcal{S} : \ell^2(\mathbb{N}_0) \rightarrow \ell^2(\mathbb{N}_0)$  given by

$$(a_0, a_1, a_2, \dots) \xrightarrow{\mathcal{S}} (0, a_0, a_1, a_2, \dots).$$

## » Right Shift on $\ell^2(\mathbb{N}_0)$

Let  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

$\ell^2(\mathbb{N}_0) := \{(a_0, a_1, a_2, \dots) \mid \forall i \in \mathbb{N}_0, a_i \in \mathbb{C}; \sum_{i=0}^{\infty} |a_i|^2 < \infty\}$ .

$\ell^2(\mathbb{N}_0)$  is an Hilbert space with the following inner product:

$$\langle (a_n)_{n \in \mathbb{N}_0}, (b_n)_{n \in \mathbb{N}_0} \rangle = \sum_{n=0}^{\infty} a_n \bar{b}_n.$$

Consider the map  $\mathcal{S} : \ell^2(\mathbb{N}_0) \rightarrow \ell^2(\mathbb{N}_0)$  given by

$$(a_0, a_1, a_2, \dots) \xrightarrow{\mathcal{S}} (0, a_0, a_1, a_2, \dots).$$

$\mathcal{S}$  does not have any eigenvalues but fortunately, it has a lot of invariant (closed) subspaces!



## » Right Shift on $\ell^2(\mathbb{N}_0)$

Let  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

$\ell^2(\mathbb{N}_0) := \{(a_0, a_1, a_2, \dots) \mid \forall i \in \mathbb{N}_0, a_i \in \mathbb{C}; \sum_{i=0}^{\infty} |a_i|^2 < \infty\}$ .

$\ell^2(\mathbb{N}_0)$  is an Hilbert space with the following inner product:

$$\langle (a_n)_{n \in \mathbb{N}_0}, (b_n)_{n \in \mathbb{N}_0} \rangle = \sum_{n=0}^{\infty} a_n \bar{b}_n.$$

Consider the map  $\mathcal{S} : \ell^2(\mathbb{N}_0) \rightarrow \ell^2(\mathbb{N}_0)$  given by

$$(a_0, a_1, a_2, \dots) \xrightarrow{\mathcal{S}} (0, a_0, a_1, a_2, \dots).$$

$\mathcal{S}$  does not have any eigenvalues but fortunately, it has a lot of invariant (closed) subspaces!

For each  $n \in \mathbb{N}_0$ ,

$\mathcal{M}_n := \{(a_n)_{n \in \mathbb{N}_0} \in \ell^2(\mathbb{N}_0) \mid a_0 = a_1 = \dots = a_{n-1} = 0\}$ . Then

$$\mathcal{S}\mathcal{M}_n = \mathcal{M}_{n+1} \subset \mathcal{M}_n.$$

## » Right Shift on $\ell^2(\mathbb{N}_0)$

Let  $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$ .

$\ell^2(\mathbb{N}_0) := \{(a_0, a_1, a_2, \dots) \mid \forall i \in \mathbb{N}_0, a_i \in \mathbb{C}; \sum_{i=0}^{\infty} |a_i|^2 < \infty\}$ .

$\ell^2(\mathbb{N}_0)$  is an Hilbert space with the following inner product:

$$\langle (a_n)_{n \in \mathbb{N}_0}, (b_n)_{n \in \mathbb{N}_0} \rangle = \sum_{n=0}^{\infty} a_n \bar{b}_n.$$

Consider the map  $\mathcal{S} : \ell^2(\mathbb{N}_0) \rightarrow \ell^2(\mathbb{N}_0)$  given by

$$(a_0, a_1, a_2, \dots) \xrightarrow{\mathcal{S}} (0, a_0, a_1, a_2, \dots).$$

$\mathcal{S}$  does not have any eigenvalues but fortunately, it has a lot of invariant (closed) subspaces!

For each  $n \in \mathbb{N}_0$ ,

$\mathcal{M}_n := \{(a_n)_{n \in \mathbb{N}_0} \in \ell^2(\mathbb{N}_0) \mid a_0 = a_1 = \dots = a_{n-1} = 0\}$ . Then

$$\mathcal{S}\mathcal{M}_n = \mathcal{M}_{n+1} \subset \mathcal{M}_n.$$

Are these all the invariant subspaces on  $\ell^2(\mathbb{N}_0)$ ?

## » Hardy Space

### Definition (The Hardy Hilbert Space)

Let  $\mathbb{D} := \{z \in \mathbb{C} : |z| < 1\}$ . We define

$$H^2(\mathbb{D}) = \left\{ f: \mathbb{D} \rightarrow \mathbb{C} \mid f(z) = \sum_{n=0}^{\infty} a_n z^n, z \in \mathbb{D}; (a_n)_{n \in \mathbb{N}_0} \in \ell^2(\mathbb{N}_0) \right\}$$

For  $f = \sum_{n=0}^{\infty} a_n z^n$  and  $g = \sum_{n=0}^{\infty} b_n z^n$  both in  $H^2(\mathbb{D})$ , we define

$$\langle f, g \rangle = \left\langle \sum_{n=0}^{\infty} a_n z^n, \sum_{n=0}^{\infty} b_n z^n \right\rangle := \sum_{n=0}^{\infty} a_n \bar{b}_n.$$

$H^2(\mathbb{D})$  is an Hilbert space with the given inner product.

## » Shift on the Hardy Space

Define  $S : H^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$  given by

$$S(a_0 + a_1z + \dots + a_nz^n + \dots) = a_0z + a_1z^2 + \dots + a_nz^{n+1} + \dots$$

This is the same as

$$[Sf](z) = zf(z) \text{ for each } f \in H^2 \text{ and each } z \in \mathbb{D}.$$

We will call it the shift on the Hardy space.

## » Shift on the Hardy Space

Define  $S : H^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$  given by

$$S(a_0 + a_1z + \dots + a_nz^n + \dots) = a_0z + a_1z^2 + \dots + a_nz^{n+1} + \dots$$

This is the same as

$$[Sf](z) = zf(z) \text{ for each } f \in H^2 \text{ and each } z \in \mathbb{D}.$$

We will call it the shift on the Hardy space. Consider the map  $U : H^2(\mathbb{D}) \rightarrow \ell^2(\mathbb{N}_0)$  given by

$$z_i \xrightarrow{U} e_i = \left( 0, 0, \dots, \underbrace{1}_{i\text{th entry}}, 0, \dots \right)$$

for each  $i \in \mathbb{N}_0$ . Thus,  $U$  is a unitary transformation and  $U^*SU = S$ . Consequently, the invariant subspaces of  $S$  and the invariant subspaces of  $\mathcal{S}$  are in one-to-one correspondence.

Some Examples in  $H^2(\mathbb{D})$

## » Some Invariant Subspaces of $H^2(\mathbb{D})$

Let  $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subset \mathbb{D}$ . Consider the subspace

$$\mathcal{M} = \{f \in H^2(\mathbb{D}) \mid f(\alpha_i) = 0 \text{ for all } 1 \leq i \leq n\}.$$

This forms a closed invariant subspace of shift  $S$  on the Hardy Space,  $1 \notin \mathcal{M}$  and  $p(z) = \prod_{i=1}^n (\alpha_i - z) \in \mathcal{M}$  and  $p \neq 0$ .

## » Some Invariant Subspaces of $H^2(\mathbb{D})$

Let  $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \subset \mathbb{D}$ . Consider the subspace

$$\mathcal{M} = \{f \in H^2(\mathbb{D}) \mid f(\alpha_i) = 0 \text{ for all } 1 \leq i \leq n\}.$$

This forms a closed invariant subspace of shift  $S$  on the Hardy Space,  $1 \notin \mathcal{M}$  and  $p(z) = \prod_{i=1}^n (\alpha_i - z) \in \mathcal{M}$  and  $p \neq 0$ .

Let  $\{\alpha_1, \alpha_2, \dots, \alpha_n, \dots\} \subset \mathbb{D}$  be an infinite subset. Like the previous case, it is easily seen that

$$\mathcal{M} = \{f \in H^2(\mathbb{D}) \mid f(\alpha_i) = 0 \text{ for all } i \in \mathbb{N}\}$$

is an closed invariant subspace of  $H^2(\mathbb{D})$ . But is it the case that

$$\{0\} \subsetneq \mathcal{M} \subsetneq H^2(\mathbb{D})?$$

NAIVE GUESS:  $p(z) = \prod_{i=1}^{\infty} (\alpha_i - z)$ . But is  $p \in H^2(\mathbb{D})$ ?



## » Inner functions

## » Inner functions

## Definition (Inner Function)

Let  $f: \mathbb{D} \rightarrow \mathbb{C}$  and let  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ . Then  $f$  is said to be an *inner function* if  $f$  is a bounded holomorphic function and the function  $\tilde{f}$  defined almost everywhere on  $\mathbb{T}$  by

$$\tilde{f}(e^{i\theta}) = \lim_{r \rightarrow 1} f(re^{i\theta})$$

has unit modulus almost everywhere.

## » Inner functions

### Definition (Inner Function)

Let  $f: \mathbb{D} \rightarrow \mathbb{C}$  and let  $\mathbb{T} = \{z \in \mathbb{C} : |z| = 1\}$ . Then  $f$  is said to be an *inner function* if  $f$  is a bounded holomorphic function and the function  $\tilde{f}$  defined almost everywhere on  $\mathbb{T}$  by

$$\tilde{f}(e^{i\theta}) = \lim_{r \rightarrow 1} f(re^{i\theta})$$

has unit modulus almost everywhere.

### Finite Blaschke Product

Let  $\alpha_1, \alpha_2, \dots, \alpha_n \in \mathbb{D}$  and  $\beta \in \mathbb{R}$ . Then the finite Blaschke product is the function given by

$$B(z) = e^{i\beta} \prod_{k=1}^n \frac{z - \alpha_j}{1 - \bar{\alpha}_j z}.$$

## » Infinite Blaschke Product

Theorem (Blaschke, F. Riesz)

Let  $(\alpha_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathbb{D} \setminus \{0\}$ . Then the infinite product

$$B(z) = \prod_{n=1}^{\infty} \frac{|\alpha_n|}{\alpha_n} \frac{\alpha_n - z}{1 - \bar{\alpha}_n z}$$

converges uniformly on compact subsets of  $\mathbb{D}$

$$\iff \sum_{i=1}^{\infty} (1 - |\alpha_i|) < \infty.$$

In either cases,  $B$  defines an inner function.

## » Infinite Blaschke Product

Theorem (Blaschke, F. Riesz)

Let  $(\alpha_n)_{n \in \mathbb{N}}$  be a sequence in  $\mathbb{D} \setminus \{0\}$ . Then the infinite product

$$B(z) = \prod_{n=1}^{\infty} \frac{|\alpha_n|}{\alpha_n} \frac{\alpha_n - z}{1 - \bar{\alpha}_n z}$$

converges uniformly on compact subsets of  $\mathbb{D}$

$$\iff \sum_{i=1}^{\infty} (1 - |\alpha_i|) < \infty.$$

In either cases,  $B$  defines an inner function.

For a proof of this, take a look at [Rud74], [Hof07] or [Mas09].

# Beurling's Theorem

»  $L^2(\mathbb{T})$ Definition (Lebesgue Measure on  $\mathbb{T}$ )

A subset  $A \subset \mathbb{T}$  is called measurable subset of  $\mathbb{T}$  if  $\{t \in (-\pi, \pi] : e^{it} \in A\}$  is Lebesgue measurable subset of  $(-\pi, \pi]$ . The Lebesgue measure  $m_{\mathbb{T}}$  on the measurable subset of  $\mathbb{T}$  is given by

$$\lambda_{\mathbb{T}}(A) = \frac{\lambda(\{t \in (-\pi, \pi] : e^{it} \in A\})}{2\pi}$$

where  $\lambda$  is the Lebesgue measure on  $(-\pi, \pi]$ .

»  $L^2(\mathbb{T})$ Definition (Lebesgue Measure on  $\mathbb{T}$ )

A subset  $A \subset \mathbb{T}$  is called measurable subset of  $\mathbb{T}$  if  $\{t \in (-\pi, \pi] : e^{it} \in A\}$  is Lebesgue measurable subset of  $(-\pi, \pi]$ . The Lebesgue measure  $m_{\mathbb{T}}$  on the measurable subset of  $\mathbb{T}$  is given by

$$\lambda_{\mathbb{T}}(A) = \frac{\lambda(\{t \in (-\pi, \pi] : e^{it} \in A\})}{2\pi}$$

where  $\lambda$  is the Lebesgue measure on  $(-\pi, \pi]$ .

Then  $L^2(\mathbb{T})$  is a Hilbert space and if we define  $\zeta : \mathbb{T} \rightarrow \mathbb{T}$  to be function given by  $\zeta(e^{it}) = e^{it}$  for each  $e^{it} \in \mathbb{T}$  then  $\{\zeta^n : n \in \mathbb{Z}\}$  forms an orthonormal basis for  $L^2(\mathbb{T})$ . (Proof: Refer [Axl20] or [Rud74]).



## » Hardy Spaces on $\mathbb{T}$

For  $f \in L^2(\mathbb{T})$ , define  $\hat{f}(n) = \langle f, \zeta^n \rangle_{L^2(\mathbb{T})} = \int_{\mathbb{T}} f \zeta^{-n} d\lambda_{\mathbb{T}}$ .

## » Hardy Spaces on $\mathbb{T}$

For  $f \in L^2(\mathbb{T})$ , define  $\hat{f}(n) = \langle f, \zeta^n \rangle_{L^2(\mathbb{T})} = \int_{\mathbb{T}} f \zeta^{-n} d\lambda_{\mathbb{T}}$ .

### Definition (Hardy Spaces on $\mathbb{T}$ )

We define the Hardy space  $H^2(\mathbb{T})$  to be given by

$$H^2(\mathbb{T}) := \left\{ f \in L^2(\mathbb{T}) : \hat{f}(n) = 0 \text{ for each } n \leq -1 \right\}.$$

Easy to see:  $H^2(\mathbb{T})$  is a closed subspace of  $L^2(\mathbb{T})$ .

## » Hardy Spaces on $\mathbb{T}$

For  $f \in L^2(\mathbb{T})$ , define  $\hat{f}(n) = \langle f, \zeta^n \rangle_{L^2(\mathbb{T})} = \int_{\mathbb{T}} f \zeta^{-n} d\lambda_{\mathbb{T}}$ .

### Definition (Hardy Spaces on $\mathbb{T}$ )

We define the Hardy space  $H^2(\mathbb{T})$  to be given by

$$H^2(\mathbb{T}) := \left\{ f \in L^2(\mathbb{T}) : \hat{f}(n) = 0 \text{ for each } n \leq -1 \right\}.$$

Easy to see:  $H^2(\mathbb{T})$  is a closed subspace of  $L^2(\mathbb{T})$ . Furthermore, if  $f \in H^2(\mathbb{D})$  then the function  $\tilde{f}: \mathbb{T} \rightarrow \mathbb{C}$

$$\tilde{f}(e^{it}) = \lim_{r \rightarrow 1^-} f(re^{it})$$

for each  $e^{it} \in \mathbb{T}$  is  $H^2(\mathbb{T})$  function, and, furthermore, we have that

$$\langle f, g \rangle_{H^2(\mathbb{D})} = \left\langle \tilde{f}, \tilde{g} \right\rangle_{H^1(\mathbb{T})}$$

for each  $f, g \in H^2(D)$  and  $\tilde{f}, \tilde{g} \in H^2(\mathbb{T})$  are their corresponding boundary functions.

»  $\theta H^2(\mathbb{D})$  are invariant subspaces!

»  $\theta H^2(\mathbb{D})$  are invariant subspaces!

$\theta H^2(\mathbb{D})$  are invariant subspaces of  $H^2(\mathbb{D})$  where  $\theta$  is inner.

Let  $\theta$  be an inner function.

»  $\theta H^2(\mathbb{D})$  are invariant subspaces!

$\theta H^2(\mathbb{D})$  are invariant subspaces of  $H^2(\mathbb{D})$  where  $\theta$  is inner.

Let  $\theta$  be an inner function.

Consider the operator  $M_\theta : H^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$  given by  $[M_\theta(f)](z) = \theta(z)f(z)$  for each  $f \in H^2(\mathbb{D})$  and  $z \in D$ .

»  $\theta H^2(\mathbb{D})$  are invariant subspaces!

$\theta H^2(\mathbb{D})$  are invariant subspaces of  $H^2(\mathbb{D})$  where  $\theta$  is inner.

Let  $\theta$  be an inner function.

Consider the operator  $M_\theta : H^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$  given by  $[M_\theta(f)](z) = \theta(z)f(z)$  for each  $f \in H^2(\mathbb{D})$  and  $z \in D$ .  
 $M_\theta$  is an isometry.

»  $\theta H^2(\mathbb{D})$  are invariant subspaces!

$\theta H^2(\mathbb{D})$  are invariant subspaces of  $H^2(\mathbb{D})$  where  $\theta$  is inner.

Let  $\theta$  be an inner function.

Consider the operator  $M_\theta : H^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$  given by  $[M_\theta(f)](z) = \theta(z)f(z)$  for each  $f \in H^2(\mathbb{D})$  and  $z \in D$ .

$M_\theta$  is an isometry.

To show it is invariant, consider the following:

$$S(\theta H^2(\mathbb{D})) = z(\theta H^2(\mathbb{D})) = \theta(z H^2(\mathbb{D})) \subset \theta H^2(\mathbb{D})$$



## » $\theta H^2(\mathbb{D})$ are invariant subspaces!

$\theta H^2(\mathbb{D})$  are invariant subspaces of  $H^2(\mathbb{D})$  where  $\theta$  is inner.

Let  $\theta$  be an inner function.

Consider the operator  $M_\theta : H^2(\mathbb{D}) \rightarrow H^2(\mathbb{D})$  given by  $[M_\theta(f)](z) = \theta(z)f(z)$  for each  $f \in H^2(\mathbb{D})$  and  $z \in D$ .

$M_\theta$  is an isometry.

To show it is invariant, consider the following:

$$S(\theta H^2(\mathbb{D})) = z(\theta H^2(\mathbb{D})) = \theta(z H^2(\mathbb{D})) \subset \theta H^2(\mathbb{D})$$

### Theorem (Beurling's Theorem)

*Every nonzero (closed) invariant subspace of  $H^2(\mathbb{D})$  is of the form  $\theta H^2(\mathbb{D})$  for some inner function  $\theta$ .*

## » Sketch of Proof of Beurling's Theorem

### Theorem (von Neumann-Wold decomposition)

Let  $T$  be a bounded linear operator on a Hilbert space  $\mathcal{H}$  and  $\mathcal{W} = \mathcal{H} \ominus T\mathcal{H}$ .

1.  $T^m\mathcal{H} \perp T^n\mathcal{H}$  for every  $m, n \geq 0$  and  $m \neq n$ .
2.  $\mathcal{W}_\infty = \bigcup_{n \geq 0} T^n\mathcal{H}$  then  $T\mathcal{W}_\infty$  and  $T|_{\mathcal{W}_\infty}$  is unitary.
3.  $\mathcal{W}_0 = \bigoplus_{n \geq 0} T^n\mathcal{W}$  then  $T\mathcal{W}_0 \subset \mathcal{W}_0$  and there is no nonzero invariant subspace of  $\mathcal{W}_0$  upon which the restriction of  $T$  is unitary.
4.  $\mathcal{H} = \mathcal{W}_0 \oplus \mathcal{W}_\infty$ .

## » Sketch of Proof of Beurling's Theorem

### Theorem (von Neumann-Wold decomposition)

Let  $T$  be a bounded linear operator on a Hilbert space  $\mathcal{H}$  and  $\mathcal{W} = \mathcal{H} \ominus T\mathcal{H}$ .

1.  $T^m\mathcal{H} \perp T^n\mathcal{H}$  for every  $m, n \geq 0$  and  $m \neq n$ .
2.  $\mathcal{W}_\infty = \bigcup_{n \geq 0} T^n\mathcal{H}$  then  $T\mathcal{W}_\infty$  and  $T|_{\mathcal{W}_\infty}$  is unitary.
3.  $\mathcal{W}_0 = \bigoplus_{n \geq 0} T^n\mathcal{W}$  then  $T\mathcal{W}_0 \subset \mathcal{W}_0$  and there is no nonzero invariant subspace of  $\mathcal{W}_0$  upon which the restriction of  $T$  is unitary.
4.  $\mathcal{H} = \mathcal{W}_0 \oplus \mathcal{W}_\infty$ .

For a proof, see [Nag+10].

» Sketch Part 1:  $\mathcal{SM} \subsetneq \mathcal{M}$

» Sketch Part 2:  $(\mathcal{M} \ominus S\mathcal{M})$  has an inner function  $\theta$

# » Sketch Part 3: Use Wold Decomposition

# » Final Remarks: Singular Inner Function & Factorization

## » References

- [Axl15]   Sheldon Axler. *Linear algebra done right*. Springer, 2015.
- [Axl20]   Sheldon Axler. *Measure, integration & real analysis*. Springer Nature, 2020.
- [GMR16]   Stephan Ramon Garcia, Javad Mashreghi, and William T Ross. *Introduction to model spaces and their operators*. Vol. 148. Cambridge University Press, 2016.
- [Hof07]   Kenneth Hoffman. *Banach spaces of analytic functions*. Courier Corporation, 2007.
- [Mas09]   Javad Mashreghi. *Representation theorems in Hardy spaces*. Vol. 74. Cambridge University Press, 2009.



## » References (cont.)

- [Nag+10] Béla Sz Nagy et al. *Harmonic analysis of operators on Hilbert space*. Springer Science & Business Media, 2010.
- [Ric04] Stefan Richter. *An Elementary Introduction To What I Do For My Research*. 2004. URL: [https://web.math.utk.edu/~richter/Research\\_elementary.pdf](https://web.math.utk.edu/~richter/Research_elementary.pdf).
- [Rud74] Walter Rudin. *Real and complex analysis*. Second. McGraw-Hill Series in Higher Mathematics. McGraw-Hill Book Co., New York-Düsseldorf-Johannesburg, 1974, pp. xii+452.

## » References (cont.)

[Sar21] Meredith Sargent. *Shift Invariant Subspaces, An Introduction*. 2021. URL: <https://meresarmath.wordpress.com/2021/09/30/shift-invariant-subspaces-an-introduction/>.

Thank You! Questions?