

ALGEBRA

Factors and Zeros of Polynomials

Let $p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$ be a polynomial. If $p(a) = 0$, then a is a *zero* of the polynomial and a solution of the equation $p(x) = 0$. Furthermore, $(x - a)$ is a *factor* of the polynomial.

Fundamental Theorem of Algebra

An n th degree polynomial has n (not necessarily distinct) zeros. Although all of these zeros may be imaginary, a real polynomial of odd degree must have at least one real zero.

Quadratic Formula

If $p(x) = ax^2 + bx + c$, and $0 \leq b^2 - 4ac$, then the real zeros of p are $x = (-b \pm \sqrt{b^2 - 4ac})/2a$.

Special Factors

$$x^2 - a^2 = (x - a)(x + a)$$

$$x^3 - a^3 = (x - a)(x^2 + ax + a^2)$$

$$x^3 + a^3 = (x + a)(x^2 - ax + a^2)$$

$$x^4 - a^4 = (x - a)(x + a)(x^2 + a^2)$$

Binomial Theorem

$$(x + y)^2 = x^2 + 2xy + y^2$$

$$(x - y)^2 = x^2 - 2xy + y^2$$

$$(x + y)^3 = x^3 + 3x^2y + 3xy^2 + y^3$$

$$(x - y)^3 = x^3 - 3x^2y + 3xy^2 - y^3$$

$$(x + y)^4 = x^4 + 4x^3y + 6x^2y^2 + 4xy^3 + y^4$$

$$(x - y)^4 = x^4 - 4x^3y + 6x^2y^2 - 4xy^3 + y^4$$

$$(x + y)^n = x^n + nx^{n-1}y + \frac{n(n-1)}{2!}x^{n-2}y^2 + \cdots + nxy^{n-1} + y^n$$

$$(x - y)^n = x^n - nx^{n-1}y + \frac{n(n-1)}{2!}x^{n-2}y^2 - \cdots \pm nxy^{n-1} \mp y^n$$

Rational Zero Theorem

If $p(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0$ has integer coefficients, then every *rational zero* of p is of the form $x = r/s$, where r is a factor of a_0 and s is a factor of a_n .

Factoring by Grouping

$$acx^3 + adx^2 + bcx + bd = ax^2(cx + d) + b(cx + d) = (ax^2 + b)(cx + d)$$

Arithmetic Operations

$$ab + ac = a(b + c)$$

$$\frac{a}{b} + \frac{c}{d} = \frac{ad + bc}{bd}$$

$$\frac{a + b}{c} = \frac{a}{c} + \frac{b}{c}$$

$$\left(\frac{a}{b}\right)\left(\frac{c}{d}\right) = \left(\frac{a}{b}\right)\left(\frac{c}{d}\right) = \frac{ac}{bd}$$

$$\left(\frac{a}{b}\right)\left(\frac{b}{c}\right) = \frac{a}{c}$$

$$\frac{a}{\left(\frac{b}{c}\right)} = \frac{ac}{b}$$

$$a\left(\frac{b}{c}\right) = \frac{ab}{c}$$

$$\frac{a - b}{c - d} = \frac{b - a}{d - c}$$

$$\frac{ab + ac}{a} = b + c$$

Exponents and Radicals

$$a^0 = 1, \quad a \neq 0$$

$$(ab)^x = a^x b^x$$

$$a^x a^y = a^{x+y}$$

$$\sqrt{a} = a^{1/2}$$

$$\frac{a^x}{a^y} = a^{x-y}$$

$$\sqrt[n]{a} = a^{1/n}$$

$$\left(\frac{a}{b}\right)^x = \frac{a^x}{b^x}$$

$$\sqrt[n]{a^m} = a^{m/n}$$

$$a^{-x} = \frac{1}{a^x}$$

$$\sqrt[n]{ab} = \sqrt[n]{a} \sqrt[n]{b}$$

$$(a^x)^y = a^{xy}$$

$$\sqrt[n]{\frac{a}{b}} = \frac{\sqrt[n]{a}}{\sqrt[n]{b}}$$

FORMULAS FROM GEOMETRY


Triangle

$$h = a \sin \theta$$

$$\text{Area} = \frac{1}{2}bh$$

(Law of Cosines)


$$c^2 = a^2 + b^2 - 2ab \cos \theta$$



Right Triangle

(Pythagorean Theorem)


$$c^2 = a^2 + b^2$$



Equilateral Triangle


$$h = \frac{\sqrt{3}s}{2}$$

$$\text{Area} = \frac{\sqrt{3}s^2}{4}$$




Parallelogram

$$\text{Area} = bh$$



Trapezoid


$$\text{Area} = \frac{h}{2}(a + b)$$



Circle

$$\text{Area} = \pi r^2$$

$$\text{Circumference} = 2\pi r$$




Sector of Circle

(θ in radians)

$$\text{Area} = \frac{\theta r^2}{2}$$

$$s = r\theta$$




Circular Ring

(p = average radius,

w = width of ring)

$$\begin{aligned} \text{Area} &= \pi(R^2 - r^2) \\ &= 2\pi pw \end{aligned}$$




Sector of Circular Ring

(p = average radius,

w = width of ring,

θ in radians)


$$\text{Area} = \theta pw$$



Ellipse

$$\text{Area} = \pi ab$$


$$\text{Circumference} \approx 2\pi \sqrt{\frac{a^2 + b^2}{2}}$$



Cone

(A = area of base)


$$\text{Volume} = \frac{Ah}{3}$$



Right Circular Cone

$$\text{Volume} = \frac{\pi r^2 h}{3}$$


$$\text{Lateral Surface Area} = \pi r \sqrt{r^2 + h^2}$$



Frustum of Right Circular Cone

$$\text{Volume} = \frac{\pi(r^2 + rR + R^2)h}{3}$$


$$\text{Lateral Surface Area} = \pi s(R + r)$$



Right Circular Cylinder

$$\text{Volume} = \pi r^2 h$$


$$\text{Lateral Surface Area} = 2\pi rh$$



Sphere

$$\text{Volume} = \frac{4}{3}\pi r^3$$

$$\text{Surface Area} = 4\pi r^2$$




Wedge

(A = area of upper face,

B = area of base)


$$A = B \sec \theta$$



TRIGONOMETRY


Definition of the Six Trigonometric Functions

Right triangle definitions, where $0 < \theta < \pi/2$.




$$\begin{aligned}\sin \theta &= \frac{\text{opp}}{\text{hyp}} & \csc \theta &= \frac{\text{hyp}}{\text{opp}} \\ \cos \theta &= \frac{\text{adj}}{\text{hyp}} & \sec \theta &= \frac{\text{hyp}}{\text{adj}} \\ \tan \theta &= \frac{\text{opp}}{\text{adj}} & \cot \theta &= \frac{\text{adj}}{\text{opp}}\end{aligned}$$

Circular function definitions, where θ is any angle.



$$\begin{aligned}\sin \theta &= \frac{y}{r} & \csc \theta &= \frac{r}{y} \\ \cos \theta &= \frac{x}{r} & \sec \theta &= \frac{r}{x} \\ \tan \theta &= \frac{y}{x} & \cot \theta &= \frac{x}{y}\end{aligned}$$



Reciprocal Identities

$$\begin{aligned}\sin x &= \frac{1}{\csc x} & \sec x &= \frac{1}{\cos x} & \tan x &= \frac{1}{\cot x} \\ \csc x &= \frac{1}{\sin x} & \cos x &= \frac{1}{\sec x} & \cot x &= \frac{1}{\tan x}\end{aligned}$$

Quotient Identities

$$\tan x = \frac{\sin x}{\cos x} \quad \cot x = \frac{\cos x}{\sin x}$$

Pythagorean Identities

$$\begin{aligned}\sin^2 x + \cos^2 x &= 1 \\ 1 + \tan^2 x &= \sec^2 x & 1 + \cot^2 x &= \csc^2 x\end{aligned}$$

Cofunction Identities

$$\begin{aligned}\sin\left(\frac{\pi}{2} - x\right) &= \cos x & \cos\left(\frac{\pi}{2} - x\right) &= \sin x \\ \csc\left(\frac{\pi}{2} - x\right) &= \sec x & \tan\left(\frac{\pi}{2} - x\right) &= \cot x \\ \sec\left(\frac{\pi}{2} - x\right) &= \csc x & \cot\left(\frac{\pi}{2} - x\right) &= \tan x\end{aligned}$$

Even/Odd Identities

$$\begin{aligned}\sin(-x) &= -\sin x & \cos(-x) &= \cos x \\ \csc(-x) &= -\csc x & \tan(-x) &= -\tan x \\ \sec(-x) &= \sec x & \cot(-x) &= -\cot x\end{aligned}$$

Sum and Difference Formulas

$$\begin{aligned}\sin(u \pm v) &= \sin u \cos v \pm \cos u \sin v \\ \cos(u \pm v) &= \cos u \cos v \mp \sin u \sin v \\ \tan(u \pm v) &= \frac{\tan u \pm \tan v}{1 \mp \tan u \tan v}\end{aligned}$$

Double-Angle Formulas

$$\begin{aligned}\sin 2u &= 2 \sin u \cos u \\ \cos 2u &= \cos^2 u - \sin^2 u = 2 \cos^2 u - 1 = 1 - 2 \sin^2 u \\ \tan 2u &= \frac{2 \tan u}{1 - \tan^2 u}\end{aligned}$$

Power-Reducing Formulas

$$\begin{aligned}\sin^2 u &= \frac{1 - \cos 2u}{2} \\ \cos^2 u &= \frac{1 + \cos 2u}{2} \\ \tan^2 u &= \frac{1 - \cos 2u}{1 + \cos 2u}\end{aligned}$$

Sum-to-Product Formulas

$$\begin{aligned}\sin u + \sin v &= 2 \sin\left(\frac{u+v}{2}\right) \cos\left(\frac{u-v}{2}\right) \\ \sin u - \sin v &= 2 \cos\left(\frac{u+v}{2}\right) \sin\left(\frac{u-v}{2}\right) \\ \cos u + \cos v &= 2 \cos\left(\frac{u+v}{2}\right) \cos\left(\frac{u-v}{2}\right) \\ \cos u - \cos v &= -2 \sin\left(\frac{u+v}{2}\right) \sin\left(\frac{u-v}{2}\right)\end{aligned}$$

Product-to-Sum Formulas

$$\begin{aligned}\sin u \sin v &= \frac{1}{2}[\cos(u-v) - \cos(u+v)] \\ \cos u \cos v &= \frac{1}{2}[\cos(u-v) + \cos(u+v)] \\ \sin u \cos v &= \frac{1}{2}[\sin(u+v) + \sin(u-v)] \\ \cos u \sin v &= \frac{1}{2}[\sin(u+v) - \sin(u-v)]\end{aligned}$$

DERIVATIVES AND INTEGRALS

Basic Differentiation Rules

- $\frac{d}{dx}[cu] = cu'$
- $\frac{d}{dx}[u \pm v] = u' \pm v'$
- $\frac{d}{dx}[uv] = uv' + vu'$
- $\frac{d}{dx}\left[\frac{u}{v}\right] = \frac{vu' - uv'}{v^2}$
- $\frac{d}{dx}[c] = 0$
- $\frac{d}{dx}[u^n] = nu^{n-1}u'$
- $\frac{d}{dx}[x] = 1$
- $\frac{d}{dx}[|u|] = \frac{u}{|u|}(u'), \quad u \neq 0$
- $\frac{d}{dx}[\ln u] = \frac{u'}{u}$
- $\frac{d}{dx}[e^u] = e^u u'$
- $\frac{d}{dx}[\log_a u] = \frac{u'}{(\ln a)u}$
- $\frac{d}{dx}[a^u] = (\ln a)a^u u'$
- $\frac{d}{dx}[\sin u] = (\cos u)u'$
- $\frac{d}{dx}[\cos u] = -(\sin u)u'$
- $\frac{d}{dx}[\tan u] = (\sec^2 u)u'$
- $\frac{d}{dx}[\cot u] = -(\csc^2 u)u'$
- $\frac{d}{dx}[\sec u] = (\sec u \tan u)u'$
- $\frac{d}{dx}[\csc u] = -(\csc u \cot u)u'$
- $\frac{d}{dx}[\arcsin u] = \frac{u'}{\sqrt{1-u^2}}$
- $\frac{d}{dx}[\arccos u] = \frac{-u'}{\sqrt{1-u^2}}$
- $\frac{d}{dx}[\arctan u] = \frac{u'}{1+u^2}$
- $\frac{d}{dx}[\operatorname{arccsc} u] = \frac{-u'}{|u|\sqrt{u^2-1}}$
- $\frac{d}{dx}[\sinh u] = (\cosh u)u'$
- $\frac{d}{dx}[\cosh u] = (\sinh u)u'$
- $\frac{d}{dx}[\tanh u] = (\operatorname{sech}^2 u)u'$
- $\frac{d}{dx}[\coth u] = -(\operatorname{csch}^2 u)u'$
- $\frac{d}{dx}[\operatorname{sech} u] = -(\operatorname{sech} u \tanh u)u'$
- $\frac{d}{dx}[\operatorname{csch} u] = -(\operatorname{csch} u \coth u)u'$
- $\frac{d}{dx}[\sinh^{-1} u] = \frac{u'}{\sqrt{u^2+1}}$
- $\frac{d}{dx}[\cosh^{-1} u] = \frac{u'}{\sqrt{u^2-1}}$
- $\frac{d}{dx}[\tanh^{-1} u] = \frac{u'}{1-u^2}$
- $\frac{d}{dx}[\coth^{-1} u] = \frac{u'}{1-u^2}$
- $\frac{d}{dx}[\operatorname{sech}^{-1} u] = \frac{-u'}{u\sqrt{1-u^2}}$
- $\frac{d}{dx}[\operatorname{csch}^{-1} u] = \frac{-u'}{|u|\sqrt{1+u^2}}$

Basic Integration Formulas

- $\int kf(u) du = k \int f(u) du$
- $\int [f(u) \pm g(u)] du = \int f(u) du \pm \int g(u) du$
- $\int du = u + C$
- $\int u^n du = \frac{u^{n+1}}{n+1} + C, \quad n \neq -1$
- $\int \frac{du}{u} = \ln|u| + C$
- $\int e^u du = e^u + C$
- $\int a^u du = \left(\frac{1}{\ln a}\right)a^u + C$
- $\int \sin u du = -\cos u + C$
- $\int \tan u du = -\ln|\cos u| + C$
- $\int \cot u du = \ln|\sin u| + C$
- $\int \sec u du = \ln|\sec u + \tan u| + C$
- $\int \csc u du = -\ln|\csc u + \cot u| + C$
- $\int \sec^2 u du = \tan u + C$
- $\int \csc^2 u du = -\cot u + C$
- $\int \sec u \tan u du = \sec u + C$
- $\int \csc u \cot u du = -\csc u + C$
- $\int \frac{du}{\sqrt{a^2 - u^2}} = \arcsin \frac{u}{a} + C$
- $\int \frac{du}{u\sqrt{u^2 - a^2}} = \frac{1}{a} \operatorname{arcsec} \frac{|u|}{a} + C$