

Robust Regression

유제진

Department of Statistics, University

INDEX

1. Check function
2. Building QR
3. Outlier Protection on Median
4. QR on Engel Data
5. QR on Wage Data

QUANTILE REGRESSION

Quantile

- r -th Quantile of random variable Y with Cdf F_Y

$$Q_Y(\tau) = F_Y^{-1}(\tau) = \inf\{y : F_Y(y) \geq \tau\}, \tau \in (0, 1)$$

- Here, we define the loss function as

$$\rho_\tau(y) = y * (\tau - I(y < 0)) \text{ where } I : \text{Indicator Function}$$

Also known as "Check Function".

QUANTILE REGRESSION

Quantile

- CDF Function : function returns probabilities of X being smaller than or equal to some value x .


$$Pr(X \leq x) = F(x)$$

- The inverse of CDF = Quantile Function

$$F^{-1}(p) = x$$

where x would return some value P .

QUANTILE REGRESSION



- Gumbel Distribution

$$F(x) = e^{-e^{-x}}$$

\longleftrightarrow

$$F^{-1}(p) = -\ln(-\ln p)$$

QUANTILE REGRESSION

Generalized Inverse Distribution Function

- ▶ Not every function has an inverse.
- ▶ So condition "Monotonically increasing" is needed.
- ▶ Also, to be a function, one to one condition need to be satisfied.
- ▶ CDF satisfy this condition.
- ▶ But what happens when it comes to discrete random variable CDF?
- ▶ Discrete random variable CDF is not continuous, but increasing.

QUANTILE REGRESSION

Generalized Inverse Distribution Function

- ▶ Generalized Inverse Distribution only requires non-decreasing condition.

$$Q_Y(\tau) = F_Y^{-1}(\tau) = \text{Inf}\{y : F_Y(y) \geq \tau\}, \tau \in (0, 1)$$

- ▶ For given probability value τ ,
- ▶ Look for some y that results in $F(y)$ returning value greater or equal then τ .
- ▶ But since there could be multiple values of y that meet this condition.
e.g. $F(y) \geq 0$ is true for any y
- ▶ So use Infimum to take the smallest among y .

QUANTILE REGRESSION

Derivation of Check Function

- ▶ Find "Location" x^* relative to a distribution or set of data F .
- ▶ mean

$$L_F(\bar{x}) = \int_R (x - \bar{x})^2 dF(x)$$

mean minimized the expected squared residual.

- ▶ L_F is for loss function determined by F .
- ▶ To show x^* minimized any function begins with,
- ▶ Demonstrating the function's value does not decrease when x^* is changed by a little bit.
- ▶ Such a value is called a critical point of the function.

QUANTILE REGRESSION

Derivation of Check Function

- What kind of loss function Λ would result in a percentile $F^{-1}(\alpha)$ being a critical point?

$$\begin{aligned} L_F(F^{-1}(\alpha)) &= \int_R \Lambda(x - F^{-1}(\alpha)) dF(x) \\ &= \int_0^1 \Lambda(F^{-1}(u) - F^{-1}(\alpha)) du \end{aligned}$$

- For this to be critical point, derivative must be zero.

QUANTILE REGRESSION

Derivation of Check Function

$$\begin{aligned} 0 &= L'_F(x^*) = L_F(F^{-1}(\alpha)) = - \int_0^1 \Lambda'(F^{-1}(u) - F^{-1}(\alpha)) du \\ &= - \int_0^\alpha \Lambda'(F^{-1}(u) - F^{-1}(\alpha)) du - \int_\alpha^1 \Lambda'(F^{-1}(u) - F^{-1}(\alpha)) du \end{aligned}$$

- ▶ On the left hand side, argument of Λ' is negative.
- ▶ On the right hand side, argument of Λ' is positive.
- ▶ Other than that, we have little control over the values of these integrals.
- ▶ Because F could be any distribution function.

QUANTILE REGRESSION

Derivation of Check Function

- ▶ Now, make Λ' depend only on the sign of its argument, otherwise it must be constant.
- ▶ This implies Λ will be piecewise linear, potentially with different slopes to the left and right of zero.
- ▶ Moreover, Rescaling Λ by a constant will not change its properties.
- ▶ So may feel free to set the left hand slope to -1.

QUANTILE REGRESSION

Derivation of Check Function

- ▶ Let $\tau > 0$ be the right hand slope.
- ▶ Then final equation simplifies to

$$0 = \alpha - \tau(1 - \alpha) \implies \tau = \frac{\alpha}{1 - \alpha}$$

- ▶ To conclude,

$$\Lambda(x) = \begin{cases} -x \\ \alpha \\ 1 - \alpha \end{cases} x \implies \Lambda(x) = \begin{cases} -(1 - \alpha)x & x \leq 0 \\ \alpha x & x \geq 0 \end{cases}$$

DRAWING CHECK FUNCTION

Check Function

```
► rho <- function(u) {u * (tau - ifelse(u <
  0,1,0) )}
tau <- .25; curve(rho,-2,2,lty=1,lwd=3)
tau <- .50; curve(rho,
  -2,2,lty=2,col="blue",add=T,lwd=3)
tau <- .90; curve(rho,
  -2,2,lty=3,col="red",add=T, lwd=3)
abline(v=0,lty=5,col="gray",lwd=3)
legend("bottomleft",c(".25",".5",".9"),
  lty=1:3,col=c("black","blue","red"),cex=.6)
```

DRAWING CHECK FUNCTION

Data Plot




Figure 1: Loss of check function

BUILDING QUANTILE REGRESSION

Median

- Consider simple $\{y_1 \cdots y_n\}$

$$\min_{\mu} \left\{ \sum_{i=1}^n |y_i - \mu| \right\} \iff \min_{\mu, a, b} \left\{ \sum_{i=1}^n a_i + b_i \right\}$$

subject to $a_i, b_i \geq 0$ and $y_i - \mu = a_i - b_i \quad \forall 1, \dots, n$

- Turn into Linear Programming Problem.
- To illustrate, consider a lognormal sample.

BUILDING QUANTILE REGRESSION

Median

► Pre-built Function

```
n=101
set.seed(32420)
y = rlnorm(n)
median(y)
[1] 1.03435
```


BUILDING QUANTILE REGRESSION

Median

► Solve by Linear Programming

```
library(lpSolve)
A1 = cbind(diag(2*n), 0)
A2 = cbind(diag(n), -diag(n), 1)
r = lp("min", c(rep(1, 2*n), 0),
        rbind(A1, A2), c(rep(">=", 2*n),
                          rep("=", n)), c(rep(0, 2*n), y))
tail(r$solution, 1)
[1] 1.03435
```

BUILDING QUANTILE REGRESSION

Quantile

- Now Change the equation into quantile regression,

$$\min_{\mu, a, b} \left\{ \sum_{i=1}^n \tau a_i + (1 - \tau) b_i \right\}$$

subject to $a_i, b_i \geq 0$ and $y_i - [\beta_0^\tau + \beta_1^\tau x_i] = a_i - b_i \quad \forall 1, \dots, n$

BUILDING QUANTILE REGRESSION

Quantile

► Pre-built Function

```
tau = .3
quantile(y,tau)
30% 0.5775248
```

► Solve by Linear Programming

```
A1 = cbind(diag(2*n), 0)
A2 = cbind(diag(n), -diag(n), 1)
r = lp("min", c(rep(tau,n), rep(1-tau,n), 0),
      rbind(A1, A2), c(rep(">=", 2*n),
      rep("=", n)), c(rep(0, 2*n), y))
tail(r$solution, 1)
30% 0.5775248
```

BUILDING QUANTILE REGRESSION

Quantile

- ▶ Other than above programming, there are several more approaches.
- ▶ If interested, check out more on this page.
[Quantile Regression Computation: From the Inside and the Outside](#)
- ▶ Instead building your own function, we can use R package "quantreg".

OUTLIER PROTECTION ON MEDIAN REGRESSION

Basic Setting

- ▶ Import necessary packages.

```
library(MASS)
library(quantreg)
set.seed(32420)
```

- ▶ Same seed will be used repeatedly.
- ▶ $n_1 = 1000$ and $n_2 = 200$ for this trial.

OUTLIER PROTECTION ON MEDIAN REGRESSION

Generate Data Sample

- Generate n_1 of Good Observation.

```
data1 <- mvrnorm(n=n1,mu=c(0,0),Sigma =  
matrix(c(1,0.8,0.8,1),ncol=2))
```

- Generate n_2 of Bad Observation.

```
data2 <- mvrnorm(n=n2,mu=c(1.5,-1.5), Sigma  
= .2*diag(c(1,1)))
```

OUTLIER PROTECTION ON MEDIAN REGRESSION

Generate Data Sample

- Bind the Data and turn it into data.frame.

```
data <- rbind(data1, data2)
data <- data.frame(data)
```

- Distinguish the Good and Bad using Indicator vector.

```
names(data) <- c("X", "Y")
ind <- c(rep(1, n1), rep(2, n2))
```

OUTLIER PROTECTION ON MEDIAN REGRESSION

Drawing Scatter Plot

- Draw plot of the generated Data Sample.

```
plot(Y ~ X, data, pch=c("x", "o")[ind],  
      col=c("black", "red")[ind],  
      main=paste("N1 =", n1, "N2 =", n2))
```


OUTLIER PROTECTION ON MEDIAN REGRESSION

Data Plot




Figure 2: Scatter Plot

OUTLIER PROTECTION ON MEDIAN REGRESSION

Data Plot

- Fit and Draw Quantile Regression of $\tau=0.5$.

```
r1 <- rq(Y~X,data=data,tau=0.5)
abline(r1)
summary(r1)
```

- See Summary if you need.

OUTLIER PROTECTION ON MEDIAN REGRESSION

Drawing Quantile Regression Plot




Figure 3: Plot with QR

OUTLIER PROTECTION ON MEDIAN REGRESSION

Comparison Data Plot

- Draw Ordinary Least Square.

```
abline(lm(Y~X,data),lty=2, col="red")
```

- Draw Ordinary Least Square on Good.

```
abline(lm(Y~X,data,subset= 1:n1),
       lty=1,col='blue'))
```

- Draw Topleft Index.

```
legend("topleft",c("L1","ols","ols on good"),
      inset=0.02, lty=c(1,2,1),
      col=c("black","red","blue"),cex=.9)
```

OUTLIER PROTECTION ON MEDIAN REGRESSION

Data Plot




Figure 4: Final Comparison Plot

OUTLIER PROTECTION ON MEDIAN REGRESSION

Comparison Data Plot

- ▶ Produce 4 simulation plot with difference amount of bad observation.
- ▶ By observing several graph, we can guess how much outlier can be bear-ed on median regression.
- ▶ In this simulation, 1000 good observations could bear till 371 bad observations.

OUTLIER PROTECTION ON MEDIAN REGRESSION

Merge into One Function

```
qresult <- function(n1,n2){
  data <- mvrnorm(n=n1,mu=c(0,0),
    Sigma = matrix(c(1,0.8,0.8,1),ncol=2))
  data <- rbind(data,mvrnorm(n=n2,mu=c(1.5,-1.5),
    Sigma = .2*diag(c(1,1))))
  data <- data.frame(data) names(data) <- c("X","Y")
  ind <- c(rep(1,n1),rep(2,n2))
  plot(Y~X,data,pch=c("x","o")[ind], col=c("black","red")[ind],
    main=paste("N1 =",n1,"N2 =",n2))
  summary(r1 <- rq(Y~X,data=data,tau=0.5))
  abline(r1)
  abline(lm(Y~X,data),lty=2, col="red")
  abline(lm(Y~X,data,subset= 1:n1),lty=1,col='blue')
  legend("topleft",c("L1","ols","ols on good"),
    inset=0.02, lty=c(1,2,1),
    col=c("black","red","blue"),cex=.9)}
```

OUTLIER PROTECTION ON MEDIAN REGRESSION

Data Plot




Figure 5: Final Comparison Plot

OUTLIER PROTECTION ON MEDIAN REGRESSION

Data Plot




Figure 6: Final Comparison Plot

OUTLIER PROTECTION ON MEDIAN REGRESSION

Heteroscedasticity



- ▶ Simply to check, Heteroscedasticity, check out the Quantile Regression line.
- ▶ If they look like having different slope, probably there will be Heteroscedasticity.

Predictive Interval

- ▶ For different quantiles, there will be different predictive intervals obviously.
- ▶ By plotting quantiles, the accuracy of OLS interval can be checked.



OUTLIER PROTECTION ON MEDIAN REGRESSION

Heteroscedasticity



OUTLIER PROTECTION ON MEDIAN REGRESSION

Heteroscedasticity - Residual Plot



OUTLIER PROTECTION ON MEDIAN REGRESSION

Predictive Intervals





- Summary Plot of the Intercept and Coefficient.

```
plot(summary(rq(Y~X, data1,  
               tau=2:98/100)))
```

- The horizontal line is the OLS estimate.
- Dashed lines for confidence interval for OLS estimate.

OUTLIER PROTECTION ON MEDIAN REGRESSION

Predictive Intervals



QUANTILE REGRESSION ON ENGEL DATA

Engel's Law

- ▶ As income rises, the proportion of income spent on food falls, even if absolute expenditure on food rises.
- ▶ Empirical Law based on observation.
- ▶ This example shows expenditures on food as a function of income for 19th century Belgian households.

QUANTILE REGRESSION ON ENGEL DATA

Plot Data

- ▶ Engel Data is pre-included in the R package 'quantreg'.

```
data(engel)
```

- ▶ Draw the Scatter plot.

```
plot(foodexp~income, engel,  
      cex=.5,xlab="Household Income",  
      ylab="Food Expenditure")
```

- ▶ Draw the OLS on the plot.

```
abline(lm(foodexp~income,engel))
```


QUANTILE REGRESSION ON ENGEL DATA

Data Plot




Figure 10: Scatter Plot and OLS

QUANTILE REGRESSION ON ENGEL DATA

Plot Quantile Regression

- ▶ Draw Median Regression on the plot.

```
abline(rq(foodexp~income, engel, tau=.5),  
col="blue")
```

- ▶ Draw the 10-20-75-90 Quantile Regression on the plot.

```
taus <- c(.1, .25, .75, .90)  
for( i in 1:length(taus)){  
  abline(rq(foodexp~income, engel, tau=taus[i]), )  
    col="gray") }
```

QUANTILE REGRESSION ON ENGEL DATA

Data Plot




Figure 11: Quantile Regression

QUANTILE REGRESSION ON ENGEL DATA

Plot Summary

- Summary Plot of the Intercept and Coefficient.

```
plot(summary(rq(foodexp~income, engel,  
               tau=2:98/100)))
```

- The horizontal line is the OLS estimate.
- Dashed lines for confidence interval for OLS estimate.

QUANTILE REGRESSION ON ENGEL DATA

Data Plot




Figure 12: Summary Plot of Quantile Regression

QUANTILE REGRESSION ON ENGEL DATA

Log Transformation Case of Engel Data




Figure 13: Summary Plot of Log10 Quantile Regression

QUANTILE REGRESSION ON WAGE DATA

Wage Data from ISLR

- ▶ R package 'ISLR' contains only data, for use of the book "Introduction to Statistical Learning with applications in R".
- ▶ Load Wage Data from the package 'ISLR'.

```
library(ISLR)  
data(Wage)
```

- ▶ Rest of procedure will be same as the previous case.

QUANTILE REGRESSION ON WAGE DATA

Data Plot




Figure 14: Scatter Plot and OLS

QUANTILE REGRESSION ON WAGE DATA

Data Plot




Figure 15: Quantile Regression

QUANTILE REGRESSION ON WAGE DATA

Data Plot




Figure 16: Summary Plot of Quantile Regression

QUANTILE REGRESSION ON WAGE DATA

Log Transformation Case of Wage Data




Figure 17: Summary Plot of Log10 Quantile Regression