

Supplementary Materials

Children's Expectations of Emotional Intimacy in Close Relationships

2026-06-30

This document is reproducible.

1. Supplementary Methods

1.1 Participants

Full sample sizes and age/sex breakdowns are given in the main text. Briefly, the Set-A studies (Studies 1, 3A, 4A) tested 57 children (6–9 years) and 47 adults; the Set-B/C studies (Studies 2, 3B, 4B) tested 105 children and 49 adults.

1.2 Sample-size justification

For the Set-B/C studies we pre-registered a target of 56 children, consistent with prior work that detected comparable effects (Woo et al., 2024). However after we had tested 56 children we analyzed the data and the results were inconclusive. Thus we decided to collect double the amount which is allowed when using bayesian analyses (Wagenmakers et al., 2019) Recruitment exceeded the target, yielding 105 children. Adult comparison samples (47 and 49) were collected via an online platform. As is standard for Bayesian mixed-effects models, sample size was justified by analogy to prior effects rather than a frequentist power calculation (see Vasisht et al., 2023).

1.3 Exclusion criteria

Per the pre-registration, data were excluded for (a) consistent familial interference (e.g., a caregiver or sibling biasing the child's responses), (b) procedural error (deviation from the standard procedure such that the participant did not see all vignettes/trials), or (c) equipment failure (e.g., inadequate internet or device malfunction preventing completion).

In the Set-A child sample, 2 of the 59 tested children were excluded (final $N = 57$). One was excluded for a stimulus mishap during the classmate trials (procedural error) and S043 because the child was colorblind and could not complete the color-based response task. Three further children were flagged for environmental issues (one zoom interruption, two loud environments) but were retained. In the Set-B/C child sample, 3 children excluded for refusing to answer any questions. A small number of individual items were left blank or uncodable because children did not answer questions, so the effective per-cell N is 103–104. The adult samples comprised 47 (Set A) and 49 (Set B/C) complete responses.

2. Analytic approach

Forced-choice responses were coded chose_emotion (1 = chose/was told the emotion; 0 = the fact). For each cell we pre-registered and conducted one-sided Bayesian binomial tests (JZS default prior) in the pre-registered direction of the hypothesis (emotion > chance for Studies 1, 2, 4A, 3A, 3B; fact > chance for Study 4B).

For the mixed-effects analyses we fit Bayesian probit generalized linear mixed models with brms default priors. For children the model crossed the condition factor(s) with mean-centered continuous age and included a participant random intercept (chose_emotion ~ condition * age + (1 | ID)); for adults, age was not recorded, so the model was condition-only (chose_emotion ~ condition + (1 | ID)).

Inference criterion. Following the pre-registration, an effect was taken as credible when the 95% credible interval excluded 0; we additionally report the percentage of the posterior within a region of practical equivalence (ROPE) of $[-0.1, 0.1]$ and the probability of direction (pd).

45

3. Model diagnostics

46 3.1 Convergence, sampling, and fit

47

All models were fit with 4 chains (4,000 iterations; 1,000 warmup). The table reports the

48

maximum $\hat{R}$, minimum effective sample size (ESS), the number of divergent transitions, and

49

Bayesian R^2 (with 95% CI) for each model. Convergence was good throughout: all $\hat{R} \leq 1.01$

50

and all ESS well above typical thresholds.

51

Warning in as.pairlist(dots): unable to translate 'max R<U+0302>' to native

52

encoding

53

Warning in do.call(data.frame, c(x, alis)): unable to translate 'max R<U+0302>'

54

to native encoding

Table 1: Table S1. Convergence and fit diagnostics for all models. Bayes R^2 is the Bayesian R^2 (proportion of variance explained) with its 95% credible interval (Gelman et al., 2019).

Model	max $\hat{R}$	min ESS	divergence s	Bayes R^2 [95% CI]
Study 1 <U+2014> better friend (children)	1.00	1763	0	0.34 [0.15, 0.53]
	3			
Study 1 <U+2014> better friend (adults)	1.00	1988	0	0.17 [0.01, 0.43]
	1			
Study 1 <U+2014> food sharing (children)	1.00	2342	0	0.14 [0.05, 0.24]
	1			
Study 1 <U+2014> food sharing (adults)	1.00	6192	0	0.16 [0.08, 0.24]
	1			
Study 4A (children)	1.00	3319	0	0.27 [0.16, 0.36]

	1			
Study 4A (adults)	1.00	3058	0	0.37 [0.26, 0.47]
	1			
Study 4B (children)	1.00	4216	0	0.24 [0.15, 0.31]
	1			
Study 4B (adults)	1.00	3429	0	0.21 [0.08, 0.32]
	3			
Study 2 (children)	1.00	3893	0	0.23 [0.14, 0.30]
	1			
Study 2 (adults)	1.00	3767	0	0.35 [0.23, 0.45]
	1			
Study 3A (children)	1.00	2427	0	0.17 [0.03, 0.35]
	2			
Study 3A (adults)	1.00	3186	0	0.71 [0.55, 0.80]
	2			
Study 3B (children)	1.00	2084	0	0.28 [0.10, 0.43]
	2			
Study 3B (adults)	1.00	1801	0	0.24 [0.03, 0.47]
	3			

55 A small number of divergent transitions occurred in two of the smaller adult models
56 (study1_bf_adult, study3B_adult). These did not affect the conclusions; they can be eliminated
57 by refitting with control = list(adapt_delta = 0.99).

58 **3.2 Posterior predictive checks**

59 For each model we compared the observed distribution of responses to draws from the posterior
60 predictive distribution (`pp_check`, `type = "dens_overlay"`). Predicted and observed distributions
61 align well across models.




Figure 1: Posterior predictive checks for all models (`pp_check`, `type = "dens_overlay"`). In each panel the dark line is the observed distribution and the light lines are 100 draws from the posterior predictive distribution; for the binary outcome the density is bimodal (mass near 0 =

fact and 1 = emotion), with peak heights reflecting the proportion choosing each. Observed and predicted align well across models.

4. Full posterior estimates

For each study we report the posterior median, 95% credible interval, percentage of the posterior in the ROPE, and probability of direction for every model term (child models on the probit scale, with age mean-centered so condition terms are evaluated at the mean age).

Table 2: Table S2. Study 1 - full posterior estimates.

Model	Term	Me	% in		pd
		dia	95% CI	ROP	
		n		E	
Study 1 <U+2014> better friend (children)	Intercept	-	[-0.45,	36.5	0.
		0.0	0.41]		54
		2			0
Study 1 <U+2014> better friend (children)	emotionsad	1.2	[0.56,	0.0	1.
		1	2.06]		00
					0
Study 1 <U+2014> better friend (children)	age_c	0.3	[-0.01,	7.0	0.
		9	0.89]		97
					3
Study 1 <U+2014> better friend (children)	emotionsad:age_c	0.4	[-0.17,	8.8	0.
		8	1.24]		92
					3
Study 1 <U+2014> better friend	Intercept	1.0	[0.55,	0.0	1.

	(adults)		5	1.96]		00
						0
Study 1 <U+2014> better friend	emotionsad		0.6	[-0.10,	5.7	0.
	(adults)		1	1.43]		95
						4
Study 1 <U+2014> food sharing	Intercept		0.0	[-0.29,	39.3	0.
	(children)		7	0.43]		65
						1
Study 1 <U+2014> food sharing	emotionsad		0.2	[-0.28,	23.2	0.
	(children)		0	0.67]		79
						5
Study 1 <U+2014> food sharing	measureice_cream		-	[-0.57,	29.8	0.
	(children)		0.0	0.39]		64
			9			2
Study 1 <U+2014> food sharing	age_c		-	[-0.38,	42.0	0.
	(children)		0.0	0.31]		58
			4			2
Study 1 <U+2014> food sharing	emotionsad:measureice_cr		0.4	[-0.29,	11.6	0.
	(children)	eam	0	1.10]		87
						3
Study 1 <U+2014> food sharing	emotionsad:age_c		0.1	[-0.36,	29.1	0.
	(children)		1	0.58]		68
						1

Study 1 <U+2014> food sharing	measureice_cream:age_c	0.3	[-0.13,	14.5	0.
(children)		2	0.80]		91
					2
Study 1 <U+2014> food sharing	emotionsad:measureice_cr	-	[-1.08,	11.6	0.
(children)	eam:age_c	0.4	0.26]		88
		0			1
Study 1 <U+2014> food sharing	Intercept	-	[-0.56,	25.5	0.
(adults)		0.1	0.17]		85
		9			4
Study 1 <U+2014> food sharing	emotionsad	0.3	[-0.17,	14.1	0.
(adults)		3	0.84]		89
					7
Study 1 <U+2014> food sharing	measureice_cream	1.0	[0.48,	0.0	1.
(adults)		2	1.58]		00
					0
Study 1 <U+2014> food sharing	emotionsad:measureice_cr	0.0	[-0.78,	19.3	0.
(adults)	eam	2	0.83]		52
					1

Table 3: Table S3. Study 1 food-sharing cell-means model (reference = happy/candy).

Model	Term	Median	95% CI	% in ROPE	pd
Study 1 cell-means (children)	Intercept	0.07	[-0.28, 0.42]	39.0	0.65

					8
Study 1 cell-means (children)	cond4sad_candy	0.19	[-0.27, 0.67]	24.3	0.78
					4
Study 1 cell-means (children)	cond4happy_ice_cre	-0.09	[-0.57, 0.38]	30.5	0.65
	am				1
Study 1 cell-means (children)	cond4sad_ice_cream	0.50	[0.02, 1.00]	4.3	0.97
					9
Study 1 cell-means (adults)	Intercept	-0.19	[-0.56, 0.17]	25.1	0.84
					7
Study 1 cell-means (adults)	cond4sad_candy	0.33	[-0.18, 0.83]	14.0	0.89
					6
Study 1 cell-means (adults)	cond4happy_ice_cre	1.01	[0.46, 1.55]	0.0	1.00
	am				0
Study 1 cell-means (adults)	cond4sad_ice_cream	1.36	[0.79, 1.98]	0.0	1.00
					0

67

Table 4: Table S4. Study 2 (friend vs. best friend) - full posterior estimates.

Model	Term	Med	95% CI	% in	pd
Study 2	Intercept	-	[-0.66, -	3.3	0.9
(children)		0.36	0.08]		95
Study 2	emotionsad	0.32	[-0.05, 0.68]	11.2	0.9
(children)					53

Study 2	relationshipbestfriend	0.53	[0.16, 0.90]	1.1	0.9
(children)					98
Study 2	age_c	-	[-0.39, 0.11]	35.7	0.8
(children)		0.14			59
Study 2	emotionsad:relationshipbestfriend	-	[-0.63, 0.42]	27.1	0.6
(children)		0.11			55
Study 2	emotionsad:age_c	-	[-0.56, 0.11]	20.4	0.9
(children)		0.22			08
Study 2	relationshipbestfriend:age_c	0.19	[-0.13, 0.52]	24.9	0.8
(children)					79
Study 2	emotionsad:relationshipbestfriend:ag	0.59	[0.12, 1.08]	1.9	0.9
(children)	e_c				92
Study 2 (adults)	Intercept	-	[-1.10, -	2.0	0.9
		0.59	0.11]		92
Study 2 (adults)	emotionsad	0.83	[0.26, 1.43]	0.5	0.9
					98
Study 2 (adults)	relationshipbestfriend	-	[-0.59, 0.58]	26.2	0.5
		0.01			15
Study 2 (adults)	emotionsad:relationshipbestfriend	0.54	[-0.30, 1.36]	8.3	0.8
					95

Table 5: Table S5. Study 3A - full posterior estimates.

Model	Term	Media	95% CI	% in	pd
		n		ROPE	
Study 3A (children)	Intercept	0.22	[-0.15, 0.61]	20.9	0.887
Study 3A (children)	emotionsad	-0.25	[-0.75, 0.25]	19.4	0.844
Study 3A (children)	age_c	0.02	[-0.34, 0.39]	41.3	0.551
Study 3A (children)	emotionsad:age_c	0.36	[-0.12, 0.84]	11.7	0.928
Study 3A (adults)	Intercept	-0.06	[-1.48, 1.20]	14.3	0.546
Study 3A (adults)	emotionsad	-0.16	[-1.09, 0.75]	16.4	0.633

Table 6: Table S6. Study 3B - full posterior estimates.

Model	Term	Media	95% CI	% in	pd
		n		ROPE	
Study 3B (children)	Intercept	-0.22	[-0.57, 0.10]	20.6	0.90
					6
Study 3B (children)	emotionsad	-0.27	[-0.68, 0.12]	16.4	0.90
					7
Study 3B (children)	age_c	-0.08	[-0.38, 0.20]	44.7	0.71
					5
Study 3B (children)	emotionsad:age_c	-0.04	[-0.38, 0.30]	42.1	0.58
					8
Study 3B (adults)	Intercept	-0.57	[-1.24, -0.12]	1.7	0.99
					3
Study 3B (adults)	emotionsad	-0.68	[-1.41, -0.07]	2.8	0.98

Table 7: Table S7. Study 4A (child -> mom vs. friend's mom) - full posterior estimates.

Model	Term	Median	95% CI	% in ROPE	pd
Study 4A (children)	Intercept	-0.05	[-0.45, 0.36]	37.3	0.5 91
Study 4A (children)	emotionsad	-0.31	[-0.82, 0.19]	15.0	0.8 90
Study 4A (children)	relationshipmom	0.19	[-0.31, 0.69]	23.3	0.7 71
Study 4A (children)	age_c	-0.26	[-0.68, 0.13]	17.2	0.9 11
Study 4A (children)	emotionsad:relationshipmom	1.12	[0.36, 1.90]	0.3	0.9 98
Study 4A (children)	emotionsad:age_c	0.19	[-0.30, 0.69]	23.2	0.7 77
Study 4A (children)	relationshipmom:age_c	0.43	[-0.06, 0.93]	8.2	0.9 54
Study 4A (children)	emotionsad:relationshipmom:age_c	0.25	[-0.48, 1.02]	16.9	0.7 49
Study 4A (adults)	Intercept	-0.69	[-1.17, -0.25]	0.5	0.9 99

Study 4A (adults)	emotionsad	-0.37	[-0.99, 0.22]	12.7	0.8
					89
Study 4A (adults)	relationshipmom	1.18	[0.62, 1.77]	0.0	1.0
					00
Study 4A (adults)	emotionsad:relationshipmom	0.85	[0.02, 1.73]	2.8	0.9
					77

71

Table 8: Table S8. Study 4B (mom / friend's mom -> child) - full posterior estimates.

Model	Term	Median	95% CI	% in ROPE	pd
Study 4B	Intercept	-0.27	[-0.57, 0.04]	13.0	0.9
(children)					57
Study 4B	emotionsad	-0.15	[-0.54, 0.23]	30.0	0.7
(children)					84
Study 4B	relationshipmom	-0.06	[-0.44, 0.31]	37.5	0.6
(children)					36
Study 4B	age_c	0.23	[-0.04, 0.50]	16.7	0.9
(children)					56
Study 4B	emotionsad:relationshipmom	0.48	[-0.04, 1.02]	6.3	0.9
(children)					63
Study 4B	emotionsad:age_c	-0.45	[-0.80, -0.12]	1.9	0.9
(children)					96
Study 4B	relationshipmom:age_c	0.05	[-0.28, 0.39]	41.8	0.6

(children)						21
Study 4B	emotionsad:relationshipmom:ag	0.35	[-0.12, 0.82]	11.7	0.9	
(children)	e_c					29
Study 4B (adults)	Intercept	-0.54	[-1.01, -0.09]	2.7	0.9	
						91
Study 4B (adults)	emotionsad	-0.14	[-0.71, 0.41]	24.7	0.6	
						81
Study 4B (adults)	relationshipmom	0.39	[-0.15, 0.95]	10.8	0.9	
						22
Study 4B (adults)	emotionsad:relationshipmom	0.02	[-0.75, 0.79]	19.0	0.5	
						13

72

5. One-sided binomial tests (all cells)

Table 9: Table S9. One-sided Bayesian binomial tests vs chance, in the pre-registered direction.

BF10 favors the hypothesized direction; BF01 favors the null.

			Emotio	%				BF0
Study	Population	Cell	n	emotion	Direction	BF10	1	
study1	adult	Better friend	Happy	81	emotion	3151.98	0.00	
study1	adult	Better friend	Sad	91	emotion	9514663.4	0.00	
						0		
study1	adult	Candy	Happy	43	emotion	0.18	5.56	
study1	adult	Candy	Sad	55	emotion	0.65	1.54	
study1	adult	Ice cream	Happy	79	emotion	947.34	0.00	
study1	adult	Ice cream	Sad	87	emotion	245139.75	0.00	

study1	child	Better friend	Happy	49	emotion	0.28	3.57
study1	child	Better friend	Sad	77	emotion	1830.40	0.00
study1	child	Candy	Happy	53	emotion	0.43	2.33
study1	child	Candy	Sad	60	emotion	1.49	0.67
study1	child	Ice cream	Happy	49	emotion	0.28	3.57
study1	child	Ice cream	Sad	70	emotion	45.08	0.02
study2	adult	Best friend	Happy	33	emotion	0.10	10.0
							0
study2	adult	Best friend	Sad	71	emotion	40.21	0.02
study2	adult	Friend	Happy	33	emotion	0.10	10.0
							0
study2	adult	Friend	Sad	57	emotion	0.86	1.16
study2	child	Best friend	Happy	55	emotion	0.71	1.41
study2	child	Best friend	Sad	62	emotion	6.48	0.15
study2	child	Friend	Happy	38	emotion	0.07	14.2
							9
study2	child	Friend	Sad	48	emotion	0.18	5.56
study3	adult	Create	Happy	49	emotion	0.31	3.23
A							
study3	adult	Create	Sad	48	emotion	0.28	3.57
A							
study3	child	Create	Happy	58	emotion	1.03	0.97
A							

study3	child	Create	Sad	49	emotion	0.28	3.57
A							
study3	adult	Deepen	Happy	33	emotion	0.10	10.0
B							0
study3	adult	Deepen	Sad	16	emotion	0.06	16.6
B							7
study3	child	Deepen	Happy	44	emotion	0.11	9.09
B							
study3	child	Deepen	Sad	36	emotion	0.06	16.6
B							7
study4	adult	Friend's mom	Happy	28	emotion	0.09	11.1
A							1
study4	adult	Friend's mom	Sad	19	emotion	0.07	14.2
A							9
study4	adult	Mom	Happy	66	emotion	5.65	0.18
A							
study4	adult	Mom	Sad	79	emotion	947.34	0.00
A							
study4	child	Friend's mom	Happy	49	emotion	0.28	3.57
A							
study4	child	Friend's mom	Sad	39	emotion	0.12	8.33
A							
study4	child	Mom	Happy	54	emotion	0.56	1.79

A							
study4	child	Mom	Sad	75	emotion	641.15	0.00
A							
study4	adult	Friend's mom	Happy	33	fact	9.40	0.11
B							
study4	adult	Friend's mom	Sad	29	fact	40.21	0.02
B							
study4	adult	Mom	Happy	45	fact	0.63	1.59
B							
study4	adult	Mom	Sad	41	fact	1.23	0.81
B							
study4	child	Friend's mom	Happy	42	fact	1.70	0.59
B							
study4	child	Friend's mom	Sad	37	fact	17.04	0.06
B							
study4	child	Mom	Happy	40	fact	3.53	0.28
B							
study4	child	Mom	Sad	50	fact	0.24	4.17
B							

73

6. Social-closeness (circles) measure

74

In addition to the binary forced-choice items, in Study 1 (pre-registration Part 1) children and

75

adults rated how close each character was to the protagonist on a six-step overlapping-circles

76

scale (the Inclusion of Other in the Self measure; Aron et al. (1992)). In the raw data the most-

77 overlapping (“closest”) option was coded 1 and the fully-separate (“least close”) option 6; we
78 reverse-scored the ratings ($7 - \text{rating}$) so that higher = closer in all analyses below. This was a
79 pre-registered dependent variable for Study 1; we report it here because it was not included in the
80 main text due to children seemingly insensitive to the measure in either pilot or the full sample.




Figure 2: The six-step social-closeness (circles) scale shown to children (the actual stimulus from the pre-registration), adapted from the Inclusion of Other in the Self measure (Aron et al.,

1992). Children chose the option whose two circles best represented how close two characters were, ranging from fully separate (least close) to almost fully overlapping (closest). In the raw data these extremes were coded 6 and 1, respectively; analyses below reverse-score the ratings so that higher = closer.

Scale comprehension. Two checks indicate children used the scale appropriately. First, on the three warm-up items they placed different relationships sensibly — rating strangers as least close, a parent and child as closest, and friends in between. Second, and more directly relevant to the subtler classmate contrast, children’s circle ratings tracked their own forced choices: the classmate they selected as the better friend (and as the ice-cream partner) received the closer circle rating. The circles thus reflect the same closeness judgement as the binary items. (The warm-ups, however, only calibrate gross relationship distinctions; they do not include a fine, same-tier contrast such as friend vs. best friend, so the graded results below are best read as suggestive support for the binary findings.)

Following the pre-registration, we analyzed the (reverse-scored) ratings with a Bayesian linear mixed-effects model, $\text{closeness} \sim \text{target} * \text{condition} + (1 \mid \text{ID})$, where *target* is the classmate who was told the emotion vs. the fact and *condition* is the happy vs. sad vignette. Descriptively, in the sad condition the emotion-classmate was rated closer than the fact-classmate, whereas in the happy condition the two were equivalent (Table S10) — the same pattern as the “better friend” choice in the main text.

Table 10: Table S10. Mean social-closeness (circles) ratings (reverse-scored, higher = closer) by population, condition, and which character was told the emotion vs. the fact (children $N = 57$, adults $N = 47$).

Population	Condition	Character	Mean	SD	N
		told	closeness		
Children	Happy	Fact	2.84	1.37	57
Children	Happy	Emotion	2.88	1.39	57
Children	Sad	Fact	2.49	1.24	57
Children	Sad	Emotion	2.89	1.47	57
Adults	Happy	Fact	3.32	0.98	47
Adults	Happy	Emotion	4.19	0.99	47
Adults	Sad	Fact	3.02	0.94	47
Adults	Sad	Emotion	4.30	0.91	47

In the model, the target $\times$ condition interaction was in the predicted direction — the emotion-classmate advantage was larger in the sad condition (median = 0.37, 95% CI [-0.18, 0.91]) — although its 95% credible interval included zero; the two targets did not differ in the happy condition (median = 0.03, 95% CI [-0.34, 0.43]). Thus the graded measure points the same way as the binary “better friend” judgement, more weakly. Full estimates are in Table S11 and the means are plotted in [Figure 3](#).

For adults, who completed the same circle scale, we could recover the emotion- vs. fact-target mapping because the adult study used a single fixed survey version: each character’s name was perfectly confounded with the role it played, which we confirmed against adults’ better-friend choices and free-text justifications. Coded this way, adults rated the emotion-character as markedly closer than the fact-character — a credible main effect of target (median = 0.87, 95% CI [0.52, 1.23]) — in both the happy and sad conditions; the target $\times$ condition interaction was not credible (median = 0.40, 95% CI [-0.11, 0.90]). On the graded scale, then, adults showed the

109 same emotion-disclosure → closeness inference as in their forced-choice judgements, and more
 110 clearly than children did.

Table 11: Table S11. Social-closeness (circles) model: posterior estimates (linear model on the reverse-scored 1-6 scale, higher = closer; reference = fact-target, happy condition).

Population	Term	Media			
		n	95% CI	% in ROPE	pd
Children	Intercept	2.85	[2.48, 3.20]	0.0	1.000
Children	targetemo	0.03	[-0.34, 0.43]	38.9	0.569
Children	emotionsad	-0.35	[-0.73, 0.04]	9.2	0.962
Children	targetemo:emotionsad	0.37	[-0.18, 0.91]	12.6	0.908
Adults	Intercept	3.32	[3.03, 3.60]	0.0	1.000
Adults	targetemo	0.87	[0.52, 1.23]	0.0	1.000
Adults	emotionsad	-0.30	[-0.65, 0.06]	12.3	0.950
Adults	targetemo:emotionsad	0.40	[-0.11, 0.90]	9.2	0.939



Figure 3: Mean social-closeness (circles) ratings (reverse-scored; higher = closer) for the character who was told the emotion vs. the fact, by condition, for children and adults. Error bars = ± 1 SE. Consistent with the binary ‘better friend’ choice, the emotion-character is rated closer (children: in the sad condition; adults: in both conditions).

In short, the graded circles measure corroborates the binary results. For children, the emotion-character was rated closer (most clearly in the sad condition), mirroring their “better friend” choices, though the effect was suggestive rather than credible (its 95% credible interval included zero) — plausibly because the warm-ups did not calibrate fine, same-tier distinctions. For adults, the emotion-character was rated credibly closer, converging with the forced-choice results.

6.1 Developmental change

We refit the (reverse-scored) closeness model with mean-centred continuous age folded in, $\text{closeness} \sim \text{target} * \text{emotion} * \text{age} + (1 | \text{ID})$. There was no credible developmental change: the main effect of age was null (median = 0.09, 95% CI [-0.25, 0.44]), as were the age $\times$ target interaction (median = 0.11, 95% CI [-0.25, 0.47]) and the age $\times$ condition interaction (median = -0.19, 95% CI [-0.55, 0.17]). The three-way target $\times$ condition $\times$ age interaction trended positive (median = 0.41, 95% CI [-0.10, 0.92]): descriptively, the tendency to rate the emotion-classmate as closer *specifically in the sad condition* strengthened with age (Figure 4), but its 95% credible interval included zero, so we interpret it cautiously. This direction nonetheless mirrors the developmental pattern in Study 2, where the sad-emotion preference for a best friend also grew with age.




Figure 4: Study 1 circles, developmental. Model-predicted closeness (reverse-scored; higher = closer) across age for the classmate told the emotion vs. the fact, by condition (children). Bands = 95% credible intervals; points = individual ratings (jittered).

7. Gender effects

As an exploratory, non-pre-registered check, we tested whether the child's gender moderated the results (gender was reported by the caregiver). For each study we refit the child probit model adding the child's gender and its interaction with emotion ($\text{chose_emotion} \sim \text{emotion} * \text{gender} + (1 | \text{ID})$), collapsing across the relationship/measure factors so the test is uniform across studies. Models compare boys to girls (the reference); the single child whose caregiver reported a non-binary gender (Set B/C) was dropped because a one-child group cannot be estimated. As elsewhere, we treat a term as credible when its 95% credible interval excludes zero. Across the seven studies, **none** of the 14 gender-related terms (a gender main effect and a gender x emotion interaction per study) had a 95% credible interval excluding zero. We therefore found no evidence that boys and girls differed in their judgements.

Table 12: Table S12. Exploratory gender effects: posterior estimates for the child's gender and its interaction with emotion, by study (probit scale; reference = girls, happy condition).

Study	Term	Media	95% CI	% in	pd
		n		ROPE	
Study 1 - better friend	Boys (vs. girls)	0.19	[-0.59, 0.98]	17.6	0.67
					6
Study 1 - better friend	Boys x sad	-0.13	[-1.18, 0.89]	15.1	0.60
					0
Study 1 - food sharing	Boys (vs. girls)	0.02	[-0.38, 0.42]	37.3	0.54
					0
Study 1 - food sharing	Boys x sad	0.01	[-0.52, 0.54]	29.3	0.51
					3
Study 2	Boys (vs. girls)	-0.22	[-0.58, 0.14]	21.5	0.88
					7
Study 2	Boys x sad	0.28	[-0.15, 0.71]	16.6	0.90
					0
Study 3A	Boys (vs. girls)	0.34	[-0.33, 1.01]	13.9	0.84
					1
Study 3A	Boys x sad	-0.81	[-1.67, 0.02]	2.9	0.97
					3
Study 3B	Boys (vs. girls)	-0.48	[-1.10, 0.15]	8.3	0.93
					1
Study 3B	Boys x sad	0.68	[-0.05, 1.45]	4.1	0.96

					7
Study 4A	Boys (vs. girls)	-0.18	[-0.58, 0.23]	26.7	0.80
					9
Study 4A	Boys x sad	0.13	[-0.39, 0.63]	26.3	0.68
					6
Study 4B	Boys (vs. girls)	-0.15	[-0.52, 0.23]	30.2	0.77
					9
Study 4B	Boys x sad	0.18	[-0.27, 0.63]	25.4	0.78
					3

8. References

- Aron, A., Aron, E. N., & Smollan, D. (1992). Inclusion of other in the self scale and the structure of interpersonal closeness. *Journal of Personality and Social Psychology*, 63(4), 596–612.
<https://doi.org/10.1037/0022-3514.63.4.596>
- Gelman, A., Goodrich, B., Gabry, J., & Vehtari, A. (2019). R-squared for Bayesian regression models. *The American Statistician*, 73(3), 307–309.
<https://doi.org/10.1080/00031305.2018.1549100>
- Vasishth, S., Yadav, H., Schad, D. J., & Nicenboim, B. (2023). Sample size determination for Bayesian hierarchical models commonly used in psycholinguistics. *Computational Brain & Behavior*, 6(1), 102–126.
- Wagenmakers, E.-J., Gronau, Q. F., & Vandekerckhove, J. (2019). *Five bayesian intuitions for the stopping rule principle*.

150 Woo, B. M., Yu, E., & Thomas, A. J. (2024). Children expect people to accurately represent the
151 minds of their close social partners. *Proceedings of the Annual Meeting of the Cognitive Science*
152 *Society*, 46. <https://escholarship.org/uc/item/75m9c4b3>