

Batch details	Chennai-Sept-2025
Team Members	• Adlin Raja D • Adrian Jacob Rosario • Ksheerabdhinath M K G • Mohamed Thahar • Ashok S • Manideep Doddapaneni
Domain of Project	Health Care
Proposed project title	Machine Learning-Based Early Prediction of Medication Dosage Adjustment Risk in Hospitalized Diabetic Patients
Group Number	Group 1
Team Leader	Adrian Jacob Rosario
Mentor Name	Siddharth Koshta

Date: 04/03/2026

SL NO	Particulars	Page No
1	Overview	3
2	Business problem statement (GOALS)	3
3	Topic survey in depth	4
4	Critical assessment of topic survey	6
5	Methodology to be followed	5
6	References	9

Machine Learning-Based Early Prediction of Medication Dosage Adjustment Risk in Hospitalized Diabetic Patients

OVERVIEW

Hospitalized diabetic patients often experience fluctuating glucose levels and rapidly changing clinical conditions, requiring frequent medication dosage adjustments. Delayed or improper modifications can result in adverse drug reactions (ADRs), poor glycemic control, and increased complications. This project proposes the development of a machine learning classification model to predict patients who are more likely to require medication dosage adjustments during hospitalization. Early predictions can assist clinicians in prioritizing high-risk patients for medication review, enabling timely interventions and improving patient safety and treatment outcomes.

BUSINESS PROBLEM STATEMENT (GOALS)

1. Business Problem Understanding

Hospitalized diabetic patients frequently require medication dosage modifications due to unstable blood glucose levels and changing physiological conditions. In many cases, these adjustments are delayed because clinicians must monitor large volumes of patient data while managing multiple patients simultaneously. Delayed or incorrect dosage adjustments can lead to adverse drug reactions, poor glycemic control, prolonged hospital stays, and increased healthcare costs.

2. Business Objective

The objective of this project is to develop a machine learning classification model capable of predicting which hospitalized diabetic patients are more likely to require medication dosage adjustments during their hospital stay. Early prediction of high-risk patients can support clinicians in prioritizing medication review and monitoring, thereby reducing adverse drug reactions and improving treatment outcomes.

3. Approach

This project will utilize patient clinical and demographic data to train a supervised machine learning classification model. The dataset will be preprocessed and explored to identify relevant predictors influencing dosage adjustments. Various classification algorithms such as Logistic Regression, Random Forest, and

Decision Trees will be evaluated. Model performance will be assessed using metrics such as accuracy, sensitivity, precision, and F1 score to ensure reliable prediction of patients requiring dosage modification.

4. Conclusions

The proposed system can act as a clinical decision-support tool to help healthcare providers identify high-risk patients who may require dosage adjustments. This can lead to earlier clinical intervention, improved glycemic control, reduction in adverse drug reactions, and enhanced patient safety. Additionally, the model can provide insights into clinical and demographic factors influencing dosage modification, supporting data-driven clinical decision-making.

TOPIC SURVEY IN BRIEF

1. Problem Understanding

Diabetes management in hospitalized patients is complex due to fluctuating blood glucose levels caused by stress, illness, medication interactions, and changes in dietary intake. These fluctuations often require frequent adjustments in medication dosage to maintain optimal glycemic control. However, identifying patients who require dosage modifications in a timely manner is challenging for clinicians managing multiple patients simultaneously. Delayed dosage adjustments may result in hyperglycemia, hypoglycemia, or adverse drug reactions, ultimately affecting patient safety and increasing hospitalization costs.

2. Current Solutions to the Problem

Currently, dosage adjustments are primarily based on manual clinical monitoring and physician judgment. Clinicians review patient records, laboratory reports, and glucose readings before making dosage decisions. While this approach is clinically effective, it is time-consuming and may lead to delays in identifying patients who require immediate medication modification. Some hospitals use basic clinical decision-support systems that generate alerts based on glucose thresholds, but these systems often lack predictive capabilities.

3. Proposed Solution to the Problem

The proposed solution is to develop a machine learning-based predictive model capable of identifying patients who are likely to require medication dosage adjustments during their hospital stay. By analyzing patient demographic data, clinical variables, and laboratory parameters, the model can identify patterns associated with dosage modifications. This predictive approach allows clinicians to intervene earlier and prioritize high-risk patients for medication review. Such a system can serve as an intelligent clinical decision-support tool that improves patient monitoring and treatment efficiency.

4. Reference to the Problem

Recent studies in healthcare analytics have demonstrated the potential of machine learning models in predicting adverse drug reactions, hospital readmissions, and patient risk stratification. These models use clinical data to identify patterns that may not be easily detected through traditional analysis methods. Applying similar techniques to predict medication dosage adjustments in diabetic patients can significantly improve clinical decision-making and patient safety.

CRITICAL ASSESSMENT OF TOPIC SURVEY

Although current clinical monitoring methods help physicians adjust medication dosages, they largely rely on manual evaluation of patient data and do not provide predictive insights. Existing clinical decision-support systems often generate alerts only after abnormal glucose levels occur. There is a gap in predictive tools that can proactively identify patients who are likely to require dosage adjustments. This project addresses this gap by developing a machine learning model capable of early risk prediction, enabling clinicians to intervene sooner and improve patient safety.

METHODOLOGY TO BE FOLLOWED

The project will follow a structured data science workflow consisting of the following steps.

Step 1: Data Collection

Patient clinical and demographic data related to hospitalized diabetic patients will be collected from the dataset. Variables may include patient demographics, laboratory results, medication information, and hospitalization records.

Step 2: Data Preprocessing

The dataset will be cleaned to handle missing values, remove inconsistencies, and encode categorical variables. Feature scaling and normalization may also be applied to improve model performance.

Step 3: Exploratory Data Analysis (EDA)

Exploratory analysis will be conducted to understand the distribution of variables, identify correlations, and detect patterns associated with medication dosage adjustments. Visualizations such as histograms, boxplots, and correlation heatmaps will be used.

Step 4: Model Development

Several machine learning classification algorithms will be implemented, like:

- Logistic Regression
- Decision Tree
- Random Forest

The models will be trained using the prepared dataset to predict whether a patient is likely to require medication dosage adjustment.

Step 5: Model Evaluation and Interpretation

The models will be evaluated using performance metrics such like:


- Accuracy
- Precision
- Recall (Sensitivity)
- F1 Score

The best-performing model will be selected and interpreted to identify key predictors influencing dosage adjustment decisions

ML PIPELINE

Data Collection





REFERENCES

Clore, J., Cios, K., DeShazo, J., & Strack, B. (2014). Diabetes 130-US Hospitals for Years 1999-2008 [Dataset]. UCI Machine Learning Repository. <https://doi.org/10.24432/C5230J>.

Strack B, DeShazo JP, Gennings C, Olmo JL, Ventura S, Cios KJ, Clore JN. Impact of HbA1c measurement on hospital readmission rates: analysis of 70,000 clinical database patient records. *Biomed Res Int*. 2014;2014:781670. doi: 10.1155/2014/781670. Epub 2014 Apr 3. PMID: 24804245; PMCID: PMC3996476.

<https://archive.ics.uci.edu/dataset/296/diabetes+130-us+hospitals+for+years+1999-2008>