

Klippel-Trenaunay Syndrome Complicated by Kasabach-Merritt Syndrome During Pregnancy

A case report and review of the literature

Authors

Mourad Elfaham¹, Aya Attia¹, Mohamed Waheed¹, Mohamed Essam¹, Amgad Gamal¹, Salma Saad¹, Mariam Sarhan¹, Marwa Mourad¹, Mohamed Hegazy¹, Farah Said¹, Sarah Tarek¹, Sara Ibrahim Abdelkader^{2*}, Ahmed Azab², Mohammed Ghanem², Rahma Alaa Abdelhafez², Nour Atif², Maha Moemen², Belgin Ahmed Nagah², Ashraf Nabhan^{1,3}

Affiliation

¹ Faculty of Medicine, Ain Shams University, Cairo, Egypt.

² Faculty of Medicine, Galala University, Attaka, Suez, Egypt.

³ Faculty of Medicine, MTI University, Cairo, Egypt. 01069663999

* Corresponding author: Sara Ibrahim Abdelkader. Faculty of Medicine, Galala University, Attaka, Suez, Egypt. Email: sara.elzeftawy@gu.edu.eg, ORCID: [0009-0002-5877-4722](https://orcid.org/0009-0002-5877-4722)

Word count: 2500

Abstract

The co-occurrence of Klippel-Trenaunay syndrome and Kasabach-Merritt syndrome in pregnancy is extremely rare and represents a critical obstetric emergency with significant maternal and fetal risks, demanding early recognition and immediate multidisciplinary intervention.

We report the case of a 28-year-old pregnant woman with known Klippel-Trenaunay syndrome who developed Kasabach-Merritt phenomenon during pregnancy. The fetus suffered severe

growth restriction and absent end-diastolic flow on umbilical artery Doppler. A cesarean section was performed at 35 weeks of gestation.

This case highlights the complexity of managing pregnancy in women with Klippel–Trenaunay syndrome complicated by Kasabach Merritt syndrome and emphasizes the importance of multidisciplinary surveillance and timely delivery.

Keywords: Klippel–Trenaunay syndrome; angioosteohypertrophy syndrome; Kasabach Merritt syndrome; fetal growth restriction; high-risk pregnancy

Introduction

Klippel-Trenaunay syndrome (KTS) is a rare congenital disorder affecting an estimated 2–5 per 100,000 individuals. Its features include a triad of capillary malformations, venous and lymphatic anomalies, and limb overgrowth[1,2]. It was first described in 1900 as Naevus vasculosus osteohypertrophicus by two French physicians, Maurice Klippel and Paul Trenaunay. It was previously considered similar to the less common Parkes Weber syndrome, distinguished from KTS by the presence of arteriovenous malformations. [3,4].

KTS hallmark features predominantly affect one lower limb (85%) causing a discrepancy in limb length, although bilateral involvement may occur in a minority of cases (12.5%). However, it is a generalized disorder and can involve multiple body regions including the pelvis, abdomen, head and neck, back, pelvis, and urinary bladder. Capillary malformations are the most common presentation (98%), followed by varicosities or venous malformations (72%), and limb hypertrophy (67%) [5].

Pregnancy poses considerable challenges in patients with KTS, as hormonal and hemodynamic changes exacerbate underlying vascular abnormalities increasing the risk of thromboembolism, hemorrhage, and coagulopathy. The emergence of coagulopathy, in particular, may indicate the development of Kasabach-Merritt syndrome (KMS). [6]

Kasabach-Merritt syndrome (KMS), first described in 1940, is a rare and potentially fatal condition defined by the triad of thrombocytopenia, microangiopathic hemolytic anemia, and consumptive coagulopathy occurring in association with vascular abnormalities [7]. In the setting of KTS, pregnancy and labor may precipitate or exacerbate KMS; the concurrent occurrence of both conditions during the peripartum period constitutes a critical obstetric emergency carrying a reported mortality rate of 30–40% [8].

To date, only three cases of KTS complicated by KMS in pregnancy have been reported in the literature [9–11]. We hereby describe the clinical course and successful multidisciplinary management of a pregnancy in a woman with KTS complicated by fetal growth restriction and peripartum acute KMS. Through this case, we aim to describe the clinical challenges inherent to this rare case, outline effective management strategies, and emphasize the critical

role of early recognition and interdisciplinary collaboration in achieving favorable maternal and neonatal outcomes.

Case Presentation

A 28-year-old multigravida presented at 30+1 weeks' gestation with clinically significant hematuria and was admitted on December 16th to Ain Shams University Hospital, Egypt. Her medical history was notable for Klippel–Trenaunay syndrome (KTS), diagnosed at two years of age. She had an extensive KTS phenotype, with disease involvement encompassing both lower limbs, the back, pelvis, urinary bladder, and breast tissue — a distribution reflecting a multisystem vascular dysplasia. Her surgical history included varicose vein surgery seven years prior and orthopedic fixation with plates and screws approximately 18 months before this admission. Her past obstetric history was significant. Her first pregnancy, seven years prior, ended in neonatal death. Her second pregnancy was complicated by severe preeclampsia, postpartum wound hematoma, and sepsis necessitating ICU admission. She also had a history of one spontaneous miscarriage. In the current pregnancy, she received two doses of antenatal corticosteroids on December 14th and 15th.

On admission, vital signs were stable with a blood pressure 120/80 mmHg, temperature 36.2°C, and heart rate 68 bpm. Physical examination revealed port-wine stains over the back and gluteal region, with discrepancy in lower limb length bilaterally. Blood typing identified B-positive blood group, and viral serology was negative for hepatitis B, hepatitis C, and HIV. Her complete blood count (CBC) demonstrated microcytic hypochromic anemia (hemoglobin 7.0 g/dL, hematocrit 23.0%) and thrombocytopenia (128,000/ μ L). She received six units of packed red blood cells (pRBCs) to correct her declining hemoglobin. Obstetric ultrasonography revealed an estimated fetal weight below the 10th percentile, consistent with fetal growth restriction (FGR). There was an absent end-diastolic flow on umbilical artery Doppler.

Urological consultation was obtained for management of severe hematuria, with initial measures comprising intravenous tranexamic acid and insertion of a three-way indwelling silicone urinary catheter maintained for one week.

On December 18th, pelviabdominal MRI demonstrated a viable single intrauterine pregnancy in cephalic presentation with unremarkable placental location and attachment. Compression of the left ovarian and left common iliac veins by the gravid uterus was identified, with associated venous dilatation and slowing of flow. Extensive dilated, tortuous tubular lesions consistent with varicosities were observed throughout the pelvic, perivesical, perineal, and gluteal regions — more prominent on the left — exhibiting a classic “bag of worms” appearance. Similar lesions were identified in the subcutaneous tissue of both breasts. The urinary bladder contained blood. The liver was mildly enlarged with vascular compression, while the gallbladder, spleen, and kidneys were unremarkable. Minimal bilateral pleural effusion was also noted.

On December 21st, a pelviabdominal ultrasonography, color Doppler of bilateral lower limb veins, and duplex arterial studies were performed. Ultrasonography demonstrated hepatosplenomegaly, soft tissue lesions within the urinary bladder, and bilateral refluxing parametrial venous channels, with a single viable intrauterine fetus confirmed. Lower limb vascular studies revealed a normal venous and arterial system with no evidence of deep vein thrombosis.

Serial CBCs over the four days following transfusion showed slight improvement in hemoglobin, reaching 9.4 g/dL on December 22nd, while platelet count remained persistently low, ranging between 82,000 and 104,000/ μ L. Iron studies revealed elevated unsaturated iron-binding capacity (UIBC) and reduced transferrin saturation. Lactate dehydrogenase (LDH) was elevated at 454 IU/L. Urinalysis on the same day demonstrated hematuria (2+), pyuria, and mild proteinuria (1+); microscopic examination showed >100 red blood cells per high-power field (RBCs/HPF), 40–45 pus cells/HPF, increased epithelial cells (2+), and uric acid crystals (2+). Coagulation studies on December 27th and 28th revealed mildly elevated prothrombin time (PT), while serum creatinine remained low.

Contrast-enhanced pelvic MRI on January 1st demonstrated extensive venous malformations involving the cervix, vagina, anal canal, perineum, and bilateral labia, with additional lesions in the gluteal region, thighs, anterior abdominal wall, and intravesically.

The patient was monitored daily with regular assessment of vital parameters, fetal kick counts, vaginal bleeding, uterine contractions, and signs of membrane rupture. She received one unit of pRBCs on January 2nd and a further unit on January 5th.

At 35 weeks' gestation, a multidisciplinary team comprising obstetrics, hematology, anesthesia, and neonatology convened and recommended delivery by emergency cesarean section in view of deteriorating fetal Doppler parameters and cumulative maternal risks. On January 18th, after preoperative preparation with 2gm ceftriaxone, cesarean delivery was performed under general anesthesia via a vertical midline abdominal incision. Intraoperatively, significant varicosities were encountered over the anterior abdominal wall, lower uterine segment, and the uterovesical interface. A lower uterine segment incision was made for delivery of the fetus and placenta; the uterine wall was subsequently closed in two layers. The placenta was delivered complete, the perineum was intact, and hemostasis was secured. No intraperitoneal drains were placed; a subcutaneous drain was inserted and maintained for seven days. Skin closure was achieved with interrupted mattress sutures. One packed RBCs, one FFP, 12 units pf platelets and 2 gm tranexamic acid were infused intraoperatively to control oozing.

A female neonate weighing 1,900 g was delivered with Apgar scores of 7 and 9 at one and five minutes, respectively. Given the low birth weight, routine vaccination was deferred. The newborn did not require respiratory support.

Postoperatively, the patient was admitted to the ICU following identification of hemoperitoneum. Her pharmacological regimen included low-molecular-weight heparin (LMWH), tranexamic acid, ceftriaxone, pantoprazole, intravenous paracetamol, and a charcoal-based herbal preparation.

On postoperative day one, mild hematuria persisted and laboratory investigations revealed declining hemoglobin and platelet counts alongside markedly elevated LDH, alkaline phosphatase, creatine kinase, and bilirubin. LMWH was withheld, and the patient was commenced on pRBCs and fresh frozen plasma (FFP) in a 1:1 ratio, receiving a total of ten units of each over seven days. Additional laboratory results are presented in ?@tbl-2

On postoperative day two, hemoglobin and platelet count deteriorated rapidly, falling from 9.6 g/dL and 218,000/ μ L to 7.6 g/dL and 104,000/ μ L, respectively, necessitating ongoing blood product replacement.

On postoperative day three, transvaginal ultrasonography (TVUS) revealed marked hemoperitoneum with a localized lower uterine segment hematoma measuring approximately 12 cm in maximum diameter. A decision to continue conservative management was made by the attending consultant and by the head of the unit. The pharmacological regimen was revised to include LMWH, tranexamic acid, dexamethasone, magnesium sulfate, cefepime, omeprazole, clindamycin, and intravenous paracetamol.

On postoperative day four, coagulation studies revealed hypofibrinogenemia (fibrinogen 1.4 g/L) and markedly elevated D-dimer (35.2 μ g/mL FEU), findings consistent with a diagnosis of Kasabach-Merritt syndrome (KMS).

Urinalysis on day six demonstrated persistent hematuria (2+), mild pyuria, and mild proteinuria (1+), with microscopic examination showing >100 RBCs/HPF, 13–15 pus cells/HPF, and increased epithelial cells (1+). By day seven, hemoglobin had stabilized at 10.1 g/dL and remained stable over the subsequent three days, though thrombocytopenia persisted at 64,000/ μ L.

Following clinical stabilization, the patient was discharged on postoperative day ten. At her outpatient review on day nineteen, she was in satisfactory general condition with a soft, lax abdomen; skin sutures were removed and the abdominal wound demonstrated optimal healing.

?@tbl-1 shows the full clinical timeline in chronological order.

Table 1: Patient timeline

Date	Timeline
30 +1 weeks' gestation (Dec 16, 2025)	Admission of a case of KTS, presenting with hematuria; anemia and thrombocytopenia; fetus <10th percentile, received tranexamic acid and 6 units PRBCs.
30+1 to 35 weeks' gestation (Dec 18, 2025 – Jan 17, 2026)	MRI: extensive venous malformations (pelvis, thigh, abdomen). Close maternal/fetal monitoring. Intermittent transfusions slight hemoglobin improvement; fluctuating platelets; prolonged PT.

Date	Timeline
35 weeks (Jan 18, 2026)	Emergency cesarean section for fetal compromise → female neonate (1900 g). Mother admitted to ICU, hemoperitoneum managed conservatively.
Postoperative course	Day 1: Transfusion support (total 10 units PRBC + FFP over 7 days) Day 2: Hb 9.6→7.6 g/dL; platelets $218 \times 10^3 \rightarrow 104 \times 10^3 / \mu\text{L}$ Day 3: Conservative management, LMWH resumed, tranexamic acid initiated Day 4: Hypofibrinogenemia (1.4 g/L), elevated D-dimer (35.2 $\mu\text{g/mL}$) consistent with Kasabach–Merritt syndrome Day 7: Hb stabilized (10.1 g/dL); persistent thrombocytopenia ($64 \times 10^3 / \mu\text{L}$) Day 10: Discharged in stable condition Day 19: Outpatient follow-up: fair general condition; sutures removed

Table 2: Laboratory results throughout the clinical course

Parameters/Timing	December 16 (Day of admission)	De- cem- ber 22	De- cem- ber 28	POD 1	POD 2	POD 4	POD 6	Refer- ence range
Hb (g/dL)	7.0	9.4	10.2	9.6	7.6	8.6	10.0	12.0-15.0
Platelets (per μL)	128,000	97,000	129,000	218,000	104,000	75,000	64,000	150,000–410,000
PT (seconds)	-	-	12.4	40.70	-	-	17.7	11.7–15.3
Total bilirubin (mg/dL)	-	-	-	2.8	1.8	-	1.4	1.8-3.5
ALP (IU/L)	-	-	-	135	115	-	108	34–104
LDH (IU/L)	-	454	-	424	357	-	449	140–271

POD: Post operative day, Hb: Hemoglobin, HTC: Hematocrit, PT: Prothrombin time, ALP: Alkaline phosphatase, LDH : Lactate dehydrogenase.

Review of literature

A systematic search of MEDLINE, Embase, and Scopus identified only three reported cases of KTS complicated by peripartum KMS. KTS was diagnosed prior to pregnancy in two cases [9,10], whereas in the third it had been misdiagnosed as acromegaly and was not recognized until the final trimester [11]. All cases presented with classic KTS features — multiple varicosities, limb hypertrophy, and hemangiomas — with one additionally demonstrating port-wine stains [11]. Despite treatment, postpartum laboratory parameters failed to normalize, indicating refractory coagulopathy consistent with KMS.

All three cases had marked vulvar varicosities necessitating cesarean delivery, with intraoperative or postoperative hemorrhage requiring multiple transfusions; two required massive transfusion protocols [9,11]. Management followed a multidisciplinary approach in all cases, incorporating anticoagulation and blood product support. Two cases received corticosteroids and antifibrinolytics [9,10]. One patient developed life-threatening hemorrhage requiring postpartum hysterectomy, complicated further by DIC, sepsis, peritonitis, and hemoperitoneum necessitating emergency laparotomy [11]. All patients required prolonged postoperative hospitalization. None of the neonates demonstrated clinical or laboratory evidence of KTS.

Summary of the three cases is shown in [Table 3](#)

Table 3: Literature review

Author (Year)	Preg- nancy	Key findings	KMS	Management	Out- come
Neubert (1995) [9]	22 years- old primi- gravida, 37 weeks' gesta- tion	known KTS, hemangiomas, limb hypertrophy, massive vulvovaginal varicosities	Post-op KMS: thrombocytope- nia, decreased fibrinogen, prolonged PT/PTT, increased fibrin split products	CS at 39 wks; hematoma evacuation; ICU; transfusion; heparin + aminocaproic acid	Dis- charged day 20; healthy neonate

Au- thor (Year)	Preg- nancy	Key findings	KMS	Management	Out- come
Kan- mani (2021) [10]	24 years- old, primi- gravida, 29 weeks' gesta- tion	known KTS, bleeding gums, Limb hypertrophy, varicosities (vulvar, lower limb)	Antenatal KMS: severe thrombocytopenia (limited laboratory investigation reported)	Steroids + platelet transfusion; CS at 35 wks; post-op anticoagulation and corticosteroids	Dis- charged day 20; healthy neonate
Zhang (2019) [11]	26 years old. Postpar- tum.	KTS diagnosed in the third trimester of pregnancy. Admitted on day 10 postpartum with severe postpartum hemorrhage, hemodynamic instability, ecchymoses, massive limb hypertrophy, varicosities	Post-op KMS: thrombocytope- nia, ↓ fibrinogen, ↑ PT/APTT, increased fibrin split products, increased D-dimer	CS at 39 wks; hysterectomy; embolization; laparotomy; anticoagulation	Dis- charged day 42; healthy neonate

Discussion

The occurrence of Klippel-Trenaunay syndrome complicated by Kasabach-Merritt syndrome during pregnancy is exceedingly rare, with only three such cases reported in the literature prior to this report [9–11]. The present case thus represents the fourth documented instance of this life-threatening combination, and the first to describe its successful management in the context of concomitant fetal growth restriction and peripartum acute KMS.

Diagnosis and baseline complexity

KTS patients may present with complications such as cellulitis, massive vaginal or visceral bleeding and other vascular complications as severe hematuria and hematochezia (fewer than 5%). Other less common but serious manifestations include deep vein thrombosis (5.8%), pulmonary embolism (2.3%) and severe PPH (11%)[12]. Our patient had clinically significant hematuria which was the sentinel presentation leading to admission, a manifestation attributable to intravesical venous malformations confirmed on both MRI and contrast-enhanced pelvic imaging. Imaging studies are crucial for assessing the extent of vascular malformations

near the spinal cord and pelvic structures, with MRI being the preferred imaging modality for accurately evaluating lesions size, extent, and anatomical involvement which is critical for planning the anesthesia and delivery type. [13,14].

Our patient underwent first-trimester ultrasound and MRI at 30 weeks. Findings collectively reflecting the hemodynamic burden imposed by the gravid state on an already compromised vascular system. This aligns with prior observations that pregnancy-associated hormonal and hemodynamic changes exacerbate the vascular pathology inherent to KTS, amplifying risks of thromboembolism and hemorrhage [15].

Antenatal management

Initial antenatal management was guided by two principal concerns: controlling hematuria and optimizing fetal and maternal status ahead of inevitable preterm delivery. Hematuria was managed conservatively with intravenous tranexamic acid and continuous bladder irrigation via a three-way urinary catheter, achieving partial resolution without surgical intervention. Serial CBCs demonstrated persistent thrombocytopenia throughout the antenatal period despite transfusion support, accompanied by elevated LDH — early biochemical signals of evolving consumptive coagulopathy. These antenatal laboratory aberrations, in retrospect, likely reflected subclinical KMS activity preceding its overt peripartum manifestation.

Fetal complications, including intrauterine growth restriction (IUGR), were reported in a few cases. However, no significant fetal effects were reported in the majority of patients with KTS, indicating that its impact on fetal development is generally limited[16]. In our case, Fetal surveillance revealed IUGR with absent end-diastolic umbilical artery flow, indicating significant uteroplacental insufficiency. The administration of two doses of antenatal corticosteroids at 30 weeks represented an important adjunct in preparing for anticipated preterm delivery. The decision to deliver at 35 weeks was reached through multidisciplinary consensus, balancing the risk of ongoing fetal compromise against the substantial maternal surgical risks posed by the extensive pelvic varicosities. This mirrors the management philosophy reported in all previously published cases, where cesarean delivery was ultimately necessitated by vulval or pelvic varicosities precluding safe vaginal birth [[9]][10][11]

Intraoperative findings and delivery

Cesarean section was performed under general endotracheal anesthesia, with the choice of a vertical midline incision allowing adequate visualization and surgical access in the setting of extensive anterior abdominal wall and lower uterine segment varicosities. Hemostasis was secured without recourse to uterine artery ligation, balloon tamponade, or hysterectomy, distinguishing the intraoperative course of this case favorably from that of a previously published case report [11].

A healthy female neonate was delivered demonstrated no clinical features of KTS, supporting the current understanding that KTS does not follow a straightforward vertical transmission pattern, respectively. Consistent with all three previously reported cases[[9]][10][11].

Peripartum KMS and postoperative course

The postoperative course was complicated by hemoperitoneum and lower uterine segment hematoma. The definitive diagnosis of KMS was established on day four, supported by hypofibrinogenemia and markedly elevated D-dimer along with the severe thrombocytopenia — laboratory findings fulfilling the diagnostic criteria of consumptive coagulopathy in the context of underlying vascular abnormalities [7].

Management of peripartum KMS in this case required careful navigation of the competing hemostatic risks inherent to the syndrome: thrombocytopenia and consumptive coagulopathy predispose to hemorrhage, while the vascular malformations underlying KMS simultaneously confer thrombotic risk. Using anticoagulation-antifibrinolytic combination reflects the dual-pathway pathophysiology of KMS and is consistent with the approach described in two of the three previously reported cases [[9]][10]. Corticosteroid therapy was incorporated given its proposed role in reducing vascular lesion activity and platelet trapping. The patient ultimately received a massive transfusion — alongside infective prophylaxis.

A pattern consistent with the refractory coagulopathy observed in all previously reported cases, where postpartum laboratory parameters failed to normalize despite active treatment [9–11]. The prompt recognition and initiation of therapy are critical to reduce mortality in KMS. The patient eventually made a complete recovery on day ten, representing a favorable outcome relative to the severe postoperative morbidity — including sepsis, peritonitis, DIC, and hysterectomy — reported in the most complicated previously published case[11].

Multidisciplinary care and broader implications

The successful outcome in this case was underpinned by early multidisciplinary involvement across obstetrics, hematology, anesthesia, neonatology, urology, and radiology — a consensus echoed across all published cases and reflective of the inherent complexity of managing concurrent KTS and KMS in pregnancy. Early recognition of KMS, even in its subclinical antenatal form, appears critical: the persistent antenatal thrombocytopenia and elevated LDH in our patient, though not initially diagnostic, warranted heightened vigilance and likely informed the decision to deliver at 35 rather than a later gestational age.

This case further highlights the importance of preconception counseling in women with KTS. The published cross-sectional data [15] documenting significantly elevated risks of DVT, pulmonary embolism, and severe postpartum hemorrhage in KTS patients provides epidemiological context for the individual complexity illustrated here. Our patient’s cumulative obstetric history reflects the substantial reproductive morbidity that KTS may confer across successive pregnancies and underscores the need for individualized, specialist-led preconception assessment in this population.

Conclusion

Pregnancy in women with Klippel–Trenaunay syndrome complicated by Kasabach–Merritt syndrome is extremely rare and poses significant diagnostic and therapeutic challenges. Early

suspicion and diagnosis with intensive antenatal surveillance and multidisciplinary management are crucial to navigate the complex balance between thrombosis and bleeding and to optimize maternal and perinatal outcomes in such high-risk pregnancy and childbirth.

Abbreviation

Klippel-Trenaunay syndrome (KTS)

Kasabach Meritt Syndrome (KMS)

Magnetic resonance imaging (MRI)

Ultrasonography (US)

Transvaginal ultrasound (TVUS)

Intensive care unit (ICU)

Low-molecular-weight heparin (LMWH)

Lactate dehydrogenase (LDH)

Complete blood picture (CBC)

Disseminated Intravascular Coagulation (DIC)

Multidisciplinary team (MDT)

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

A written consent for publication has been obtained from the patient.

Availability of data and materials

The data and materials supporting the conclusions of this article are available as a supplementary materials.

Competing interests

The authors declare that they have no competing interests.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Authors' contributions

SIA, ME, AFN contributed to writing the first draft of the manuscript. All the authors revised the manuscript critically for important intellectual content. All authors approved the final version of the manuscript.

Acknowledgements

We sincerely thank the healthcare providers at Ain Shams University Hospital of Obstetrics and Gynecology for their dedication and commitment to patient care in the operating room. We deeply appreciate their contributions and the valuable role they play in ensuring high-quality surgical care.

References

1. International Society for the Study of Vascular Anomalies. ISSVA Classification for Vascular Anomalies. <https://www.issva.org/UserFiles/file/ISSVA-Classification-2018.pdf>; 2018.
2. Pavone P, Marino L, Cacciaguerra G, Di Nora A, Parano E, Musumeci G, et al. [Klippel-trenaunay syndrome, segmental/focal overgrowth malformations: A review](#). *Children (Basel)*. 2023;10:1421.
3. McKusick VA. Frederick parkes weber—1863-1962. *JAMA* [Internet]. 1963;183. Available from: <http://dx.doi.org/10.1001/jama.1963.63700010018015>
4. Sikova DVD, Pavlova LjT, Laskoska MTV, Nikolovska ST, Biljanovska NC. Naevus varicosus osteohypertrophicus. *Springer US*; 1999. p. 535–40. Available from: http://dx.doi.org/10.1007/978-1-4615-4857-7_80

5. Jacob AG, Driscoll DJ, Shaughnessy WJ, Stanson AW, Clay RP, Gloviczki P. Klippel-trénaunay syndrome: Spectrum and management. Mayo Clinic Proceedings [Internet]. 1998;73:28–36. Available from: [http://dx.doi.org/10.1016/S0025-6196\(11\)63615-X](http://dx.doi.org/10.1016/S0025-6196(11)63615-X)
6. Blei F. Klippel-Trenaunay Syndrome: Clinical Manifestations, Diagnosis, and Management. UpToDate. <https://www.uptodate.com/contents/klippel-trenaunay-syndrome-clinical-manifestations-diagnosis-and-management>; 2024.
7. Mahajan P, Margolin J, Iacobas I. Kasabach-merritt phenomenon: Classic presentation and management options. Clinical Medicine Insights: Blood Disorders [Internet]. 2017;10:1179545X1769984. Available from: <http://dx.doi.org/10.1177/1179545X17699849>
8. Tian Y, Yuan Y, Wei L, Xu Z, Li L, Li L. [Kaposiform hemangioendothelioma: Diagnosis and treatment](#). Pediatric Investigation. 2026;
9. Neubert GA, Golden MA, Rose NC. [Kasabach-merritt coagulopathy complicating klippel-trenaunay-weber syndrome in pregnancy](#). Obstetrics & Gynecology. 1995;85:831–3.
10. Kanmani K, Meena M, Narmadha D, A. GV. [Klippel trenaunay syndrome—an obstetric challenge](#). International Journal of Reproduction, Contraception, Obstetrics and Gynecology. 2021;10:2081–4.
11. Zhang J, Wang K, Mei J. Late puerperal hemorrhage of a patient with klippel–trenaunay syndrome. Medicine [Internet]. 2019;98:e18378. Available from: <http://dx.doi.org/10.1097/MD.00000000000018378>
12. Chang R, Zhang J, Zhang F, Wen L, Jiang Y. Pregnancy in women with klippel–trenaunay syndrome complicated by uterine vascular malformation: A case report and literature review. Cureus [Internet]. 2025; Available from: <http://dx.doi.org/10.7759/cureus.90065>
13. Howlett DC, Roebuck DJ, Frazer CK, Ayers B. The use of ultrasound in the venous assessment of lower limb klippel-trenaunay syndrome. European Journal of Radiology [Internet]. 1994;18:224–6. Available from: [http://dx.doi.org/10.1016/0720-048x\(94\)90340-9](http://dx.doi.org/10.1016/0720-048x(94)90340-9)
14. Asghar F, Aqeel R, Farooque U, Haq A, Taimur M. Presentation and management of klippel-trenaunay syndrome: A review of available data. Cureus [Internet]. 2020; Available from: <http://dx.doi.org/10.7759/cureus.8023>
15. Silva Correia IF, Hussain M, Johnson J-A. Obstetric management for pregnant women with klippel–trenaunay syndrome: A UK case report and review of the literature. International Journal of Gynecology & Obstetrics [Internet]. 2024;168:484–6. Available from: <http://dx.doi.org/10.1002/ijgo.15889>

16. Akay E, Uzun Gül AD, Türkoğlu A. [Recurrent preeclampsia in a pregnant woman with klippel-trenaunay syndrome: Two cesarean deliveries and multiple extremity involvement – case report and literature review](#). BMC Pregnancy and Childbirth. 2025;25:407.