

Klippel-Trenaunay Syndrome Complicated by Kasabach-Merritt Syndrome During Pregnancy

A case report and review of the literature

Authors

Mourad Elfaham¹, Aya Attia¹, Mohamed Waheed¹, Mohamed Essam¹, Salma Saad¹, Mariam Sarhan¹, Marwa Mourad¹, Mohamed Hegazy¹, Farah Said¹, Sarah Tarek¹, Amgad Gamal¹, Sara Ibrahim Abdelkader^{2*}, Ahmed Azab², Mohammed Ghanem², Rahma Alaa Abdelhafez², Nour Atif², Maha Moemen², Belgin Ahmed Nagah², Ashraf Nabhan^{1,3}

Affiliation

¹ Faculty of Medicine, Ain Shams University, Cairo, Egypt.

² Faculty of Medicine, Galala University, Attaka, Suez, Egypt.

³ Faculty of Medicine, MTI University, Cairo, Egypt. 01069663999

* Corresponding author: Sara Ibrahim Abdelkader. Faculty of Medicine, Galala University, Attaka, Suez, Egypt. Email: sara.elzeftawy@gu.edu.eg, ORCID: [0009-0002-5877-4722](https://orcid.org/0009-0002-5877-4722)

Word count: 2500

Abstract

The co-occurrence of Klippel-Trenaunay syndrome and Kasabach-Merritt syndrome in pregnancy is extremely rare and represents a critical obstetric emergency with significant maternal and fetal risks, demanding early recognition and immediate multidisciplinary intervention.

We report the case of a 28-year-old pregnant woman with known Klippel-Trenaunay syndrome who developed Kasabach-Merritt phenomenon during pregnancy. The fetus suffered severe

growth restriction and absent end-diastolic flow on umbilical artery Doppler. A cesarean section was performed at 35 weeks of gestation.

This case highlights the complexity of managing pregnancy in women with Klippel–Trenaunay syndrome complicated by Kasabach Merritt syndrome and emphasizes the importance of multidisciplinary surveillance and timely delivery.

Keywords: Klippel–Trenaunay syndrome; angioosteohypertrophy syndrome; Kasabach Merritt syndrome; fetal growth restriction; high-risk pregnancy

Introduction

Klippel-Trenaunay syndrome (KTS) is a rare congenital disorder affecting an estimated 2–5 per 100,000 individuals. Its features include a triad of capillary malformations, venous and lymphatic anomalies, and limb overgrowth. [1,2]

KTS was first described in 1900 as *Naevus vasculosus osteohypertrophicus* by two French physicians, Maurice Klippel and Paul Trenaunay. It was previously considered similar to the less common Parkes Weber syndrome, distinguished from KTS by the presence of arteriovenous malformations. [3,4]. KTS is currently classified within the PIK3CA-Related Overgrowth Spectrum (PROS), a group of overgrowth syndromes with overlapping phenotypic features that share mutations in the PIK3CA gene. KTS, also referred to as capillary-lymphatic-venous malformation, is characterized by capillary malformations, soft tissue or bony hypertrophy, and varicose veins or venous malformations. The presence of at least two of these features suggests the diagnosis, which is supported by imaging. Magnetic resonance imaging, Color Doppler studies, and pelviabdominal ultrasounds may all be used to visualize vascular malformations associated with KTS.[5] KTS hallmark features predominantly affect one lower limb (85%) causing a discrepancy in limb length, although bilateral involvement may occur in a minority of cases (12.5%). However, it is a generalized disorder and can involve multiple body regions including the pelvis, abdomen, head and neck, back, pelvis, and urinary bladder. Capillary malformations are the most common presentation (98%), followed by varicosities or venous malformations (72%), and limb hypertrophy (67%) [6].

Pregnancy poses considerable challenges in patients with KTS, as hormonal and hemodynamic changes exacerbate underlying vascular abnormalities increasing the risk of thromboembolism, hemorrhage, and coagulopathy. The emergence of coagulopathy, in particular, may indicate the development of Kasabach-Merritt syndrome (KMS). [7]

Kasabach-Merritt syndrome (KMS), first described in 1940, is a rare and potentially fatal condition defined by the triad of thrombocytopenia, microangiopathic hemolytic anemia, and consumptive coagulopathy occurring in association with vascular abnormalities [8]. In the setting of KTS, pregnancy and labor may precipitate or exacerbate KMS; the concurrent occurrence of both conditions during the peripartum period constitutes a critical obstetric emergency carrying a reported mortality rate of 30–40% [9].

To date, only three cases of KTS complicated by KMS in pregnancy have been documented in the literature [10–12]. We report a fourth case, describing the clinical course and successful multidisciplinary management of a pregnant woman with KTS complicated by fetal growth restriction and peripartum acute KMS. Through this case, we aim to illuminate the clinical challenges inherent to this rare complication, outline effective management strategies, and emphasize the critical role of early recognition and interdisciplinary collaboration in achieving favorable maternal and neonatal outcomes.

Case Presentation

A 28-year-old multigravida (P2+1, both prior deliveries by cesarean section, one living child) presented at 30+1 weeks' gestation with clinically significant hematuria and was admitted on December 16th to Ain Shams University Hospital, Egypt. The clinical timeline is detailed in Table 1.

Her background was notable for Klippel-Trenaunay syndrome (KTS), diagnosed at age two, with involvement of both lower limbs, the back, pelvis, and urinary bladder. Her surgical history included varicose vein surgery seven years prior and orthopedic fixation with plates and screws approximately 18 months before this admission.

Her obstetric history was significant. Her first pregnancy, seven years prior, ended in neonatal death. Her second pregnancy was complicated by severe preeclampsia, postpartum wound hematoma, and sepsis necessitating ICU admission. She also had a history of spontaneous miscarriage not managed surgically. In the current pregnancy, she received two doses of antenatal corticosteroids on December 14th and 15th.

On admission, vital signs were stable: blood pressure 120/80 mmHg, temperature 36.2°C, and heart rate 68 bpm. Physical examination revealed port-wine stains over the back and gluteal region, with discrepancy in lower limb length bilaterally.

A complete blood count (CBC) demonstrated microcytic hypochromic anemia (hemoglobin 7.0 g/dL, hematocrit 23.0%) and thrombocytopenia (128,000/ μ L). She received six units of packed red blood cells (pRBCs) to correct her declining hemoglobin. Blood typing identified B-positive blood group, and viral serology was negative for hepatitis B, hepatitis C, and HIV.

Obstetric ultrasonography revealed an estimated fetal weight below the 10th percentile, consistent with intrauterine growth restriction (IUGR), and absent end-diastolic flow on umbilical artery Doppler. Urological consultation was obtained for management of severe hematuria, with initial measures comprising intravenous tranexamic acid and insertion of a three-way indwelling silicone urinary catheter maintained for one week.

On December 18th, pelviabdominal MRI demonstrated a viable single intrauterine pregnancy in cephalic presentation with unremarkable placental location and attachment. Compression

of the left ovarian and left common iliac veins by the gravid uterus was identified, with associated venous dilatation and slowing of flow. Extensive dilated, tortuous tubular lesions consistent with varicosities were observed throughout the pelvic, perivesical, perineal, and gluteal regions — more prominent on the left — exhibiting a classic “bag of worms” appearance. Similar lesions were identified in the bilateral breast subcutaneous tissue. The urinary bladder contained blood products. The liver was mildly enlarged with vascular compression, while the gallbladder, spleen, and kidneys were unremarkable. Minimal bilateral pleural effusion was also noted.

On December 21st, pelviabdominal ultrasonography, color Doppler of bilateral lower limb veins, and duplex arterial studies were performed. Ultrasonography demonstrated hepatosplenomegaly, soft tissue lesions within the urinary bladder, and bilateral refluxing parametrial venous channels, with a single viable intrauterine fetus confirmed. Lower limb vascular studies revealed a normal venous and arterial system with no evidence of deep vein thrombosis.

Serial CBCs over the four days following transfusion showed progressive improvement in hemoglobin, reaching 9.4 g/dL on December 22nd, while platelet count remained persistently low, ranging between 82,000 and 104,000/ μ L. Iron studies revealed elevated unsaturated iron-binding capacity (UIBC) and reduced transferrin saturation. Lactate dehydrogenase (LDH) was elevated at 454 IU/L. Urinalysis on the same day demonstrated hematuria (2+), pyuria, and mild proteinuria (1+); microscopic examination showed >100 red blood cells per high-power field (RBCs/HPF), 40–45 pus cells/HPF, increased epithelial cells (2+), and uric acid crystals (2+). Coagulation studies on December 27th and 28th revealed mildly elevated prothrombin time (PT), while serum creatinine remained low.

Contrast-enhanced pelvic MRI on January 1st demonstrated extensive venous malformations involving the cervix, vagina, anal canal, perineum, and bilateral labia, with additional lesions in the gluteal region, thighs, anterior abdominal wall, and intravesically.

The patient was monitored daily with regular assessment of vital parameters, fetal kick counts, vaginal bleeding, uterine contractions, and signs of membrane rupture. She received one unit of pRBCs on January 2nd and a further unit on January 5th.

At 35 weeks' gestation, a multidisciplinary team comprising obstetrics, hematology, anesthesia, and neonatology convened and recommended delivery by emergency cesarean section in view of deteriorating fetal Doppler parameters and cumulative maternal risks. On January 18th, cesarean delivery was performed under general endotracheal anesthesia via a vertical midline abdominal incision. Intraoperatively, significant varicosities were encountered over the anterior abdominal wall, lower uterine segment, and the uterovesical interface. A lower uterine segment incision was made for delivery of the fetus and placenta; the uterine wall was subsequently closed in two layers. The placenta was delivered complete, the perineum was intact, and hemostasis was secured. No intraperitoneal drains were placed; a subcutaneous drain was inserted and maintained for seven days. Skin closure was achieved with interrupted mattress sutures.

A female neonate weighing 1,900 g was delivered with Apgar scores of 7 and 9 at one and five minutes, respectively. Given the low birth weight, routine vaccination was deferred, and neither observation nor respiratory support was required.

Postoperatively, the patient was admitted to the ICU following identification of hemoperitoneum. Her pharmacological regimen included low-molecular-weight heparin (LMWH), tranexamic acid, ceftriaxone, pantoprazole, intravenous paracetamol, and a charcoal-based herbal preparation.

On postoperative day one, mild hematuria persisted and laboratory investigations revealed declining hemoglobin and platelet counts alongside markedly elevated LDH, alkaline phosphatase, creatine kinase, and bilirubin. LMWH was withheld, and the patient was commenced on pRBCs and fresh frozen plasma (FFP) in a 1:1 ratio, receiving a total of ten units of each over seven days. Additional laboratory results are presented in @tbl-2.

On postoperative day two, hemoglobin and platelet count deteriorated rapidly, falling from 9.6 g/dL and 218,000/ μ L to 7.6 g/dL and 104,000/ μ L, respectively, necessitating ongoing blood product replacement.

On postoperative day three, transvaginal ultrasonography (TVUS) revealed marked hemoperitoneum with a localized lower uterine segment hematoma measuring approximately 12 cm in maximum diameter. Conservative management was elected by the attending consultant and departmental head. The pharmacological regimen was revised to include resumed LMWH, tranexamic acid, calcium, dexamethasone, magnesium sulfate, cefepime, omeprazole, clindamycin, and intravenous paracetamol.

On postoperative day four, coagulation studies revealed hypofibrinogenemia (fibrinogen 1.4 g/L) and markedly elevated D-dimer (35.2 μ g/mL FEU), findings consistent with a diagnosis of Kasabach-Merritt syndrome (KMS).

Urinalysis on day six demonstrated persistent hematuria (2+), mild pyuria, and mild proteinuria (1+), with microscopic examination showing >100 RBCs/HPF, 13–15 pus cells/HPF, and increased epithelial cells (1+). By day seven, hemoglobin had stabilized at 10.1 g/dL and remained stable over the subsequent three days, though thrombocytopenia persisted at 64,000/ μ L.

Following clinical stabilization, the patient was discharged on postoperative day ten. At her outpatient review on day nineteen, she was in satisfactory general condition with a soft, lax abdomen; skin sutures were removed and the abdominal wound demonstrated optimal healing.

Table 1: Patient timeline

Date	Timeline
30 +1 weeks' gestation (Dec 16, 2025)	Admission of a case of KTS, presenting with hematuria; anemia and thrombocytopenia; fetus <10th percentile, received tranexamic acid and 6 units PRBCs.
30+1 to 35 weeks' gestation (Dec 18, 2025 – Jan 17, 2026)	MRI: extensive venous malformations (pelvis, thigh, abdomen). Close maternal/fetal monitoring. Intermittent transfusions slight hemoglobin improvement; fluctuating platelets; prolonged PT.
35 weeks (Jan 18, 2026)	Emergency cesarean section for fetal compromise → female neonate (1900 g). Mother admitted to ICU, hemoperitoneum managed conservatively.
Postoperative course	Day 1: Transfusion support (total 10 units PRBC + FFP over 7 days) Day 2: Hb 9.6→7.6 g/dL; platelets $218 \times 10^3 \rightarrow 104 \times 10^3 / \mu\text{L}$ Day 3: Conservative management → LMWH resumed + tranexamic acid Day 4: Hypofibrinogenemia (1.4 g/L), elevated D-dimer (35.2 $\mu\text{g/mL}$) → consistent with Kasabach–Merritt syndrome Day 7: Hb stabilized (10.1 g/dL); persistent thrombocytopenia ($64 \times 10^3 / \mu\text{L}$) Day 10: Discharged in stable condition Day 19: Outpatient follow-up: fair general condition; sutures removed

Table 2: Relevant abnormal laboratory results throughout the clinical course.

Pa- rame- ters/ Tim- ing	De- cem- ber 16 (Day of admis- sion)	De- cem- ber 22	De- cem- ber 28	POD 1	POD 2	POD 4	POD 6	POD 7	Refer- ence range
Hb (g/dL)	7.0	7.6	10.2	9.6	7.6	8.6	10.0	10.0	12.0- 15.0
HCT (%)	23.0	23.5	33.1	20.0	21.8	26.8	30.8	-	36-46
Platelets (/ μL)	128,000	97,000	129,000	218,000	104,000	75,000	64,000	64,000	150,000–410,000

Pa- rame- ters/Tim- ing	De- cem- ber 16 (Day of admis- sion)	De- cem- ber 22	De- cem- ber 28	POD 1	POD 2	POD 4	POD 6	POD 7	Refer- ence range
PT (sec- onds)	-	-	12.4	40.70	-	-	17.7	-	11.7–15.3
aPTT (sec- onds)	-	-	-	-	-	-	48.90	-	26-40
Fib- rino- gen (g/L)	-	-	-	-	-	1.4	-	-	2.00- 4.00
D- dimer (µg/mL FEU)	-	-	-	-	-	35.2	-	-	0-0.55
Total biliru- bin (mg/dL)	-	-	-	2.8	1.8	-	1.4	-	1.8-3.5
ALP (IU/L)	-	-	-	135	115	-	108	-	34–104
LDH (IU/L)	-	454	-	424	357	-	449	-	140–271
BUN (mg/dL)	-	-	-	-	-	-	24	-	5–23
Serum creati- nine (mg/dL)	-	-	0.35	-	-	0.30	-	-	0.6–1.2

POD: Post operative day, Hb: Hemoglobin, HTC: Hematocrit, PT: Prothrombin time, aPTT: Activated partial thromboplastin time, ALP: Alkaline phosphatase, LDH : Lactate dehydrogenase, BUN: Blood Urea Nitrogen.

Review of literature

A systematic search of MEDLINE, Embase, and Scopus identified only three reported cases of KTS complicated by peripartum KMS. KTS was diagnosed prior to pregnancy in two cases [10,11], whereas in the third it had been misdiagnosed as acromegaly and was not recognized until the final trimester [12]. All cases presented with classic KTS features — multiple varicosities, limb hypertrophy, and hemangiomas — with one additionally demonstrating port-wine stains [12]. Despite treatment, postpartum laboratory parameters failed to normalize, indicating refractory coagulopathy consistent with KMS.

All three patients had marked vulvar varicosities necessitating cesarean delivery, with intra-operative or postoperative hemorrhage requiring multiple transfusions; two required massive transfusion protocols [10,12]. Management followed a multidisciplinary approach in all cases, incorporating anticoagulation and blood product support. Two cases received corticosteroids and antifibrinolytics [10,11]. One patient developed life-threatening hemorrhage requiring postpartum hysterectomy, complicated further by DIC, sepsis, peritonitis, and hemoperitoneum necessitating emergency laparotomy [12]. All patients required prolonged postoperative hospitalization. None of the neonates demonstrated clinical or laboratory evidence of KTS.

Detailed summary of the three cases is shown in Table 3

Table 3: Literature review

Au- thor (Year)	Preg- nancy	Key findings	KMS	Management	Out- come
Neu- bert (1995) [10]	22 years- old primi- gravida, 37 weeks’ gesta- tion	known KTS, hemangiomas, limb hypertrophy, massive vulvovaginal varicosities	Post-op KMS: thrombocytope- nia, decreased fibrinogen, prolonged PT/PTT, increased fibrin split products	CS at 39 wks; hematoma evacuation; ICU; transfusion; heparin + aminocaproic acid	Dis- charged day 20; healthy neonate

Au- thor (Year)	Preg- nancy	Key findings	KMS	Management	Out- come
Kan- mani (2021) [11]	24 years- old, primi- gravida, 29 weeks' gesta- tion	known KTS, bleeding gums, Limb hypertrophy, varicosities (vulvar, lower limb)	Antenatal KMS: severe thrombocytopenia (limited laboratory investigation reported)	Steroids + platelet transfusion; CS at 35 wks; post-op anticoagulation and corticosteroids	Dis- charged day 20; healthy neonate
Zhang (2019) [12]	26 years old. Postpar- tum.	KTS diagnosed in the third trimester of pregnancy. Admitted on day 10 postpartum with severe postpartum hemorrhage, hemodynamic instability, ecchymoses, massive limb hypertrophy, varicosities	Post-op KMS: thrombocytope- nia, ↓ fibrinogen, ↑ PT/APTT, increased fibrin split products, increased D-dimer	CS at 39 wks; hysterectomy; embolization; laparotomy; anticoagulation	Dis- charged day 42; healthy neonate

Discussion

The occurrence of KTS complicated by KMS during pregnancy is exceedingly rare, and, to our knowledge, only three such cases have been reported in the literature. [10–12]

Morbidity in KTS is driven predominantly by thromboembolism and hemorrhage, and pregnancy may substantially increase these risks. Nevertheless, robust evidence guiding pregnancy management in this population remains scarce. A cross-sectional study enrolling sixty adult women with KTS demonstrated significantly increased rates of pregnancy-associated deep vein thrombosis, pulmonary embolism, and severe postpartum hemorrhage relative to the general population, with nearly half of participants reporting concurrent exacerbation of KTS-related symptoms [13]. These findings contrast with those of a comparative study in which pregnant women with KTS showed no significant difference in thromboembolic or bleeding risk compared with their nulligravid counterparts [14]; however, the authors acknowledged that women with more severe KTS may have been selectively discouraged from pregnancy, introducing potential selection bias.

Fetal complications, including intrauterine growth restriction (IUGR), were reported in a few cases, including our case. However, no significant fetal effects were reported in the majority of patients with KTS, indicating that its impact on fetal development is generally limited.[15]

Patients may present with a wide variety of manifestations either primary or due to complications such as pain, bleeding, cellulitis, massive vaginal bleeding, visceral bleeding, and bleeding gums, and other vascular complications such as severe hematuria and hematochezia (fewer than 5%). Our case presented with clinically significant hematuria. Other less common but serious manifestations include deep vein thrombosis (5.8%), pulmonary embolism (2.3%), severe PPH (11%), and potential preeclampsia. [15,16]

Patients commonly present with persistent bleeding and thrombocytopenia refractory to blood products. Our patient developed post-operative hematoma and hemoperitoneum which were managed conservatively with success. Specific therapy was initiated once Kasabach-Merritt syndrome was suspected and the patient eventually made a complete recovery. The prompt recognition and initiation of therapy are critical to reduce mortality in KMS. Delayed recognition was associated with more complicated clinical courses, as in the KTS patient at 39 weeks who was not previously diagnosed with KTS, and did not receive LMWH, losing 1500 mL of blood postoperatively and 4500 mL of blood through vaginal bleeding on day 10 postpartum, necessitating a hysterectomy and abdominal gauze packing.[12]

Diagnosis of KMS is confirmed by lab findings of thrombocytopenia, hypofibrinogenemia, elevated D-dimer, and prolonged coagulation times. The absence of thrombocytopenia may indicate refractory coagulopathy rather than KMS, although atypical presentations should be suspected in presence of hemangiomas.[16]

The management of KTS becomes more complex when complicated by Kasabach Merritt syndrome (KMS). Imaging studies are crucial for assessing the extent of vascular malformations near the spinal cord and pelvic structures, with MRI being the preferred imaging modality for accurately evaluating lesion size, extent, and anatomical involvement which is critical for planning the anesthesia and delivery type. Doppler ultrasound is effective for diagnosing venous malformations. Regular maternal and fetal assessment is essential due to risk of placental involvement [17,18]. Our patient underwent first-trimester ultrasound and MRI at 30 weeks. MRI demonstrated extensive venous malformations involving the pelvic, perivesical, perineal, gluteal, thigh, and anterior abdominal regions. Ultrasound revealed hepatosplenomegaly, soft tissue lesions in the urinary bladder, bilateral refluxing parametrial venous channels, and no evidence of deep vein thrombosis.

Deciding the mode of delivery in a pregnant woman with KTS requires a multidisciplinary team (MDT). In most reported cases, cesarean section was opted for due to the presence of vulval varicosities.[19] In our case, emergency cesarean section under general anesthesia was recommended by an MDT involving obstetrics, hematology, anesthesia, and neonatology due to both fetal and maternal risks. Vertical midline abdominal incision revealed significant varicosities in the anterior abdominal wall, over the lower uterine segment, and in the interface

between the uterus and the bladder. Comprehensive preparations were made before and after delivery, including the administration of tranexamic acid and LMWH.

Prophylactic anticoagulation with LMWH should be considered throughout pregnancy given the elevated risk of venous thromboembolism (VTE) [20], with regular coagulopathy monitoring and serial imaging. Postpartum management may involve anticoagulation, platelet transfusion, cryoprecipitate, fresh frozen plasma and antifibrinolytic agents such as TXA. Postpartum follow up is essential as KMS can develop or worsen after delivery and patients may require large volumes of blood products to manage the ongoing coagulopathy, but the condition is often refractory, showing minimal to no improvement despite transfusion support. [13]

Conclusion

Pregnancy in women with Klippel–Trenaunay syndrome complicated by Kasabach–Merritt syndrome is extremely rare and poses significant diagnostic and therapeutic challenges. Early suspicion and diagnosis with intensive antenatal surveillance and multidisciplinary management are crucial to navigate the complex balance between thrombosis and bleeding and to optimize maternal and perinatal outcomes in such high-risk pregnancy and childbirth.

Abbreviation

Klippel-Trenaunay syndrome (KTS)

Kasabach Meritt Syndrome (KMS)

Magnetic resonance imaging (MRI)

Ultrasonography (US)

Transvaginal ultrasound (TVUS)

Intensive care unit (ICU)

Low-molecular-weight heparin (LMWH)

Lactate dehydrogenase (LDH)

Blood Urea Nitrogen (BUN)

Prothrombin time (PT)

Partial Thromboplastin Time (PTT)

Complete blood picture (CBC)

Disseminated Intravascular Coagulation (DIC)

Multidisciplinary team (MDT)

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

A written consent for publication has been obtained from the patient.

Availability of data and materials

The data and materials supporting the conclusions of this article are available as a supplementary materials.

Competing interests

The authors declare that they have no competing interests.

Funding

This research received no specific grant from any funding agency in the public, commercial or not-for-profit sectors.

Authors' contributions

SIA, ME, AFN contributed to writing the first draft of the manuscript. All the authors revised the manuscript critically for important intellectual content. All authors approved the final version of the manuscript.

Acknowledgements

We sincerely thank the healthcare providers at Ain Shams University Hospital of Obstetrics and Gynecology for their dedication and commitment to patient care in the operating room. We deeply appreciate their contributions and the valuable role they play in ensuring high-quality surgical care.

References

1. International Society for the Study of Vascular Anomalies. ISSVA Classification for Vascular Anomalies. <https://www.issva.org/UserFiles/file/ISSVA-Classification-2018.pdf>; 2018.
2. Pavone P, Marino L, Cacciaguerra G, Di Nora A, Parano E, Musumeci G, et al. Klippel-trenaunay syndrome, segmental/focal overgrowth malformations: A review. *Children (Basel)*. 2023;10:1421.
3. McKusick VA. Frederick parkes weber—1863-1962. *JAMA* [Internet]. 1963;183. Available from: <http://dx.doi.org/10.1001/jama.1963.63700010018015>
4. Sikova DVD, Pavlova LjT, Laskoska MTV, Nikolovska ST, Biljanovska NC. Naevus varicosus osteohypertrophicus. Springer US; 1999. p. 535–40. Available from: http://dx.doi.org/10.1007/978-1-4615-4857-7_80
5. Horbach S, Lokhorst M, Oduber C, Middeldorp S, Post J van der, Horst C van der. Complications of pregnancy and labour in women with klippel-trénaunay syndrome: A nationwide cross-sectional study. *BJOG: An International Journal of Obstetrics & Gynaecology* [Internet]. 2017;124:1780–8. Available from: <http://dx.doi.org/10.1111/1471-0528.14698>
6. Jacob AG, Driscoll DJ, Shaughnessy WJ, Stanson AW, Clay RP, Gloviczki P. Klippel-trénaunay syndrome: Spectrum and management. *Mayo Clinic Proceedings* [Internet]. 1998;73:28–36. Available from: [http://dx.doi.org/10.1016/S0025-6196\(11\)63615-X](http://dx.doi.org/10.1016/S0025-6196(11)63615-X)
7. Blei F. Klippel-Trenaunay Syndrome: Clinical Manifestations, Diagnosis, and Management. UpToDate. <https://www.uptodate.com/contents/klippel-trenaunay-syndrome-clinical-manifestations-diagnosis-and-management>; 2024.
8. Mahajan P, Margolin J, Iacobas I. Kasabach-merritt phenomenon: Classic presentation and management options. *Clinical Medicine Insights: Blood Disorders* [Internet]. 2017;10:1179545X1769984. Available from: <http://dx.doi.org/10.1177/1179545X17699849>

9. Tian Y, Yuan Y, Wei L, Xu Z, Li L, Li L. [Kaposiform hemangioendothelioma: Diagnosis and treatment](#). *Pediatric Investigation*. 2026;
10. Neubert GA, Golden MA, Rose NC. [Kasabach-merritt coagulopathy complicating klippel-trenaunay-weber syndrome in pregnancy](#). *Obstetrics & Gynecology*. 1995;85:831–3.
11. Kanmani K, Meena M, Narmadha D, A. GV. [Klippel trenaunay syndrome—an obstetric challenge](#). *International Journal of Reproduction, Contraception, Obstetrics and Gynecology*. 2021;10:2081–4.
12. Zhang J, Wang K, Mei J. Late puerperal hemorrhage of a patient with klippel–trenaunay syndrome. *Medicine [Internet]*. 2019;98:e18378. Available from: <http://dx.doi.org/10.1097/MD.00000000000018378>
13. Silva Correia IF, Hussain M, Johnson J-A. Obstetric management for pregnant women with klippel–trenaunay syndrome: A UK case report and review of the literature. *International Journal of Gynecology & Obstetrics [Internet]*. 2024;168:484–6. Available from: <http://dx.doi.org/10.1002/ijgo.15889>
14. Marvin EK, Schoch JJ, Nguyen H, Anderson KR, Driscoll DJ, Rose CH, et al. Venous thromboembolic and bleeding complications among pregnant women with klippel-trenaunay syndrome. *Journal of the American Academy of Dermatology [Internet]*. 2019;81:1277–82. Available from: <http://dx.doi.org/10.1016/j.jaad.2019.04.018>
15. Akay E, Uzun Gül AD, Türkoğlu A. [Recurrent preeclampsia in a pregnant woman with klippel-trenaunay syndrome: Two cesarean deliveries and multiple extremity involvement – case report and literature review](#). *BMC Pregnancy and Childbirth*. 2025;25:407.
16. Chang R, Zhang J, Zhang F, Wen L, Jiang Y. Pregnancy in women with klippel–trenaunay syndrome complicated by uterine vascular malformation: A case report and literature review. *Cureus [Internet]*. 2025; Available from: <http://dx.doi.org/10.7759/cureus.90065>
17. Howlett DC, Roebuck DJ, Frazer CK, Ayers B. The use of ultrasound in the venous assessment of lower limb klippel-trenaunay syndrome. *European Journal of Radiology [Internet]*. 1994;18:224–6. Available from: [http://dx.doi.org/10.1016/0720-048x\(94\)90340-9](http://dx.doi.org/10.1016/0720-048x(94)90340-9)
18. Asghar F, Aqeel R, Farooque U, Haq A, Taimur M. Presentation and management of klippel-trenaunay syndrome: A review of available data. *Cureus [Internet]*. 2020; Available from: <http://dx.doi.org/10.7759/cureus.8023>
19. Keepanasseril A, Keerthana K, Keepanasseril A, Maurya DK, Kadambari D, Sistla S. Pregnancy in women with klippel-trenaunay syndrome: Report of three pregnancies in a single patient and review of literature. *Obstetric Medicine [Internet]*. 2017;10:177–82. Available

from: <http://dx.doi.org/10.1177/1753495X17719181>

20. Marvin EK, Schoch JJ, Nguyen H, Anderson KR, Driscoll DJ, Rose CH, et al. Venous thromboembolic and bleeding complications among pregnant women with klippel-trenaunay syndrome. *Journal of the American Academy of Dermatology* [Internet]. 2019;81:1277–82. Available from: <http://dx.doi.org/10.1016/j.jaad.2019.04.018>