
LC Classical Mechanics and Relativity 2 Lecture Notes

Year 1 Semester 2

Ash Stewart

MSci Physics with Particle Physics and Cosmology

School of Physics and Astronomy
University of Birmingham

Table of Lectures

Lecture 8: Special Relativity I: Foundations and Time Dilation	2
Lecture 9: Special Relativity II: Length Contraction	6
Lecture 10: Special Relativity III: Lorentz Transformations	9
Lecture 12: Special Relativity IV: More Space-Time	12
Lecture 13: Special Relativity V:	15
Lecture IDK: Special Relativity VII: Kinematics Examples, Doppler and Natural Units	16

Tue 17 Feb 2026 11:00

Lecture 8 - Special Relativity I: Foundations and Time Dilation

1 Introduction

In this lecture:

- Recap from CMR1
- Einstein's two postulates
- Time dilation
- Galilean Transformations

Crucially, there are no special or absolute frames of reference - laws of motion are only sensible when we consider a frame of reference. The only slightly special frame is the one in which we are currently stationary.

Special relativity provides a theory of relative motion between inertial frames of reference. An inertial frame is one which is not accelerating relative to the other frames being considered.

2 Frames of Reference

We will generally consider two related frames of reference, denoted Σ and Σ' . Σ is our “stationary” frame, i.e. the frame taken by an observer sat on earth. Σ' is our “moving frame”, relative to the stationary observer and is moving with constant speed v .

Coordinates in the stationary frame are (x, y, z, t) , while in the moving frame they add a prime, so are given by: (x', y', z', t') .

We want to compare observations of position/experience of time/velocity in the moving frame with observations of the same categories in the stationary frame.

The frame in which an object is stationary is denoted its rest frame Σ_0 .

3 Galilean Transformations

“Galilean Invariance”: The laws of physics are invariant in all inertial (non-accelerating) frames.

“Galilean Transformation”: Time is invariant and universal in all frames.

In Classical Mechanics, we treat time as invariant, but this stops being true in special relativity. We can transform between frames in such a manner that time is kept constant, this is a Galilean Transformation, but this is not true by default.

Consider our two reference frames Σ and Σ' , where the latter moves with speed v relative to the former. Our coordinates in (x, y) become coordinates in (x', y') .

An object moves at speed u' in the x -direction in reference frame Σ' . If we assume that time is invariant (as we're still currently doing classical physics without having introduced special relativity), then:

$$t = t'$$

And as the motion is entirely in the x-axis:

$$y = y'$$

After time t :

$$x = x' + \text{distance moved by } \Sigma' \text{ in time } t \text{ relative to } \Sigma$$

$$x = x' + vt$$

The object velocity is defined as:

$$\begin{aligned} u &= \frac{dx}{dt} = \frac{d}{dt}(x' + vt) \\ &= \frac{dx'}{dt} + v = u' + v \end{aligned}$$

This holds if and only if time is the quantity we treat as being invariant. This agrees with our classical understanding.

3.1 Where does this break down?

Lets assume that the moving object is a photon, moving in Σ' with velocity c' , hence, according to the previous derivation:

$$c = c' + v$$

Therefore observers in different frames will measure different values for the speed of light...oh no!

If we assume Galilean invariance, the laws of physics are the same in all reference frames, and yet the speed of light is included as a constant in many laws (i.e. electromagnetism). Therefore observers must measure the same speed of light in all reference frames.

This is a contradiction - we cannot assume that both time and the laws of physics are invariant.

3.2 Experimental Verification

The earth travels around the sun extremely quickly, and the sun is travelling even faster around the galactic centre. Therefore, the earth is moving in space.

If Galilean relativity is correct, we would measure the speed of light in one direction as $c + u$ and measure the speed of light in the other direction as $c - u$.

This was tested by the Michelson-Morley experiment, where incoming light was split by a half-silvered mirror. Half the light travels in one direction, and half the light is split off by 90° . They reflect off a pair of mirrors and are recombined in a splitter to be observed.

If the speed of light was different in different directions, the two beams would be out of phase and we would observe an interference pattern on combination. We would expect to see phase difference that varies with time, i.e. turning from destructive to constructive etc.

This was not observed and the interference pattern generated was constant. Therefore, the results showed no variation in the speed of light.¹

4 The Solution - Einstein's Special Relativity

To fix this contradiction, Einstein came up with two postulates:

1. The laws of physics are the same in any inertial frame of reference.
 - This is the same as Galilean invariance and is easily believable.
2. The speed of light is constant in every frame of reference.
 - This was revolutionary and is much less intuitive.

The last postulate is difficult to understand intuitively, but makes everything work if we accept it as true!

¹This is the same idea used in the LIGO experiment to discover gravitational waves, except this was used to show that the path length changed, and not the speed of light. If the path lengths changed, this was due to a gravitational wave (ripple in space time) propagating to the earth and interfering with the measurement.

4.1 Proving Time Dilation

Consider an observer on the earth in frame Σ . This observer watches two rockets both travelling side by side away from the earth in the x -direction with speed v . They are both travelling in x , with a constant y -difference y_0 between them.

Suppose the upper rocket A fires a laser beam directly at B (i.e. directly in the y -direction downwards).

Frame Σ : This is the rest frame of the earth (and the observer on earth) with coordinates (x, y) .

Frame Σ' : This is the rest frame of the rockets, with coordinates (x', y') .

In the rockets rest frame Σ' , the rockets are stationary and the time taken for a laser pulse to travel between them is:

$$t_0 = \frac{y_0}{c}$$

Or equivalently:

$$y_0 = ct_0$$

In the earth's rest frame Σ , the observer on the earth doesn't just see the light travelling in the y -direction. It also sees the light moving in the x -direction, as the whole Σ' frame containing the rockets relativistically move away.

Effectively, we have:




Figure 1.1

B moves distance x horizontally while it waits for the laser to hit it, and the length of the path of the laser beam in time t is given by D . If the time taken for light to reach B in Σ is t , then:

$$D = ct$$

As the speed of light is constant in all frames. As the rockets are moving away:

$$x = vt$$

This gives us a Pythagorean triangle, where:

$$D^2 = x^2 + y^2 = x^2 + y_0^2$$

And substituting in:

$$c^2 t^2 = v^2 t^2 + c^2 t_0^2$$

$$t^2 (c^2 - v^2) = t_0^2 c^2$$

$$t = t_0 \sqrt{\frac{c^2}{c^2 - v^2}}$$

Dividing through by c_2 :

$$t = t_0 \sqrt{\frac{1}{1 - \frac{v^2}{c^2}}}$$

Or:

$$t = t_0 \gamma, \quad \gamma = \frac{1}{\sqrt{1 - \frac{v^2}{c^2}}}$$

$\gamma > 1$ for all object speeds that don't exceed the speed of light (which is required), so the time observed in the frame Σ is longer than the "proper time" observed by the rocket.

5 The Twin Paradox

Consider two twins, an astronaut and a physics professor. The astronaut goes on a long trip close to the speed of light and reunites with his brother. Which brother is older?

Special relativity says that because there is no absolute frame of reference, the brother travelling away also sees the same problem in reverse, i.e. he sees the physics professor brother travelling away in the opposite direction at equal and opposite speed, i.e. they are the same age.

In practice, the astronaut is non-inertial and special rel isn't sufficient, so we can't solve it directly.

Thu 19 Feb 2026 15:00

Lecture 9 - Special Relativity II: Length Contraction

1 Proving Length Contraction

Consider a horizontal laser cavity. We fire a laser from the left of the cavity (point A) to the right of the cavity (point B). The laser bounces off a mirror at point B and bounces back.

The cavity has length L_0 as measured in the rest frame Σ_0 . In this frame:

$$t_0 = \frac{2L_0}{c}, \quad \text{Where } t_0 \text{ is the time taken for the } A \rightarrow B \rightarrow A \text{ journey in } \Sigma_0$$

Now suppose the cavity moves at speed v relative to a stationary observer in a different frame (Σ).

The observer sees the cavity as having length L , how long does the $A \rightarrow B \rightarrow A$ journey take as observed by a stationary observer in Σ .

$$t = t_{A \rightarrow B} + t_{B \rightarrow A}$$

The cavity is moving away from the observer, so the laser beam doesn't just travel L when going from $A \rightarrow B$. It must travel L plus the distance the whole cavity has moved away in this time (as B is moving away from the pulse). This extra distance is $vt_{A \rightarrow B}$, hence the distance travelled when going from A to B is $L + vt_{A \rightarrow B}$.

As c is constant in all frames, we can say:

$$c = \frac{L + vt_{A \rightarrow B}}{t_{A \rightarrow B}}$$

$$t_{A \rightarrow B}(c - v) = L$$

$$t_{A \rightarrow B} = \frac{L}{c - v}$$

And now on the return journey from $B \rightarrow A$, as it travels back towards A , the cavity is moving in the same direction as the light, so A "catches up" to the pulse and results in a smaller required distance to be travelled, $L - vt_{B \rightarrow A}$

$$c = \frac{L - vt_{B \rightarrow A}}{t_{B \rightarrow A}} \implies t_{B \rightarrow A} = \frac{L}{c + v}$$

And returning to the total time:

$$\begin{aligned} t &= t_{A \rightarrow B} + t_{B \rightarrow A} \\ t &= \frac{L}{c - v} + \frac{L}{c + v} \\ &= L \left(\frac{1}{c - v} + \frac{1}{c + v} \right) \\ &= L \left(\frac{c + v + c - v}{(c - v)(c + v)} \right) \\ &= L \frac{2c}{c^2 - v^2} \\ &= \frac{2L}{c} \frac{c^2}{c^2 - v^2} \\ &= \frac{2L}{c} \frac{1}{1 - \frac{v^2}{c^2}} \end{aligned}$$

$$= \frac{2L}{c} \gamma^2$$

We now apply time dilation, which says that $t = \gamma t_0$, hence:

$$\gamma t_0 = \frac{2L}{c} \gamma^2$$

$$t_0 = \frac{2L}{c} \gamma$$

From the rest frame, we know that $t_0 = 2L_0/c$, so:

$$\frac{2L_0}{c} = \frac{2L\gamma}{c}$$

$$L_0 = L\gamma$$

As $\gamma > 1$, $\forall (u < c)$, the proper length measured from the rest frame will always be greater than the length measured by an observer, hence length is contracted for a moving object.

2 Example

In a particle accelerator, protons are accelerated to a measly $0.9c$. These protons pass through a tunnel of length 2km as viewed from the a laboratory rest frame at the accelerator.

Recall that:

$$t = \gamma t_0$$

$$L = \frac{L_0}{\gamma}$$

$$\gamma = \frac{1}{\sqrt{1 - \beta^2}}, \quad \text{where: } \beta = u/c$$

2.1 How long would the journey take, according to the rest frame?

Viewed in the lab frame, and $v = d/t$ so:

$$t = \frac{2 \times 10^3 \text{m}}{0.9 \times 3 \times 10^8} = 7.4 \times 10^{-6} \text{s} = 7.4 \mu\text{s}$$

No relativity needed!

2.2 How long would the journey take, according to the proton's frame?

We do need to consider relativity here, and use time dilation:

$$t = \gamma t_0$$

$$\gamma = \frac{1}{\sqrt{1 - 0.9^2}} = 2.3$$

$$t = \gamma t_0 \implies t_0 = \frac{t}{\gamma} = \frac{7.4 \mu\text{s}}{2.3} = 3.2 \mu\text{s}$$

2.3 How long is the tunnel, according to the proton's frame?

The rest frame now corresponds to the tunnel. It has proper length in its rest frame of 2km.

As viewed by the protons in their rest frame, it is the tunnel that is moving and is rushing towards them at $0.9c$. This is therefore a length contraction problem.

$$L = \frac{L_0}{\gamma} = \frac{2 \text{km}}{2.3} = 870 \text{m}$$

2.4 Checking Values

To check, we can reuse $v = d/t$ but this time in the proton rest frame.

$$d = 870\text{m}, \quad t = 3.2 \times 10^{-6}\text{s}$$

Hence

$$v = \frac{870\text{m}}{3.2 \times 10^{-6}\text{s}} = 2.7 \times 10^8 \text{ms}^{-1} = 0.9c \text{ as required!}$$

3 Example II: Cosmic Rays

The highest energy cosmic rays are protons with massive energies $E \sim 10^{20}\text{eV}$, compared to the LHC with $E \sim 10^{12}\text{eV}$

This corresponds to a $\gamma = 10^{11}$, and our galaxy is $\sim 10^{20}\text{m}$ across.¹

As viewed on earth, these cosmic protons travel at $v \approx c$, hence,

$$t \approx \frac{d}{v} \approx \frac{10^{20}}{3 \times 10^8} = 3 \times 10^{11}\text{s} \approx 10^4 \text{years}$$

As viewed by the protons:

$$t_0 = \frac{t}{\gamma} = \frac{3 \times 10^{11}}{10^{11}} \approx 3\text{s}$$

Which is starkly different!

4 Lorentz Transformations

In general, transforming between coordinates in two different frames can get extremely messy, more than can easily be handled in simple applications of length contraction or time dilation.

The Lorentz Transformations provide a general set of coordinate transformations between Σ and Σ' , i.e. $(x, y, z, t) \rightarrow (x', y', z', t')$.

These transformations must be:

1. Symmetric about a change in sign of u , i.e. the transformation from $\Sigma \rightarrow \Sigma'$ with u and the transformation from $\Sigma' \rightarrow \Sigma$ with $-u$ must be the same.
2. They must be linear as otherwise a constant speed in one would not be constant in the other (and therefore not inertial frames).
3. If relative velocity is only in the x direction, the transformations must be independent of y and z .

These transformations are (assuming motion only in x):

$$\begin{aligned} x &= \gamma(x' + vt') \\ y &= y' \\ z &= z' \\ t &= \gamma\left(t' + \frac{x'v}{c^2}\right) \end{aligned}$$

And:

$$\begin{aligned} x' &= \gamma(x - vt) \\ y' &= y \\ z' &= z \\ t' &= \gamma\left(t - \frac{xv}{c^2}\right) \end{aligned}$$

We do not need to know the derivations, but do need to know the results.

¹ $\gamma = E/mc^2$ - will be expanded on in later lectures.

Tue 24 Feb 2026 11:00

Lecture 10 - Special Relativity III: Lorentz Transformations

Recap of the first two relativity lectures:

- Recap from Semester 1 Special Rel:
 - Background
 - Einstein's postulates.
 - Time dilation.
 - Lorentz contraction.
- Galilean transformations and when they fail.
- Lorentz transformations for space and time.

1 Galilean vs. Lorentz Transformations

In a standard setup with two frames, Σ and Σ' , where Σ' moves with speed v relative to Σ .

In Σ , we have coordinates (x, y, t) , in Σ' we have coordinates (x', y', t') .

In a Galilean transformations, we kept time invariant across both frames, so $t = t'$. This lead to $x' = x - vt \implies u' = u - v$ but raised a contradiction with the speed of light as observers in different frames would take two measurements of the speed of light.

In the second lecture, we consider the speed of light as being invariant across all frames instead of time to resolve the contradiction. This gave us the Lorentz Transformations:

$$x' = \gamma(x - vt)$$

$$t' = \gamma\left(t - \frac{xv}{c^2}\right)$$

Where $y = y', z = z'$.

As we cannot use $u' = u - v$ due to the contradiction with the speed of light, we in this lecture will consider:

- Lorentz transformations for velocity.
- Space-time, Lorentz invariance, causality.

2 Lorentz Transformation for Velocity

We can revisit the question from the first lecture, but in the Lorentz transformation perspective rather than a Galilean perspective.

Again, we have our two standard frames Σ, Σ' . An object is moving with speed u' in frame Σ' . We want to find an expression for the velocity of the object as viewed from Σ , given by u . The prime frame moves at speed v wrt the rest frame and the objects motion is entirely along the x axis.

By definition, we have:

$$u = \frac{dx}{dt} = \frac{dx}{dt'} \frac{dt'}{dt} = \frac{\left(\frac{dx}{dt'}\right)}{\left(\frac{dt}{dt'}\right)} \quad (1)$$

And:

$$\frac{dx}{dt'} = \gamma \left(\frac{dx'}{dt'} + v \right) = \gamma (u' + v) \quad (2)$$

$$\frac{dt}{dt'} = \gamma \left(1 + \frac{v}{c^2} \frac{dx'}{dt'} \right) = \gamma \left(1 + \frac{u'v}{c^2} \right) \quad (3)$$

And substituting (3) and (2) into (1):

$$u = \frac{\gamma(u' + v)}{\gamma \left(1 + \frac{u'v}{c^2} \right)} \Rightarrow \boxed{u = \frac{u' + v}{1 + \frac{u'v}{c^2}}}$$

To get the reverse transformation, we invoke forward-backward symmetry, i.e. swap $x \rightarrow x'$, $t \rightarrow t'$ etc, where $v \rightarrow -v'$:

$$u' = \frac{u - v}{1 - \frac{uv}{c^2}}$$

2.1 Limiting Cases

Firstly, we'll look at the non-relativistic case, i.e. where $u, v \ll c$:

$$u = \frac{u' + v}{1 + \frac{u'v}{c^2}} = \frac{u' + v}{1 + \text{small}} \approx u' + v$$

This is the classical Galilean approach, which is what we'd expect for non-relativistic behaviour.

What about an ultra-relativistic limit where $u, u' \rightarrow c$ with $v \ll c$. Now we have:

$$u = \frac{u' + \text{small}}{1 + \text{small}} \approx u'$$

As the speed of light is invariant, speeds close to the speed of light do not change between reference frames.

2.2 Example

An observer on a rocket sees a second rocket moving away from them at $u_1 = 0.8c$. Another observer on a planet sees the first rocket moving at $0.7c$. Assuming all motion is in the same direction, how fast does the planetary observer see the second rocket?

- From Σ' (the first rocket's rest frame), the second rocket moves with $0.8c$.
- From Σ (the planetary rest frame), the first rocket moves with $0.7c$, and the second rocket moves with some unknown speed u .

We apply the formula:

$$u = \frac{u' + v}{1 + \frac{u'v}{c^2}}$$

In Σ' , $u' = 0.8c$. The two frames move with relative velocity equal to the change in speed of the first rocket from $\Sigma \rightarrow \Sigma'$: $v = 0.7c$.

In frame Σ we are trying to do:

$$\begin{aligned} u &= \frac{0.8c + 0.7c}{1 + \frac{0.8c \times 0.7c}{c^2}} \\ &= \frac{1.5c}{1.56} = 0.96c \end{aligned}$$

3 Space-Time and Lorentz Invariants

From length contraction and time dilation, we can see that in special relativity, intervals in time and space are not fixed but vary from frame-to-frame.

This means that the order of events may even be different for different observers.

There are some quantities however which are invariant across frames, such as the speed of light. One of these is space-time.

3.1 Space-Time Invariance

As usual, we have Σ and Σ' with relative speed v . The axes of these frames coincide at $t = 0, x = 0$ and $t' = 0, x' = 0$.

At $t = t' = 0$, there is a flash of light at the origin which propagates isotropically.

In Σ , the light forms a spherical shell of radius Δr at time Δt , and in Σ' it reaches a radial distance $\Delta r'$ at time $\Delta t'$.

After time ΔT in Σ we have $\Delta r = c\Delta T$. Similarly in Σ' we have $\Delta r' = c\Delta t'$.

Squaring these we have:

$$\begin{aligned}(\Delta r)^2 &= c^2 \Delta t^2 = \Delta x^2 + \Delta y^2 + \Delta z^2 \\ (\Delta r')^2 &= c^2 \Delta t'^2 = \Delta x'^2 + \Delta y'^2 + \Delta z'^2\end{aligned}$$

Hence:

$$c^2 \Delta t^2 - \Delta x^2 - \Delta y^2 - \Delta z^2 = 0 = c^2 \Delta t'^2 - \Delta x'^2 - \Delta y'^2 - \Delta z'^2$$

This means that, **for light**:

$$c^2 \Delta t^2 - \Delta x^2 - \Delta y^2 - \Delta z^2 = 0$$

We call this quantity Δs^2 :

What about for objects moving at a speed less than the speed of light? We want to apply the Lorentz transformations to $\Delta s'^2$ (again treating the “boost” as being in x):

$$\begin{aligned}\Delta s'^2 &= c'^2 \Delta t'^2 - \Delta x'^2 - \Delta y'^2 - \Delta z'^2 \\ &= c^2 \gamma^2 \left(\Delta t - \frac{v}{c^2} \Delta x \right)^2 - \gamma^2 (\Delta x - v \Delta t)^2 - \Delta y^2 - \Delta z^2 \\ &= c^2 \gamma^2 \left(\Delta t^2 + \frac{v^2}{c^4} \Delta x^2 - 2 \frac{v}{c^2} \Delta t \Delta x \right) - \gamma^2 (\Delta x^2 + v^2 \Delta t^2 - 2v \Delta x \Delta t) - \Delta y^2 - \Delta z^2 \\ &= \Delta t^2 (c^2 \gamma^2 - v^2 \gamma^2) + \Delta x^2 \left(\frac{v^2 \gamma^2}{c^2} - \gamma^2 \right) + \Delta t \Delta x (-2v \gamma^2 + 2v \gamma^2) - \Delta y^2 - \Delta z^2 \\ &= \Delta t^2 \gamma^2 (c^2 - v^2) + \Delta x^2 \gamma^2 \left(\frac{v^2}{c^2} - 1 \right) - \Delta y^2 - \Delta z^2 \\ &= \Delta t^2 \frac{c^2 - v^2}{1 - \frac{v^2}{c^2}} + \Delta x^2 \frac{\left(\frac{v^2}{c^2} - 1 \right)}{1 - \frac{v^2}{c^2}} - \Delta y^2 - \Delta z^2 \\ \Delta s'^2 &= \boxed{c^2 \Delta t^2 - \Delta x^2 - \Delta y^2 - \Delta z^2 = \Delta s^2}\end{aligned}$$

Hence Δs^2 is invariant under the Lorentz transformations regardless of speed.

Tue 03 Mar 2026 11:00

Lecture 12 - Special Relativity IV: More Space-Time

Recap from last lecture:

- Lorentz Transformations for space and time.
- Lorentz Transformations for velocity/velocity addition.
- A Lorentz Invariant: The space-time interval is invariant across frames of reference:

$$\delta S^2 = c^2 \Delta t^2 - \Delta r^2$$

In this lecture:

- Minkowski space and causality.
- Relativistic energy and momentum.

1 Space-Time Invariance

Last lecture we showed that:

$$c^2 \Delta t^2 - \Delta r^2 = c^2 \Delta t'^2 - \Delta r'^2 = \quad (= 0 \text{ for light})$$

And we define the space-time interval Δs as $\Delta S^2 = c^2 \Delta t^2 - r^2$. There is some relationship between space and time that keeps this quantity constant across frames. What does this quantity actually *mean*, however?

In a 3D set of axes, we have:

$$\Delta r^2 = \Delta x^2 + \Delta y^2 + \Delta z^2$$

And we have defined:

$$\Delta s^2 = c^2 \Delta t^2 - \Delta r^2 = c^2 \Delta t^2 - (\Delta x^2 + \Delta y^2 + \Delta z^2)$$

This looks almost like a 4D space (where one axis is time), but with some strange negative signs appearing? We don't have the equipment yet to properly understand the negative signs, but can be in future courses.

2 Minkowski Space (Space-Time Diagrams)

In 2D, we consider one spatial coordinate and one time coordinate, where the x axis represents our spatial coordinate and the vertical axis is given by ct .

Consider some kind of light-emitting event happening at the origin. This light travels at a fixed speed, so we end up with the path of light at 45 degrees on this space-time diagram:




Figure 4.1

These are given by the lines $c^2t^2 = \pm x^2$ and are called “light cones”, as they form cones in higher dimensions. The positive velocity axis is called the “future cone”, the path taken by light emerging from event A. The negative velocity axis is called the “past cone”, and represents the path taken by light emerging at A.




Figure 4.2: Light cones in a 3D space (two spatial, one velocity)

Inside the future cone, events may be caused by the event happening at A. Things travelling at a velocity $< c$ can reach them.

The region inside the past cone gives regions in spacetime where events could have caused A.

Events outside the two cones cannot influence, nor can they be influenced by A. They cannot be causally connected in any way to the event at A.

We now have two events. A is defined as before, and a new event B separated from A by distance $\Delta \underline{r}$ and time Δt .

What is the condition that B lies within the light cone given by A? We require:

$$|\Delta \underline{r}| \leq c|\Delta t|$$

Or in invariant form:

$$\Delta S^2 = c^2 \Delta t^2 - \Delta r^2 \geq 0$$

We call ΔS the space-time interval, or invariant distance.

There are three possible cases:

- $\Delta S^2 > 0$, “time-like intervals”. This is where $|\Delta \underline{r}| \leq c|\Delta t|$ and B is inside A’s light cone. $\Delta S \in \mathbb{R}$. Here, a signal can pass between A and B and the two events may be causally connected (where A causes B).
- $\Delta S^2 = 0$. Here, $|\Delta \underline{r}| = c|\Delta t|$. This is where B sits on A’s light cone, and A/B may be causally connected. A signal can only travel between the two events if the signal is at velocity c . This is called a “light-like interval”
- $\Delta S^2 < 0$, “space-like intervals”. This is where $|\Delta \underline{r}| > c|\Delta t|$. $\Delta S = \sqrt{c^2 \Delta t^2 - \Delta r^2} \in i\mathbb{R}$. No signal can pass between A and B, not even light. No causal connection is possible.

3 Causality and Time Order

It is possible for observers in two different frames to disagree on the order in which events have taken place. If we have two events with t_1, t_2 in Σ and t'_1, t'_2 in Σ' , there are possible cases where:

- $t_2 > t_1$.
- $t'_1 > t'_2$.

This may seem uncomfortable - does this cause a breakdown in causality?

Suppose we have a frame Σ with two events, $(x_1, t_1), (x_2, t_2)$, such that $\Delta t = t_2 - t_1 > 0$.

Frame Σ' moves with speed v relative to Σ and in Σ' the same two events have coordinates (x'_1, t'_1) and (x'_2, t'_2) . For the observers to disagree on causality, we have $\Delta t' t'_2 - t'_1 < 0$. We begin by applying the Lorentz Transformation to $\Delta t'$:

$$\begin{aligned} \Delta t' &= \gamma \left(t_2 - \frac{v}{c^2} x_2 \right) - \gamma \left(t_1 - \frac{v}{c^2} x_1 \right) < 0 \\ \implies \gamma(t_2 - t_1) + \frac{\gamma v}{c^2} (-x_2 + x_1) &< 0 \\ \Delta t - \frac{v}{c^2} \Delta x &< 0 \end{aligned}$$

I.e. $\Delta t < \frac{v}{c^2} \Delta x$ or $c\Delta t < \frac{v}{c} \Delta x$. We know that $v/c < 1$, hence $c\Delta t < \Delta x$. This means that:

$$\Delta S = \sqrt{c^2 \Delta t^2 - \Delta x^2}$$

Which is the square root of a negative number, hence this gives a space-like interval. This means that while we can define two points in space-time where two observers will disagree about their order, the space-time invariant is imaginary hence these events cannot be causally linked. This means that observers disagreeing on event order does not break causality.

Tue 10 Mar 2026 11:00

Lecture 13 - Special Relativity V:

Recap of last lecture:

- Minkowski Space (space-time)
- Time-like and space-like intervals.
- Ensuring that this new model does not violate causality.
- Simultaneity.

Tue 24 Mar 2026 11:00

Lecture IDK - Special Relativity VII: Kinematics Examples, Doppler and Natural Units

1 Example I

As viewed in a lab frame, a cosmic ray proton of energy E_p hits a stationary proton. What is the minimum threshold energy of E_p that allows the reaction:

$$p + p \rightarrow p + p + \pi$$

To take place, given $m_p = 938\text{MeV}/c^2$ and $m_\pi = 104\text{MeV}/c^2$?

Let the cosmic ray proton be proton one with E_p, p_p and the stationary proton be proton 2 with m_p .

We can consider energy and momentum in the lab frame, initially (before the collision):

$$(\Sigma E)_{\text{initial}}^{\text{lab}} = E_p + M_p c^2 \quad \text{and} \quad (\Sigma p)_{\text{initial}}^{\text{lab}} = p_p \quad (1)$$

For a threshold energy, the produced particles have no relative motion (i.e. at rest in the centre of mass frame). This means that all available relativistic energy goes into particle production with no excess.

Hence in the centre of mass frame finally, we have two stationary protons and one stationary pion. This is the lowest energy state we can have in the centre of mass frame, with all these particles still existing.

$$(\Sigma E)_{\text{final}}^{\text{CMS}} = (2m_p + m_\pi)c^2 \quad \text{and} \quad (\Sigma p)_{\text{final}}^{\text{CMS}} = 0 \quad (2)$$

We can then apply the energy momentum invariant:

$$s = [(\Sigma E)^2 - (\Sigma p)^2]_{\text{final}}^{\text{lab}} = [(\Sigma E)^2 - (\Sigma p)^2]_{\text{CMS}}^{\text{final}} \quad (3)$$

Inserting 1, 2 into 3:

$$\begin{aligned} (E_p + m_p c^2)^2 - p_p^2 c^2 &= (2m_p + m_\pi)^2 c^4 \\ \Rightarrow E_p^2 + m_p^2 c^4 + m_p c^2 E_p - p_p^2 c^2 &= (2m_p + m_\pi)^2 c^4 \end{aligned}$$

For a single particle, $E_p^2 - p_p^2 c^2 = m_p^2 c^4$, hence:

$$\begin{aligned} 2m_p^2 c^4 + 2m_p c^2 E_p &= (2m_p + m_\pi)^2 c^4 \\ E_p &= \frac{(2m_p + m_\pi)^2 c^4 - 2m_p^2 c^4}{2m_p c^2} \end{aligned}$$

And we have known masses m_p, m_π hence (noting that the masses are in units energy per speed of light squared, so the c s cancel out):

$$E_p = \frac{(2 \times 938 + 104)^2 - 2 \times 938^2}{2 \times 938} = 1228\text{MeV}$$

2 Example II

In an LHC experiment, a B_0 meson ($b\bar{d}$) with mass $5\text{GeV}/c^2$ is produced in a lab frame momentum of $50\text{GeV}/c$. The meson has proper decay time $2 \times 10^{-12}\text{s}$. How far does the experiment observe the meson fly before decay?

3 Relativistic Longitudinal Doppler Effect

We want to build an analogue for the sonic doppler effect for light. This is especially important in astronomy, i.e. tracking the movement of cosmic spectral lines to determine speeds of distant sources.

We will only consider the longitudinal effect, i.e. the movement of bodies directly towards or away from an observer in their line of sight. We will not (yet) consider transverse effects in this course.

This comes from two components:

- The standard doppler effect (as for sonic waves).
- Relativistic time dilation.