
LC Introduction to Particle Physics and Cosmology Lecture Notes

Year 1 Semester 2

Ash Stewart

MSci Physics with Particle Physics and Cosmology

School of Physics and Astronomy
University of Birmingham

Table of Lectures

Lecture 1: Particle Physics I: The Standard Model	2
Lecture 2: Particle Physics II: Luminosity and Particle Signatures	5
Lecture 3: Particle Physics III: Particle/Matter Interactions	10
Lecture 4: Particle Physics IV: Vertexing and Tracking Systems	14
Lecture 5: Particle Physics V: Calorimetry	18
Lecture 6: Particle Physics VI: Calorimetry II and PID	22
Lecture 7: Cosmology I: A Brief Introduction to Cosmology	25
Lecture 8: Cosmology II: Expansion of the Universe and Key Principles	29
Lecture 9: Cosmology III: Friedmann Equation	32
Lecture 10: Cosmology IV: Friedmann Equation, Hubble and Redshift Combined	35
Lecture 11: Cosmology V: Cosmological Models	39
Lecture 12: Cosmology VI: Measurements, Universe Age and Deceleration	42
Lecture 13: Cosmology VII: Cosmological Constant, CMB and Big Bang Problems	45

Thu 22 Jan 2026 16:00

Lecture 1 - Particle Physics I: The Standard Model

1 Course Introduction

Course Structure

- Particle Physics: 6 lectures in weeks 1 to 6.
 1. Introduction and the standard model.
 2. Experimental measurements.
 3. Interactions with matter
 4. Tracking detectors I
 5. Tracking detectors II
 6. Calorimeters and Particle Identification.
- Cosmology: 4 lectures in weeks 7 to 11.

Course Aims

- Overview of current methods in Particle Physics experiments.
- An emphasis placed on the questions and challenges.

For example, the LHC has already been programmed with experiments all the way up to 2041. Therefore, any detectors we design today only become relevant in over a decade, which makes good detector design decisions incredibly important. This course will equip us to understand what drives those design choices.

The course is assessed by a single one hour long exam, weighted half particle physics and half cosmology.

Recommended Texts

- Detectors for particle radiation (2nd edition), K. Kleinknecht (1998)
- Particle Physics, Martin and Shaw.
- High Energy Physics, D. H. Perkins (2nd through 4th editions)
- Feynman Lectures.
- Modern Particle Physics, M. Thomson.

2 Matter Particles

Fermions all have quantum spin $1/2$. Spin is a purely inherent quantum property (like mass or charge) and has no classical representation, but is analogous to angular momentum. They are subject to Fermi-Dirac statistics, which means that no identical fermion in a system of fermions can have the same quantum number as any other. Fermions are divided into two types, quarks and leptons.

2.1 Quarks

There are three generations of quarks:

First Generation

- Up Quark (u), mass of $\approx 0.001\text{GeV}$
- Down Quark (d), mass of $\approx 0.001\text{GeV}$

Second Generation

- Charm Quark (c), mass of $\approx 1.3\text{GeV}$
- Strange Quark (s) mass of $\approx 95\text{MeV}$

Third Generation

- Top Quark (t), mass of $\approx 175\text{GeV}$
- Bottom Quark (b), mass of $\approx 4.3\text{GeV}$

“up-type quarks”, u, c, t have electromagnetic charge $+2/3$ and “down-type quarks”, d, s, b , have electromagnetic charge $-1/3$ (charges in units of e). Quarks do not ever exist alone in isolation.

2.2 Leptons

There are three generations of leptons, given by the electron e^- , muon, μ^- and tau τ^- . These all have charge of -1 (in units of e). These have masses (in MeV) of approximately 0.5, 105, 1800.

These also have associated neutrinos, the electron neutrino, the mu neutrino and the tau neutrino, ν_e, ν_μ, ν_τ . These are not massless, but have a tiny mass many orders of magnitude smaller than their corresponding non-neutrino counterparts. These are neutrally charged.

Leptons are not subject to the strong interaction.

2.3 Hadrons

Quarks do not exist in isolation, but form bound states subject to the strong force called hadrons. There are two types of hadrons - baryons and mesons.

Baryons are formed from three quarks, which may be the same or different $q_1 q_2 q_3$.

Mesons are formed from a quark-antiquark pair, $q_1 \bar{q}_2$. Examples of baryons include the proton and the neutron, given by uud and udd .

3 Forces

As far as we know, there are four fundamental forces:

- Gravity
- Electromagnetism
- Strong
- Weak

For considering particle interactions, we disregard gravity as it becomes incredibly weak for small masses. Creating a complete theory that incorporates all four is an open question in physics. It's okay to neglect it, but it is unsatisfying.

We consider these forces as arising by the exchange (between two particles subject to a force between them) of particles called bosons. These have spin-1, so are called “gauge bosons”. These are subject to Bose-Einstein statistics, which does not impose the same restriction as Fermi-Dirac for quantum numbers in a system.

For EM: The exchange particle is a photon, γ . This is represented on a Feynman diagram as a wiggly line. They are massless and couple to electric charge.

For Weak: The exchange particle is a $W^\pm$ or Z^0 boson. This is represented by a wiggly line or a dotted straight line. They are not massless, and have masses of approx. 80 and 90GeV respectively.

For Strong: The exchange particle is a gluon g . This is represented by a series of curls on a Feynman diagram. They are massless. They couple to “colour charge” which is just another quantum number analogous to electric charge. Just like electric charge has values $\pm$, colour charge has values we denote r, g, b


Quarks are subject to the strong, electromagnetic and weak interactions

Leptons are not subject to the strong interaction, but the e, μ, τ are subject to EM (ν is not as it is neutrally charged), and all are subject to the weak interaction. This makes neutrinos very difficult to detect as they are only affected by the weak interaction.

4 Feynman Diagrams


Feynman diagrams are space (y-axis), time (x-axis) diagrams to show allowed interactions between particles.

Consider a simple example of electron-positron annihilation. They travel towards each other, meeting and annihilating into either a photon or a Z boson. This is called a ‘time-like exchange’. The boson then decays and we see pair production of two muons (one μ^- muon and one μ^+ antimuon).



Arrows in Feynman diagrams convey “fermion flow”. This means that for a matter particle, the arrows aligns to the time axis. For antimatter particles, they antialign. Some conservation laws (i.e. charge) apply at the vertex level, while others only apply across whole processes.

We now consider a space-like exchange where the exchange is aligned with the vertical (space) axis. An antimuon scatters off a positron like such:



5 Luminosity

We can determine the rate of interactions with the following:

$$W = \mathcal{L}\sigma$$

Where $W(\text{s}^{-1})$ is interaction rate, $\mathcal{L}(\text{cm}^{-2}\text{s}^{-1})$ represents the luminosity (an attribute of the accelerator being used), and $\sigma(\text{cm}^{-2})$ is the cross section, representing the underlying physics of the interaction.

These are investigated in greater detail in Lecture 03.

Mon 02 Feb 2026 12:00

Lecture 2 - Particle Physics II: Luminosity and Particle Signatures

1 Cross Sections

Consider a proton-proton interaction, producing some unknown particle X :

$$pp \rightarrow ppX$$

We have said that the rate of interaction is given by:

$$W = \mathcal{L}\sigma$$

Where $\mathcal{L}$ in $\text{cm}^{-2}\text{s}^{-1}$ is (coarsely) a parameter of the accelerator, describing its ability to produce collisions, and σ in cm^{-2} is a measure of interaction probability. Even though the particles are point-like, we treat them as having an effective area, and the magnitude of that area dictates how likely an interaction is to take place.

In this interaction, we have two protons (modelled as solid balls) passing immediately next to each other (one travelling clockwise and one counter-clockwise) around the accelerator. Assuming they pass immediately next to each other, and we model them as having radius 10^{-15}m , we have a separation between the centres of each proton as $2 \times 10^{-15}\text{m}$, therefore a cross section of:

$$\pi (2 \times 10^{-15}\text{m})^2 \sim 0.125 \times 10^{-28}\text{m}^2$$

To move this to a less annoying length scale, we define a new unit, the barn:

$$1\text{barn} \equiv 10^{-28}\text{m}^2 = 10^{-24}\text{cm}^2$$

In reality, this model may approximate a cross section, but it's not accurate. In reality, there's a much wider variation:




Figure 2.1

Here we have two x-axes running in parallel - the black axis is the momentum in a lab frame (as it hits some fixed target) while the red axis is the corresponding “centre of mass energy”. How do we relate these two?

We want to know what the maximum mass of the particle we can generate is. In the lab frame, this requires us to take the momentum of the incoming and generated particle into account. We then have to take the final momentum of the system into account to conserve momentum, as the whole system must continue moving in the direction of motion for conservation.

Translating into the frame of reference given by the system centre of mass gives us a system where the two masses can be thought of as approaching each other with equal and opposite momenta. Since the total momentum is zero, the objects (incoming particle and the target) can theoretically hit each other and come to a complete stop. Because the system does not have to keep moving after the collision, all of the energy in this frame is free to be converted into the mass of a new particle.

While this may not be accurate in practice, it gives us a hard upper maximum for the possible energy available for production.

We can show relativistically that the energy in the centre of mass frame (labelled $\sqrt{s}$, the energy available for particle production) is:

$$E_{\text{com}} = \sqrt{s} \propto \sqrt{p_{\text{lab}}}$$

This can be seen in the final values of the x-axis, which (both starting approx. 1) are 10^8 and 10^4 . This tells us that we reach diminishing returns with a fixed target collider - increasing energies by 8 orders of magnitude only increases the energy available for particle production by 4 orders. This is an inherent inefficiency of a fixed target collider.

This is much cheaper as its easier to align, we just fire a beam at a block of (for example) lead. It also makes it quite easy to change the target material. Changing materials in a colliding-beam collider (where two beams are fired in opposite directions, one clockwise and one anticlockwise, and collide with each other), i.e. to fire lead nuclei requires an extensive recalibration process.

In a dual-beam collider, only a very small proportion of the accelerated material from each beam actually interact with each other. In a fixed target collider, the target is much more dense, so we see a higher rate of interactions.

In summary, the advantages of a fixed target collider are:

1. Easier to collide.
2. Easier to change the target.

3. Very high density

2 Luminosity

2.1 Fixed Target Case

Consider a fixed target collider. We want to build an expression for the luminosity of this setup.

We have some incoming flux of particles (per second per unit area), J , incident on the block of material with density ρ , thickness t and mass of one nucleus m_A . The beam, modelled by a cylinder, illuminates some circular portion of the block, with area A .

Consider our interaction rate W . This is given by (where V is the volume of a cylinder from the illuminated beam, of thickness equal to the target):

$$W = \underbrace{JA}_{\substack{\text{incident particles} \\ \text{per unit time}}} \times \underbrace{\frac{\rho}{m_A} V}_{\substack{\text{total number of} \\ \text{target particles}}} \times \underbrace{\frac{\sigma}{A}}_{\substack{\text{probability of} \\ \text{interaction}}}$$

$$W = JA \times \frac{\rho}{m_A} At \times \frac{\sigma}{A}$$

$$W = J \times \frac{\rho}{m_A} At \times \sigma$$

$$W = \frac{J\rho At}{m_A} \sigma$$

Comparing to $W = \mathcal{L}\sigma$ gives the luminosity as:

$$\mathcal{L} = \frac{J\rho At}{m_A}$$

2.2 Colliding Beam Case

In a colliding beam case, the derivation is more (and too) complex. It is equal to:

$$\mathcal{L} = \frac{f_{\text{rep}} n_b N_1 N_2}{4\pi\sigma_x\sigma_y}$$

Where:

- f_{rep} is the repetition frequency, i.e the rate of the beam passing the collision point.
- n_b is the number of bunches in the beam.
 - A beam can be thought of as as string of pearls, rather than a single discrete constant beam - i.e. clusters of particles “bunches”, followed by empty space between them.
- N_1, N_2 are the number of particles per bunch for each beam
- σ_x, σ_y are the dimensions of the beam in the x- and y-direction, not a cross section as previously.

3 Examples of Detectors

We have some interaction point producing a spray of particles, and surround this with a series of different detector layers. Each produced particle will trigger a different subset of these layers, defining a unique signature we can use to identify produced particles.

Broadly, in some generic detector, we have:




Figure 2.2

The tracking system is non-destructive. Formed of layers of silicon, a charged particle will ionise small portions of each layer. This can be turned into a signal. Neutral hadrons and photons will pass straight through, but charged particles will leave a deposit of charge and pass through unaffected.

We then move to destructive layers. Particles leave tree-like structures as they pass through these layers and create a shower of particles.

3.1 Decays

Most decays take place over a very low time scale, $< 10^{-8}$ s and produce a final state made up of some subset of the following: $\gamma, \pi^+, \pi^-, \kappa^+, \kappa^-, p, n, \pi^0, e^+, e^-, \mu^+, \mu^-$.

So, in order to detect some exotic particle, we assume that it will either persist long enough to be detected itself, or decay into some subset of these known particles which will reach out detector.

Consider a parent particle A , for example $B^0 (\bar{b}d)$ decaying into two child particles B, C , given by a “J Psi” ($c\bar{c}$), J/ψ and a “K Short”, $K_s^0 (d\bar{s})$:

$$B \rightarrow J/\psi \quad K_s^0$$

$$\bar{b}d \rightarrow c\bar{c} \quad d\bar{s}$$

This decay has a lifetime of 10^{-12} s, via the weak interaction due to the change in quark flavour. The J Psi decays into e^+e^- or $\mu^+\mu^-$ via the EM interaction very rapidly in 10^{-21} s (its lifetime is governed by the strong interaction, which it may also use to decay via, even though we detect it via the EM decay path). The K Short decays into $\pi^+\pi^-$ or $\pi^0\pi^0$ again via the weak interaction with lifetime in 10^{-10} s.

The range of a particle is given by:

$$\text{Range} = \beta\gamma c\tau$$

Where τ is a time scale (lifetime), c is the speed of light, γ is the Lorentz Factor and β scales the range based on the speed actually being travelled. We also have:

$$E = \gamma m$$

$$p = \beta\gamma m$$

And familiarly:

$$E^2 - p^2 = m^2$$

If the B^0 has energy 20GeV and mass 5GeV, we have $\gamma = 4$ and this gives a range of $\approx 1\text{mm}$. This is so small we will never observe it directly. The K_s^0 however has range $\approx 30\text{cm}$, so is detectable.

Crucially:

- SI lifetimes: $\sim 10^{-21} - 10^{-24}\text{s}$
- EM lifetimes: $\sim 10^{-16} - 10^{-20}\text{s}$
- WI lifetimes: $\sim 10^{-12}\text{s}$

Thu 05 Feb 2026 16:00

Lecture 3 - Particle Physics III: Particle/Matter Interactions

In order to look at specific detectors and how they work, we need to consider how particles interact with matter. We'll consider categories of particles and their standard interactions, and use this to build a model for how we can build a detector.

We've looked at this previously:




Figure 3.1

For example:


- Charged particles produce ionisation in the non-destructive tracking system, leaving deposits we can detect as signal.
- Electrons and photons leave distinctive signatures (shape of the shower of particles) in the electromagnetic calorimeters.
- Hadrons leave deposits in the electromagnetic calorimeter, but they dump all their energy (and are mostly identifiable by) the hadronic calorimeter.
- Muons make it through all of the previous layers, and are picked up at the very end by the outermost muon system.

1 Charged Particles

1.1 Ionisation

This is the process that powers the tracking system. A charged particle, h comes into the system (it could be a hadron or a lepton). It exchanges a photon with an electron in the system and excites the electron, ejecting it from the atom.

Strictly:



We can try to characterise the rate of energy loss of the charged particle. The average rate of energy loss per unit distance is:

$$-\left\langle \frac{dE}{dl} \right\rangle \propto \ln E$$

Where E is energy, and the sign is negative as energy is being lost here.

For example, a real detector here might be two charged plates with a high voltage sandwiching a gas mixture. As the gas mixture is ionised, the ionised portion drifts towards one of the electrodes where it induces a detectable current.

1.2 Bremsstrahlung

Translates to “braking radiation”. Consider a free electron radiating a photon. This is impossible for a free particle (as if we consider the electron’s rest frame, emission would violate conservation of energy). We therefore need a source of external interference, in this case matter.

An electron is accelerated by nuclear change as it passes through material and is scattered. This scattering causes bremsstrahlung photon emission. The average rate of energy loss is given by:

$$-\left\langle \frac{dE}{dx} \right\rangle \propto \frac{E}{m^2} \propto \frac{E}{X_0}$$

An electron will then generate more bremsstrahlung than a muon, due to its much smaller mass. X_0 is called radiation length, and is covered in future lectures.

1.3 Cherenkov Radiation

Consider a charged particle moving through a (non-vacuum) material. It emits photons at some angle θ_c if the particle is moving faster than the speed of light in the medium (note this does not violate relativity, as the speed of light in a medium is less than the speed of light in a vacuum).

These emitted photons cause a coherent wavefront to form around the particle’s trajectory, forming a cone around the direction of travel. The angle θ_c is given by:

Geometrically, after time t , the emitted photon has travelled ct/n and the particle vt , hence:

$$\cos \theta_c = \frac{ct/n}{vt} = \frac{c}{nv} = \frac{1}{n\beta}$$

Where n is the refractive index of the material. This is analogous to shock waves forming when an object goes faster than the speed of sound.

A planar detector will take a single cross-section through this cone, detecting rings around the point the charged particle passed through the material. By measuring this ring, we can work out the speed of the

particle, and use this along with a measured momentum (in a tracking detector) to work out the mass of the particle.

Again:

$$-\left\langle \frac{dE}{dx} \right\rangle \propto z^2 \sin^2 \theta_c$$

Where z is the particle charge in units of $|e|$. It is important to note that this is a very small energy loss for the particle. It may emit $10^3 \gamma/\text{cm}$, and only lose a few keV/cm

2 Photons

2.1 Photoelectric Effect

We have a photon strike a atom, transferring energy and forcing an electron to be ejected. The max kinetic energy of the electron is given by the photon energy minus some amount of work to eject it:

$$E_{kmax} = hf - \phi$$

Where ϕ , the work function, is the energy required to liberate one electron from the atom's surface.


2.2 Compton Scattering

See QM1. A photon scatters off a quasi-free electron in an atom. The photon and electron are scattered with a change in energy:

$$\gamma + e^- \rightarrow \gamma' + e^{-'}$$

Thomson scattering is a low-energy form of scattering where the energies do not change, and Rayleigh scattering is a very low energy limit, where the interaction takes place between a photon and multiple atomic electrons.

2.3 Pair Production



A photon interacting with a nucleus can (leaving the nucleus unscathed) produce a particle and the corresponding antiparticle, typically a positron and an electron. This has a minimum photon energy of $E_\gamma > 2m_e c^2$ to ensure conservation of energy isn't violated and must occur in the presence of matter (for the same reason as Bremsstrahlung). The photon is not present in the final state.

However, say a photon has precisely the energy required to create a pair. This would (as it stands) create a pair of electrons with zero momentum (or very small momentum). However, a photon always has a momentum given by its energy, so momentum is not conserved. If the pair travel to attempt to resolve this, they now have some kinetic energy too, which means the photon must have a higher energy, and hence a higher momentum. This creates mismatch, we cannot conserve both energy and momentum in this situation.

By this interaction taking place in the presence of a nucleus, the nucleus can absorb some recoil (via exchange of a virtual photon) to ensure conservation of energy and momentum are satisfied.

This must take place in a Coulomb field to contribute this photon. The present nucleus has charge $z|e|$.

Lets consider the cross section of this interaction:

$$\sigma_{\text{pprod}} \propto \frac{7}{9} \frac{A}{N_A} \frac{1}{X_0}$$

$$^1 m_e c^2 = 0.511 \text{ MeV}$$

X_0 is the radiation length. It is a complex property of the material but is proportional to $1/z^2$.

3 Neutrinos

Neutrinos are really difficult to detect, as they are only effected by the weak interaction and have no ionisation/pair production etc - they are effectively non-interacting. We detect them by detecting the products of a weak force interaction.

For example:

- A muon neutrino can exchange a W boson with a proton, which becomes a neutron. The muon neutrino becomes a muon and the up quark in the proton becomes a down quark in the neutron.
 - We can detect the muon that's produced as the neutrino passes through the detector and interacts.
 - We build a massive multi-tonne detector that puts a large amount of mass in the way of the neutrino, often water.
 - We do this in the hope that it will interact with some of the matter and produce the more-detectable muon.
- The neutrino can scatter off a nucleus.
 - A small amount of energy is exchanged with the nucleus, depositing a small amount of heat energy in the detector's matter.
 - Nothing changes/decays/is produced etc other than a small amount of heat.
 - We build a very sensitive detector capable of detecting this very small amount of heat.

4 Hadrons

Hadrons are really complex in their interactions. They can have inelastic nuclear interactions, with large energy deposits at a small number of sites (compared to an ECal shower where we'd expect to see many smaller deposits). This arises as a result of the nucleus becoming excited or breaking apart entirely and producing a chain of secondary hadrons/particles and a change in the nucleus.

These secondary particles can go on to interact again. These are complex and messy objects which can fragment off to cause many secondary impacts.

5 Conclusion

In summary:

- The key interactions we care about are ionisation, Bremsstrahlung and Cherenkov radiation for charged particles.
- For photons, we care about pair production.
- For hadrons, hadron showers are messy and complex. We add inelastic nuclear interactions on top of an electromagnetic component from ionisation and everything becomes rather tricky.

Mon 09 Feb 2026 12:00

Lecture 4 - Particle Physics IV: Vertexing and Tracking Systems

In our familiar layered model:




Figure 4.1

The first subdetector type is the initial tracking system which provides a non-destructive estimate of a particle's trajectory.

1 Tracking Systems

Purpose: To determine the trajectory of charged particles (usually in the presence of a magnetic field), in order to infer the momentum of the particle. This happens by ionisation, whereby the particle leaves small deposits of charge in either a gas or layers of a semiconductor sensor.

Position: As the following layers are destructive (i.e. calorimeters will absorb the particle entirely in order to make an energy measurement, while a muon system will block any other particle), we need to place the tracking system first for it to be effective.

We also want it placed near the primary interaction point. Since we want to use position measurements in the tracking system to extrapolate backwards and determine the origin point (where the collision took place), keeping them as close as possible to this origin point reduces the overall uncertainty of the track.

For a long path, a small uncertainty in position points propagates to a much larger uncertainty in track the further away we are from the initial collision point.

1.1 Material Budget

- In an ideal world, the tracking detector would have a perfectly massless and lightweight material, in order to reduce the risk of scattering. However, we need some mass in order to measure ionisation, so it's a constant trade off between the two.
- We want as small a number of radiation lengths X_0 s within / upstream of the tracking system as possible.
- Some material is unavoidable, for example in the LHC there needs to be a conductive shield around the beam to prevent the large magnetic fields from inducing currents that would interfere with the sensitive measurement equipment. It also separates the highly pure vacuum of the beam pipe from the slightly less pure vacuum of the outer portion containing the LHCs electronics.

Radiation Length This is an inherent property of each material and is a measure of energy loss. A particle of energy E_0 passes through a distance of one radiation length and loses a factor of energy $1/e$.

$$E(x) = E_0 \exp\left(\frac{-x}{X_0}\right)$$

For example:

- For Cu: $X_0 = 15\text{mm}$.
- For Be: $X_0 = 35.2\text{cm}$.

The processes by which energy is lost depends on the particle species in question. For example, an electron loses energy by Bremsstrahlung much more rapidly compared to a muon, due to the differences in mass (average energy loss $\propto 1/m^2$). To define radiation length, we use an electron as a scale.

This lets us talk about “material budget”, as adding more material causes a higher uncertainty. For each particle path we care about, we want the fewest radiation lengths possible.

2 Measuring Trajectories

For a particle passing through a tracking system, we reconstruct its trajectory by measuring individual energy deposits (called “hits”) caused by ionisation in a large volume (typically $1\text{m} \times 1\text{m} \times 1\text{m}$)

This volume is made up often of many layers, so we can gain an idea of the particles position as it passes through each layer and use this to extrapolate. A tracking system has a resolution of a few $100\mu\text{m}$ and may leave $10 - 100$ hits.

In a “vertex detector”, we use a smaller system to reconstruct tracks at/around the interaction point with a higher precision - typically aiming for $\sim 10\mu\text{m}$. It is typically a silicon detector (layers of silicon detector sheets) and generally records fewer hits (< 10).

In a “tracking detector” we use a much larger system commonly further away from the primary interaction point. In a vertex system, there's no magnetic field so while we can use it to extrapolate back to find the PIP, we cannot use it to estimate momentum.

The LHC-b experiment for example has both a vertexer very close to the PIP (to determine the PIP location), and a much more substantial tracking system behind a magnet further away (to determine the deflected track in a magnetic field and hence the momentum.)

3 Measuring Momentum

The motion of a charged particle in a magnetic arises due to the Lorentz force and is proportional to charge:

$$\underline{F} = q(\underline{E} + \underline{v} \times \underline{B})$$

Where $\underline{F}$ is the force, $\underline{E}$ is the electric field, $\underline{v}$ is the particle velocity and $\underline{B}$ is the magnetic field. Assuming we are only dealing with a magnetic field, and taking magnitudes, we have:

$$\frac{p}{\text{GeV}} = 0.3 \times \frac{B}{\text{T}} \times \frac{q}{|e|} \times \frac{r}{\text{m}}$$

Where the magnetic field produces curvature of radius r in the plane perpendicular to the B -field. Motion parallel to $\underline{B}$ is unchanged.

In 3D, and assuming we are contained within the magnetic field, the particle will follow a helix of constant radius of curvature. This assumes that there are no non-conservative forces (i.e. scattering). In the ideal case, there is no work done, meaning in our force:

$$\underline{F} = q\underline{v} \times \underline{B}$$

The force and the velocity are perpendicular, so $\underline{F} \cdot \underline{v} = 0$.

3.1 Quantitatively

Consider a particle path with a constant radius of curvature R . Consider three hits, with vertical separation between the top and the bottom being $L/2$. The distance between the top/bottom hit and the origin is l , and the final distance between this straight line and the deflected path is the “sagitta” s .




Figure 4.2

We know $L/2$ (the vertical separation between hits) as we've build the detector so know the resolution, and we want to measure the sagitta, the deviation from a straight line.

We know that:

$$p = 0.3BqR$$

And from the diagram:

$$R = l + s$$

$$l = R \cos\left(\frac{\theta}{2}\right)$$

Putting these together:

$$s = R\left(1 - \cos\frac{\theta}{2}\right)$$

Assuming the bending is gentle in a detector, so θ and s are small. Hence we apply a small angle approximation:

$$\cos\frac{\theta}{2} \approx 1 - \left(\frac{\theta}{2}\right)^2 \frac{1}{2!} + \left(\frac{\theta}{2}\right)^4 \frac{1}{4!} + \dots$$

Taking the first order:

$$s \approx \left(\left(\frac{\theta}{2} \right)^2 \frac{1}{2!} \right) = \frac{R\theta^2}{8}$$

We can also use the small angle approximation for sine:

$$\sin \frac{\theta}{2} = \frac{L}{2R}$$

For small theta:

$$\sin \frac{\theta}{2} \approx \frac{\theta}{2}$$

$$\theta \approx \frac{L}{R}$$

So:

$$s = \frac{L^2}{8R}$$

Hence, finally, we have:

$$p = 0.3Bq \frac{L^2}{8s}$$

3.2 Uncertainties

We want to find the uncertainty on momentum, σ_p/p :

$$\frac{\sigma_p}{p} = \frac{\sigma_s}{s} = \frac{8p}{0.3BqL^2} \sigma_s$$

However we want to use our uncertainties on individual hits, x_1, x_2, x_3 etc. We have (and will not derive):

$$\frac{\sigma_p}{p} = \frac{\sigma_{xy} P}{0.3BL^2} \sqrt{\frac{720}{N+4}}$$

Mon 09 Feb 2026 12:00

Lecture 5 - Particle Physics V: Calorimetry

Now that we've finished talking about determining the principal interaction point and the momentum of produced particles in a tracking detector, we move onto calorimeters:




Figure 5.1

1 Materials and Energy Loss

Energy loss in a material is described by the Bethe formula, which gives mean energy loss by ionisation for unit path length:

$$\left\langle -\frac{dE}{dx} \right\rangle = K Z^2 \frac{1}{\beta^2} \left[\frac{1}{2} \ln \frac{2m_e c^2 \beta^2 \gamma^2 W_{\max}}{I^2} - \beta^2 - \frac{\delta(\beta\gamma)}{2} \right]$$

This is:

- Valid for $\beta\gamma < 1000$ within a few percent precision.
- Includes dependence on the medium, I, Z, A etc.

The good news is that we do not need to know this formula. We do however need to know some key features:

- $\frac{dE}{dx} \propto \frac{1}{\beta^2}$ below some minimal value of $\frac{dE}{dx}$
- $\frac{dE}{dx} \propto \ln(\beta^2 \gamma^2)$ above this minimal value of $\frac{dE}{dx}$. This is called “relativistic rise”.
- This minima happens at approximately $\beta\gamma \sim 3 - 4$.

- At large $\beta\gamma$, polarisation of the medium causes saturation (effectively a plateau).

Broadly, we expect to see a fall up until some minimal value, then a relativistic rise, and then a plateau:




Figure 5.2

2 Calorimeters

The purpose of a calorimeter is to determine the total energy of an incident particle. It does so by absorbing the particle entirely, creating a measurable signal which is proportional to the energy of the incident particle. As it is destructive, it is always located after the non-destructive tracking system. It has no need for the B-field of the tracking detector, and may be called a “calo”, “ECAL” or “HCAL”.

There are two types of calorimeter:

- Electromagnetic Calorimeter: ECAL.
- Hadronic Calorimeter: HCAL.

We characterise the length of an ECAL in terms of radiation length X_0 , typically $15X_0 - 30X_0$. It works by bremsstrahlung, and emitted bremsstrahlung photons causing pair production and more bremsstrahlung emission etc.

A HCAL works on nuclear hadronic interactions, so we instead define an interaction length λ_{int} , typically $5\lambda_{\text{int}} - 8\lambda_{\text{int}}$.

They don't measure the total energy of a particle if particles produced in the process pass straight through, so not all the energy in an ECAL will be captured, as some will pass straight through. Muons, neutrinos and sometimes pions pass straight through and escape, they are not (may not for a pion) be detected by the calorimeters.

An ideal calorimeter has an output signal “response” which is proportional to the energy of the input particle. We would ideally like a directly proportional linear relationship, but this may not always be realistic.

If we already know mass (from Cherenkov rings, to be discussed later in the course) and momentum from the tracking detector, why do we need a calo when we can just use $E^2 - p^2 = m^2$ to determine energy?

- Cherenkov rings and tracking systems only work with a charged particle. The calculation method is not sensitive enough, as the Cherenkov+tracker setup cannot detect γ , ν or neutral hadrons.
- Having a direct energy measurement adds another constraint to our system - the more information we can get the better. A direct energy measurement improves resolution and is more likely to be accurate than determining it from two other uncertain quantities.
- At very high momentum, (with little Coulomb scattering) the relative uncertainty of the transverse momentum of a tracking system is given by:

$$\frac{\sigma_{p_{\perp}}}{p_{\perp}} \propto \frac{p_{\perp}}{BL^2}$$

Therefore at a higher momentum, we have a higher uncertainty. At high momenta, there is a smaller deflection, so a high momentum track will just pass rapidly through the field with very little deflection and is therefore much more difficult to measure accurately - the tracking system degrades at higher momenta.

For a calorimeter, we have nicer behaviour at high momenta:

$$\frac{\sigma_E}{E} = \frac{a}{\sqrt{E}} \oplus b \oplus \frac{c}{E}$$

Where:

- a arises statistical fluctuations inherent to the measurement.
- b arises from calibration effects (i.e. from non-linearity)
- c arises from electronic noise.
- $\oplus$ means to “add in quadrature”, $(p \oplus q \oplus r) = \sqrt{p^2 + q^2 + r^2}$
- Trackers are relatively slow as they require ionisation clouds drifting towards a collection point and involve quite a computationally complex problem to recognise the patterns and reconstruct the track. A calorimeter relies on “scintillation” to convert energy to a response, which is much faster.

This is important for triggering. Every 25ns, we need to decide whether or not to keep an event (as there is nowhere near enough storage/processing power) to perform detailed track reconstruction for every single event. We need a calorimeter to participate in this.

2.1 EM Calorimeters

There are two types of electromagnetic calorimeters, sampling or homogeneous. A sampling calorimeter is layered, with a layer of absorber producing a shower of particles (i.e. lead) in front of a collector layer with a scintillator that generates a signal when particles pass through.

HCALs are always sampling-type calorimeters, as it takes a lot more mass to slow down/break apart a hadron compared to a lighter charged particle.

Lets build a simple model of an electromagnetic shower that might be detected by a calorimeter:

We have a high energy photon (so we can mostly disregard scattering etc) enter the detector. This pair produces (i.e. an electron and a positron) which emits photons via bremsstrahlung emission. These emitted photons can then go on to pair produce themselves, generating (i.e.) an electron and a positron again, which emit more photons via bremsstrahlung, etc.

The maximum energy is deposited when the average particle energy (of particles developing in the shower) is the “critical energy”.

Here, a photon has entered from the left, and the first few radiation lengths have left the material intact.




Figure 5.3: Damage to a copper block that has had energy dumped into it via this process.

After the first few radiation lengths, we have the “critical point”, where a large amount of energy has been dumped. This point is defined as the point where the probability of bremsstrahlung and ionisation are equal.

The shower deposits a constant amount of energy per X_0 travelled (by definition of radiation length). Each step in path length, in terms of X_0 has a doubling number of particles and a halving individual particle energy. The angle of photon emission is fairly small, so the shower is narrow. This is unlike a hadronic calorimeter that has a wider shower. The width of the shower is given by the multiple scattering of the e^+ and e^- .

When the energy of the particle is less than the critical energy, ionisation dominates the interactions and the shower development stops rapidly. When the particle energy is $> E_c$, pair production and bremsstrahlung dominate.

After depth t (in units of X_0), the number of particles is 2^t , and the average particle energy is $E_0/2^t$.

The shower stops developing after $E = \frac{E_0}{2^t} < E_c$. This happens when $E = 2^t = E_0/E_c$. Hence:

$$t_{\max} \log 2 = \log \left(\frac{E_0}{E_c} \right)$$

$$t_{\max} = \frac{\log (E_0/E_c)}{\log 2}$$

This is what defines how deep our calorimeter needs to be in order to collect the energies we are expecting to require.

Mon 16 Feb 2026 16:00

Lecture 6 - Particle Physics VI: Calorimetry II and PID

1 EM Calorimetry: Shower Shape

The energy deposited vs the depth in material can be expressed as:

$$\frac{dE}{dt} = E_0 b \frac{(bt)^{a-1} e^{-bt}}{\Gamma(a)}$$

Where t is expressed as a length in units of radiation length $t = x/X_0$ and $\Gamma(a)$ is the standard gamma function $\Gamma(x) = \int_0^\infty t^{x-1} e^{-t} dt$. a, b are both constants.

For small t (early in the detector): t^{a-1} dominates as $e^{-bt} \approx 1$

For large t (later in the detector): e^{-bt} dominates.

Considering a plot of $\frac{dE}{dt}$ against t , we initially have polynomial growth, reaching a peak. We then from that peak decay exponentially with an infinite tail.

The maximum point of the shower is given at $t = t_{\max}$ and is achieved by further differentiating dE/dt and is given by:

$$t_{\max} = \frac{a-1}{b} = \ln\left(\frac{E_0}{E_c}\right) + C_{\gamma \text{ or } e^-}$$

For a photon, $C_\gamma = 0.5$ and for an electron, $C_e = -1$. Hence, the shower maximum is at a smaller value of t for an electron/positron compared to photon.

Crucially, we want to find the total energy deposited by the particle. We get that by integrating dE/dt :

$$\int_0^{\text{calo thickness}} \frac{dE}{dt} dt$$

While the tail is an infinitely long decaying exponential, we only care about the portion of the tail which is inside the calorimeter. Any energy that would be deposited past this point is lost:




Figure 6.1

If the calorimeter is too thin, we need to make a large correction to adjust for this and we therefore have a bigger uncertainty. We want to ensure that the calorimeter is thick enough to capture the vast majority of deposited energy with minimal loss.

1.1 Resolution

The resolution for the calorimeter varies heavily based on the material used to construct it. Typically, however:

- ECAL resolution: 2 – 10%
- HCAL resolution: 50 – 100%

Different materials will have different characteristic radiation lengths,

For example:

- W has $X_0 = 0.35\text{cm}$, $\lambda_{\text{int}} = 9.9\text{cm}$, $R_M = 0.93\text{cm}$
- Pb has $X_0 = 0.56\text{cm}$, $\lambda_{\text{int}} = 17.6\text{cm}$, $R_M = 1.6\text{cm}$
- Fe has $X_0 = 1.8\text{cm}$, $\lambda_{\text{int}} = 16.7\text{cm}$, $R_M = 1.7\text{cm}$
- Cu has $X_0 = 1.4\text{cm}$, $\lambda_{\text{int}} = 15.3\text{cm}$, $R_M = 1.8\text{cm}$

R_M is the “Moliere Radius” and is the radial width in which 90% of the shower’s energy deposit is confined. Note that we have different design considerations for an HCAL vs an ECAL, for example the size of an HCAL is generally much larger than that of an ECAL.

2 Particle Identification (PID)

We can now measure a particle’s momentum and energy, and we now want to determine the mass of the particle. Surely we can do that with:

$$E^2 - p^2 = m^2$$

Unfortunately not...the resolution (especially for energy on a calo) is pretty poor, so we would end up with an unreasonably large uncertainty on our mass, too large to do reasonable particle identification with. Instead, we:

- Use information from the whole ensemble of detectors.
- Use hypothesis tests: proposing a specific particle type and then using all the information we have (i.e. momentum) see if this is a reasonably likely outcome.
- For example, we can use information on whether or not the particle reached the muon chambers to help narrow down what it could be. Alternatively, what is the pattern of energy deposit observed in the ECAL or HCAL?
- We can also use dedicated PID, such as:
 - A time of flight detector, determining the particle velocity (hence mass, with momentum measurements from a tracker).
 - Cherenkov detectors (not useful at high momenta, i.e $>100\text{GeV}$): As a recap, the particle emits a cone of photons at angle θ_c which depends on the refractive index of the material and β . Measuring this cone allows us to determine θ_c hence β hence v .

Here is an example of a detector setup:




Figure 6.2

We have covered the vertex locator, magnet, tracking stations T1, T2, T3, the HCAL, ECAL and (while we haven't yet touched on them), we have 5 muon systems.

We are yet to discuss RICH1 and RICH2. These are called "Ring Imaging Cherenkov Detectors". These cones of Cherenkov emissions are focussed using a magnet onto a plane, where they form rings. The radii of these rings can be measured and the detector geometry can be used to determine the Cherenkov angle. The two detectors work similarly, but cover different ranges of measurable momenta.




Figure 6.3

This Cherenkov angle can therefore be used to discriminate against particles, up to a limit of momentum, past which they become fairly useless.

NOTE: There was a practice exam question here that would be good to come back to during revision.

End of Particle Physics

Mon 23 Feb 2026 12:00

Lecture 7 - Cosmology I: A Brief Introduction to Cosmology

Primary Textbook: “An Introduction to Modern Cosmology”, Andrew Liddle. Each lecture will have a chapter or two as recommended reading from this book and is available as an ePDF from the library.

1 The origin of Cosmology

Comes from the Ancient Greek word *κοσμος*, “kosmos” meaning order. Cosmology is a branch of astronomy concerned with studying the origin and evolution of the universe. We won’t really consider anything on a smaller scale than clusters of galaxies.

The earliest form of this is *celestial mechanics* which goes back to Ancient Greek philosophers such as Aristotle and Ptolemy.

1.1 Geocentric Ptolemaic System

This was the prevailing theory until the 16th century where the earth was at the centre of the universe and other stars/planet etc revolved around the earth in “epicycles”. Here, the planets sat on rotating spheres, which orbited the earth.




Figure 7.1

The epicycles explain retrograde motion. Observational power was too limited to get more accurate measurements, and they noticed that some objects appeared to go backwards as they were moving forwards overall in the orbit. We can solve this by placing another smaller circular orbit superimposed on the larger orbit.

We now know that this is not accurate, but it explained the motion at the time.

1.2 Heliocentric System

Copernicus in 1543 radically changed the understanding of orbits. He suggested that the sun, not the earth, was near the centre of the universe. The planets, including the earth revolved in circles around the sun, with other stars orbiting much further away.

In order to match the observed motion, we still need epicycles in this initial version of the theory.

This was the foundation of modern astronomy, and lead to:

- Kepler discovering elliptical orbits. This got rid of the need for epicentres and described the motion of planets with the data collected at the time.
- Galilei discovering the moons of Jupiter and phases of Venus. Before Galilei, these telescopes were not sufficiently powerful to observe these phases. The existence of these moons provided additional evidence for the heliocentric system as we now had a celestial body which did not orbit the Earth. This suggested that the Earth wasn't special, which was one of the beliefs that drove the geocentric model.
- Newton's theory of universal gravity in 1678.

This theory, and the implications it had on humanity's role in the universe was deeply unpopular with religious leaders and this caused opposition to its widespread acceptance.

This was a better way to think about the solar system (which was about the limit of what was practically observable). Copernicus therefore needed to provide evidence to justify this. These later from Kepler, Galilei and Newton acted as this evidence.

2 From Herschel to Hubble

In order to go from scales on a solar system level to a cosmological level, we need more powerful measurement equipment to be able to gather as much data as possible on much larger scales. William and Caroline Herschel pioneered this by creating the most powerful telescopes to date in 1877.

They probed deeper and deeper past the solar system, into the "deep sky". They investigated the contents of the Milky Way to build a catalogue of objects and try to find the edge of the galaxy.

This was in an attempt to work out whether or not there was some kind of edge to the universe. They discovered diffuse objects ("nebulae") beyond the milky way, and began to be able to resolve individual stars with improved technology.

This lead to the question of whether these nebulae (named "island galaxies" at the time) belong as components of the milky way or existed in their own right as galaxies.

2.1 Shapley-Curtis Debate

In the 1920s, the "Great Debate of Astronomy" investigated whether these spiral shape nebulae were small object on the outskirts of the Milky Way (advocated for by Harlow Shapley) or were independent small galaxies outside the gravitational attraction of the Milky Way (advocated for by Heber Curtis).

In 1924, Edwin Hubble established that these bright nebulae were not part of the Milky Way and were instead of extragalactic origin. This told us that the observable universe extended beyond the Milky Way.

He discovered Cepheid Variable Stars, which are stars of periodic varying brightness. The period of pulsation is linked to intrinsic luminosity, and they act as distance indicators based on the perceived brightness and period on Earth.

These observations estimated the distance of these CVSes away from the Earth, and proved that they were too distant to sit within the Milky Way. This profoundly changed our understanding of the scale of brightness.

These CVSes are called "standard candles" which are astronomical objects with known intrinsic luminosity, M . We observe some different apparent brightness m , and can measure the pulsation period to determine the actual intrinsic luminosity.

Using M (determined from period) and m (measured) directly, we can use this formula:

$$5 \log D = m - M - 10$$

$$\implies D = 10^{(m-M-10)/5}$$

This made it apparent that these distances were much larger than our understanding of the size of the Milky Way, by in excess of 2 orders of magnitude.

We can also know the intrinsic luminosity of a Type 1a supernovae, when a star explodes and collapses into a white dwarf.

2.2 Cosmic Ladder

We can form the “Cosmic Distance Ladder”. Since traditional distance measurements become infeasible as scales get larger, we need to use different ranges for different distances. For example:

- Cepheid Variable Stars are not that bright, so stop being useful at longer ranges.
- Supernovae are brighter, so stay useful for longer.




Figure 7.2: The Cosmic Distance Ladder

3 Expansion of the Universe

In 1929, Hubble analysed the emission spectra of galaxies. As a source moves, wavelengths (hence the positions of spectral lines) are impacted by redshift:

$$z = \frac{\lambda_o - \lambda_e}{\lambda_e} \approx \frac{v}{c} \quad \text{approximation valid for non-relativistic limits.}$$

From this, noting the changes of the spectral lines vs known non-redshifted calibration points on earth allowed for the velocity of each galaxy to be determined.

Plotting v against r yields a straight line, suggesting $v \propto r$. Hence galaxies further away are moving faster. The sign of the redshift (i.e. is it indeed redshift and not blueshift) confirms that they are moving away from us.

This is very strong evidence that the universe is not static and is actually expanding. This gives us a dramatically new understanding of the universe, where it grows in size as time goes on.

4 Key Observations

We have a number of key objectives, and want to build a theory that fits these and makes predictions. Most of these observations are made using electromagnetic radiation, although additional information is provided by particle physics (and more recently gravitational wave astronomy).

Some of these key observations are:

- The universe is expanding (Hubble).
- There is an abundance of light elements (nucleosynthesis).
- Large-scale structure (LSS).
- Cosmic microwave background (CMB).
- Galaxy rotation curves (implying the existence of dark matter).
- Acceleration of the rate of expansion of the universe (dark energy).

4.1 Nucleosynthesis

Stars are very good at producing heavy elements via nuclear fusion. Here, stars start with lighter elements (H, He) and fuse upwards through increasing mass numbers up to Iron. Heavier elements require a stellar explosion to form.

It turns out that young stars have an abundance of light elements. These start from gas clouds that have collapsed gravitationally. In this very early stage, they started out with some contamination of heavier lighter elements (Helium isotopes and other Hydrogen isotopes) and couldn't just start with Hydrogen. They need some of these other elements to kickstart the process.

There has to be some process that took place before stars were formed that provided these elements to kickstart star formation. Stars could not have formed without this.

Theorists (George Gamov 1948) hypothesised reversing the expanding universe, i.e. stepping back in time to a much smaller, denser, hotter early universe. This would have formed a plasma-like state which could have provided these elements.

Nuclear fusion requires a very large energy, i.e. nuclear binding energy for these interactions is $\sim 1\text{MeV}$. This plasma state lasted about 3 minutes, and allows for the formation up to Hydrogen-4. Heavier elements are still formed in stars.

4.2 Large Scale Structure

The fundamental building block of the universe is galaxies. These cluster themselves together into filaments. While on a local scale, we see very large gaps between these, on a large enough scale (i.e. zoomed out enough) we see that the universe is very smooth on average. We say that, on cosmological scales, the distribution of matter is homogeneous and isotropic (has identical physical properties in all directions with uniform structure).

4.3 Cosmic Microwave Background

There is a nearly isotropic low frequency radiation background that penetrates the entire universe. It has an almost perfect black body spectrum with $T \sim 2.73\text{K}$. This is in the microwave region.

4.4 Discovery of Dark Matter

Fritz Zwicky and Vera Rubin discovered strange velocities in galaxy clusters and in the rotational curves of galaxies. This suggests that there is some extra mass that only interacts via gravity.

This is also potentially (dark energy) a driving force for the increasing rate of the expansion of the universe. It implies that we may need an extra particle added to the standard model to explain what actually makes this up.

Thu 26 Feb 2026 14:00

Lecture 8 - Cosmology II: Expansion of the Universe and Key Principles

1 Expansion of the Universe

Data from supernovae found that the universe was not expanding at a constant rate. The rate of expansion is increasing and accelerating.

The best current model we have is the Hot Big Bang Model:




Figure 8.1

This is where the universe was created from an initial singularity, resulting in the big bang - an explosion.

Space itself is expanding and the universe is shaped by gravity as a fundamental force. The fundamental particles were created first in a dense and hot universe. As the universe expanded, it cooled down allowing for the propagation of CMB photons and the creation of meaningful cosmic structure.

1.1 Big Bang Cosmology

For the rest of the course we will:

- Explore big-bang centric cosmology, in the context of Newtonian gravity.

- We'll build and discuss the main equations that model the evolution of the universe. Using these we'll explore:
 - The history of the universe.
 - The composition of the universe.
 - The eventual fate of the universe.
- We'll discuss key observations and any problems where these do not fit the models we're applying them to.

2 Key Principles

2.1 The Cosmological Principle

The distribution of matter is:

- **Homogeneous**, meaning that the universe looks the same at every point. There is no special point in the universe, and the average matter density is the same everywhere.
- **Isotropic**, meaning the universe looks the same in all directions

However this isn't perfectly fulfilled in our universe. It's a valid approximation on sufficiently large scales $> 100\text{Mpc}$, but it isn't perfect.

2.2 Distance Measures - The Parsec

The parsec is a unit of distance useful on cosmological scales:

$$1\text{pc} = 3.086 \times 10^{16}\text{m}$$

Suppose an observer on Earth measures a very distant star. We take two measurements six months apart. As the Earth is moving relative to the star, it appears to us that the star has shifted position (in the Earth's rest frame). The total observed angular shift is $2p$, where p is the parallax angle — the angle at the star in the right triangle formed by the Sun, Earth, and star, with 1 AU as the opposite side.

The parsec is defined as the distance at which $p = 1$ arcsecond (4.848×10^{-6} rad), giving the relationship:

$$d [\text{pc}] = \frac{1}{p [\text{arcsec}]}$$

2.3 Redshift

This has already been covered in other modules, but to recap the fractional shift is:

$$z = \frac{\lambda_0 - \lambda_e}{\lambda_e} \approx \frac{v}{c}$$

for non-relativistic sources where $v \ll c$.

3 Expansion of the Universe

Hubble found that there was a linear relationship between the recessional velocity of a galaxy and its distance.

$$v = H_0 r$$

Where $H \approx 68\text{kms}^{-1}\text{Mpc}^{-1}$. Note that the parameter H varies with time. H_0 represents the value at the current epoch.

This isn't exact, and does not hold for nearby galaxies where other forces dominate over the expansion of the universe. These galaxies have some peculiar velocity which forms the total velocity:

$$v_{\text{total}} = v_{\text{pec}} + v_H$$

For distant galaxies, the recessional velocity dominates. This is called Hubble flow.

3.1 Comoving Coordinates

As the universe expands, the physical (proper) distance between two galaxies grows. This means that, in a coordinate system centred on the earth, the coordinates of other galaxies are increasing.

Instead, we define a new coordinate system where the coordinate system itself expands alongside the universe. This means that galaxies remain at fixed coordinates as the universe expands, and the coordinate grid expands at the same rate as the universe. The distance between galaxies in these coordinates is therefore constant.

This rate of increase is given by the universal expansion rate (a scale factor) $a(t)$. Due to homogeneity, this scale factor is only time-dependant, as the rate of expansion is the same everywhere in the universe.

This means that we can relate proper distance $r(t)$ and the fixed comoving distance x with:

$$r(t) = a(t)x$$

Mon 02 Mar 2026 12:00

Lecture 9 - Cosmology III: Friedmann Equation

1 Equation of Motion for the Universe

Goal: To derive an equation of motion for the dynamics of the Universe.

Assumptions:

- We will assume Newtonian Gravity.
- We will assume The Cosmological Principle holds on the scales we're considering.
- We will assume uniform expansion.
- We will assume no dissipation. This means that total energy for a particle of mass m will be conserved.
- As a result of these assumptions, we will be using comoving coordinates.

We'll be using a non-vectorised form of the comoving coordinates:

$$r(t) = a(t)x$$

Assume at some time t_1 we have some spherical volume of radius $r(t_1)$, containing mass M . As time progresses and the universe expands, this radius will increase. This is given by some generic $r(t)$ and volume $V(t)$

As a result of our assumptions, we assume that we have uniform mass density given by $\rho(t)$ for some t :

$$\rho(t) = \frac{M}{V(t)}$$

Consider a particle of mass m on the boundary of our initial t_1 universe. This particle will have some gravitational potential energy, and some kinetic energy as it moves on expansion:

$$\text{total energy} = \text{potential energy} + \text{kinetic energy}$$

$$U = V + T$$

For potential energy:

$$V = \frac{-GMm}{r(t)} = -\frac{4\pi}{3}Gm\rho(t)r^2(t) \quad \text{assuming a spherical universe.}$$

For kinetic energy:

$$T = \frac{1}{2}mv^2 = \frac{1}{2}m\left(\frac{dr(t)}{dt}\right)^2 = \frac{1}{2}m\dot{r}^2$$

Hence for total energy:

$$U = T + V = \frac{1}{2}m\dot{r}^2 - \frac{4\pi}{3}Gm\rho(t)r^2(t)$$

In doing this, we've had to implicitly assume that we can treat the mass M distributed throughout the sphere as if it were a point mass at the centre of the sphere. This gives us an ordinary differential equation in terms of $r(t)$.

1.1 Comoving Coordinates

We now want to transform this into comoving coordinates.

$$r(t) = \frac{d}{dt}(a(t)x) = \dot{a}(t)x$$

$$\implies U = \frac{-4\pi}{3}Gm\rho(t)a^2(t)x^2 + \frac{1}{2}m\dot{a}^2(t)x^2$$

$$\frac{2U}{ma^2(t)x^2} = \left(\frac{\dot{a}(t)}{a(t)}\right)^2 - \frac{8\pi G}{3}\rho(t)$$

Let $\frac{2U}{mx^2} \equiv -kc^2$, hence¹:

$$\boxed{\left(\frac{\dot{a}(t)}{a(t)}\right)^2 = \frac{8\pi G}{3}\rho(t) - \frac{kc^2}{a(t)^2}}$$

This is called the Friedmann Equation. This is now an ODE in terms of scale factor and not in terms of radius. Before we can solve it however, we need to deal with the $\rho(t)$ term.

Considering the second term, we have:

$$[k] = \frac{1}{L^2}$$

This k has different implications on the geometry of the universe, and we consider 3 cases:

- $k = 0$: A flat universe.
- $k > 0$: A positive-curvature (closed) universe with spherical geometry, analogous to the surface of a sphere.
- $k < 0$: A negative-curvature (open) universe with hyperbolic geometry, analogous to the surface of a saddle.

Current observations suggest that our universe is very close to the $k = 0$ case, so we will disregard the second term in first year cosmology.

1.2 The Fluid Equation

In order to solve the Friedmann Equation, we need to build something to describe $\rho(t)$, called the “fluid equation”. We will assume the following:

- We will assume that the universe is a perfect/ideal fluid. This means that it is characterised by a density ρ and an isotropic pressure p .
- We will assume that the 1st Law of Thermodynamics holds:

$$dE = -pdV + TdS$$

- We will assume that the process is adiabatic, $dS = 0$.

Considering a comoving unit volume, where:

$$x = 1 \implies r(t) = a(t)$$

Hence:

$$V(t) = \frac{4\pi}{3}a^3(t)$$

The internal energy of our fluid is given by:

$$E = mc^2 = \frac{4\pi}{3}\rho(t)a^3(t)c^2$$

¹This seems very random, but all we are doing is setting a constant equal to another constant, so it does work!

$$\begin{aligned}\frac{dE}{dt} &= \frac{4\pi}{3}c^2 \frac{d}{dt} (\rho(t)a^3(t)) \\ &= \frac{4\pi c^2}{3} (\dot{\rho}(t)a^3(t) + \rho(t)3a^2(t)\dot{a}(t))\end{aligned}$$

And:

$$V(t) = \frac{4\pi}{3}a^3(t) \implies \frac{dV}{dt} = 4\pi a^2(t)\dot{a}(t)$$

Hence:

$$\begin{aligned}\frac{4\pi c^2}{3} (\dot{\rho}(t)a^3(t) + \rho(t)3a^2(t)\dot{a}(t)) + p4\pi a^2(t)\dot{a}(t) &= 0 \\ \dot{\rho}(t) + 3\frac{\dot{a}(t)}{a(t)} \left(\rho(t) + \frac{p}{c^2} \right) &= 0\end{aligned}$$

This is now much easier to solve!

Thu 05 Mar 2026 14:00

Lecture 10 - Cosmology IV: Friedmann Equation, Hubble and Redshift Combined

We have the Friedmann Equation from last lecture:

$$\left(\frac{\dot{a}(t)}{a(t)}\right)^2 = \frac{8\pi G}{3}\rho(t) - \frac{kc^2}{a(t)^2}$$

Where $a(t)$ is the scale factor relating the comoving scaled coordinates to the initial coordinate system. As observations suggest our universe has little to no curvature, hence $k \approx 0$, we will generally ignore the second term at this level.

We also assume:

- Newtonian Gravity
- Conservation of Energy
- The Cosmological Principle

We also have the fluid equation:

$$\dot{\rho}(t) + 3\frac{\dot{a}(t)}{a(t)}\left(\rho(t) + \frac{p(t)}{c^2}\right) = 0$$

As the universe is not empty, we need to describe how the matter density changes with time. This is the fluid equation. We also need an equation of state, to link the density and the pressure of the fluid.

Depending on the type of material being considered, this may give multiple different equations of state. These are the equations for a matter-dominated (non-relativistic) and a radiation dominated universe (relativistic).

For a matter dominated universe, we have:

$$p = 0$$

For a radiation dominated universe, we have:

$$p = \frac{\rho c^2}{3}$$

1 Acceleration Equation

We want some kind of equation to describe the acceleration of the scale factor, i.e $\ddot{a}(t)$.

We can get this through a combination of:

1. The Friedmann Equation → differentiate it with respect to time.
2. Simplify with the fluid equation.
3. Use the Friedmann equation again to get the acceleration equation in the form:

$$\frac{\ddot{a}(t)}{a(t)} = \frac{-4\pi G}{3}\left(\rho(t) + \frac{3p}{c^2}\right)$$

Note the sign here, if the contents of the brackets are positive, then there's an overall global negative sign, which suggests that the whole expression is negative. This suggests that the rate of universal expansion is slowing... this contradicts observations.

For now, we accept this, but we'll make a modification to the Friedmann equation to correct for this next week.

1.1 Hubble and Friedmann

How are Hubble's observation and Friedmann's equation connected?

We have the Friedmann eqn:

$$\left(\frac{\dot{a}(t)}{a(t)}\right)^2 = \frac{8\pi G}{3}\rho(t) - \frac{kc^2}{a(t)^2}$$

And Hubble's observation:

$$\mathbf{v}_h = H \cdot \mathbf{r}$$

Where $\mathbf{v}_h$ represents the radial recessional velocity only, excluding peculiar motion. But we want to find a form of the Hubble observation which includes a term in a . We consider the recessional velocity, starting from our definition of comoving coordinates:

$$\begin{aligned}\mathbf{r}(t) &= a(t)\mathbf{x} \\ \mathbf{v}_H &= \frac{d\mathbf{r}}{dt} = \dot{a}(t)\mathbf{x} = \dot{a}(t)\frac{\mathbf{r}(t)}{a(t)} = \frac{\dot{a}(t)}{a(t)}\mathbf{r}(t)\end{aligned}$$

And comparing with Hubble's law gives:

$$H(t) \cdot \mathbf{r} = \frac{\dot{a}(t)}{a(t)}\mathbf{r}$$

$$H(t) = \frac{\dot{a}(t)}{a(t)}$$

While commonly called the Hubble Constant, because it's constant wrt space, it's a function of time which means it's not a constant and is properly referred to as the Hubble Parameter. H_0 is $H(t = t_0)$ where t_0 denotes right now.

2 Expansion and Redshift

Consider two galaxies separated by a (cosmologically) small distance dr . A photon is emitted at one galaxy with λ_{em} and travels towards an observer on the second galaxy.

According to Hubble's law, there will be a small recessional velocity between the two (disregarding peculiar velocity):

$$dv = H dr = \frac{\dot{a}(t)}{a(t)} dr$$

The very small change in wavelength will be:

$$d\lambda = \lambda_{\text{obs}} - \lambda_{\text{em}}$$

And in the non-relativistic limit:

$$\frac{d\lambda}{\lambda_{\text{em}}} \approx \frac{dv}{c}$$

The time dt for the photon to travel distance dr is:

$$dt = \frac{dr}{c}$$

Hence, substituting $dr = c dt$:

$$\frac{d\lambda}{\lambda_{\text{em}}} = \frac{dv}{c} = \frac{\dot{a}(t)}{a(t)} \frac{dr}{c} = \frac{\dot{a}(t)}{a(t)} dt = \frac{1}{a} \frac{da}{dt} dt$$

$$\frac{d\lambda}{\lambda_{\text{em}}} = \frac{da}{a}$$

This is a separable differential equation. Integrating both sides from emission to observation:

$$\begin{aligned} \int_{\lambda_{\text{em}}}^{\lambda_{\text{obs}}} \frac{d\lambda}{\lambda} &= \int_{a_{\text{em}}}^{a_{\text{obs}}} \frac{da}{a} \\ \ln\left(\frac{\lambda_{\text{obs}}}{\lambda_{\text{em}}}\right) &= \ln\left(\frac{a_{\text{obs}}}{a_{\text{em}}}\right) \\ \Rightarrow \boxed{\lambda \propto a} \end{aligned}$$

This means that, as the universe expands, the wavelength of these photons changes as well.

Going back to the definition of z :

$$1 + z = \frac{\lambda_{\text{obs}}}{\lambda_{\text{em}}} = \frac{a(t_{\text{obs}})}{a(t_{\text{em}})}$$

Effectively, photons have their wavelength stretched proportional to the scale factor of the universes growth between emission and observation. As the universe grows, wavelengths become stretched and energy decreases.

3 Simple Cosmological Models

In order to solve the Friedmann equation, we make assumptions about the contents and geometry of the universe, i.e. $\rho(t)$ and k .

For our universe, where $k \approx 0$, we assume $k = 0$ at a first year level as this is what is currently favoured by observations. We assume a flat universe.

We need to build an expression for ρ . Ideally, we'd use a mixture model for an environment which has both relativistic and non-relativistic particles. In practice, we'd use a model that combines these, however in order to get something we can easily solve analytically at our level, we assume that one is true.

For relativistic particles, we have (and will investigate in more depth in future lectures):

$$p_r(t) = \frac{\rho c^2}{3}$$

If we assume non-relativistic particles, we have:

$$p_m(t) = 0$$

In practice, we don't have to choose between these. A real model of a universe has both present, so we would need to consider a mixture module. This gives us two different contributions, one for matter and one for radiation, $\rho_r(t), \rho_m(t)$.

However, the broader we make our assumptions the more difficult analytical solutions become. We therefore will for now assume that only one of the two holds.

4 Model I: Flat, Matter-Dominated Universe

With $k = 0, p = 0$ the fluid equation now becomes:

$$\dot{\rho}(t) = \frac{3\dot{a}(t)}{a(t)}\rho(t) = 0$$

This becomes:

$$\frac{1}{a^3(t)} \frac{d}{dt} (\rho(t)a^3(t)) = 0$$

This is directly soluble, hence:

$$\frac{d}{dt} (\rho(t)a^3(t)) = 0$$

And hence:

$$\rho(t) = \frac{\text{constant}}{a^3(t)}$$

We need to use a boundary condition to solve for the constant. Generally, we define the scale factor at the current epoch as one to normalise.

$$a(t_0) = 1 = a_0$$

This gives us the density at the current epoch as:

$$\rho(t_0) = \frac{\rho_0}{a^3(t)}$$

This is a standard volumetric dilution, with the density decreasing as the universe expands. This is as-expected for standard matter particles.

den

Mon 09 Mar 2026 12:00

Lecture 11 - Cosmology V: Cosmological Models

1 Cosmological Models

1.1 Flat, Matter Only Universe Continued

We have the Friedmann equation:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho(t)$$

And using our equation of state for a matter only universe:

$$\begin{aligned} &= \frac{8\pi G}{3} \frac{\rho_0}{a^3} \\ \Rightarrow \dot{a}^2 &= \frac{8\pi G}{3} \rho_0 \frac{1}{a} \\ \Rightarrow \dot{a} &= \sqrt{\frac{8\pi G}{3} \rho_0} \sqrt{\frac{1}{a}} \end{aligned}$$

And hence to get to $a(t)$ we can separate variables:

$$\begin{aligned} \int \sqrt{a} da &= \sqrt{\frac{8\pi G}{3} \rho_0} \int dt \\ \frac{2}{3} a^{3/2} + C &= \sqrt{\frac{8\pi G \rho_0}{3}} t \end{aligned}$$

We need a boundary condition to determine the constant. We again say that the scale factor at $t = 0$ is zero.

$$a(t = 0) = 0 \Rightarrow C = 0$$

$$a(t) = (6\pi G \rho_0)^{1/3} t^{2/3}$$

And it is common to rewrite this using $a(t_0) \equiv a_0 = 1$. This gives us (by dividing through by a_0 , as dividing by one changes nothing):

$$a(t) = \left(\frac{t}{t_0}\right)^{2/3}$$

This causes power law growth for all t .

2 Flat, Radiation Only Universe

Now, we keep $k = 0$ but the equation of state is now $p = \rho c^2/3$. The fluid equation now gives us:

$$\begin{aligned} \dot{\rho} + \frac{3\dot{a}}{a} \left(\rho + \frac{\rho c^2}{3c^2} \right) &= 0 \\ \dot{\rho} + \frac{\dot{a}}{a} (4\rho) &= 0 \end{aligned}$$

This is almost directly equivalent to the matter only case, except the factor of 4:

$$\frac{1}{a^4} \frac{d}{dt} (\rho a^4) = 0$$

Hence:

$$\rho(t) = \frac{\rho_0}{a^4(t)}$$

This extra factor of $1/a$ causes an apparent loss of energy. This is due to redshift as the universe expands.

We can express ρ as energy per unit volume:

$$\rho = \frac{E}{V} = \frac{h\nu}{V} = \frac{hc}{V\lambda}$$

This gives us the same $1/a^3$ due to volume dilution, but also the extra $1/a$ due to the $1/\lambda$ term as the wavelength of photons becomes stretched out.

As before, we want to solve the Friedmann equation, so we substitute in our new ρ :

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} \frac{\rho_0}{a^4}$$

Following the same steps we get:

$$a(t) = \left(\frac{t}{t_0}\right)^{1/2}$$

This is still power law growth, but where the universe grows more slowly compared to the matter-only case.

3 Mixture Model

These two models are unrealistic for a true universe that contains both matter and radiation. We can form a more general $\rho(t)$ which is a sum of the two species.

We still assume a flat $k = 0$ universe. This gives us:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3} (\rho_m(t) + \rho_r(t))$$

We can do this by considering limits, i.e. what happens when:

- $r_m \ll r_r$
- $r_m \gg r_r$

These are universes that are dominated by one species or the other, but not solely.

3.1 Case I: $\rho_r \gg \rho_m$

Since $\rho(t) = \rho_m(t) + \rho_r(t)$, in this limit it approximates to just the radiation component. When this happens, $a \propto \sqrt{t}$. In this phase where radiation dominates, we have:

$$\rho_r(t) = \frac{\rho_{r,0}}{a^4(t)} \implies \rho_r(t) \propto \frac{1}{t^2}$$

And hence:

$$\rho_m(t) = \frac{\rho_{m,0}}{a^3(t)} \implies \rho_m(t) \propto \frac{1}{t^{3/2}}$$

3.2 Case II: $\rho_m \gg \rho_r$

In this limit, the matter component dominates, and we have $a \propto t^{2/3}$. In this phase where matter dominates, we have:

$$\rho_m(t) = \frac{\rho_{m,0}}{a^3(t)} \implies \rho_m(t) \propto \frac{1}{t^2}$$

And:

$$\rho_r(t) \propto \frac{1}{t^{8/3}}$$

3.3 Point of Equipartition

At some point in this evolution from a radiation to matter dominated world, there will be some crossover point where $\rho_m = \rho_r$. This is called the point of equipartition.

4 Curvature - Non-Flat Matter-Only Universe

We've already seen what happens with the $k = 0$ case of a flat universe. When $k < 0$ we have a hyperbolic geometry and $k > 0$ spherical geometry (and continue to assume that we are only dealing with a matter only universe)

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho_m(t) - \frac{kc^2}{a^2(t)}$$

Crucially, is the unbound nature of the expansion wrt time still true for these new geometries?

If $k < 0$ (Hyperbolic):

Here, the negative sign of k causes the final second term to be positive. This means that the expansion is still unbound. As:

$$H^2(t) = \left(\frac{\dot{a}}{a}\right)^2 > 0$$

This causes a larger magnitude of $H^2(t)$ hence the expansion is at a more rapid rate.

Thu 12 Mar 2026 14:00

Lecture 12 - Cosmology VI: Measurements, Universe Age and Deceleration

1 Hyperbolic Geometry Cont.

At late times, the matter density decreases and the curvature term dominates. This gives us:

$$\left(\frac{\dot{a}}{a}\right)^2 \approx \frac{-kc^2}{a^2}$$

This results in $\dot{a} \propto t$ where the universe expands linearly with time. This is called free expansion.

In both flat and hyperbolic geometries we have unbound infinite growth, with the hyperbolic geometry expanding faster than a universe with the flat geometry.

2 Spherical Geometry

Here, $k > 0$. Again we have the Friedmann equation:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho(t) - \frac{kc^2}{a^2}$$

Now, the second term remains negatively signed. This means that as the universe expands and the curvature term dominates, the expansion rate slows down and eventually stops. The universe then begins to contract, leading to a big crunch.

There is some point in time in which the two terms can cancel leading to $H(t) = 0$. This is an unstable equilibrium that will eventually collapse under its own gravity.

3 Observational Parameters

Instead of guessing at values of parameters, we want to make an attempt at measuring true values from the universe itself. There are a few key parameters we'll consider:

- The Hubble constant at the current epoch.
- Density parameters, we denote these $\Omega_x(t)$. The x may vary depending on the density being considered, i.e. mass density, energy density etc.
- Deceleration parameter, q_0 . This describes the rate at which the expansion of the universe is changing (regardless of acceleration or deceleration).

3.1 Hubble Constant at Current Epoch

We can measure this using velocities and distances. Velocities can be ascertained using redshift and distances using standard candles. Current measurements are between units of:

$$68 - 72 \text{ km s}^{-1} \text{ Mpc}^{-1}$$

The smaller this value is, the smaller the expansion, the higher value the more rapid the expansion.

We can say:

$$t_h \propto \frac{1}{H_0}$$

Where t_h is called the hubble time. This time provides a very rough estimate for the age of the universe. We can look at stars and use our understanding of the stellar evolution to determine a minimum possible age of the universe.

If we compute the hubble time for a flat matter dominated universe this underestimates the age of the universe. Not all stars can therefore be adequately explained by this model. To fix this, we can adjust our understanding of the curvature and species-makeup of the universe.

If we move from a flat universe to a hyperbolic one, the hubble time actually moves closer to the actual age. This fails for spherical geometry which gives an even younger universe.

This rules out a spherical geometry as a possible geometry for our current use. As adjusting to a hyperbolic universe would require changing other values to compensate (and this is indicating against), the current favoured solution is the flat universe.

If the Friedmann equation is solved properly, without our limit argument, a much better argument for the stellar age is generated.

3.2 Densities

It is preferable for us to determine densities as fractions of some value. In order to define our density values, they need to be relative to some important value. This value is called the critical density:

$$\Omega_x(t) = \frac{\rho_x(t)}{\rho_c(t)}$$

This critical density is the density value that we require in order keep $k = 0$. To determine this we look at the Friedmann Equation:

$$H^2(t) = \frac{8\pi G}{3}\rho(t) \implies \boxed{\rho_c(t) = \frac{3H^2(t)}{8\pi G}}$$

We can use these densities to simplify the Friedmann Eqn:

$$\begin{aligned} \left(\frac{\dot{a}}{a}\right)^2 &= H^2 = \frac{8\pi G}{3}\rho_c\Omega - \frac{kc^2}{a^2} \\ &= H^2\Omega - \frac{kc^2}{a^2} \\ 1 &= \Omega - \frac{kc^2}{a^2H^2} \\ \Omega(t) - 1 &= \frac{kc^2}{a^2(t)H^2(t)} \end{aligned}$$

It is then common to define a curvature density parameter $-\Omega_k(t)$ such that:

$$\begin{aligned} \Omega(t) - 1 &= -\Omega_k(t) \\ \Omega(t) + \Omega_k(t) &= 1 \end{aligned}$$

For a critically dense universe, the second term is zero and this means that the first term (all of the relative energy components) sums to one.

3.3 Deceleration Parameter

We can consider a Taylor expansion of $a(t)$ around the current epoch $a(t_0)$ (in the current epoch):

$$a(t_0) + \dot{a}(t_0)(t - t_0) + \frac{\ddot{a}}{2}(t - t_0)^2$$

And dividing through by the current epoch scale factor:

$$\begin{aligned}\frac{\dot{a}(t_0)}{a(t_0)} &\approx 1 + \left. \frac{\dot{a}(t_0)}{a(t_0)} \right|_{t=t_0} + \left. \frac{\ddot{a}(t_0)}{a(t_0)} \right|_{t=t_0} (t - t_0)^2 \\ &= 1 + H_0(t - t_0) - q_0 H_0^2 (t - t_0)^2\end{aligned}$$

Where q_0 is defined by this, and the negative sign is historical choice. This gives:

$$q_0 \equiv - \frac{\ddot{a}(t_0)}{a(t_0)} \frac{1}{H_0^2}$$

This was defined on the historical assumption of a decelerating expansion which has proven incorrect. The sign conventions are therefore weird.

For a matter dominated universe:

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3} \rho(t)$$

And at the present epoch:

$$\begin{aligned}&= -\frac{4\pi G}{3} \rho_0 \implies q_0 = \frac{4\pi G}{3} \frac{\rho_0}{H_0^2} \\ &= \frac{4\pi G}{3} \frac{\Omega_0 \rho_c}{H_0^2} \\ &= \frac{4\pi G}{3} \frac{\Omega_0 3H_0^2}{8\pi G H_0^2} = \frac{\Omega_0}{2} > 0\end{aligned}$$

This is positive for the deceleration constant, so the universe is...decelerating in its expansion. We know that this isn't accurate, so there's a missing piece.




Figure 12.1

We can fix this by adding a cosmological constant, shown in the dotted line that we will encounter next lecture.

Mon 16 Mar 2026 12:00

Lecture 13 - Cosmology VII: Cosmological Constant, CMB and Big Bang Problems

1 Cosmological Constant

The “missing piece” is called the Cosmological Constant Λ . This allows us to add a third term to the RHS of the Friedmann Equation:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}\rho(t) - \frac{kc^2}{a^2(t)} + \frac{\Lambda}{3}$$

Why we can do this is summed up by “general relativity says so”...

This extra term allows us to create a static (non-stable) universe with no expansion. This forms an unstable solution where any deviation will trigger a collapse. Different values give different results and we have the conditions $\Lambda \in \mathbb{R}$ and $\Lambda > 0$.

With this added term, a rederivation of the acceleration equation gives:

$$\frac{\ddot{a}}{a} = -\frac{4\pi G}{3}\left(\rho(t) + \frac{3p}{c^2}\right) + \frac{\Lambda}{3}$$

This first term is negative and the latter is positive, so depending on the value of the cosmological constant. This new term acts against the collapsing gravitational impacts so a sufficiently large constant can prevent this shrink, or even opposite it and cause an acceleration.

We want to model this as being some kind of fluid, and modify our $\rho(t)$ term:

$$\rho(t) + \rho_m(t) + \rho_r(t) \rightarrow \rho(t) + \rho_\Lambda$$

We can do so with:

$$\rho_\Lambda = \frac{\Lambda}{8\pi G}$$

Hence our Friedmann equation becomes:

$$\left(\frac{\dot{a}}{a}\right)^2 = \frac{8\pi G}{3}(\rho(t) + \rho_\Lambda) + \frac{kc^2}{a^2(t)}$$

We can use our fluid equation since we model this as a fluid and this causes a pressure term:

$$p_\Lambda = -c^2\rho_\Lambda$$

This effectively causes a negative pressure. In order to explain this weird repulsive behaviour, it makes sense that the sign of the pressure is flipped to flip the final impact.

There is however a massive discrepancy (causing an open question) between the order of magnitude of the measured Λ versus the predicted one.

As we’ve defined a density, we can now also define an Omega parameter:

$$\Omega_\Lambda(t) = \frac{\rho_\Lambda}{\rho_c(t)} = \frac{\Lambda}{8\pi G} \frac{8\pi G}{3H^2(t)} = \frac{\Lambda}{3H^2(t)}$$

Note that the constant is not time dependant, but the omega parameter is due to the hubble parameter.

We can now modify the Friedmann equation to finally say:

$$\Omega_m(t) + \Omega_r(t) + \Omega_\Lambda(t) + \Omega_k(t) = 1$$

2 Universe Evolution

This gives us a new equipartition point with the three fluids (matter, radiation, Λ). A universe eventually will be dominated by the cosmological constant. Depending on the relative combinations of these parameters, there is many many more possible outcomes for the universe.

Current best estimates is about 70% cosmological constant contributions and 30% matter and radiation contributions (combined) with no curvature contribution due to a flat universe.

Current estimates give:

- 5% baryonic matter.
- 26% dark matter.
- 69% dark energy.

This crucially gives $\Omega_0 + \Omega_{\Lambda,0} \approx 1$, hence we are a critically dense universe (or close to critically dense).

3 Cosmic Microwave Background

3.1 Why is the Big Bang “Hot”?

The strongest evidence is the Cosmic Microwave Background. This measures an isotropic almost-perfect black body spectrum with a temperature of:

$$T_0 = 2.7255\text{K}$$

It fits a Plank spectrum almost perfectly. By Wein’s law, we can know the temperature based on the energy spectrum distribution and vice-versa.

The radiation density is given by the Stefan-Boltzmann Law:

$$\epsilon_r \equiv \rho_r c^2 = \alpha T^4$$

where:

$$\alpha = \frac{\pi^2 k^2 k_B^4}{15 h^3 c^3}$$

With a value given in the formula sheet. Since we know that the ρ_r is related to the scale factor (scaling inversely with the scale factor) hence we know that the temperature has scaled down as the universe expands.

This tells us that:

$$\rho_r \propto a^{-4} \implies \frac{\alpha T^4}{c^2} \propto a^{-4} \implies \frac{T}{T_0} \propto \frac{1}{a}$$

As $a \rightarrow 0$, $T \rightarrow \infty$, hence the big bang was extremely hot.

4 Origins of the CMB

If we take a very old time when $a = a_0/1,000,000$, the temperature of the CMB photons is therefore $T(a) = 3,000,000\text{K}$.

This gives a mean energy much larger than the ionisation energy of hydrogen. Therefore, the hydrogen atoms could not exist at this epoch. This gave a primordial soup of ionised plasma with nuclei, electrons and photons.

Once the universe cools and expands, the photons are insufficiently energetic and allow the first neutral hydrogen ions to form (recombination). These photons become decoupled, where they can freely travel without interaction and can escape.

We calculate this properly in Y3, but the last scattering (on average) happened when the universe was about 300,000 years old. We cannot electromagnetically see the epochs that precede this, as the photons wouldn’t have been able to reach an observer.

The CMB is extremely important because it provides:

- Very strong evidence for the big bang.
- Constraints on the density parameters and geometry of the universe.

- Density fluctuations cause under- and over-densities causing “anisotropies” which provided the seeds for large scale structure forming.

5 Problems with the Big Bang

- Flatness Problem
 - The current universe is close to being critically dense and hence being flat.
 - However the flat universe is unstable, i.e. small perturbations to the total density cause more and more curvature quite quickly.
 - As the critical density varies with time and scale factor, the universe initial conditions must have been very very close (even closer) to being a flat universe at the big bang.
 - There is no explanation why this should be the case.
- Monopole Problem
 - Initially, at very high temperatures, all the forces would have combined into one primordial superforce.
 - All grand unified theories (GUTs) suggest that a magnetic monopole should be able to exist.
 - We know that this is not possible.
- Horizons Problem
 - The universe is isotropic and homogenous.
 - For this to happen, at some time any two sources must have been in causal contact (i.e. able to exchange momentum and energy with each other).
 - We¹ can calculate that this isn't the case.

6 Inflation

We can fix this by having a very large exponential expansion given by another cosmological constant when the universe was $\sim 10^{-36}$ years old.

End of Module.

¹Physicists in general...not us