

MATH 316 Lecture 7

Ashtan Mistal

May 20 2021

1 Boundary value Problems

Is there a similar setup for BVPs/ Let's consider 3 different BVPs:

1. P1: $y'' + \lambda y = 0$, for $x \in [0, L]$, with $y(0) = 0 = y(L)$
2. P2: $y'' + \lambda y = 0$, for $x \in [0, L]$, with $y'(0) = 0 = y'(L)$
3. P3: $y'' + \lambda y = 0$, for $x \in [0, L]$, with $y(0) = y(L)$ and $y'(0) = y'(L)$

Any value of λ for which P1 (P2 or P3) has a non-zero solution is called an **eigenvalue** of P1 (P2 or P3) and the corresponding solution is called an **eigenfunction** of P1 (P2 or P3).

Exercise: find the eigenvalues and eigenfunctions of problems P1, P2 and P3.

1.1 Solving BVP (P1,P2,P3)


P1:

$$y'' + \lambda y = 0, \text{ and } y(0) = 0 = y(L)$$

λ : Eigenvalue. There are three categories that we have to investigate each time we solve such a problem:

1. If λ is negative ($\lambda < 0$):

- $\lambda = -\mu^2 \rightarrow y'' - \mu^2 y = 0$
- $r^2 - \mu^2 = 0 \rightarrow r_1 = \mu, r_2 = -\mu \rightarrow y(x) = C_1 e^{\mu x} + C_2 e^{-\mu x}$
- Note that $\cosh(\mu x) = \frac{e^{\mu x} + e^{-\mu x}}{2}$ and $\sinh(\mu x) = \frac{e^{\mu x} - e^{-\mu x}}{2} \rightarrow$
- $\hookrightarrow y(x) = A \sinh(\mu x) + B \cosh(\mu x)$



- ---
- Note that we have the boundary conditions $y(0) = 0 \rightarrow B = 0$ and $y(L) = 0 \rightarrow A \sinh(\mu L) = 0$
- $\Rightarrow A = 0$ and $\Rightarrow y(x) = 0$ which is trivial

2. If λ is zero: $y'' = 0 \rightarrow y(x) = Ax + B$

- $y(0) = 0 \rightarrow B = 0$
- $y(L) = 0 \rightarrow AL = 0 \rightarrow A = 0 \rightarrow y(x) = 0$
- Therefore it's a trivial solution.

3. If $\lambda > 0$: $\lambda = \mu^2 \rightarrow y'' + \mu^2 y = 0$

- $r^2 + \mu^2 = 0 \rightarrow r = \pm i\mu$
- $y(x) = A \sin(\mu x) + B \cos(\mu x)$
- $y(0) = 0 \rightarrow B = 0$
- $y(L) = 0 \rightarrow A \sin(\mu L) = 0 \rightarrow \begin{matrix} A = 0(\text{trivial}) \\ \sin(\mu L) = 0 \rightarrow \mu L = n\pi \end{matrix}$ therefore $\mu = \frac{n\pi}{L}$
- Eigenvalue: $\lambda = \left(\frac{n\pi}{L}\right)^2$
- Eigenfunction: $y_n(x) = A_n \sin\left(\frac{n\pi}{L}x\right)$

1.2 P2: $y'' + \lambda y = 0$

$$y'(0) = 0 = y'(L)$$

$$r^2 + \lambda = 0$$

1. If $\lambda > 0$, $\lambda = \mu^2$

- $r^2 + \mu^2 = 0 \rightarrow r = \pm i\mu$
- $y(x) = A \sin(\mu x) + B \cos(\mu x)$
- Sub boundary conditions:
- $y'(0) = A = 0$ and $y'(L) = 0 \rightarrow -B\mu \sin(\mu L) = 0$

$B = 0$ which is trivial

- This gives us two solutions: $\underbrace{\sin(\mu L) = 0}_{\mu = \frac{n\pi}{L}}$

- $\Rightarrow \lambda = \left(\frac{n\pi}{L}\right)^2$ is the eigenvalue
- $y_n(x) = B \cos\left(\frac{n\pi}{L}x\right)$ is the eigenfunction

2. If $\lambda < 0$: $\rightarrow \lambda = -\mu^2$

- $r^2 - \mu^2 = 0 \rightarrow r = \pm\mu \rightarrow y(x) = C_1 e^{\mu x} + C_2 e^{-\mu x} \Rightarrow y(x) = A \sinh(\mu x) + B \cosh(\mu x)$
- Substituting the boundary conditions into $y(x) = A \sinh(\mu x) + B \cosh(\mu x)$, we get:
- $y'(0) = 0 \rightarrow A = 0$
- $y'(L) = 0 \rightarrow -\mu B \sinh(\mu L) = 0 \rightarrow B = 0$ (which means that $y(x) = 0$ which is trivial)

3. If $\lambda = 0 \rightarrow y'' = 0 \rightarrow y = Ax + B$

- $y'(0) = 0 \rightarrow A = 0$ and $y'(L) = 0 \rightarrow A = 0$: $\rightarrow y(x) = B$
- $\lambda = 0$ is an eigenvalue $\rightarrow y(x) = 1$ is the eigenfunction
- For P2 problems: eigenvalues are: $0, \frac{n^2\pi^2}{L^2}$ and eigenfunctions are: $1, \cos\left(\frac{n\pi}{L}x\right)$

1.3 P3: $y'' + \lambda y = 0$

Periodic boundary conditions: $y(0) = y(L)$

1. If $\lambda > 0$: $\lambda = \mu^2$

- $r = \pm\mu i \rightarrow y(x) = A \sin(\mu x) + B \cos(\mu x)$
- $y(0) = y(L) \rightarrow B = A \sin(\mu L) + B \cos(\mu L)$
- $y'(0) = y'(L) \rightarrow A\mu = A\mu \cos(\mu L) - B\mu \sin(\mu L)$

$$\begin{aligned} r = \pm\mu i &\rightarrow y(x) = A \sin(\mu x) + B \cos(\mu x) \\ y(0) = y(L) &\rightarrow B = A \sin(\mu L) + B \cos(\mu L) \\ y'(0) = y'(L) &\rightarrow A\mu = A\mu \cos(\mu L) - B\mu \sin(\mu L) \end{aligned} \quad \left\{ \begin{array}{l} \times \frac{A}{B} \\ \rightarrow A = \frac{A^2}{B} \sin(\mu L) + A \cos(\mu L) \\ \rightarrow A = A \cos(\mu L) - B \sin(\mu L) \end{array} \right. \rightarrow \frac{A^2}{B} + B \sin(\mu L) = 0$$

-
- We get the last term by using the previous two terms (on the right) and cancelling out.
- $\sin(\mu L) = 0 \rightarrow \mu L = n\pi$. $A \neq 0$ and $B \neq 0$ and therefore $\mu = \frac{n\pi}{L}$
- Eigenvalues: $\lambda = \left(\frac{n\pi}{L}\right)^2$

- Eigenfunction is $y_n(x) = A_n \sin(\frac{n\pi}{L}x) + B_n \cos(\frac{n\pi}{L}x)$

2. If $\lambda < 0$: $\lambda = -\mu^2$

- $y(x) = A \sinh(\mu x) + B \cosh(\mu x)$
- $y(0) = y(L) \rightarrow B = A \sinh(\mu L) + B \cosh(\mu L)$
- $y'(0) = y'(L) \rightarrow A = A \cosh(\mu L) - B \sinh(\mu L)$
- Multiplying $B = A \sinh(\mu L) + B \cosh(\mu L)$ by:

$$-1 \quad y(x) = A \sinh(\mu x) + B \cosh(\mu x) \quad \textcircled{1} \quad B = A \sinh(\mu L) + B \cosh(\mu L) \quad \times \frac{A}{B} \rightarrow A = \frac{A^2}{B} \sinh(\mu L) + A \cosh(\mu L)$$

- Okay this is going way too fast... screenshots it is.


(ii) If $\lambda < 0$: $\lambda = -\mu^2$ $y(x) = A \sinh(\mu x) + B \cosh(\mu x)$ $\textcircled{1}$

$$y(0) = y(L) \rightarrow B = A \sinh(\mu L) + B \cosh(\mu L) \quad \times \frac{A}{B} \rightarrow A = \frac{A^2}{B} \sinh(\mu L) + A \cosh(\mu L)$$

$$y'(0) = y'(L) \rightarrow \textcircled{2} \quad A = A \cosh(\mu L) - B \sinh(\mu L)$$

$$\textcircled{1} - \textcircled{2} \rightarrow \left(\frac{A^2}{B} + B \right) \sinh(\mu L) = 0 \quad \times$$

No solution \rightarrow No eigenvalues



3. $\lambda = 0$:

- View screenshot below.

(iii) $\lambda = 0$: $y'' = 0 \rightarrow y = Ax + B$

$$\left. \begin{array}{l} y(0) = B \text{ \& } y(L) = AL + B \rightarrow A = 0 \\ y'(0) = y'(L) = A \end{array} \right\} \rightarrow y(x) = B \quad \checkmark$$

$\lambda = 0$ is an eigenvalue

$y(x) = 1$ is an eigenfunction

2 Fourier Series

Fourier series arise in 3 different situations of relevance to this course: 1. Simple boundary value problems, e.g. P1-P32. Partial differential equations that describe heat flow, waves and diffusion (more later). 3. Some initial value problems with less simple periodic forcing, e.g. we are very unlikely to have exactly: $f(t) = F_0 \cos(\omega t)$, in any real system, but might have a periodic forcing function.

For what follows, let the interval in P1-P3 be the interval $[a, b] = [-L, L]$. The key idea is that an arbitrary function, $f(t)$, defined on $[-L, L]$ can be represented in the following form:

$$f(t) = \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi t}{L}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi t}{L}\right) \quad (1)$$

Note that these are the eigenfunctions of problem P3. Outside of the interval, because each function above has period $2L$, the above series must converge to a periodic extension of $f(t)$ of period $2L$.

Two immediate questions:

1. Can all functions $f(t)$ be represented in this way, i.e. which functions?
2. How do we find the coefficients a_n and b_n ?

Definition: If the series on the right-hand side of (1) converges to a function $f(t)$ then this is called the Fourier series of $f(t)$.

Comments:

Firstly, in order for $f(t)$ to have Fourier series representation (1), that is valid for all t it is necessary that $f(t)$ is periodic, with period $2L$, i.e.

$$f(t + 2L) = f(t) \quad \forall t$$

Secondly, suppose that $f(t)$ has a Fourier series representation (1). Then a_n and b_n are determined straightforwardly. See below for a_n :

1. Multiply (1) by $\cos(\frac{m\pi t}{L})$
2. Integrate both sides of the equation between $[-L, L]$:

$$\int_{-L}^L f(t) \cos \frac{m\pi t}{L} dt = \int_{-L}^L \left(\frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos \frac{n\pi t}{L} + \sum_{n=1}^{\infty} b_n \sin \frac{n\pi t}{L} \right) \cos \frac{m\pi t}{L} dt$$

Note that:

$$\begin{aligned} \int_{-L}^L \cos \frac{n\pi t}{L} \cos \frac{m\pi t}{L} dt &= \begin{cases} 0 & m \neq n \\ L & m = n \end{cases} \\ \int_{-L}^L \cos \frac{n\pi t}{L} \sin \frac{m\pi t}{L} dt &= 0 \\ \int_{-L}^L \sin \frac{n\pi t}{L} \sin \frac{m\pi t}{L} dt &= \begin{cases} 0 & m \neq n \\ L & m = n \end{cases} \end{aligned}$$

Therefore, interchanging summation and integration:

$$\begin{aligned} \int_{-L}^L f(t) \cos \frac{m\pi t}{L} dt &= a_m \int_{-L}^L \cos \frac{m\pi t}{L} \cos \frac{m\pi t}{L} dt = a_m L \\ a_m &= \frac{1}{L} \int_{-L}^L f(t) \cos \frac{m\pi t}{L} dt \end{aligned}$$

For the coefficients b_n a similar procedure is possible (exercise).
Thus, we finally have:

$$\begin{aligned}a_0 &= \frac{1}{L} \int_{-L}^L f(t) dt \\a_m &= \frac{1}{L} \int_{-L}^L f(t) \cos \frac{m\pi t}{L} dt \quad m = 1, 2, 3, \dots \\b_m &= \frac{1}{L} \int_{-L}^L f(t) \sin \frac{m\pi t}{L} dt \quad m = 1, 2, 3, \dots\end{aligned}$$

which are known as the **Euler-Fourier series**.

2.1 Example 1

Assume that the function $f(t)$, defined by $f(t) = \begin{cases} t & -L \leq t < 0 \\ 0 & 0 \leq t < L \end{cases}$ with $f(t + 2L) = f(t)$, has a fourier series. Sketch the function and find the fourier series.

Solution:

$$\begin{aligned}f(t) &= \frac{a_0}{2} + \sum_{n=1}^{\infty} a_n \cos\left(\frac{n\pi t}{L}\right) + \sum_{n=1}^{\infty} b_n \sin\left(\frac{n\pi t}{L}\right) \\a_0 &= \frac{1}{L} \int_{-L}^L f(t) dt = \frac{1}{L} \int_{-L}^0 t dt = \frac{-L}{2} \\a_n &= \frac{1}{L} \int_{-L}^L f(t) \cos\left(\frac{n\pi t}{L}\right) dt = \frac{1}{L} \int_{-L}^0 t \cos\left(\frac{n\pi t}{L}\right) dt\end{aligned}$$

Using integration by parts, with $u = t$ ($du = dt$) and $dv = \cos(\frac{n\pi t}{L})dt$, with $v = \frac{L}{n\pi} \sin(\frac{n\pi t}{L})$, we get the following:

$$\begin{aligned}a_n &= \frac{1}{L} \left[\frac{tL}{n\pi} \frac{\sin(n\pi t)}{L} \right]_{-L}^0 - \int_{-L}^0 \frac{L}{n\pi} \frac{\sin(n\pi t)}{L} dt \\&= \frac{1}{n\pi} \left[\frac{L}{n\pi} \cos\left(\frac{n\pi t}{L}\right) \right]_{-L}^0 = \frac{L}{n^2\pi^2} (1 - \cos(n\pi))\end{aligned}$$