



+91-9527070000

Call Us Now- 24/7 Customer Support

SERVICES

Learn more about the services offered at SPECT LAB Pune, India. We offer the most advanced medical imaging services available in India.

THYROID SCAN

PREREQUISITE	2-4 days prior to test off Antythroid medication. Fasting not required
TIME REQUIRED	1.5 Hr \n Scanning Duration: 20 mins. Scanning Duration 20 Min.
PROCEDURE	Tc99m Pertechnetate is injected intravenously, blood flow, blood pool and after 20 minutes static images will be acquired for anterior neck.
INDICATIONS	Determination of thyroid size, function and position. Evaluation of functional status of thyroid nodules. Differentiating Thyroiditis from Grave's disease. Evaluation of Thyroid and neck masses. Quantitative thyroid uptake. Detection of ectopic thyroid eg. Lingual thyroid. To detect restroternal extension in MNG or large goiters.

COLLOID LIVER SCAN

PREREQUISITE	No Prequisite
TIME REQUIRED	Scan Time 30 – 45 minutes .
PROCEDURE	Tc 99m Phytate is injected intravenously and following that images in multiple projections (anterior, anterior with costal marker, posterior, right and left laterals and right and left oblique) were acquired of the abdomen.
INDICATIONS	To assess hepatocellular dysfunction in diffuse liver disease liver disease. Suspected chronic liver disease (7cirrhosis). Assess hepatic function in chronic alcoholics. Evaluation of splenomegaly.

GASTRIC EMPTYING

PRE PROCEDURE PREPARATION	Fasting for 4 hrs
METHOD OF ADMINISTRATION	Oral.
PROCEDURE IMAGING TIME	1.5 to 2 hrs
TOTAL TIME ALONG WITH REPORTING	immediate
POST PROCEDURE INSTRUCTIONS	Avoid contact with small children less than 5 years and pregnant lady for 1 day. Stop breast feeding for 24 hours and encourage patient to pass urine more frequently.

DTPA RENOGRAPH (GLOMERULAR FUNCTION) / EC

(MAG-3) RENOGRAM (GLOM+ TUBULAR FUNCTION)

PREREQUISITE	Good Hydration.
TIME REQUIRED	Scan Time 45mins – 2 hours.
PROCEDURE	Tc99m DTPA is injected IV after ensuring good hydration of the patient. I.V. lasix is given after 10 mins of tracer injection. Sequential dynamic images are obtained for 31 minutes followed by post void and 1hour delayed image.
INDICATIONS	Evaluation of differential renal function. Evaluation of obstructive uropathy. Evaluation of unequal sized kidney / ? agenesis of kidney. Indirect radio nuclide cystography of VU reflux. Follow up renal transplantation.

BONE SCAN

PREREQUISITE	No Prerequisite.
TIME REQUIRED	Scan Time 2-3hr minutes .
PROCEDURE	20 mCi of Tc 99m M.D.P. is administered intravenously and after 2.5 hours anterior and posterior images is acquired of the whole skeleton.
INDICATIONS	Screen for bone metastases in patient with malignancy. Infection (osteomyelitis vs. cellulitis). Evaluation of suspected fracture and stress fractures. Evaluation of avascular necrosis and vascularity of bone graft. Evaluates response to chemo, or radiation treatment for cancer. Evaluation of prosthesis for loosening or infection. Evaluation of nonspecific X-ray/ MRI bone signals. Metabolic bone disease.

LYMPHOSCINTIGRAPHY

PRE PROCEDURE PREPARATION	None
METHOD OF ADMINISTRATION	Interdigital with no side or after effects.
PROCEDURE IMAGING TIME	30 to 60 mins.
INJECTION TO IMAGING TIME	Immediate, followed by 40-min delayed post walk image.
TOTAL TIME ALONG WITH REPORTING	2 – 3 hrs.
POST PROCEDURE INSTRUCTIONS	No specific .

I-131 WHOLE BODY SCAN

PREREQUISITE	3-4 weeks post total thyroidectomy, Off medication post surgery. Avoid Iodine intake 4-5 days.

PRIOR THE STUDY	3-4 hrs fasting.
TIME REQUIRED	Scan Time 30 – 45 minutes.
PROCEDURE	Day1 I 131 solution is given orally. Day 2 I 131 Whole body scan is performed.
INDICATIONS	Detection of residual tissue in thyroid bed and local or distant thyroid mets after total thyroidectomy for Ca thyroid. Assessment of response to I- 131 therapy for Ca thyroid.

HEPATOBIILIARY SCINTRAPHY (HIDA SCAN)

PREREQUISITE	No Prerequisite.
TIME REQUIRED	Scan Time 2 – 24 hours
PROCEDURE	2 mCi 99m Tc Mebrofenin is injected intravenously after skipping one normal feed. Dynamic images are obtained for 20 minutes. Second set of dynamic images are obtained post feed for 20 minutes. Delayed static images may be acquired till 24hours if required.
INDICATIONS	Evaluation of suspected acute cholecystitis. Congenital abnormalities of biliry tract disorders. Evaluation of commonbile duct obstruction. detection of bile extravassion (biliry leaks).

PARATHYROID SCAN

PREREQUISITE	No Prerequisite.
TIME REQUIRED	Scan Time 30 – 45 minutes.
PROCEDURE	4 mCi of 99mTc(O4-) is injected intravenously and sequential dynamic images and static blood pool and 20th minute are acquired for anterior neck. Without moving the patient 20 mCi of 99mTc Sestamibi is injected intravenously and static images at 20th minute (Early) and 45 mins (delayed) are acquired for the ant. neck.
INDICATIONS	Evaluation of primary or secondary hyperparathyroidism. Parathyroid adenomas or hyperplasia prior to surgical removal.

STRESS & REST MYOCARDIAL PERFUSION IMAGING (THALLIUM SCAN)

PREREQUISITE	Off beta blockers 48 prior the study. 4hrs fasting. No medication on the day of scan.
TIME REQUIRED	Scan Time 30 – 45 minutes .

PROCEDURE	Patient has to undergo physiological / pharmacological stress. 99mTc Sestamibi is injected when patient achieves 85% THR . 45 mins post injection scan is performed. Almost after 1hour 99mTc Sestamibi is again injected to the patient in the resting position(SOS post nitrate) . 45 mins post injection scan is performed.
INDICATIONS	Diagnosis of coronary artery disease. Prognosis of known coronary artery disease. Risk stratification in known or suspected CAD. Suspected restenosis after CABG. Evaluation of PTCA / stent, anti – angina drugs. Pre – operative cardiac risk assessment. Assessment of myocardial viability.

POST CAPTOPRIL RENOGRAM (SUSPECTED RENAL ART. STENOSIS)

PREREQUISITE	Good Hydration. Off medications 2-3 days prior study.
TIME REQUIRED	Scan Time 45mins for each scan.
PROCEDURE	A) Baseline 4 mCi of Tc DTPA is injected intravenously after ensuring good hydration of the patient. Dynamic images are obtained for 31 minutes . B) Post Captopri I Patient is given Tab. Enalapril & similar imaging protocol is followed 2 hours after the Baseline study.
INDICATIONS	Evaluation of reno – vascular hypertension.

DMSA RENAL SCAN (CORTICAL FUNCTION)

PREREQUISITE	Good Hydration.
TIME REQUIRED	Scan Time 15mins.
PROCEDURE	Tc99m DMSA is injected intravenously after ensuring good hydration of the patient. Static images in posterior, LPO & RPO views are acquired at 2.5 hrs post injection.
INDICATIONS	Diagnosis of acute pyelonephritis. Diagnosis of renal scarring.

BRAIN SPECT (CEREBRAL PERFUSION SCAN)

PREREQUISITE	No PreRequisite.
TIME REQUIRED	Scan Time 30mins.
PROCEDURE	Tc99m ECD in injected intravenously & immediate spect images of skull is acquired.
INDICATIONS	Alzheimer's disease. Differential diagnosis of various forms of dementia. Detection of epileptogenic foci.

VENTRICULAR SHUNT PATENCY STUDY (CISTERNOGRAPHY)

PREREQUISITE	No PreRequisite.
TIME REQUIRED	Scan Time 30mins.
PROCEDURE	In order to determine the position of the leak a 99m Tc DTPA material is injected into the subarachnoid space via a lumbar approach.and dynamic and static views will acquired.
INDICATIONS	Study of CSF dynamics.

RBC BLOOD POOL SCAN/GI BLEED SCAN

PREREQUISITE	No PreRequisite.
TIME REQUIRED	Scan Time 45mins – 24 hrs.
PROCEDURE	Invivo RBC Labeling Method 20 minutes after the intravenous administration of stannous pyrophosphate (cold Kit), 10 mCi of 99m Tc (O4-) is injected intravenously by positioning the patient supine covering the anterior abdomen & Perfusion images (2sec/fr x 60 frames) followed by dynamic images (1min/fr x 60 frames). There after delayed static images are acquired. If required TOMO images may be acquired.
INDICATIONS	Liver hemangiomas. Detection and localization of gastrointestinal bleeding.

MECKEL'S DIVERTICULUM IMAGING

PREREQUISITE	2-4 hrs fasting.
TIME REQUIRED	Scan Time 45mins.
PROCEDURE	99mTc is injected intravenously under the camera and dynamic images are acquired for 30 minutes immediately thereafter of the anterior abdomen, followed by static images of the LAO , RAO views and post void static image of the abdomen.
INDICATIONS	Detection and localization of Meckel's diverticulum.

MILK SCAN (GE REFLUX)

PREREQUISITE	No PreRequisite.
TIME	Scan Time 45mins

REQUIRED	
PROCEDURE	99mTc-DTPA is fed orally to the patient mixed in 5 ml milk. & then unlabelled 50 ml milk is given to wash the esophagus and sequential dynamic images are acquired for 30 minutes (10sec/frame) immediately thereafter. Static anterior & posterior abdomen image was taken after 1 hour.
INDICATIONS	Detection of GE reflux in infants having LRTI.

MIBG SCAN

PREREQUISITE	PreRequisite No PreRequisite.
TIME REQUIRED	Scan Time 45mins.
PROCEDURE	Orally one tea spoon twice a day of Colloidal Iodine solution is started 2 day prior to injection and throughout the study. ¹³¹ I MIBG (Meta Iodo Benzyl Guanidine) is injected intravenously and whole body views of the upper torso (Anterior and Posterior) are acquired at 24 & 48,72 hours.5mins .
INDICATIONS	Pheochromocytoma. Neuroblastoma. 1)TESTICULAR SCAN Differentiating epididymitis from torsion. 2)Salivary gland scan Stenosis of salivary gland for dryness of mouth. 3)Lymphoscintigraphy To differentiate lymphoedema from venous pathology.

MUGA SCAN

PREREQUISITE	No PreRequisite.
TIME REQUIRED	Scan Time 45mins.
TECHNIQUE IN VIVO RBC LABELING METHOD	Twenty minutes after the intravenous administration of stannous pyrophosphate, 25 mCi of 99m Tc (O4-) is MUGA SCAN

DACRYOSCINTIGRAPHY

PREREQUISITE	No Prequisite.
TIME REQUIRED	Scan Time 30 – 45 minutes
TECHNIQUE	With patient lying supine under the gamma camera,100microCi of 99mTc O4- was instilled at the outer canthus of both the eyes with the help of a micropipette.The patient was instructed to blink frequently when sequential dynamic images were acquired for both the eyes immediately for 15 mins (128X128 matrix at the rate of 15 Secs/frame). Static images were acquired at 15 mins , 1 hour & at 2 hours post instillation
INDICATION	For assessing the patency of nasolacrymal duct in patients having epiphora

GASTRIC EMPTYING SCAN

PREREQUISITE	No PreRequisite.
PROCEDURE	For a gastric emptying study, a patient eats a meal in which the solid component of the meal (for example, scrambled eggs), the liquid component of the meal (for example, water), or both, are mixed with a small amount of radioactive material. A scanner is placed over the patient's stomach to monitor the amount of radioactivity in the stomach for several hours after the test meal is eaten. As the radioactively-labeled food empties from the stomach, the amount of radioactivity in the stomach decreases. The rate at which the radioactivity leaves the stomach reflects the rate at which food is emptying from the stomach.
INDICATION	Gastroparesis.

TESTICULAR TORSION SCAN

PREREQUISITE	No PreRequisite
TIME REQUIRED	Scan Time 1-2h.
PROCEDURE	99mTc is injected intravenously under the camera and dynamic images are acquired for 15 minutes immediately thereafter of the anterior abdomen. Static image taken after 30min.

DIRECT CYSTOGRAM

PREREQUISITE	No PreRequisite
TECHNIQUE	2 mci 99mTc-DTPA is administered into the bladder via supra pubic puncture under aseptic conditions followed by an appropriate volume of sterile normal saline. The filling and voiding images are acquired continuously with the patient lying supine followed by post void static image.
INDICATION	To rule out VU Reflux

PET-CT IMAGING PROCEDURE(FDG)

PRE REQUISITE	6 hrs of fasting. Blood sugar level below 200mg/dl.
	- You would be then directed to a room where a small dosage of 18 F – FDG / Sodium Fluoride will be injected in your vein. - After injection, you would be asked to quietly relax and rest in a separate room at our Centre for about an hour before you are taken up for the scan. - During this resting period, unless recommended by our staff, no

PROCEDURE	attendants or relatives would be permitted to stay with you in this separate waiting room. - You are advised to stay calm, restrict physical movements and avoid talking to others while present in the room. - In case you feel any vomiting sensation or giddiness, inform the staff immediately by pressing the call bell. - You may be given oral contrast to drink. You will have to finish drinking it in around 30 minutes. This helps to get a better image quality. However, don't force yourself to drink it in case you are feeling nauseatic. You would be required to drink plenty of water after the injection and till you are in the waiting room. - You will be asked to void (pass urine) before starting the scanning procedure. Take care to ensure that you don't spill urine on any of your body parts. - You would be asked to lie down flat on your back on the bed of the PET-CT scanner and are expected to remain still during the entire scanning period of 15-20 minutes. - Your body would pass through a dough-nut like structure for obtaining the scan. Please let us know if you are claustrophobic or anxious. - You would be informed about the beginning and end of the scanning procedure by our staff. - After the completion of the scanning, you will be asked to sit in another room for some time. The Nuclear staff in the mean time will evaluate your PET-CT images and if needed may repeat your whole body scan. - Your PET-CT scan would be finally evaluated by our experienced Nuclear Medicine Physician.
PET AND PET/CT SCANS ARE PERFORMED TO	- Detect cancer. - Determine whether a cancer has spread in the body. - Assess the effectiveness of a treatment plan, such as cancer therapy. - Determine if a cancer has returned after treatment. - Determine blood flow to the heart muscle.

RADIO IODINE THERAPY

PREREQUISITE	Patient should not eat or drink after midnight on the day of the procedure. If you have been taking anti-thyroid medications, you must stop at least three days before the therapy is given. Frequently, the anti-thyroid medication is stopped for five to seven days before therapy. Radioactive Iodine I-131 (also called Radioiodine I-131) therapy is a treatment for an overactive thyroid, a condition called hyperthyroidism. Radioactive iodine I-131 may also be used to treat thyroid cancer. Low dose of iodine used for hyperthyroidism and high dose used for thyroid cancer patient required admission after high dose therapy.
POST	Use private toilet facilities, if possible, and flush twice after each use. - Bathe daily and wash hands frequently. - Drink a normal amount of fluids. - Use disposable eating utensils or wash your utensils separately from others. - Sleep alone and avoid prolonged intimate contact for three or four days. Brief periods of close contact, such as handshaking and hugging, are permitted. - Launder your linens, towels, and clothes daily at home, separately. No special

PROCEDURE	cleaning of the washing machine is required between loads. • Do not prepare food for others that requires prolonged handling with bare hands. • If you are breast-feeding, you must stop several days before to ensure that milk production has also stopped. • You should avoid becoming pregnant from six months to one year after treatment.
PRE REQUISITE	You are requested to report one hour before the scheduled appointment time. You should not eat and drink for at least 6 hours before your scheduled time of appointment. You are, however, permitted to drink plain water, but should strictly avoid coffee, tea, fruit juices, coconut water, prasad, chocolates, chewing gum, etc. Radioactive iodine I-131 may also be used to treat thyroid cancer. You should not undertake any intense and strenuous physical activity or exercise for 24 hours before the scan appointment.
ON THE DAY OF SCAN	You should carry all your relevant medical records, reports, film and/or CD of Xrays, CT, MRI, PET-CT or any other investigation done till date. These are important in the interpretation of your scheduled PET-CT scan by the doctor. • You should take along an adult attendant. • You should avoid children and pregnant women accompanying you. • You should wear loose and comfortable clothing and preferably carry an additional set. • You should not wear any metallic objects, jewellery or valuables. You should leave all such items at home or may keep them with the person accompanying you. • You may be required to remain at our Centre for at least 4 hours from the scheduled time of appointment for the PET-CT scan.
PROCEDURE	Registration and billing for the PET/CT. • After registration You would be then directed to a room where a small dosage of 18 F – FDG / Sodium Fluoride will be injected in your vein. • You should avoid children and pregnant women accompanying you. • After injection, you would be asked to quietly relax and rest in a separate room at our Centre for about an hour before you are taken up for the scan. • During this resting period, unless recommended by our staff, no attendants or relatives would be permitted to stay with you in this separate waiting room. You are advised to stay calm, restrict physical movements and avoid talking to others while present in the room. • You may be given oral contrast to drink. You will have to finish drinking it in around 30 minutes. This helps to get a better image quality. However, don't force yourself to drink it in case you are feeling nauseatic. You would be required to drink plenty of water after the injection and till you are in the waiting room. • You should avoid children and pregnant women accompanying you. You will be asked to void (pass urine) before starting the scanning procedure. Take care to ensure that you don't spill urine on any of your body parts. • You would be asked to lie down flat on your back on the bed of the PET-CT scanner and are expected to remain still during the entire scanning period

	of 15-20 minutes. Your body would pass through a dough-nut like structure for obtaining the scan. Please let us know if you are claustrophobic or anxious. You would be informed about the beginning and end of the scanning procedure by our staff. • After the completion of the scanning, you will be asked to sit in another room for some time. The Nuclear staff in the mean time will evaluate your PET-CT images and if needed may repeat your whole body scan. • Your PET-CT scan would be finally evaluated by our experienced Nuclear Medicine Physician.
PET AND PET/CT SCANS ARE PERFORMED TO	Detect cancer. • Determine whether a cancer has spread in the body. • Assess the effectiveness of a treatment plan, such as cancer therapy. • Determine if a cancer has returned after treatment.

GALLIUM-68 PSMA SCAN

PREREQUISITE	• Fasting not required. • Please bring previous investigations.
TIME REQUIRED	Duration of the procedure: 2 - 3hr
PROCEDURE	• IV administration of a small radioactive agent. • There are no side effects. • Waiting period of 60 – 90 Mins. • Patient cannot leave the waiting room in this period as per government regulations. • Few patients may require a delayed image at 3 Hrs. • Attendants and relatives are not allowed to accompany the patient in waiting room. • Only disabled (bed-ridden) patients who require close monitoring and physical assistance should be accompanied by an attendant.
INDICATIONS	• To detect occult prostate cancer. • To determine its spread. • To assess treatment response. • Surveillance

GALLIUM-68 OCTREOTIDE /DOTA/SOMATOSTATIN SCAN

PREREQUISITE	No PreRequisite
TIME REQUIRED	Duration of the procedure 1-2hr
	DOTA studies are able to determine the extent of neuro endocrine tumor (NET)cancer and whether it has spread to the lymph nodes or other parts of the

TECHNIQUE	body, unlike traditional imaging technologies such as magnetic resonance imaging (MRI) and computed tomography (CT), which are often unable to detect the spread of NET cancer cells. Molecular imaging technologies help physicians select the most effective therapy for NET cancer, taking into account a tumor's unique molecular properties and whether the cancer is localized or diffuse, or spread out.
PROCEDURE	The study involves a special molecule called a monoclonal antibody developed in a laboratory and designed to bind to the somastatin receptors. This antibody is paired with a radioactive material called 68-Gallium that can be detected by a PET CT camera. When injected into the patient's bloodstream, the radioactive antibody travels and attaches to cancer cells. The PET CT then produces three-dimensional images of the tumor and its location inside the body.

ADDRESS
 Sr. No. 268, Behind Crystal Honda Showroom,
 Near Mantri Alpine and DSK Toyota, Mumbai - Banglore Highway, Bavdhan, Pune - 411 021

MOB NO.

+91-9527070000,
 +91-8975758509,
 +91-8975758566

EMAIL ADDRESS

drsolv1@gmail.com (<https://web.archive.org/web/20190909073837/mailto:drsolv1@gmail.com>)
 kiran.ayre@spectlab.com



OPENING HOURS

Monday - Sat

7 AM TO 7 PM

Sunday Closed