

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/336370234>

# Long-term Behavior of Mean-field Noisy Bounded Confidence Models with Distributed Radicals

Conference Paper · December 2019

DOI: 10.1109/CDC40024.2019.9030212

CITATIONS

2

READS

59

3 authors, including:



[Anton V Proskurnikov](#)

Politecnico di Torino

179 PUBLICATIONS 1,912 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Universal controllers in optimal disturbance rejection and tracking problems [View project](#)



New results in stability theory for nonlinear systems. [View project](#)

# Long-term Behavior of Mean-field Noisy Bounded Confidence Models with Distributed Radicals

M. A. S. Kolarijani, A. V. Proskurnikov and P. Mohajerin Esfahani

**Abstract**—In this paper, we consider the mean-field model of noisy bounded confidence opinion dynamics under exogenous influence of static radical opinions. The long-term behavior of the model is analyzed by providing a sufficient condition for exponential convergence of the dynamics to stationary state. The stationary state is also characterized by a global estimate for a sufficiently large noise. Furthermore, we consider the order-disorder transition in the model in order to identify the effect of the (relative) mass of the radicals on the critical noise level at which this transition occurs. A numerical scheme for approximating the critical noise level is provided and validated through numerical simulations of the mean-field model and the corresponding agent-based model for a particular distribution of radical opinions.

## I. INTRODUCTION

Long before the recent “boom” in the study of complex systems, the necessity of mathematical models that can capture the diversity of clustering behaviors in real-world networks was realized in mathematical sociology [1]. The problem of disclosing these mechanisms is nowadays referred to as the community cleavage problem or Abelson’s diversity puzzle [2], [3] and primarily concerned with temporal mechanisms of *opinion formation* under social influence. The interdisciplinary area of social dynamics [3]–[8] has attracted enormous attention of the research community.

One feature observed in social and biological systems is the *homophily* [9], or tendency of individuals to bond with similar ones. In other words, like-minded individuals influence a social actor “stronger” than different-minded ones. Mathematically, social influence weights should be non-constant, but rather *opinion-dependent*. That is, the underlying interaction graph is dynamic and varies as the opinions of the individuals change. A class of nonlinear models resembling this feature are the so-called *bounded confidence* models proposed as extensions of the deterministic [10] and randomized gossip-based [11] consensus algorithms for multi-agent networks. Bounded confidence models stipulate that a social actor is insensitive to opinions beyond its bounded *confidence set* (usually, this set is an open or closed ball, centered at the actor’s own opinion). A detailed survey of bounded confidence models and relevant mathematical results can be found in [8]. Bounded confidence models

exhibit convergence of the opinions to some steady values, which can reach consensus or split into several disjoint clusters. Opinions in real social groups, however, usually do not terminate at steady values, which is usually explained by two factors.

The first reason explaining opinion fluctuation is exogenous influence, which can be interpreted as some “truth” available to some individuals [12] or a position shared by a group of close-minded opinion leaders (“radicals”) [13], [14]. Typically, the exogenous signal are supposed to change slowly compared to the opinion evolution and is thus replaced by a constant; the main concern is the dependence between the constant input and the resulting opinion profile. The second culprit of persistent opinion fluctuation is uncertainty in the opinion dynamics, usually modeled as a random drift of each opinion. Whereas these models are still waiting for clear sociopsychological interpretation, they are broadly adopted in statistical physics [15]–[18] to study phase transitions in systems of interacting particles.

In spite of some progress in analysis of “noisy” bounded confidence models [19], [20], in particular, the interplay of confidence ranges and noise levels, all consequences of a noise and exogenous influence in nonlinearly coupled networks are far from being understood. Even for the classical models from [10], [11], disclosing the relation between the initial and the terminal opinion profiles remains a challenging problem (including e.g. the *2R*-conjecture [21], [22]). This motivates examination of the corresponding *mean-field* models as the number of actors  $N \rightarrow \infty$ . The arising macroscopic approximations of microscopic models describe the evolution of the *distribution* (a probability measure or a density) of opinion over some domain. In continuous-time models (considered in this paper), the density obeys a nonlinear Fokker-Planck (FP) equation. This models are known as density-based [23], continuum-agent [24], [25], Eulerian [26], [27], kinetic [28] or mean-field [22], [29].

In this paper, we advance the theory of macroscopic modeling of noisy bounded confidence dynamics by considering “radical” opinions (equivalently, exogenous influences) that are not necessarily concentrated at a single point (as in [13], [26]) but have their own distribution. Following [22], [30], [31], we consider the model on a circle by considering *periodic* boundary conditions. However, we employ an *even 2-periodic* extension of the system in order to preserve the extreme opinions explicitly. Our main theoretical result shows that for sufficiently large noises the dynamic will converge exponentially fast to a stationary state that can be made arbitrarily close to uniform distribution by increasing

M.A.S. Kolarijani and P. Mohajerin Esfahani are with Delft Center for Systems and Control, Delft University of Technology, Delft, The Netherlands. A.V. Proskurnikov is with Department of Electronics and Telecommunications, Politecnico di Torino, Turin, Italy and also with the Institute for Problems of Mechanical Engineering, Russian Academy of Sciences (IPME RAS). M.A.SharifiKolarijani@tudelft.nl; P.MohajerinEsfahani@tudelft.nl; anton.p.1982@ieee.org.

the noise level.

Exploiting the periodic nature of the system, we derive a system of quadratic ordinary differential equations (ODEs) describing the evolution of the Fourier coefficients of the solution to the FP equation. For a uniform initial data, we use the linearized ODEs to study the order-disorder transition in the system. In particular, we provide a approximation scheme for computing the critical noise level at which this transition occurs. Our analysis shows that there is a direct relation between the mass of radicals and the critical noise level. This result is then validated by numerical simulations of the mean-field partial differential equation (PDE) and the corresponding agent-based stochastic differential equations (SDEs) for a particular distribution of radicals.

**Notations.** We use  $\star$  to denote the convolution of two functions. First and second order differentiation with respect to (w.r.t.)  $x$  are denoted by  $(\cdot)_x$  and  $(\cdot)_{xx}$ , respectively. Similar notations are used for differentiation w.r.t.  $t$ .  $\mathcal{P}(X)$  is the space of probability densities on  $X = [0, 1]$ , that is, for all  $\rho \in \mathcal{P}(X)$  we have  $\int_X \rho(x) dx = 1$  and  $\rho(x) \geq 0$ .  $\mathcal{P}_e(\tilde{X})$  is the space of *even* density functions on  $\tilde{X} = [-1, 1]$  such that for all  $\rho \in \mathcal{P}_e(\tilde{X})$  we have  $\int_{\tilde{X}} \rho(x) dx = 2$ ,  $\rho(x) \geq 0$  and  $\rho(x) = \rho(-x)$ . The subscript *ep* is used to denote the subspace of *even 2-periodic* functions in the corresponding function space; e.g.,  $L_{ep}^p(\tilde{X}) \subset L^p(\tilde{X})$  is the corresponding subspace of even 2-periodic functions on  $\tilde{X} = [-1, 1]$  for which the  $p$ -th power of the absolute value is Lebesgue integrable.

## II. MODEL

The model includes  $N$  interacting *normal* agents with opinions  $x_i, i = 1, \dots, N$ , where each agent  $i$  is influenced by agents  $j$  resided in its confidence bound, that is,  $j \in \mathcal{N}_i = \{j : |x_i - x_j| \leq R\}$ . We also consider exogenous influence in form of  $N_r$  *radical* agents with static opinions  $x_{r_i}, i = 1, \dots, N_r$ , introduced to the original population of normal agents. The same bounded confidence mechanism is assumed for interaction between a normal agent and the radicals. The model also includes noise representing the uncertainties [30] or the effect of free will [32] in agents' opinions. The following system of interacting SDEs describes the evolution of normal agents' opinions

$$\begin{cases} dx_i = \frac{1}{N} \left[ \sum_{j \in \mathcal{N}_i} (x_j - x_i) + \sum_{j \in \mathcal{N}_i} (x_{r_j} - x_i) \right] dt + \sigma dW_t^i, \\ x_i(0) = x_{i0}, \end{cases} \quad (1)$$

where  $x_{i0}, i = 1, 2, \dots, N$ , are the initial opinions of the normal agents and  $\sigma > 0$  is the noise level with  $W_t^i$  representing independent Wiener processes.

Following [30], we pass to the mean-field limit ( $N \rightarrow \infty$ ) of the agent-based model described by (1). Then, the density of the normal opinions  $\rho(x, t)$  satisfies the following nonlinear FP equation

$$\begin{cases} \rho_t = [\rho (w \star (\rho + M\rho_r))]_x + \frac{\sigma^2}{2} \rho_{xx}, \\ \rho(x, 0) = \rho_0(x), \end{cases} \quad (2)$$

where  $w(x) = x \mathbf{1}_{|x| \leq R}$  is the interaction kernel. The function  $\rho(x, t)$  is the limit measure of the empirical measure  $\rho^N(dx, t) = N^{-1} \sum_{i=1}^N \delta_{x_i(t)}(dx)$ , as  $N \rightarrow \infty$  (here,  $\delta_x(dx)$  denotes a Dirac measure centered at  $x$ ). The exogenous influence of radicals is included in the mean-field model by introducing the radical opinions density function  $M\rho_r(x)$  to (2). Here,  $M = \frac{N_r}{N}$  is a constant denoting the relative mass (the zeroth moment) of the radical opinions density and  $\rho_r(x)$  is the corresponding time-invariant probability density function.

We assume that the opinions reside in the bounded interval. Without loss of generality, we take  $X = [0, 1]$  to be the opinion domain and  $\tilde{X} = [-1, 1]$  as its even extension. Notice, however, the diffusion term in (1) and (2) allows the opinions to leave  $[0, 1]$ . To avoid this, following [17], [22], [30], [31], we consider a periodic boundary condition. However, unlike the usual periodic extensions, we consider an *even 2-periodic* extension of system. In effect, we treat the same mean-field model as in [31] with an extra constraint on  $\rho_0$  (and the newly introduced  $\rho_r$ ) to be even. This particular extension has the advantage that explicitly preserves the two extreme opinion values at  $x = 0$  and  $x = 1$ .

The corresponding PDE with the even 2-periodic extension is formally expressed as

$$\begin{cases} \rho_t = (\rho G_\rho)_x + \frac{\sigma^2}{2} \rho_{xx} & \text{in } \tilde{X} \times (0, T], \\ \rho(-1, \cdot) = \rho(1, \cdot) & \text{on } \partial \tilde{X} \times [0, T], \\ \rho(x, \cdot) = \rho_0(x) & \text{on } \tilde{X} \times \{t = 0\}, \end{cases} \quad (3)$$

where

$$G_\rho(x, t) := w(x) \star (\rho(x, t) + M\rho_r(x)). \quad (4)$$

Notice that  $\rho_0$  and  $\rho_r$  are the even 2-periodic extensions of the initial density of normal opinions and the radical opinions density, respectively. Due to periodicity, the mass is preserved in (3), that is,  $\int_{\tilde{X}} \rho(x, t) dx = \int_{\tilde{X}} \rho_0(x) dx = 1$  for all  $t \geq 0$ . PDE (3) fully describes the macroscopic model considered in this paper.

*Remark 1 (Well-posedness of dynamics):* The mean-field PDE (3) has a unique weak solution  $\rho \in C^1(0, \infty; C_{ep}^2(\tilde{X}))$  with  $\rho(t) \in \mathcal{P}_e(\tilde{X})$  for all  $t \geq 0$ . See [33, Theorem 1] for the details on the required regularity on data  $\rho_0$  and  $\rho_r$ .

## III. LONG-TERM BEHAVIOR

In this section, we study the long-term behavior of the system by considering the equilibria of system. To this end, we consider the corresponding stationary equation given by

$$\frac{\sigma^2}{2} \rho_{xx} + (\rho G_\rho)_x = 0. \quad (5)$$

In this regard, notice that, unlike the autonomous systems (without radicals) considered in previous studies [31], [34], the uniform distribution  $\rho = 1$  is not an equilibrium of the system. There, the authors provide sufficient condition for exponential convergence towards uniform distribution. Here, we extend this result to our model with the exogenous influence, i.e., the radicals.

*Remark 2 (Existence of equilibrium):* A fixed-point characterization of the solution to the stationary equation (5) can

be used to derive existence result for a classical solution  $\rho \in C_{ep}^2(\tilde{X}) \cap \mathcal{P}_e(\tilde{X})$  for equation (5). See [33, Theorem 2] for the details.

Let us start by presenting two main theoretical results regarding the long-term behavior of the system. The following Proposition provides an estimate that characterizes the equilibria of the system for noises larger than a particular value.

*Proposition 3 (Estimate for equilibrium):* Assume  $\rho_s \in C_{ep}^2(\tilde{X}) \cap \mathcal{P}_e(\tilde{X})$  is a classical solution to the stationary equation (5) with  $\rho_r \in L_{ep}^2(\tilde{X}) \cap \mathcal{P}_e(\tilde{X})$ . Then, it holds that  $\|\rho_s - 1\|_{L^2} < \frac{1}{\eta} \|\rho_r\|_{L^2}$  if  $\sigma^2 > \sigma_b^2 + \eta c_b$ , where

$$\sigma_b^2 := \frac{4R}{\pi} \left( M + \frac{R}{\sqrt{3}} + 2 \right) \text{ and } c_b := \frac{4R^2 M}{\pi \sqrt{3}}. \quad (6)$$

*Proof:* See [33, Section 4.2]. ■

Proposition 3 implies that, even in presence of radical opinions, the stationary solution can be made arbitrarily close to the uniform distribution by increasing the noise level. Particularly, one notices that the minimum noise level  $\sigma_b$  is directly related to the confidence bound  $R$  and the relative mass  $M$ . Also, as the “energy”  $M \|\rho_r\|_{L^2}$  of the radicals increases, in order to counteract their effect and keep the system in a somewhat uniform state, one must increase the noise level further beyond  $\sigma_b$ .

Proposition 3, by itself, cannot be used for describing the long-term behavior of the dynamics as it only characterizes the equilibria of the system. To put this result in use, we need to also consider the stability of these equilibria. Our next theoretical result provides a sufficient condition for exponential convergence of the dynamics to stationary state for arbitrary (and sufficiently smooth) initial and radical densities.

*Proposition 4 (Exponential convergence to equilibrium):* Let  $\rho \in C^1(0, \infty; C_{ep}^2(\tilde{X}))$  be the classical solution to PDE (3), with  $\rho(t) \in \mathcal{P}_e(\tilde{X})$  for all  $t \geq 0$ . Then,  $\rho(t)$  converges to a stationary state  $\rho_s \in C_{ep}^2(\tilde{X}) \cap \mathcal{P}_e(\tilde{X})$  exponentially in  $L^2$  as  $t \rightarrow \infty$  if  $\sigma > \sigma_s$ , where  $\sigma_s > 0$  uniquely solves

$$\sigma_s^2 = \frac{4R(3+M)}{\pi} + \frac{4R^2}{\pi \sqrt{3}} \exp\left(\frac{8R(1+M)}{\sigma_s^2}\right), \quad (7)$$

for any  $\rho_0 \in L_{ep}^2(\tilde{X}) \cap \mathcal{P}_e(\tilde{X})$  and  $\rho_r \in L_{ep}^2(\tilde{X}) \cap \mathcal{P}_e(\tilde{X})$ .

*Proof:* See [33, Section 4.3]. ■

Combining the results of Propositions 3 and 4, we obtain our main theoretical result.

*Theorem 5 (Input-output stability):* Assume PDE (3) has a classical solution  $\rho \in C^1(0, \infty; C_{ep}^2(\tilde{X}))$ , with  $\rho(t) \in \mathcal{P}_e(\tilde{X})$  for all  $t \geq 0$ . If  $\sigma^2 > \max\{\sigma_b^2 + \eta c_b, \sigma_s^2\}$ , where  $\sigma_b$  and  $c_b$  are defined in (6) and  $\sigma_s$  uniquely solves (7), then it holds that

$$\|\rho(t) - 1\|_{L^2} \leq \beta e^{-\lambda t} + \frac{1}{\eta} \|\rho_r\|_{L^2}, \quad (8)$$

where  $\beta, \lambda > 0$  depend on system data and initial condition.

*Proof:* Notice that

$$\|\rho(t) - 1\|_{L^2} \leq \|\rho(t) - \rho_s\|_{L^2} + \|\rho_s - 1\|_{L^2}.$$

Then, the inequality (8) immediately follows from Propositions 3 and 4. ■

This result implies that, for sufficiently large noises, the dynamics will converge to a stationary state that can be made arbitrarily close to uniform distribution by increasing the noise level. We also note that the exact dependence of the constants  $\beta$  and  $\lambda$  on the system data  $(R, \sigma, M, \rho_r)$  and initial condition can be found in the proofs of Propositions 3 and 4, available in [33].

#### IV. ORDER-DISORDER TRANSITION

A common behavior in noisy interactive particle systems is the so-called order-disorder transition. For large values of  $\sigma$ , the effect of diffusion process can overcome the attracting forces among agents preventing the system from forming any cluster. This behavior has been analyzed and observed in several noisy bounded confidence models for opinion dynamics. Pineda et. al. used linear stability analysis in [32], [35] to compute the critical noise level above which the clustering behavior disappears for a modified version of Defuant model. The same technique of linear stability analysis was used in [22] and [30] to compute the critical noise level for a noisy Hegselmann-Krause system similar to our model, except without radicals. In particular, [22] considered the interplay between the confidence bound  $R$  and the critical noise level  $\sigma_c$  at which the system experiences the order-disorder transition.

In this section, we study the effect of the relative mass  $M$  of radicals on the critical noise level  $\sigma_c$ . We emphasize that we consider a uniform distribution for initial normal opinions. In the sequel, we make use of the *order parameter* [22]

$$\begin{cases} Q_d(x) = \frac{1}{N^2} \sum_{i,j=1}^N \mathbf{1}_{|x_i(t) - x_j(t)| \leq R}, \\ Q_c(\rho) = \iint_{X^2} \rho(x, t) \rho(y, t) \mathbf{1}_{|x-y| \leq R} dx dy, \end{cases} \quad (9)$$

to quantify the clustering behavior of the agent-based and mean-field models, respectively. The order parameter provides a measure for orderedness in opinions: for a uniform distribution of opinions - absolute disorder, we have  $Q = 0.2$ ; and, for a single cluster distribution with all agents residing in an interval of width  $R$  or less - complete order, we have  $Q = 1$ . Roughly speaking, in case of emergence of a clustered profile, the inverse of the order parameter is equal to the number of clusters.

##### A. Approximating the Critical Noise Level $\sigma_c$

In what follows, we provide a method for approximating the critical noise level  $\sigma_c$ . To do so, we exploit the periodic nature of the system and derive the ODEs describing the evolution of the Fourier coefficients of the solution  $\rho$  to (3). Then, we use the linearized ODEs to present a numerical scheme for approximating  $\sigma_c$ , without actually solving the dynamics.

1) *Fourier ODEs:* The even 2-periodic extension of the probability densities in the model allows us to consider the Fourier expansions of  $\rho$  and  $\rho_r$  expressed as  $\rho(x, t) = \sum_{n=0}^{\infty} p_n(t) \cos(\pi n x)$  and  $\rho_r(x) = \sum_{n=0}^{\infty} q_n \cos(\pi n x)$ , respectively. Inserting these expansions into (3) and setting the

inner product of the residual with elements of the basis to zero, we can obtain a system of quadratic ODEs describing the evolution of Fourier coefficients  $p_n(t)$ . For  $n = 1, \dots, N_f$ , these ODEs are expressed as

$$\dot{p}_n = c_n + b_n^T p + p^T Q_n p, \quad (10)$$

where  $p = (p_1, p_2, \dots, p_{N_f})^T$ . Note that for  $n = 0$  (the constant term in the Fourier expansion), we obtain  $\dot{p}_0 = 0$  which corresponds to the periodic nature of the system that preserves the zeroth moment. The coefficients of (10) are given by

$$c_n := 2MR f_n q_n, \quad (11)$$

$$(b_n)_k := \begin{cases} 2R f_n + \frac{MR}{2} f_{2n} q_{2n} - \frac{\pi^2 \sigma^2 n^2}{2}, & k = n, \\ nMR \left\{ \frac{q_{n+k} f_{n+k}}{n+k} + \frac{q_{|n-k|} f_{n-k}}{n-k} \right\}, & k \neq n, \end{cases} \quad (12)$$

$$(Q_n)_{k,l} := \begin{cases} nR \frac{f_k}{k}, & l = n - k > 1, \\ nR \left\{ \frac{f_k}{k} + \frac{f_{n-k}}{n-k} \right\}, & l = k - n > 1, \\ 0, & \text{Otherwise,} \end{cases} \quad (13)$$

where  $q_n, n \in \mathbb{N}$  are the Fourier coefficients of  $\rho_r$  and  $f_n := -\cos(n\pi R) + \text{sinc}(n\pi R)$ .

2) *Approximation Scheme*: Linearizing the system at  $t = 0$  (which is equivalent to disregarding the quadratic terms in (10) since the initial condition is uniform, i.e.,  $p_n(0) = 0$  for  $n \neq 0$ ), we obtain the system of linear ODEs

$$\dot{p} = c + Bp, \quad (14)$$

where the vector  $c$  and matrix  $B$  are defined accordingly using the objects  $c_n$  and  $b_n$  in (11) and (12) for  $n = 1, \dots, N_f$ .

Looking at coefficients of the quadratic ODEs (10), we notice that the noise level  $\sigma$  only appears in the diagonal entries of  $B$  such that by increasing  $\sigma$ , these diagonal entries decrease. That is, for a large enough  $\sigma$ , all eigenvalues of  $B$  are negative and the linearized system (14) is stable. This will be our first criterion for determining the critical noise level  $\sigma_c$ : the noise level above which all eigenvalues of  $B$  are negative.

In order to consider the effect of the constant linear growth rates  $c$  in (14), we further require the stationary values  $\bar{p}_n, n = 1, \dots, N_f$  of the linearized system (14) (i.e., the solutions of the equations  $c + Bp = 0$ ) to be relatively small. In order to quantify this description, we use Parseval's identity to set our second criterion as

$$\sum_{n=1}^{N_f} \bar{p}_n^2 < \gamma, \quad (15)$$

where the constant  $\gamma > 0$  determines the level of similarity between  $\rho$  and uniform distribution (disordered state). To sum up, we solve numerically for the level of noise above which the eigenvalues of  $B$  are negative and the inequality (15) holds.

## B. Numerical Simulations

In this section, we numerically study the effect of the relative mass of radicals  $M$  on the order-disorder transition. Furthermore, we use our simulation results to examine the

approximation scheme presented in Section IV-A.2. To this end, we consider a particular distribution of radical opinions, namely, a triangular one with average  $A$  and width  $2S$

$$\rho_r(x) = (S - |x - A|)/S^2 \text{ for } |x - A| \leq S. \quad (16)$$

The corresponding Fourier coefficients of  $\rho_r$  are given by

$$q_n = 2 \cos(n\pi A) \text{sinc}^2(n\pi S/2). \quad (17)$$

In all the simulations of this section, the average and width of radical distribution are fixed at  $A = 0.7$  and  $S = 0.1$ , respectively.

In order to solve the mean-field model described by PDE (3) numerically, we use the ODEs (10) to compute the coefficients of Fourier expansion of normal opinion density  $\rho$  cut at  $N_f = 128$ . In particular, we use the Fourier coefficients of the triangular  $\rho_r$  given in (17) for solving ODEs (10). To be precise, we need the Fourier coefficients  $q_n$  of  $\rho_r$  for  $1 \leq n \leq 2N_f$ , that is, twice the length of Fourier expansion of  $\rho$  (see the coefficients in (12)). For the initial condition, we consider uniform distribution  $\rho_0 = 1$  which corresponds to  $p_0 = 1$  and  $p_n(0) = 0$  for the Fourier coefficients.

For the agent-based model, the SDEs (1) are solved numerically using the Euler-Maruyama method for  $N = 500$  normal agents with time step  $\Delta t = 0.01$ . For the radical agents, we produce a random sample of size  $N_r = MN$  from the triangular distribution (16). The initial opinions of normal agents are randomly sampled from a uniform distribution on the interval  $[0, 1]$ . For complete correspondence between the agent-based and mean-field models, we also consider the effect of even 2-periodic extension in the simulations of the agent-based model.


1) *An Illustrative Example*: Fig. 1 shows the result of numerical simulation of the mean-field and agent-based models for different noise levels with the relative mass of radicals fixed at  $M = 0.15$ . Indeed, for  $\sigma$  larger than a critical level the clustering behavior almost disappears (see the lower panel corresponding to  $\sigma = 0.05$  in Fig. 1a).

In this regard, we observe that as the level of noise increases, the number of clusters in the possible clustering behavior of the system decreases. To be more precise, a higher level of noise decreases the life-time of clustering behaviors with larger number of clusters. This effect can be particularly seen in the evolution of order parameter in Fig. 1b. In this regard, notice that, for noises smaller than the critical noise level (here,  $\sigma = 0.015, 0.030, 0.045$ ) the flat areas in the order parameter in Fig. 1b correspond to a clustered behavior where the number of clusters is equal to the inverse of the order parameter. To illustrate, observe that for  $\sigma = 0.03$  and  $\sigma = 0.04$ , the system reaches a single-cluster profile around the average radical opinion  $A = 0.7$ . Notice, however, for  $\sigma = 0.03$  the system goes through a 2-cluster profile as depicted in Fig. 1b (the almost flat areas in the order parameter). For  $\sigma = 0.015$ , we observe 4-cluster and 3-cluster profiles in mean-field and agent-based models, respectively, at  $t = 10^4$ . This particular difference between mean-field and agent-based models has been also mentioned and explained by [22]. Finally, we also note that for  $M =$


0.15, the approximation scheme results in  $\sigma_c = 0.043$  for  $\gamma = 1$  and  $\sigma_c = 0.051$  for  $\gamma = 0.1$  (see (15) for influence of  $\gamma$ ).

2) *Effect of  $M$  on  $\sigma_c$* : Fig. 2 shows the order parameter derived numerically by simulating the mean-field and agent-based models. Notice how for each  $M$ , the system experiences a transition from order (clustered phase with  $Q = 1$  in the yellow strip) to the disorder (with  $Q = 0.2$  in the blue area in the upper part of the plots) as noise increases. Also, we note that the blue strip in the lower part of plots in Fig. 2 represents clustering behaviors with larger number of clusters (similar to the behavior seen for  $\sigma = 0.015$  in Fig. 1).

This result shows that as the relative mass of radicals  $M$



(a) Distribution of opinions/agents at  $t = 10^4$



(b) Evolution of order parameter (9)


Fig. 1. Numerical simulation of the mean-field (MF) and agent-based (AB) models for increasing values of noise  $\sigma$  with  $M = 0.15$  and uniform initial state. See Section IV-B.1.

increases, the corresponding critical noise level  $\sigma_c$ , above which the system is in disordered state, also increases. The dependence of  $\sigma_c$  on  $M$  is in the form of a concave function. Furthermore, for small values of  $M$ , the transition seems to be discrete, signaling a first-order transition. However, for large values of  $M$  the transition becomes blurry. This phenomenon was also observed and reported in [22] for the dependence of the critical noise level on the confidence bound  $R$ . Notice that as  $M$  increases, the required noise level for disordered behavior also increases. This increase in the noise level leads to wider clusters (see Fig. 1) which, in turn, makes differentiating the clustered (ordered) and not clustered (disordered) behaviors difficult.


Also shown in Fig. 2 (red lines) is the result of scheme provided in Section IV-A.2 for approximating the critical noise level. As can be seen, the scheme indeed provides a good approximation of the critical noise level, in particular, for  $\gamma = 1$ .

## V. CONCLUSIONS

In this paper, we considered a mean-field model for bounded confidence opinion dynamics with environmental noise. The model also included exogenous influence by



(a) Mean-field model (PDE)



(b) Agent-based model (SDE)

Fig. 2. The order parameter (9) at  $t = 10^3$  from numerical simulation of the mean-field and agent-based models starting from uniform initial distribution. For the Agent-based model, the average of order parameter over the time window  $[900, 1000]$  is reported. The plot covers the region  $\sigma \times M \in [0.01, 0.15] \times [0.01, 1]$  with step sizes  $\Delta\sigma = 0.005$  and  $\Delta M = 0.02$ . The red lines show the result of the numerical scheme described in Section IV-A.2 for approximating the critical noise level for different values of  $\gamma$  w.r.t. the second criterion (15). See Section IV-B.2.

adding a mass of radical (continuum) agents to the original population of the normal agents. Two main theoretical results were provided that characterized the long-term behavior of the system. In particular, we showed that, the effect of radical agents is restrained in systems with higher levels of noise, that is, when agents are more inclined to exercise their “free will” in exploring the opinion space. We note that the lower bound on the noise level is quite conservative, considering its theoretical nature. This can be seen by comparing the theoretical bound of Theorem 5 with the numerical one provided for the critical noise level for order-disorder transition in Section IV.

In Section IV, we also used Fourier analysis to provide a numerical scheme for approximating the critical noise level. This scheme was then validated through numerical simulations of the mean-field model and the corresponding agent-based model. We note that the proposed numerical scheme drastically reduces the effort for computing the critical noise level compared to application of standard numerical approaches for solving the PDE and the corresponding interacting SDEs. As expected (and also shown by the theoretical results), the numerical simulations revealed a higher critical noise level for a larger mass of radicals.

## REFERENCES

- [1] R. Abelson, “Mathematical models of the distribution of attitudes under controversy,” in *Contributions to Mathematical Psychology*, N. Frederiksen and H. Gulliksen, Eds. New York: Holt, Rinehart & Winston, Inc, 1964, pp. 142–160.
- [2] T. Kurahashi-Nakamura, M. Mäs, and J. Lorenz, “Robust clustering in generalized bounded confidence models,” *Journal of Artificial Societies and Social Simulation*, vol. 19, no. 4, p. 7, 2016.
- [3] N. Friedkin, “The problem of social control and coordination of complex systems in sociology: A look at the community cleavage problem,” *IEEE Control Systems Magazine*, vol. 35, no. 3, pp. 40–51, 2015.
- [4] C. Castellano, S. Fortunato, and V. Loreto, “Statistical physics of social dynamics,” *Reviews of Modern Physics*, vol. 81, pp. 591–646, 2009.
- [5] H. Xia, H. Wang, and Z. Xuan, “Opinion dynamics: A multidisciplinary review and perspective on future research,” *International Journal of Knowledge and Systems Science*, vol. 2, no. 4, pp. 72–91, 2011.
- [6] D. Acemoglu and A. Ozdaglar, “Opinion dynamics and learning in social networks,” *Dynamic Games and Applications*, vol. 1, pp. 3–49, 2011.
- [7] A. Proskurnikov and R. Tempo, “A tutorial on modeling and analysis of dynamic social networks. Part I,” *Annual Reviews in Control*, vol. 43, 2017, 65–79.
- [8] —, “A tutorial on modeling and analysis of dynamic social networks. Part II,” *Annual Reviews in Control*, vol. 45, 2018, 166–190.
- [9] M. McPherson, L. Smith-Lovin, and J. Cook, “Birds of a feather: Homophily in social networks,” *Annual Review of Sociology*, vol. 27, pp. 415–444, 2001.
- [10] U. Krause, “A discrete nonlinear and non-autonomous model of consensus formation,” *Communications in Difference Equations*, vol. 2000, pp. 227–236, 2000.
- [11] G. Deffuant, D. Neau, F. Amblard, and G. Weisbuch, “Mixing beliefs among interacting agents,” *Advances in Complex Systems*, vol. 3, no. 01n04, pp. 87–98, 2000.
- [12] R. Hegselmann and U. Krause, “Truth and cognitive division of labour: First steps towards a computer aided social epistemology,” *Journal of Artificial Societies and Social Simulation*, vol. 9, no. 3, p. 1, 2006.
- [13] —, “Opinion dynamics under the influence of radical groups, charismatic leaders and other constant signals: a simple unifying model,” *Networks & Heterogeneous Media*, vol. 10, no. 3, pp. 477–509, 2015.
- [14] Y. Zhao, L. Zhang, M. Tang, and G. Kou, “Bounded confidence opinion dynamics with opinion leaders and environmental noises,” *Computers & Operations Research*, vol. 74, pp. 205–213, 2016.
- [15] E. Ben-Naim, P. Krapivsky, and S. Redner, “Bifurcations and patterns in compromise processes,” *Physica D*, vol. 183, no. 3, pp. 190–204, 2003.
- [16] S. Grauwint and P. Jensen, “Opinion group formation and dynamics: Structures that last from nonlasting entities,” *Physical Review E*, vol. 85, p. 066113, 2012.
- [17] M. Pineda, R. Toral, and E. Hernández-García, “The noisy hegselmann-krause model for opinion dynamics,” *The European Physical Journal B*, vol. 86, no. 12, p. 490, 2013.
- [18] A. Carro, R. Toral, and M. San Miguel, “The role of noise and initial conditions in the asymptotic solution of a bounded confidence, continuous-opinion model,” *Journal of Statistical Physics*, vol. 151, pp. 131–149, 2013.
- [19] M. Huang and J. H. Manton, “Opinion dynamics with noisy information,” in *52nd IEEE Conference on Decision and Control*, 2013, pp. 3445–3450.
- [20] W. Su, G. Chen, and Y. Hong, “Noise leads to quasi-consensus of Hegselmann–Krause opinion dynamics,” *Automatica*, vol. 85, pp. 448–454, 2017.
- [21] V. D. Blondel, J. M. Hendrickx, and J. N. Tsitsiklis, “On the 2r conjecture for multi-agent systems,” in *2007 European Control Conference (ECC)*, 2007, pp. 874–881.
- [22] C. Wang, Q. Li, W. E, and B. Chazelle, “Noisy hegselmann-krause systems: Phase transition and the 2R-conjecture,” *Journal of Statistical Physics*, vol. 166, no. 5, pp. 1209–1225, 2017.
- [23] J. Lorenz, “Continuous opinion dynamics under bounded confidence: a survey,” *International Journal of Modern Physics C*, vol. 18, no. 12, pp. 1819–1838, 2007.
- [24] V. Blondel, J. Hendrickx, and J. Tsitsiklis, “Continuous-time average-preserving opinion dynamics with opinion-dependent communications,” *SIAM Journal on Control and Optimization*, vol. 48, pp. 5214–5240, 2010.
- [25] J. Hendrickx and A. Olshevsky, “On symmetric continuum opinion dynamics,” *SIAM Journal on Control and Optimization*, vol. 54, no. 5, pp. 2893–2918, 2016.
- [26] A. Mirtabatabaei, P. Jia, and F. Bullo, “Eulerian opinion dynamics with bounded confidence and exogenous inputs,” *SIAM Journal on Applied Dynamical Systems*, vol. 13, no. 1, pp. 425–446, 2014.
- [27] C. Canuto, F. Fagnani, and P. Tilli, “An Eulerian approach to the analysis of Krause’s consensus models,” *SIAM Journal on Control and Optimization*, vol. 50, pp. 243–265, 2012.
- [28] L. Boudin and F. Salvarani, “Opinion dynamics: Kinetic modelling with mass media, application to the Scottish independence referendum,” *Physica A*, vol. 444, pp. 448–457, 2016.
- [29] A. Nordio, A. Tarable, C.-F. Chiasserini, and E. Leonardi, “Belief dynamics in social networks: A fluid-based analysis,” *IEEE Transactions on Network Science and Engineering*, vol. 5, no. 4, pp. 276–287, 2018.
- [30] J. Garnier, G. Papanicolaou, and T.-W. Yang, “Consensus convergence with stochastic effects,” *Vietnam Journal of Mathematics*, vol. 45, no. 1, pp. 51–75, 2017.
- [31] B. Chazelle, Q. Jiu, Q. Li, and C. Wang, “Well-posedness of the limiting equation of a noisy consensus model in opinion dynamics,” *Journal of Differential Equations*, vol. 263, no. 1, pp. 365 – 397, 2017.
- [32] M. Pineda, R. Toral, and E. Hernández-García, “Noisy continuous-opinion dynamics,” *Journal of Statistical Mechanics: Theory and Experiment*, vol. 2009, no. 08, p. P08001, 2009.
- [33] M. A. S. Kolarijani, A. V. Proskurnikov, and P. Mohajerin Esfahani, “Macroscopic noisy bounded confidence models with distributed radical opinions,” *arXiv preprint arXiv:1905.04057*, 2019.
- [34] J. A. Carrillo, R. S. Gvalani, G. A. Pavliotis, and A. Schlichting, “Long-time behaviour and phase transitions for the mckean-vlasov equation on the torus,” *arXiv preprint arXiv:1806.01719*, 2018.
- [35] M. Pineda, R. Toral, and E. Hernandez-Garcia, “Diffusing opinions in bounded confidence processes,” *The European Physical Journal D*, vol. 62, no. 1, pp. 109–117, 2011.