

Photon Quantum Computer. Example: Mach-Zehnder interferometer.

A photon passes through the first beam splitter (BS1) and propagates via 2 different paths to another beam splitter (BS2), which directs the particle to one of the two detectors. Along each path between the two BSs, there is a phase shifter (PS i : $i = 1, 2$).




Figure 0.1: Mach-Zehnder interferometer. Pr_{D_i} is the probability of directing the particle to detector D_i .

If the initial particle is in path $|0\rangle$, it undergoes the following:

$$\begin{aligned}
 |\psi(t_0)\rangle &= |0\rangle \rightarrow \hat{U}_{\text{BS1}} |0\rangle = \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) \rightarrow \hat{U}_{\text{PS}}(\hat{U}_{\text{BS1}} |0\rangle) = \frac{1}{\sqrt{2}}(e^{i\phi_0} |0\rangle + e^{i\phi_1} |1\rangle) \\
 &= \frac{1}{\sqrt{2}} e^{i\frac{\phi_0+\phi_1}{2}} (e^{i\frac{\phi_0-\phi_1}{2}} |0\rangle + e^{-i\frac{\phi_0-\phi_1}{2}} |1\rangle) \rightarrow \\
 |\psi(t_f)\rangle &= \hat{U}_{\text{BS2}} \left(\hat{U}_{\text{PS}}(\hat{U}_{\text{BS1}} |0\rangle) \right) = \frac{1}{\sqrt{2}} e^{i\frac{\phi_0+\phi_1}{2}} \left(e^{i\frac{\phi_0-\phi_1}{2}} \frac{1}{\sqrt{2}}(|0\rangle + |1\rangle) + e^{-i\frac{\phi_0-\phi_1}{2}} \frac{1}{\sqrt{2}}(|0\rangle - |1\rangle) \right) \\
 &= \frac{1}{2} e^{i\frac{\phi_0+\phi_1}{2}} \left(2 \cos\left(\frac{\phi_0-\phi_1}{2}\right) |0\rangle + 2i \sin\left(\frac{\phi_0-\phi_1}{2}\right) |1\rangle \right) = e^{i\frac{\phi_0+\phi_1}{2}} \left(\cos\frac{\Delta\phi}{2} |0\rangle + i \sin\frac{\Delta\phi}{2} |1\rangle \right), \tag{0.1}
 \end{aligned}$$

where $\Delta\phi \equiv \phi_0 - \phi_1$. BS1 prepares the particle in a superposition of paths, PS1 and PS2 modify quantum phases and BS2 recombines the superpositions erasing the information about which path was taken from BS1.

The global phase $e^{i\frac{\phi_0+\phi_1}{2}}$, as mentioned in Section (2.3.3), is irrelevant, since it vanishes when we measure Pr_{D_i} , $i = 1, 0$:

$$\text{Pr}_{D_1} = \text{Pr}_{|\psi(t_f)\rangle}(|0\rangle) = |\langle 0|\psi(t_f)\rangle|^2 = |e^{i\frac{\phi_0+\phi_1}{2}} \cos\frac{\Delta\phi}{2}|^2 = \cos^2\frac{\Delta\phi}{2}, \tag{0.2}$$

$$\text{and } \text{Pr}_{D_2} = \text{Pr}_{|\psi(t_f)\rangle}(|1\rangle) = |\langle 1|\psi(t_f)\rangle|^2 = |e^{i\frac{\phi_0+\phi_1}{2}} \sin\frac{\Delta\phi}{2}|^2 = \sin^2\frac{\Delta\phi}{2}. \tag{0.3}$$

Notice that **this is a quantum circuit!** $\hat{U}_{\text{BS}} \equiv \hat{U}_{\text{Had}}$ and $\hat{U}_{\text{PS}} \equiv \hat{U}_{\phi} \doteq \bigoplus_{k=0}^{n-1} e^{i\phi_k} = \begin{pmatrix} e^{i\phi_0} & \\ & e^{i\phi_1} \end{pmatrix}$.

Hence, $|\psi(t_f)\rangle = \hat{U}_{\text{Had}} \hat{U}_{\phi} \hat{U}_{\text{Had}} |\psi(t_0)\rangle$.