



Numerical Analysis

Newton's method

Submitted by,

Asif Istiak; ID: 20205083

Mechanical Design Engineering

Andong National University

Submitted to,

Professor See Jo Kim

Mechanical Design Engineering

Andong National University


Contents

- **Abstract**
- **Introduction**
- **Examples**
- **Concluding Remarks**
- **Reference**

Abstract

Target:


- ❖ Define the Newton's Method
- ❖ Code explanation
- ❖ Result discussion



Introduction

Newton's Method:

- ❖ Mathematical View
- ❖ Terms of codes
- ❖ Results



[Wolfgang Bangerth's video lectures \(colostate.edu\)](http://colostate.edu)

Newton's Method

Solver Selection

$$A_k \delta U_k = F_k$$

$$A_{k,ij} = \left(\nabla \phi_i, \frac{A}{\sqrt{1+|\nabla u_{k,h}|^2}} \nabla \phi_j \right) - \left(\nabla \phi, \frac{A(\nabla u_{k,h} \cdot \nabla \phi_j)}{(1+|\nabla u_{k,h}|^2)^{3/2}} \nabla u_k \right)$$

Solver

Conjugate Gradient

A is symmetric- (derivative of a non-linear operator)

A is Positive- (convex energy function is minimized)

Newton's Method

Boundary Values of δu_k

$$u_0|_{\partial\Omega} = g$$

$$(u_k + \alpha_k \delta u_k)|_{\partial\Omega} = g$$

δu_k have zero boundary

Way of Convergence

$$U_{k+1} = U_k + \delta U_k$$

Does not follow this rule always

$$U_{k+1} = U_k + \alpha_k \delta U_k$$

Relaxing the iteration, it takes

Line search- algorithm for α_k

quadratic convergence if $\alpha_k = 1$

may not converge if $\alpha_k = 1$

slower convergence if $\alpha_k < 1$

frequently converge if $\alpha_k < 1$

α_k depends on residual function

$$R(u) = L(u) - f = 0$$

If

$$\|R(u_k + \alpha_k \delta u_k)\| \leq c \|R(u_k)\|$$

then use this α_k ; otherwise set $\alpha_k := \alpha_k/2$ and try again

$\alpha_k = 0.1$ is taken for simplicity

Compute

$$U_{k+1} = U_k + \alpha_k \delta U_k$$

Newton's Method

Implementation

1. Start with a coarse mesh
2. Do Newton iterations
3. If $\|R(u_k)\| \leq \text{tol}$ then stop
4. Save the solution on the current mesh
5. Refine the mesh
6. Go to step 2

Result:

Took a long time to run(could not visualize). Here are the results.

```
Residual: 0.00508832
Residual: 0.00457899
Residual: 0.00412073
Residual: 0.0037084
Residual: 0.00333737
***** Refined mesh 11 *****
Initial residual: 0.00342093
Residual: 0.00307852
Residual: 0.00277045
Residual: 0.00249323
Residual: 0.00224379
Residual: 0.00201932
***** Refined mesh 12 *****
Initial residual: 0.00202458
Residual: 0.00182205
Residual: 0.00163979
Residual: 0.00147577
Residual: 0.00132816
Residual: 0.00119532
***** Refined mesh 13 *****
Initial residual: 0.00115921
Residual: 0.00104326
Residual: 0.000938907
```

Conclusion

- ❖ Solver selection
- ❖ Methodology
- ❖ Conditionals

Thank You