

Joanna Masikowska

B9TB1710

In the exercise, I have given a number of events happening on average in given time span. In my case it is 1.4 earthquakes in 8 days. To find probability that at least given number of earthquakes will happen within 28 days, I will use Poisson distribution.

```
CAPS09_B9TB1710.m
1 A=1;
2 B=7;
3 C=1;
4 D=0;
5 l=(28/(B+1))*0.7*(A+1); #number of events in 28 days on average
6 k=10+C+D; #number of events in 28days
7 Y=randp(l,1,10000);
8 X=Y(Y>=k);
9 P=length(X)/length(Y)
```

Firstly, using property of Poisson distribution, I will multiply 1.4 *earthquakes/8 days* by 28/8 to get the average number of the event in 28 days. That gives me 4.9 events which I assign to variable *l*. Next, I assign the minimal number of events in 28 days, which probability I want to

find, to variable *k*. In my case, $k = 11$. Using function **randp** I generate a matrix *Y* consisting of 10000 numbers of events that happened in 28 days according to Poisson distribution. Then, I find how many numbers higher or equal to *k* are in *Y*. I do that by putting numbers $\geq k$ in the matrix *X*. Then, I calculate the probability by dividing the length of *X* by the length of *Y*. This way I get the probability that *k* or more events will happen within 28 days. After running the code a few times, I can see that the probability is around 0.011~0.013, which is small.

```
P = 0.014300
>> CAPS09_B9TB1710

P = 0.011400
>> CAPS09_B9TB1710

P = 0.012300
>> CAPS09_B9TB1710

P = 0.013600
>> CAPS09_B9TB1710

P = 0.012400
>> CAPS09_B9TB1710

P = 0.013300
>> CAPS09_B9TB1710

P = 0.011600
>> CAPS09_B9TB1710

P = 0.011800
>> CAPS09_B9TB1710

P = 0.011500
>> CAPS09_B9TB1710

P = 0.013000
>> CAPS09_B9TB1710

P = 0.013500
>> CAPS09_B9TB1710

P = 0.011100
>> CAPS09_B9TB1710

P = 0.012400
```