

**MA 102 (Mathematics II)**  
**IIT Guwahati**

Tutorial Sheet No. 5

Linear Algebra

February 21, 2019

---

1. True or False? Give justifications.

- (a) There exist distinct linear transformations  $S, T : \mathbb{V} \rightarrow \mathbb{W}$  such that  $\ker(S) = \ker(T)$  and  $\text{range}(S) = \text{range}(T)$ .
- (b) There exists a linear transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that none of  $T, T^2, T^3$  is the identity transformation but  $T^4 = I$  (identity transformation).
- (c) If  $T : \mathbb{V} \rightarrow \mathbb{W}$  is a linear transformation then  $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$  is LI in  $\mathbb{V}$  if and only if  $\{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$  is LI in  $\mathbb{W}$ .

**Solution:**

- (a) True. Consider  $S, T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $S([x, y]^\top) = [x, y]^\top$  and  $T([x, y]^\top) = [y, x]^\top$ .
- (b) True. Rotate every element of  $\mathbb{R}^2$  by 90 degrees, that is,  $T([x, y]^\top) = \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} [x, y]^\top$ .
- (c) False. If  $\{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$  is LI in  $\mathbb{W}$  then  $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$  is LI in  $\mathbb{V}$  but the converse is not true. For instance consider the  $\mathbf{0}$  transformation.

2. Determine a linear transformation from  $\mathbb{R}^3 \rightarrow \mathbb{R}^3$  such that  $\text{range}(T) = \{[x, y, z]^\top : x + 2y + z = 0\}$ . If possible give two more such linear transformations with the same range.

**Solution:** It is enough to define  $T$  on a basis.

Consider the basis  $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3\}$  of  $\mathbb{R}^3$  and define the LT  $T$  such that  $T(\mathbf{e}_1) = [2, -1, 0]^T$ ,  $T(\mathbf{e}_2) = [0, -1, 2]^T$  and  $T(\mathbf{e}_3) = \mathbf{0}$ . Then  $\text{range}(T) = \text{span}\{T(\mathbf{e}_1), T(\mathbf{e}_2)\}$ .

Note that  $\{[2, -1, 0]^T, [0, -1, 2]^T\}$  is a basis of  $S = \{[x, y, z]^T : x + 2y + z = 0\}$ .

Hence  $\text{range}(T) = \text{span}\{T(\mathbf{e}_1), T(\mathbf{e}_2)\} = S$  and  $\text{Ker}(T) = \text{span}\{\mathbf{e}_3\}$ .

If  $T(\mathbf{e}_3) = \alpha(2, -1, 0)^T + \beta(0, -1, 2)^T$  then again  $\text{range}(T)$  is same but  $\text{Ker}(T)$  is different. For the same basis by making suitable choices one can also get  $\text{Ker}(T)$  as the  $x$ -axis or the  $y$ -axis. By considering different basis of  $\mathbb{R}^3$ , one can get many more  $T$ 's.

3. If possible, find linear transformations  $S : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$  and  $T : \mathbb{R}_2[x] \rightarrow \mathbb{R}^2$  such that

- (a)  $S \circ T = I$ .
- (b)  $T \circ S = I$ .
- (c)  $\text{range}(T \circ S)$  is a line.
- (d) Neither  $S$  nor  $T$  is the zero transformation but  $S \circ T = \mathbf{0}$ .

**Solution:**

(a)  $S \circ T : \mathbb{R}_2[x] \rightarrow \mathbb{R}_2[x]$ .

Note that  $\text{range}(S \circ T) \preceq \text{range}(S)$  and  $\text{rank}(S) \leq 2$  (from the rank nullity theorem). Hence  $\text{rank}(S \circ T) \leq 2$ . An identity map from  $\mathbb{R}_2[x] \rightarrow \mathbb{R}_2[x]$  will have rank 3, hence not possible.

(b)  $T \circ S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ .

Define  $S([a, b]^\top) = a + bx$  and  $T : \mathbb{R}_2[x] \rightarrow \mathbb{R}^2$  by  $T(a_0 + a_1x + a_2x^2) = [a_0, a_1]^\top$ . Then we have  $T \circ S = I$ .

(c) Define  $S([a, b]^\top) = a + bx$  and  $T : \mathbb{R}_2[x] \rightarrow \mathbb{R}^2$  by  $T(a_0 + a_1x + a_2x^2) = [a_0, 0]^\top$ . Then  $\text{range}(T \circ S)$  is a line.

(d) Consider  $T(a_0 + a_1x + a_2x^2) = [a_0, 0]^\top$  and  $S([a, b]^\top) = bx$ .

4. Let  $\mathbb{V}, \mathbb{W}$  be finite dimensional vector spaces with ordered bases  $B$  and  $C$ , respectively. Let  $T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ . Show that  $\text{rank}(T) = \text{rank}([T]_{C \leftarrow B})$  and  $\text{nullity}(T) = \text{nullity}([T]_{C \leftarrow B})$ .

**Solution:** For  $\mathbf{v} \in \mathbb{V}$ , we have  $T\mathbf{v} = C[T\mathbf{v}]_C = C[T]_{C \leftarrow B}[\mathbf{v}]_B$ . Since  $\mathbf{v} \mapsto [\mathbf{v}]_B$  is an isomorphism, it follows that  $T\mathbf{v} = \mathbf{0} \Leftrightarrow [T]_{C \leftarrow B}[\mathbf{v}]_B = \mathbf{0}$ . Hence  $\text{nullity}(T) = \text{nullity}([T]_{C \leftarrow B})$ .

Next, let  $B$  be given by  $B = [\mathbf{v}_1, \dots, \mathbf{v}_n]$ . Then  $T\mathbf{v}_j = C[T]_{C \leftarrow B}\mathbf{e}_j$  shows that  $\text{rank}(T) = \text{rank}([T]_{C \leftarrow B})$ .

5. True or False? Give justifications.

- (a) A transformation  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  defined as  $T([x, y]^\top) = [x, y]^\top$  for  $x \neq 0$  and  $T([0, y]^\top) = [0, 0]^\top$  satisfies  $T(c[x, y]^\top) = cT([x, y]^\top)$  but is not a linear transformation.
- (b) Let  $\mathbb{V}$  and  $\mathbb{W}$  be vector spaces. Then for any  $\mathbf{v}_1, \mathbf{v}_2$  in  $\mathbb{V}$  and  $\mathbf{w}_1, \mathbf{w}_2$  in  $\mathbb{W}$ , there exists a linear transformation  $T : \mathbb{V} \rightarrow \mathbb{W}$  such that  $T(\mathbf{v}_1) = \mathbf{w}_1$  and  $T(\mathbf{v}_2) = \mathbf{w}_2$ .
- (c) Let  $\mathbb{V}$  and  $\mathbb{W}$  be  $n$ -dimensional vector spaces and  $T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$  be invertible. Then there exist ordered bases  $B$  and  $C$  of  $\mathbb{V}$  and  $\mathbb{W}$ , respectively, such that  $[T]_{C \leftarrow B} = I_n$ .

**Solution:**

(a) True.  $T(c[x, y]^\top) = cT([x, y]^\top)$  is easy to check. But for  $y \neq 0$ ,  $T([-1, y]^\top) + T([1, y]^\top) = [-1, y]^\top + [1, y]^\top = [0, 2y]^\top \neq [0, 0]^\top = T([-1, y]^\top + [1, y]^\top)$ .

(b) False. Consider  $\mathbf{v}_1 = [1, 0]^\top$  and  $\mathbf{v}_2 = [2, 0]^\top$  and  $\mathbf{w}_1 = [1, 0]^\top$  and  $\mathbf{w}_2 = [0, 1]^\top$ , then there exists no LT  $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  such that  $T(\mathbf{v}_1) = \mathbf{w}_1$  and  $T(\mathbf{v}_2) = \mathbf{w}_2$  as  $\{\mathbf{v}_1, \mathbf{v}_2\}$  is LD.

(c) True. Since  $T$  is invertible it takes every basis of  $\mathbb{V}$  to a basis of  $\mathbb{W}$ . For any basis  $B$  of  $\mathbb{V}$ , consider the ordered basis  $C := T(B)$  of  $\mathbb{W}$ . Then  $[T]_{C \leftarrow B} = I_n$ .

6. Determine a linear transformation from  $\mathbb{R}^2 \rightarrow \mathbb{R}^3$  such that  $\text{Ker}(T) = \{[x, y]^\top : 2x + y = 0\}$ .

**Solution:** Take an  $[x, y]^\top$  satisfying  $2x + y = 0$ . For example, consider  $[-1, 2]^\top$  and define an LT  $T$  such that  $T([-1, 2]^\top) = \mathbf{0}$  and  $T([1, 0]^\top) = [1, 0]^\top$ .

7. Let  $\mathbb{V}$  be a vector space and  $\dim(\mathbb{V}) = n$ . Show that there exists an LT  $T : \mathbb{V} \rightarrow \mathbb{V}$  such that  $T^j \neq \mathbf{0}$  for  $j = 1, 2, \dots, n-1$  but  $T^n = \mathbf{0}$ .

**Solution:** Let  $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$  be an ordered basis of  $\mathbb{V}$ . Define an LT  $T$  by  $T(\mathbf{v}_1) = \mathbf{0}$  and  $T(\mathbf{v}_j) = \mathbf{v}_{j-1}$  for  $j = 2, \dots, n$ . Then  $T^j \mathbf{v}_{j+1} = \mathbf{v}_1$  for  $j = 1, 2, \dots, n-1$ , and  $T^n \mathbf{v}_j = \mathbf{0}$  for  $j = 1 : n$ .

8. Let  $T : \mathcal{M}_2(\mathbb{R}) \rightarrow \mathcal{M}_2(\mathbb{R})$  be defined as  $T(A) := A - A^\top$  for all  $A \in \mathcal{M}_2(\mathbb{R})$ . Find a basis of  $\text{range}(T)$  and  $\ker(T)$ .

**Solution:** Since  $T$  maps every  $2 \times 2$  real matrix to a skew symmetric matrix and  $T\left(\begin{bmatrix} x & y \\ z & w \end{bmatrix}\right) = \begin{bmatrix} 0 & y-z \\ z-y & 0 \end{bmatrix}$ , we have  $\text{range}(T) = \text{span}\left\{\begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}\right\}$ , and  $\ker(T) = \text{span}\left\{\begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 1 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix}\right\}$ .

9. Find the change of basis matrices  $P_{C \leftarrow B}$  and  $P_{B \leftarrow C}$  for the bases  $B := [1, x, x^2]$  and  $C := [1+x, x+x^2, 1+x^2]$  of  $\mathbb{R}_2[x]$ . Consider  $p(x) := 1 + 2x - x^2$ . Find  $[p]_C$  using the change of basis matrix.

**Solution:** Note that  $P_{B \leftarrow C} = [[1+x]_B, [x+x^2]_B, [1+x^2]_B] = \begin{bmatrix} 1 & 0 & 1 \\ 1 & 1 & 0 \\ 0 & 1 & 1 \end{bmatrix}$ .

We have  $P_{C \leftarrow B} = (P_{B \leftarrow C})^{-1} = \begin{bmatrix} 1/2 & 1/2 & -1/2 \\ -1/2 & 1/2 & 1/2 \\ 1/2 & -1/2 & 1/2 \end{bmatrix}$ . Hence  $[p]_C = P_{C \leftarrow B}[p]_B = [2, 0, -1]^\top$ .

10. Let  $\mathbb{V}$  be an  $n$ -dimensional vector space with an ordered basis  $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$ . Let  $A \in \mathcal{M}_n(\mathbb{F})$  be an invertible matrix. Consider  $C := BA$ . Show that  $C$  is an ordered basis of  $\mathbb{V}$  and that the change of basis matrix is given by  $P_{B \leftarrow C} = A$ .

**Solution:** Set  $\mathbf{u}_i := BA\mathbf{e}_i$  for  $i = 1 : n$ . Then  $P_{B \leftarrow C} = [[\mathbf{u}_1]_B, \dots, [\mathbf{u}_n]_B] = [A\mathbf{e}_1, \dots, A\mathbf{e}_n] = A$ .

11. True or False? Give justifications.

- (a) Let  $\mathbf{x}$  be a nonzero vector. Then  $\mathbf{x}$  is an eigenvector of  $A$  corresponding to an eigenvalue  $\lambda$  if and only if  $\mathbf{x}$  is an eigenvector of  $A^2$  corresponding to the eigenvalue  $\lambda^2$ .

- (b) Let  $A$  be a nonzero matrix such that  $A^{31} = \mathbf{0}$ . Then  $A$  has all eigenvalues equal to 0 and  $A$  is not diagonalizable.
- (c) If  $A$  is diagonalizable then  $\text{rank}(A - cI) = \text{rank}(A - cI)^2$  for all  $c \in \mathbb{C}$ .

**Solution:**

- (a) False. If  $A\mathbf{x} = \lambda\mathbf{x}$  then  $A(A\mathbf{x}) = \lambda A\mathbf{x} = \lambda^2\mathbf{x}$  which shows that  $\mathbf{x}$  is also an eigenvector of  $A^2$  corresponding to the eigenvalue  $\lambda^2$ . But the converse is not true. For example, consider  $A = \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}$ , then  $E_0(A^2) = \mathbb{R}^2 \neq E_0(A) = \text{span}\{[1, 0]^T\}$ .
- (b) True. If  $\lambda \neq 0$  is an eigenvalue of  $A$  then  $\lambda^{31} \neq 0$  is an eigenvalue of  $A^{31}$ , which is a contradiction. Since all eigenvalues of  $A$  are equal to 0, if  $A$  is diagonalizable then  $A$  has to be the  $\mathbf{0}$  matrix, which is a contradiction.

- (c) True. If  $P^{-1}AP = \begin{bmatrix} \lambda_1 & \dots & 0 \\ \vdots & \ddots & 0 \\ 0 & \dots & \lambda_n \end{bmatrix}$ , then  $P^{-1}(A - cI)P = \begin{bmatrix} \lambda_1 - c & \dots & 0 \\ \vdots & \ddots & 0 \\ 0 & \dots & \lambda_n - c \end{bmatrix}$  and
- $$P^{-1}(A - cI)^2P = \begin{bmatrix} (\lambda_1 - c)^2 & \dots & 0 \\ \vdots & \ddots & 0 \\ 0 & \dots & (\lambda_n - c)^2 \end{bmatrix}.$$
- Hence  $\text{rank}(A - cI) = \text{rank}(P^{-1}(A - cI)P) = \text{rank}(P^{-1}(A - cI)^2P) = \text{rank}((A - cI)^2)$ .

12. Let  $\mathbb{V}, \mathbb{W}$  be  $n$  dimensional vector spaces with ordered bases  $B$  and  $C$ , respectively, and  $T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ . Show that  $T$  is invertible if and only if the matrix  $[T]_{C \leftarrow B}$  is invertible. In such a case, show that

$$([T]_{C \leftarrow B})^{-1} = [T^{-1}]_{B \leftarrow C}.$$

**Solution:** Since  $T\mathbf{v} = C[T\mathbf{v}]_C = C[T]_{C \leftarrow B}[\mathbf{v}]_B$  it follows that  $\ker(T) = \{\mathbf{0}\} \Leftrightarrow \text{null}([T]_{C \leftarrow B}) = \{\mathbf{0}\}$ . Hence  $T$  is invertible  $\Leftrightarrow [T]_{C \leftarrow B}$  is invertible. (Also follows from Q.4)

Next, for  $\mathbf{v} \in \mathbb{V}$ , we have  $[\mathbf{v}]_B = [T^{-1}(T\mathbf{v})]_B = [T^{-1}]_{B \leftarrow C}[T\mathbf{v}]_C = [T^{-1}]_{B \leftarrow C}[T]_{C \leftarrow B}[\mathbf{v}]_B$ . Hence  $[T^{-1}]_{B \leftarrow C}[T]_{C \leftarrow B} = I_n \Rightarrow ([T]_{C \leftarrow B})^{-1} = [T^{-1}]_{B \leftarrow C}$ .

13. Consider  $\mathbb{U} := \mathbb{R}^3$ ,  $\mathbb{V} := \mathcal{M}_2(\mathbb{R})$  and  $\mathbb{W} := \mathbb{R}_2[x]$  with ordered bases  $B := [\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3]$ ,  $C := \left[ \begin{bmatrix} 1 & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 1 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 1 \end{bmatrix} \right]$  and  $D := [1, x, x^2]$ , respectively. Let  $T : \mathbb{U} \rightarrow \mathbb{V}$  be given by  $T[x, y, z]^T = \begin{bmatrix} 0 & x \\ y & y + z \end{bmatrix}$  and  $S : \mathbb{V} \rightarrow \mathbb{W}$  be given by  $S \begin{bmatrix} a & b \\ c & d \end{bmatrix} = a + (b + c)x + dx^2$ . Then determine  $[S \circ T]_{D \leftarrow B}$ ,  $[S]_{D \leftarrow C}$  and  $[T]_{C \leftarrow B}$  and verify that  $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C}[T]_{C \leftarrow B}$ .
14. For each LT  $T$  on  $\mathbb{V}$ , find the eigenvalues of  $T$  and an ordered basis  $B$  of  $\mathbb{V}$  such that  $[T]_B$  is a diagonal matrix.

(a)  $\mathbb{V} := \mathbb{R}_3[x]$  and  $(Tp)(x) := xp'(x) + p''(x) - p(2)$ .

(b)  $\mathbb{V} := \mathcal{M}_2(\mathbb{R})$  and  $T\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}\right) := \begin{bmatrix} d & b \\ c & a \end{bmatrix}$ .

(c)  $\mathbb{V} := \mathcal{M}_2(\mathbb{R})$  and  $T(A) := A^\top + 2 \operatorname{Trace}(A)I_2$ .

15. Let  $T : \mathbb{R}_2[x] \longrightarrow \mathbb{R}_2[x]$  be given by  $(Tp)(x) := p(1) + p'(0)x + (p'(0) + p''(0))x^2$ . Find eigenvalues and eigenvectors of  $T$ . Also, find an ordered basis  $B$ , if it exists, of  $\mathbb{R}_2[x]$  such that  $[T]_B$  is a diagonal matrix.

\*\*\*