

Linear Algebra

Department of Mathematics
Indian Institute of Technology Guwahati

January – May 2019

MA 102 (RA, RKS, MGPP, KVK)

Matrix Operations

Topics:

- Matrix operations
- Invertible matrices
- Elementary matrices and reduction to rref
- Gauss-Jordan elimination for computing inverse of a matrix
- LU factorization

Matrices

Recall that an $m \times n$ matrix A with entries a_{ij} has m rows and n columns and can be written as

$$\begin{aligned} A &= \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \mathbf{A}_1 \\ \mathbf{A}_2 \\ \vdots \\ \mathbf{A}_m \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} = [a_{ij}]_{m \times n}, \end{aligned}$$

Matrices

Recall that an $m \times n$ matrix A with entries a_{ij} has m rows and n columns and can be written as

$$\begin{aligned} A &= \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \mathbf{A}_1 \\ \mathbf{A}_2 \\ \vdots \\ \mathbf{A}_m \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} = [a_{ij}]_{m \times n}, \end{aligned}$$

where $\mathbf{A}_i := [a_{i1}, a_{i2}, \dots, a_{in}]$ is the i -th row of A for $i = 1 : m$ and $\mathbf{a}_j := [a_{1j}, \dots, a_{mj}]^\top$ is the j -th column of A for $j = 1 : n$.

Matrices

Recall that an $m \times n$ matrix A with entries a_{ij} has m rows and n columns and can be written as

$$\begin{aligned} A &= \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \mathbf{A}_1 \\ \mathbf{A}_2 \\ \vdots \\ \mathbf{A}_m \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} = [a_{ij}]_{m \times n}, \end{aligned}$$

where $\mathbf{A}_i := [a_{i1}, a_{i2}, \dots, a_{in}]$ is the i -th row of A for $i = 1 : m$ and $\mathbf{a}_j := [a_{1j}, \dots, a_{mj}]^\top$ is the j -th column of A for $j = 1 : n$.

Transpose: The transpose of A denoted by A^\top is the $n \times m$ matrix given by $A^\top = [a_{ji}]_{n \times m}$.

Matrices

Recall that an $m \times n$ matrix A with entries a_{ij} has m rows and n columns and can be written as

$$\begin{aligned} A &= \begin{bmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \cdots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{bmatrix} = \begin{bmatrix} \mathbf{A}_1 \\ \mathbf{A}_2 \\ \vdots \\ \mathbf{A}_m \end{bmatrix} \\ &= \begin{bmatrix} \mathbf{a}_1 & \mathbf{a}_2 & \cdots & \mathbf{a}_n \end{bmatrix} = [a_{ij}]_{m \times n}, \end{aligned}$$

where $\mathbf{A}_i := [a_{i1}, a_{i2}, \dots, a_{in}]$ is the i -th row of A for $i = 1 : m$ and $\mathbf{a}_j := [a_{1j}, \dots, a_{mj}]^\top$ is the j -th column of A for $j = 1 : n$.

Transpose: The transpose of A denoted by A^\top is the $n \times m$ matrix given by $A^\top = [a_{ji}]_{n \times m}$.

Two matrices $A = [a_{ij}]$ and $B := [b_{ij}]$ are said to be **equal** (i.e., $A = B$) if A and B have the same **size** and $a_{ij} = b_{ij}$ for all i and j .

Special matrices

We denote the set of complex numbers by \mathbb{C} .

A matrix $A := [a_{ij}]$ with entries $a_{ij} \in \mathbb{C}$ (resp., $a_{ij} \in \mathbb{R}$) is said to be a complex (resp., real) matrix.

Special matrices

We denote the set of complex numbers by \mathbb{C} .

A matrix $A := [a_{ij}]$ with entries $a_{ij} \in \mathbb{C}$ (resp., $a_{ij} \in \mathbb{R}$) is said to be a complex (resp., real) matrix.

Conjugate transpose: The conjugate transpose of an $m \times n$ complex matrix $A = [a_{ij}]_{m \times n}$ is the $n \times m$ matrix denoted by A^* and is given by

$$A^* = [\bar{a}_{ji}]_{n \times m} = ([\bar{a}_{ij}]_{m \times n})^T = (\bar{A})^T,$$

where \bar{a}_{ij} is the conjugate of a_{ij} .

Special matrices

We denote the set of complex numbers by \mathbb{C} .

A matrix $A := [a_{ij}]$ with entries $a_{ij} \in \mathbb{C}$ (resp., $a_{ij} \in \mathbb{R}$) is said to be a complex (resp., real) matrix.

Conjugate transpose: The conjugate transpose of an $m \times n$ complex matrix $A = [a_{ij}]_{m \times n}$ is the $n \times m$ matrix denoted by A^* and is given by

$$A^* = [\bar{a}_{ji}]_{n \times m} = ([\bar{a}_{ij}]_{m \times n})^T = (\bar{A})^T,$$

where \bar{a}_{ij} is the conjugate of a_{ij} .

Definition: Let A be an $n \times n$ matrix. Then A is said to be

- ① symmetric if $A^T = A$,
- ② skew-symmetric if $A^T = -A$,

Special matrices

We denote the set of **complex numbers** by \mathbb{C} .

A **matrix** $A := [a_{ij}]$ with entries $a_{ij} \in \mathbb{C}$ (resp., $a_{ij} \in \mathbb{R}$) is said to be a **complex** (resp., **real**) matrix.

Conjugate transpose: The conjugate transpose of an $m \times n$ **complex matrix** $A = [a_{ij}]_{m \times n}$ is the $n \times m$ **matrix** denoted by A^* and is given by

$$A^* = [\bar{a}_{ji}]_{n \times m} = ([\bar{a}_{ij}]_{m \times n})^T = (\bar{A})^T,$$

where \bar{a}_{ij} is the **conjugate** of a_{ij} .

Definition: Let A be an $n \times n$ matrix. Then A is said to be

- ① **symmetric** if $A^T = A$,
- ② **skew-symmetric** if $A^T = -A$,
- ③ **Hermitian** if $A^* = A$,
- ④ **skew-Hermitian** if $A^* = -A$.

Special matrices (recall)

Let A be an $m \times n$ matrix with (i, j) -th entry a_{ij} . Set $p := \min(m, n)$. Then

- a_{ii} for $i = 1 : p$ are called the **diagonal entries** of A ;
- A is said to be a **diagonal matrix** if $a_{ij} = 0$ for all $i \neq j$;

Special matrices (recall)

Let A be an $m \times n$ matrix with (i, j) -th entry a_{ij} . Set $p := \min(m, n)$. Then

- a_{ii} for $i = 1 : p$ are called the **diagonal entries** of A ;
- A is said to be a **diagonal matrix** if $a_{ij} = 0$ for all $i \neq j$;
- A is said to be an **upper triangular** if $a_{ij} = 0$ for all $i > j$;
- A is said to be a **lower triangular** if $a_{ij} = 0$ for all $i < j$;

Identity matrix: An $n \times n$ diagonal matrix with all diagonal entries equal to 1 is called the **identity matrix** and is denoted by I_n or I .

Zero matrix: An $m \times n$ matrix with all entries 0 is called the **zero matrix** and is denoted by $\mathbf{O}_{m \times n}$ or simply by \mathbf{O} .

Matrix Operations

- $\mathcal{M}_{m \times n} :=$ the set of a $m \times n$ matrices.

Matrix Operations

- $\mathcal{M}_{m \times n} :=$ the set of a $m \times n$ matrices.
- To specify **real** (resp., **complex**) matrices, write $\mathcal{M}_{m \times n}(\mathbb{R})$ (resp., $\mathcal{M}_{m \times n}(\mathbb{C})$).

Matrix Operations

- $\mathcal{M}_{m \times n} :=$ the set of a $m \times n$ matrices.
- To specify **real** (resp., **complex**) matrices, write $\mathcal{M}_{m \times n}(\mathbb{R})$ (resp., $\mathcal{M}_{m \times n}(\mathbb{C})$).
- If $m = n$, we write \mathcal{M}_n for $\mathcal{M}_{m \times n}$.

Matrix Operations

- $\mathcal{M}_{m \times n} :=$ the set of a $m \times n$ matrices.
- To specify **real** (resp., **complex**) matrices, write $\mathcal{M}_{m \times n}(\mathbb{R})$ (resp., $\mathcal{M}_{m \times n}(\mathbb{C})$).
- If $m = n$, we write \mathcal{M}_n for $\mathcal{M}_{m \times n}$.
- Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be in $\mathcal{M}_{m \times n}$ and α be a scalar. Then

Matrix Operations

- $\mathcal{M}_{m \times n} :=$ the set of a $m \times n$ matrices.
- To specify **real** (resp., **complex**) matrices, write $\mathcal{M}_{m \times n}(\mathbb{R})$ (resp., $\mathcal{M}_{m \times n}(\mathbb{C})$).
- If $m = n$, we write \mathcal{M}_n for $\mathcal{M}_{m \times n}$.
- Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be in $\mathcal{M}_{m \times n}$ and α be a scalar.
Then
 - ① **Matrix addition:** $A + B := [a_{ij} + b_{ij}] \in \mathcal{M}_{m \times n}$.

Matrix Operations

- $\mathcal{M}_{m \times n} :=$ the set of a $m \times n$ matrices.
- To specify **real** (resp., **complex**) matrices, write $\mathcal{M}_{m \times n}(\mathbb{R})$ (resp., $\mathcal{M}_{m \times n}(\mathbb{C})$).
- If $m = n$, we write \mathcal{M}_n for $\mathcal{M}_{m \times n}$.
- Let $A = [a_{ij}]$ and $B = [b_{ij}]$ be in $\mathcal{M}_{m \times n}$ and α be a scalar. Then
 - 1 **Matrix addition:** $A + B := [a_{ij} + b_{ij}] \in \mathcal{M}_{m \times n}$.
 - 2 **Multiplication by a scalar:** $\alpha A := [\alpha a_{ij}] \in \mathcal{M}_{m \times n}$.
($\alpha \in \mathbb{R}$ when A and B are **real matrices**, and $\alpha \in \mathbb{C}$ when A and B are **complex matrices**)

Properties of addition and scalar multiplication

Theorem: Let $A, B, C \in \mathcal{M}_{m \times n}$ and α, β be scalars. Then

Properties of addition and scalar multiplication

Theorem: Let $A, B, C \in \mathcal{M}_{m \times n}$ and α, β be scalars. Then

- ① **Commutative Law:** $A + B = B + A$.
- ② **Associative Law:** $(A + B) + C = A + (B + C)$.

Properties of addition and scalar multiplication

Theorem: Let $A, B, C \in \mathcal{M}_{m \times n}$ and α, β be scalars. Then

- ① **Commutative Law:** $A + B = B + A$.
- ② **Associative Law:** $(A + B) + C = A + (B + C)$.
- ③ $A + \mathbf{O} = A$, where $\mathbf{O} = \mathbf{O}_{m \times n} \in \mathcal{M}_{m \times n}$.
- ④ $A + (-A) = \mathbf{O}$, where $-A = (-1)A \in \mathcal{M}_{m \times n}$.

Properties of addition and scalar multiplication

Theorem: Let $A, B, C \in \mathcal{M}_{m \times n}$ and α, β be scalars. Then

- ① **Commutative Law:** $A + B = B + A$.
- ② **Associative Law:** $(A + B) + C = A + (B + C)$.
- ③ $A + \mathbf{O} = A$, where $\mathbf{O} = \mathbf{O}_{m \times n} \in \mathcal{M}_{m \times n}$.
- ④ $A + (-A) = \mathbf{O}$, where $-A = (-1)A \in \mathcal{M}_{m \times n}$.
- ⑤ $\alpha(A + B) = \alpha A + \alpha B$.
- ⑥ $(\alpha + \beta)A = \alpha A + \beta A$.

Properties of addition and scalar multiplication

Theorem: Let $A, B, C \in \mathcal{M}_{m \times n}$ and α, β be scalars. Then

- ① **Commutative Law:** $A + B = B + A$.
- ② **Associative Law:** $(A + B) + C = A + (B + C)$.
- ③ $A + \mathbf{O} = A$, where $\mathbf{O} = \mathbf{O}_{m \times n} \in \mathcal{M}_{m \times n}$.
- ④ $A + (-A) = \mathbf{O}$, where $-A = (-1)A \in \mathcal{M}_{m \times n}$.
- ⑤ $\alpha(A + B) = \alpha A + \alpha B$.
- ⑥ $(\alpha + \beta)A = \alpha A + \beta A$.
- ⑦ $\alpha(\beta A) = (\alpha\beta)A$.
- ⑧ $1A = A$.

Matrix-vector multiplication revisited

Let $A := \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \in \mathcal{M}_{m \times n}(\mathbb{R})$ and $\mathbf{x} := [x_1, \dots, x_n]^\top \in \mathbb{R}^n$. Recall the matrix-vector multiplication

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n \in \mathbb{R}^m.$$

Matrix-vector multiplication revisited

Let $A := \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \in \mathcal{M}_{m \times n}(\mathbb{R})$ and $\mathbf{x} := [x_1, \dots, x_n]^\top \in \mathbb{R}^n$. Recall the matrix-vector multiplication

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n \in \mathbb{R}^m.$$

This shows that A act on \mathbf{x} as a well defined function

$$A: \mathbb{R}^n \longrightarrow \mathbb{R}^m, \mathbf{x} \longmapsto A\mathbf{x},$$

and satisfies following:

Matrix-vector multiplication revisited

Let $A := \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \in \mathcal{M}_{m \times n}(\mathbb{R})$ and $\mathbf{x} := [x_1, \dots, x_n]^\top \in \mathbb{R}^n$. Recall the matrix-vector multiplication

$$A\mathbf{x} = \begin{bmatrix} \mathbf{a}_1 & \cdots & \mathbf{a}_n \end{bmatrix} \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n \in \mathbb{R}^m.$$

This shows that A act on \mathbf{x} as a well defined function

$$A : \mathbb{R}^n \longrightarrow \mathbb{R}^m, \mathbf{x} \longmapsto A\mathbf{x},$$

and satisfies following:

- $A(\mathbf{x} + \mathbf{y}) = A\mathbf{x} + A\mathbf{y}$ for all \mathbf{x}, \mathbf{y} in \mathbb{R}^n .
- $A(\alpha\mathbf{x}) = \alpha A\mathbf{x}$ for all $\mathbf{x} \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$.

Matrix-vector multiplication revisited

Let $A := [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \in \mathcal{M}_{m \times n}(\mathbb{R})$ and $\mathbf{x} := [x_1, \dots, x_n]^T \in \mathbb{R}^n$. Recall the matrix-vector multiplication

$$A\mathbf{x} = [\mathbf{a}_1 \ \cdots \ \mathbf{a}_n] \begin{bmatrix} x_1 \\ \vdots \\ x_n \end{bmatrix} = x_1\mathbf{a}_1 + \cdots + x_n\mathbf{a}_n \in \mathbb{R}^m.$$

This shows that A act on \mathbf{x} as a well defined function

$$A : \mathbb{R}^n \longrightarrow \mathbb{R}^m, \mathbf{x} \longmapsto A\mathbf{x},$$

and satisfies following:

- $A(\mathbf{x} + \mathbf{y}) = A\mathbf{x} + A\mathbf{y}$ for all \mathbf{x}, \mathbf{y} in \mathbb{R}^n .
- $A(\alpha\mathbf{x}) = \alpha A\mathbf{x}$ for all $\mathbf{x} \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$.

We refer to $A : \mathbb{R}^n \longrightarrow \mathbb{R}^m, \mathbf{x} \longmapsto A\mathbf{x}$, as a **linear mapping**.

Matrix-matrix multiplication

Let $A \in \mathcal{M}_{m \times n}$ and $B := \begin{bmatrix} \mathbf{b}_1 & \cdots & \mathbf{b}_p \end{bmatrix} \in \mathcal{M}_{n \times p}$.

Definition: Define the matrix-matrix multiplication AB by

$$AB := \begin{bmatrix} A\mathbf{b}_1 & \cdots & A\mathbf{b}_p \end{bmatrix}.$$

Matrix-matrix multiplication

Let $A \in \mathcal{M}_{m \times n}$ and $B := \begin{bmatrix} \mathbf{b}_1 & \cdots & \mathbf{b}_p \end{bmatrix} \in \mathcal{M}_{n \times p}$.

Definition: Define the matrix-matrix multiplication AB by

$$AB := \begin{bmatrix} A\mathbf{b}_1 & \cdots & A\mathbf{b}_p \end{bmatrix}.$$

Justification: Define AB to be the composition of the maps

$$A : \mathbb{R}^n \longrightarrow \mathbb{R}^m, \mathbf{x} \mapsto A\mathbf{x}, \text{ and } B : \mathbb{R}^p \longrightarrow \mathbb{R}^n, \mathbf{y} \mapsto B\mathbf{y}.$$

Matrix-matrix multiplication

Let $A \in \mathcal{M}_{m \times n}$ and $B := [\mathbf{b}_1 \ \cdots \ \mathbf{b}_p] \in \mathcal{M}_{n \times p}$.

Definition: Define the matrix-matrix multiplication AB by

$$AB := [\mathbf{Ab}_1 \ \cdots \ \mathbf{Ab}_p].$$

Justification: Define AB to be the composition of the maps

$$A : \mathbb{R}^n \longrightarrow \mathbb{R}^m, \mathbf{x} \mapsto A\mathbf{x}, \text{ and } B : \mathbb{R}^p \longrightarrow \mathbb{R}^n, \mathbf{y} \mapsto B\mathbf{y}.$$

Then the linear map $AB : \mathbb{R}^p \longrightarrow \mathbb{R}^m$ is given by

$$\begin{aligned} AB\mathbf{y} &:= A(B\mathbf{y}) = A(y_1\mathbf{b}_1 + \cdots + y_p\mathbf{b}_p) \\ &= y_1\mathbf{Ab}_1 + \cdots + y_p\mathbf{Ab}_p = [\mathbf{Ab}_1 \ \cdots \ \mathbf{Ab}_p] \mathbf{y} \end{aligned}$$

for all $\mathbf{y} \in \mathbb{R}^p$

Matrix-matrix multiplication

Let $A \in \mathcal{M}_{m \times n}$ and $B := \begin{bmatrix} \mathbf{b}_1 & \cdots & \mathbf{b}_p \end{bmatrix} \in \mathcal{M}_{n \times p}$.

Definition: Define the matrix-matrix multiplication AB by

$$AB := \begin{bmatrix} A\mathbf{b}_1 & \cdots & A\mathbf{b}_p \end{bmatrix}.$$

Justification: Define AB to be the composition of the maps

$$A : \mathbb{R}^n \longrightarrow \mathbb{R}^m, \mathbf{x} \mapsto A\mathbf{x}, \text{ and } B : \mathbb{R}^p \longrightarrow \mathbb{R}^n, \mathbf{y} \mapsto B\mathbf{y}.$$

Then the linear map $AB : \mathbb{R}^p \longrightarrow \mathbb{R}^m$ is given by

$$\begin{aligned} AB\mathbf{y} &:= A(B\mathbf{y}) = A(y_1\mathbf{b}_1 + \cdots + y_p\mathbf{b}_p) \\ &= y_1A\mathbf{b}_1 + \cdots + y_pA\mathbf{b}_p = \begin{bmatrix} A\mathbf{b}_1 & \cdots & A\mathbf{b}_p \end{bmatrix} \mathbf{y} \end{aligned}$$

$$\text{for all } \mathbf{y} \in \mathbb{R}^p \implies AB = \begin{bmatrix} A\mathbf{b}_1 & \cdots & A\mathbf{b}_p \end{bmatrix}.$$

Matrix multiplication

Let $A = \begin{bmatrix} \mathbf{A}_1 \\ \vdots \\ \mathbf{A}_m \end{bmatrix} \in \mathcal{M}_{m \times n}$, $B := [\mathbf{b}_1 \ \cdots \ \mathbf{b}_p] \in \mathcal{M}_{n \times p}$. Then

$$AB = \begin{bmatrix} \mathbf{A}_1 \mathbf{b}_1 & \cdots & \mathbf{A}_1 \mathbf{b}_p \\ \vdots & \cdots & \vdots \\ \mathbf{A}_m \mathbf{b}_1 & \cdots & \mathbf{A}_m \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} \mathbf{A}_1 B \\ \vdots \\ \mathbf{A}_m B \end{bmatrix}.$$

Matrix multiplication

Let $A = \begin{bmatrix} \mathbf{A}_1 \\ \vdots \\ \mathbf{A}_m \end{bmatrix} \in \mathcal{M}_{m \times n}$, $B := [\mathbf{b}_1 \ \cdots \ \mathbf{b}_p] \in \mathcal{M}_{n \times p}$. Then

$$AB = \begin{bmatrix} \mathbf{A}_1 \mathbf{b}_1 & \cdots & \mathbf{A}_1 \mathbf{b}_p \\ \vdots & \cdots & \vdots \\ \mathbf{A}_m \mathbf{b}_1 & \cdots & \mathbf{A}_m \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} \mathbf{A}_1 B \\ \vdots \\ \mathbf{A}_m B \end{bmatrix}.$$

Thus if $A := [a_{ij}]_{m \times n}$, $B := [b_{ij}]_{n \times p}$ and $C := AB = [c_{ij}]_{m \times p}$ then

$$c_{ij} = \begin{bmatrix} a_{i1} & \cdots & a_{in} \end{bmatrix} \begin{bmatrix} b_{1j} \\ \vdots \\ b_{nj} \end{bmatrix} = \sum_{k=1}^n a_{ik} b_{kj}.$$

Matrix multiplication

Let $A = \begin{bmatrix} \mathbf{A}_1 \\ \vdots \\ \mathbf{A}_m \end{bmatrix} \in \mathcal{M}_{m \times n}$, $B := [\mathbf{b}_1 \ \cdots \ \mathbf{b}_p] \in \mathcal{M}_{n \times p}$. Then

$$AB = \begin{bmatrix} \mathbf{A}_1 \mathbf{b}_1 & \cdots & \mathbf{A}_1 \mathbf{b}_p \\ \vdots & \cdots & \vdots \\ \mathbf{A}_m \mathbf{b}_1 & \cdots & \mathbf{A}_m \mathbf{b}_p \end{bmatrix} = \begin{bmatrix} \mathbf{A}_1 B \\ \vdots \\ \mathbf{A}_m B \end{bmatrix}.$$

Thus if $A := [a_{ij}]_{m \times n}$, $B := [b_{ij}]_{n \times p}$ and $C := AB = [c_{ij}]_{m \times p}$ then

$$c_{ij} = \begin{bmatrix} a_{i1} & \cdots & a_{in} \end{bmatrix} \begin{bmatrix} b_{1j} \\ \vdots \\ b_{nj} \end{bmatrix} = \sum_{k=1}^n a_{ik} b_{kj}.$$

Note:

$$\mathbf{e}_i^\top A = \mathbf{A}_i = i\text{-th row of } A, \text{ where } \mathbf{e}_i \in \mathbb{R}^m$$

$$A \mathbf{e}_j = \mathbf{a}_j = j\text{-th column of } A, \text{ where } \mathbf{e}_j \in \mathbb{R}^n.$$

Matrix multiplication

Thoerm: Let A, B and C be matrices, and let $\alpha \in \mathbb{R}$. Then

Matrix multiplication

Thoerm: Let A, B and C be matrices, and let $\alpha \in \mathbb{R}$. Then

- ① **Associative Law:** $(AB)C = A(BC)$, if the respective matrix products are defined.
- ② **Distributive Law:**
 $A(B + C) = AB + AC$, $(A + B)C = AC + BC$,
if the respective matrix sum and matrix products are defined.

Matrix multiplication

Thoerm: Let A, B and C be matrices, and let $\alpha \in \mathbb{R}$. Then

- 1 **Associative Law:** $(AB)C = A(BC)$, if the respective matrix products are defined.
- 2 **Distributive Law:**
 $A(B + C) = AB + AC$, $(A + B)C = AC + BC$,
if the respective matrix sum and matrix products are defined.
- 3 $\alpha(AB) = (\alpha A)B = A(\alpha B)$, if the respective matrix products are defined.
- 4 $I_m A = A = A I_n$, if A is of size $m \times n$.

Block matrix

Definition: An $m \times n$ **block matrix** (or a partition matrix) is a matrix of the form

$$A := \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & \cdots & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix}$$

where each A_{ij} is a $p_i \times q_j$ **matrix** for $i = 1 : m$ and $j = 1 : n$.

Block matrix

Definition: An $m \times n$ **block matrix** (or a partition matrix) is a matrix of the form

$$A := \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & \cdots & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix}$$

where each A_{ij} is a $p_i \times q_j$ **matrix** for $i = 1 : m$ and $j = 1 : n$.

Then $\begin{bmatrix} A_{i1} & \cdots & A_{in} \end{bmatrix}$ is the i -th **block row** of A and $\begin{bmatrix} A_{1j} \\ \vdots \\ A_{mj} \end{bmatrix}$ is the j -th **block column** of A .

Block matrix

Definition: An $m \times n$ **block matrix** (or a partition matrix) is a matrix of the form

$$A := \begin{bmatrix} A_{11} & \cdots & A_{1n} \\ \vdots & \cdots & \vdots \\ A_{m1} & \cdots & A_{mn} \end{bmatrix}$$

where each A_{ij} is a $p_i \times q_j$ matrix for $i = 1 : m$ and $j = 1 : n$.

Then $\begin{bmatrix} A_{i1} & \cdots & A_{in} \end{bmatrix}$ is the i -th **block row** of A and $\begin{bmatrix} A_{1j} \\ \vdots \\ A_{mj} \end{bmatrix}$ is

the j -th **block column** of A .

Example: $\left[\begin{array}{cc|cc|c} 1 & 2 & 2 & 0 & 1 & 4 \\ 3 & 4 & 1 & 2 & 3 & 5 \\ \hline 5 & 7 & 2 & 7 & 8 & 8 \\ 3 & 4 & 1 & 9 & 2 & 2 \end{array} \right]$ has 2 block rows and 3

block columns.

Block matrix operations

Block matrix addition: Let $A := [A_{ij}]_{m \times n}$ and $B := [B_{ij}]_{m \times n}$ be block matrices such that **size of A_{ij} = size of B_{ij}** for $i = 1 : m$ and $j = 1 : n$. Then **$A + B := [A_{ij} + B_{ij}]_{m \times n}$** .

Block matrix operations

Block matrix addition: Let $A := [A_{ij}]_{m \times n}$ and $B := [B_{ij}]_{m \times n}$ be block matrices such that **size of A_{ij} = size of B_{ij}** for $i = 1 : m$ and $j = 1 : n$. Then **$A + B := [A_{ij} + B_{ij}]_{m \times n}$** .

Block matrix multiplication: Let $A := [A_{ij}]_{m \times n}$ and $B := [B_{ij}]_{n \times p}$ be block matrices. If the matrix multiplication $C_{ij} := \sum_{k=1}^n A_{ik} B_{kj}$ is well defined for $i = 1 : m$ and $j = 1 : p$ then **AB is an $m \times p$ block matrix** given by **$AB = [C_{ij}]_{m \times p}$** .

Block matrix operations

Block matrix addition: Let $A := [A_{ij}]_{m \times n}$ and $B := [B_{ij}]_{m \times n}$ be block matrices such that **size of A_{ij} = size of B_{ij}** for $i = 1 : m$ and $j = 1 : n$. Then **$A + B := [A_{ij} + B_{ij}]_{m \times n}$** .

Block matrix multiplication: Let $A := [A_{ij}]_{m \times n}$ and $B := [B_{ij}]_{n \times p}$ be block matrices. If the matrix multiplication $C_{ij} := \sum_{k=1}^n A_{ik} B_{kj}$ is well defined for $i = 1 : m$ and $j = 1 : p$ then **AB is an $m \times p$ block matrix** given by **$AB = [C_{ij}]_{m \times p}$** .

Conformal partition: If a block operation of block matrices A and B are well defined then A and B are said to be **partitioned conformably**.

Block matrix operations

Block matrix addition: Let $A := [A_{ij}]_{m \times n}$ and $B := [B_{ij}]_{m \times n}$ be block matrices such that **size of A_{ij} = size of B_{ij}** for $i = 1 : m$ and $j = 1 : n$. Then **$A + B := [A_{ij} + B_{ij}]_{m \times n}$** .

Block matrix multiplication: Let $A := [A_{ij}]_{m \times n}$ and $B := [B_{ij}]_{n \times p}$ be block matrices. If the matrix multiplication $C_{ij} := \sum_{k=1}^n A_{ik} B_{kj}$ is well defined for $i = 1 : m$ and $j = 1 : p$ then **AB is an $m \times p$ block matrix** given by **$AB = [C_{ij}]_{m \times p}$** .

Conformal partition: If a block operation of block matrices A and B are well defined then A and B are said to be **partitioned conformably**.

Example:
$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix} \begin{bmatrix} B_{11} & B_{12} & B_{13} \\ B_{21} & B_{22} & B_{23} \end{bmatrix} =$$

$$\begin{bmatrix} A_{11}B_{11} + A_{12}B_{21} & A_{11}B_{12} + A_{12}B_{22} & A_{11}B_{13} + A_{12}B_{23} \\ A_{21}B_{11} + A_{22}B_{21} & A_{21}B_{12} + A_{22}B_{22} & A_{21}B_{13} + A_{22}B_{23} \end{bmatrix}.$$

The inverse of a matrix

Definition: An $n \times n$ matrix A is said to be **invertible** if there exists a matrix B satisfying $AB = I_n = BA$. The matrix B is called an **inverse** of A .

The inverse of a matrix

Definition: An $n \times n$ matrix A is said to be **invertible** if there exists a matrix B satisfying $AB = I_n = BA$. The matrix B is called an **inverse** of A .

Fact: Inverse of an invertible matrix A is **unique**. (Exercise)

The inverse of a matrix

Definition: An $n \times n$ matrix A is said to be **invertible** if there exists a matrix B satisfying $AB = I_n = BA$. The matrix B is called an **inverse** of A .

Fact: Inverse of an invertible matrix A is **unique**. (Exercise)
We denote the inverse of A by A^{-1} .

The inverse of a matrix

Definition: An $n \times n$ matrix A is said to be **invertible** if there exists a matrix B satisfying $AB = I_n = BA$. The matrix B is called an **inverse** of A .

Fact: Inverse of an invertible matrix A is **unique**. (Exercise)
We denote the inverse of A by A^{-1} .

Note: We can talk of invertibility only of square matrices.

The inverse of a matrix

Definition: An $n \times n$ matrix A is said to be **invertible** if there exists a matrix B satisfying $AB = I_n = BA$. The matrix B is called an **inverse** of A .

Fact: Inverse of an invertible matrix A is **unique**. (Exercise)
We denote the inverse of A by A^{-1} .

Note: We can talk of invertibility only of square matrices.

- For example, the matrix $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$ is invertible since

$$\begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}.$$

The inverse of a matrix

Definition: An $n \times n$ matrix A is said to be **invertible** if there exists a matrix B satisfying $AB = I_n = BA$. The matrix B is called an **inverse** of A .

Fact: Inverse of an invertible matrix A is **unique**. (Exercise)
We denote the inverse of A by A^{-1} .

Note: We can talk of invertibility only of square matrices.

- For example, the matrix $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$ is invertible since

$$\begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}.$$

- The zero matrix \mathbf{O} is **not invertible**.

The inverse of a matrix

Definition: An $n \times n$ matrix A is said to be invertible if there exists a matrix B satisfying $AB = I_n = BA$. The matrix B is called an inverse of A .

Fact: Inverse of an invertible matrix A is unique. (Exercise)
We denote the inverse of A by A^{-1} .

Note: We can talk of invertibility only of square matrices.

- For example, the matrix $A = \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}$ is invertible since

$$\begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix} \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} = \begin{bmatrix} 3 & -5 \\ -1 & 2 \end{bmatrix} \begin{bmatrix} 2 & 5 \\ 1 & 3 \end{bmatrix}.$$

- The zero matrix \mathbf{O} is not invertible.
- If A has a zero row, then A is not invertible.

Properties of invertible matrices

Fact: Let A and B be two invertible matrices of the same size.

- 1 If $c \neq 0$ then cA is also invertible, and $(cA)^{-1} = \frac{1}{c}A^{-1}$.
- 2 The matrix AB is invertible, and $(AB)^{-1} = B^{-1}A^{-1}$.
- 3 The matrix A^T is invertible, and $(A^T)^{-1} = (A^{-1})^T$.
- 4 For any non-negative integer k , the matrix A^k is invertible, and $(A^k)^{-1} = (A^{-1})^k$.

Properties of invertible matrices

Fact: Let A and B be two invertible matrices of the same size.

- 1 If $c \neq 0$ then cA is also invertible, and $(cA)^{-1} = \frac{1}{c}A^{-1}$.
- 2 The matrix AB is invertible, and $(AB)^{-1} = B^{-1}A^{-1}$.
- 3 The matrix A^T is invertible, and $(A^T)^{-1} = (A^{-1})^T$.
- 4 For any non-negative integer k , the matrix A^k is invertible, and $(A^k)^{-1} = (A^{-1})^k$.

Let $A := \begin{bmatrix} a & b \\ c & d \end{bmatrix}$. If $ad - bc \neq 0$ then A is invertible, and

$$A^{-1} = \frac{1}{ad - bc} \begin{bmatrix} d & -b \\ -c & a \end{bmatrix}.$$

for any matrix there exist a inverse iff the determinant of the matrix is not zero.

If $ad - bc = 0$ then A is **not** invertible.

Elementary matrices

| Type | Operation | Inverse operation |
|------------|---------------------------------|---|
| <i>I</i> | $R_i \longleftarrow \alpha R_i$ | $R_i \longleftarrow \frac{1}{\alpha} R_i$ |
| <i>II</i> | $R_j \longleftarrow cR_i + R_j$ | $R_j \longleftarrow -cR_i + R_j$ |
| <i>III</i> | $R_i \leftrightarrow R_j$ | $R_i \leftrightarrow R_j$ |

Elementary matrices

| Type | Operation | Inverse operation |
|------------|---------------------------------|---|
| <i>I</i> | $R_i \longleftarrow \alpha R_i$ | $R_i \longleftarrow \frac{1}{\alpha} R_i$ |
| <i>II</i> | $R_j \longleftarrow cR_i + R_j$ | $R_j \longleftarrow -cR_i + R_j$ |
| <i>III</i> | $R_i \leftrightarrow R_j$ | $R_i \leftrightarrow R_j$ |

An **elementary matrix** is a matrix that is obtained by performing an elementary row operation on the identity matrix.

Elementary matrices

| Type | Operation | Inverse operation |
|------|-----------------------------|---------------------------------------|
| I | $R_i \leftarrow \alpha R_i$ | $R_i \leftarrow \frac{1}{\alpha} R_i$ |
| II | $R_j \leftarrow cR_i + R_j$ | $R_j \leftarrow -cR_i + R_j$ |
| III | $R_i \leftrightarrow R_j$ | $R_i \leftrightarrow R_j$ |

An elementary matrix is a matrix that is obtained by performing an elementary row operation on the identity matrix.

$$\text{Type I: } E_2(\alpha) = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \alpha & 0 \\ 0 & 0 & 1 \end{bmatrix} \Rightarrow E_2 \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ \alpha x_2 \\ x_3 \end{bmatrix}.$$

$$E_2(\alpha)A = \begin{bmatrix} \text{row}_1(A) \\ \alpha \text{row}_2(A) \\ \text{row}_3(A) \end{bmatrix} = \text{multiply 2nd row of } A \text{ by } \alpha.$$

$$(E_2(\alpha))^{-1} = E_2\left(\frac{1}{\alpha}\right).$$

Type II elementary matrices

$$\text{Type II : } E_{13}(2) := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} = I_3 + 2e_3e_1^T$$

The matrix E_{13} is obtained by performing $R_3 \leftarrow 2R_1 + R_3$ on I_3 .

Type II elementary matrices

$$\text{Type II : } E_{13}(2) := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 2 & 0 & 1 \end{bmatrix} = I_3 + 2e_3e_1^T$$

The matrix E_{13} is obtained by performing $R_3 \leftarrow 2R_1 + R_3$ on I_3 .

$$E_{13}(2) \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_2 \\ x_3 + 2x_1 \end{bmatrix} \Rightarrow E_{13}A = \begin{bmatrix} \text{row}_1(A) \\ \text{row}_2(A) \\ \text{row}_3(A) + 2\text{row}_1(A) \end{bmatrix}.$$

$(E_{13}(2))^{-1} = E_{13}(-2)$ corresponds to $R_3 \leftarrow -2R_1 + R_3$ on I_3 .

Type III elementary matrices

Type III : E_{ij} is obtained by performing $R_i \leftrightarrow R_j$ on I .

$$E_{23} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \Rightarrow E_{23} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_3 \\ x_2 \end{bmatrix} \Rightarrow E_{23}A = \begin{bmatrix} \text{row}_1(A) \\ \text{row}_3(A) \\ \text{row}_2(A) \end{bmatrix}.$$

Type III elementary matrices

Type III : E_{ij} is obtained by performing $R_i \leftrightarrow R_j$ on I .

$$E_{23} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 1 & 0 \end{bmatrix} \Rightarrow E_{23} \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} x_1 \\ x_3 \\ x_2 \end{bmatrix} \Rightarrow E_{23}A = \begin{bmatrix} \text{row}_1(A) \\ \text{row}_3(A) \\ \text{row}_2(A) \end{bmatrix}.$$

$(E_{ij})^{-1} = E_{ij}$ corresponds to row operation $R_i \leftrightarrow R_j$ on I .

Observation: Inverse of an elementary matrix is also an elementary matrix of same type.

Row operation via elementary matrices

Crux of the matter:

- Type I: Multiplying $E_i(c)$ to A giving $E_i(c)A$ amounts to performing the row operation $R_i \leftarrow cR_i$ on A .
- Type II: Multiplying $E_{ij}(c)$ to A giving $E_{ij}(c)A$ amounts to performing the row operation $R_j \leftarrow cR_i + R_j$ on A .
- Type III: Multiplying E_{ij} to A giving $E_{ij}A$ amounts to performing the row operation $R_i \leftrightarrow R_j$ on A .

Row operation via elementary matrices

Crux of the matter:

- Type I: Multiplying $E_i(c)$ to A giving $E_i(c)A$ amounts to performing the row operation $R_i \leftarrow cR_i$ on A .
- Type II: Multiplying $E_{ij}(c)$ to A giving $E_{ij}(c)A$ amounts to performing the row operation $R_j \leftarrow cR_i + R_j$ on A .
- Type III: Multiplying E_{ij} to A giving $E_{ij}A$ amounts to performing the row operation $R_i \leftrightarrow R_j$ on A .

Recall that two matrices A and B are said to be **row equivalent** if A can be transformed to B by elementary row operations.

Theorem: The matrices A and B are row equivalent $\iff B = E_k \cdots E_2 E_1 A$ for some elementary matrices E_1, E_2, \dots, E_k .

Elementary matrices and echelon form

Forward GE: $m \times n$ matrix $A \longrightarrow$ row echelon form $\text{ref}(A)$



$\text{ref}(A) = E_p \cdots E_2 E_1 A$ for some elementary matrices E_1, \dots, E_p .

Elementary matrices and echelon form

Forward GE: $m \times n$ matrix $A \longrightarrow$ row echelon form $\text{ref}(A)$



$\text{ref}(A) = E_p \cdots E_2 E_1 A$ for some elementary matrices E_1, \dots, E_p .

Gauss-Jordan: $A \longrightarrow$ reduced row echelon form $\text{rref}(A)$



$\text{rref}(A) = \hat{E}_\ell \cdots \hat{E}_2 \hat{E}_1 A$ for some elementary matrices $\hat{E}_1, \dots, \hat{E}_\ell$.

Elementary matrices and echelon form

Forward GE: $m \times n$ matrix $A \longrightarrow$ row echelon form $\text{ref}(A)$



$\text{ref}(A) = E_p \cdots E_2 E_1 A$ for some elementary matrices E_1, \dots, E_p .

Gauss-Jordan: $A \longrightarrow$ reduced row echelon form $\text{rref}(A)$



$\text{rref}(A) = \hat{E}_\ell \cdots \hat{E}_2 \hat{E}_1 A$ for some elementary matrices $\hat{E}_1, \dots, \hat{E}_\ell$.

Fact: Let $[A \mid b] \longrightarrow \text{ref}([A \mid b]) =: [U \mid d]$. Then the system $Ax = b$ and $Ux = d$ are equivalent.

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

A

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$A \xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$$

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$A \xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix}$$

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$A \xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix}$$
$$\xrightarrow{E_{32}(-1), E_{42}(-1/2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & -2 & -5 \end{bmatrix}$$

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$\begin{aligned} A &\xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix} \\ &\xrightarrow{E_{32}(-1), E_{42}(-1/2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & -2 & -5 \end{bmatrix} \xrightarrow{E_{43}(2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$\begin{aligned} A &\xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix} \\ &\xrightarrow{E_{32}(-1), E_{42}(-1/2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & -2 & -5 \end{bmatrix} \xrightarrow{E_{43}(2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

(Gaussian Elimination stops here.)

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$\begin{aligned} A &\xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix} \\ &\xrightarrow{E_{32}(-1), E_{42}(-1/2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & -2 & -5 \end{bmatrix} \xrightarrow{E_{43}(2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

(Gaussian Elimination stops here. Continue with GJE.)

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$\begin{aligned} A &\xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix} \\ &\xrightarrow{E_{32}(-1), E_{42}(-1/2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & -2 & -5 \end{bmatrix} \xrightarrow{E_{43}(2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

(Gaussian Elimination stops here. Continue with GJE.)

$$\xrightarrow{E_{34}(-3), E_{24}(-4)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$\begin{aligned} A &\xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix} \\ &\xrightarrow{E_{32}(-1), E_{42}(-1/2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & -2 & -5 \end{bmatrix} \xrightarrow{E_{43}(2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

(Gaussian Elimination stops here. Continue with GJE.)

$$\xrightarrow{E_{34}(-3), E_{24}(-4)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{23}(4)} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

Example

Find rref of $A = \begin{bmatrix} 0 & 2 & -4 & 4 \\ 1 & 0 & 2 & 0 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix}$.

$$\begin{aligned} A &\xrightarrow{E_{12}} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 2 & 2 & 1 & 7 \\ 2 & 1 & 0 & -3 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{14}(-2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 2 & -3 & 7 \\ 0 & 1 & -4 & -3 \end{bmatrix} \\ &\xrightarrow{E_{32}(-1), E_{42}(-1/2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & -2 & -5 \end{bmatrix} \xrightarrow{E_{43}(2)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 4 \\ 0 & 0 & 1 & 3 \\ 0 & 0 & 0 & 1 \end{bmatrix} \end{aligned}$$

(Gaussian Elimination stops here. Continue with GJE.)

$$\begin{aligned} &\xrightarrow{E_{34}(-3), E_{24}(-4)} \begin{bmatrix} 1 & 0 & 2 & 0 \\ 0 & 2 & -4 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_{13}(-2), E_{23}(4)} \begin{bmatrix} 1 & 0 & 0 & 0 \\ 0 & 2 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \xrightarrow{E_2(1/2)} I_4 \end{aligned}$$

Gauss-Jordan method for computing inverse

Question: Suppose $\text{rref}(A) = I_n$. Is A invertible?

Gauss-Jordan method for computing inverse

Question: Suppose $\text{rref}(A) = I_n$. Is A invertible?

Fact: Let A be an $n \times n$ matrix. If $E_p \cdots E_2 E_1 A = I_n$ then $A^{-1} = E_p \cdots E_2 E_1$, where E_1, \dots, E_p are elementary matrices.

Observation: Elementary row operations that transform A to I_n transform I_n to A^{-1} .

Gauss-Jordan method for computing inverse

Question: Suppose $\text{rref}(A) = I_n$. Is A invertible?

Fact: Let A be an $n \times n$ matrix. If $E_p \cdots E_2 E_1 A = I_n$ then $A^{-1} = E_p \cdots E_2 E_1$, where E_1, \dots, E_p are elementary matrices.

Observation: Elementary row operations that transform A to I_n transform I_n to A^{-1} .

Note: If A is invertible, then the rref of $[A \mid I_n]$ is given by

$$\text{rref}([A \mid I_n]) = [E_k \cdots E_2 E_1 A \mid E_k \cdots E_2 E_1] =$$

Gauss-Jordan method for computing inverse

Question: Suppose $\text{rref}(A) = I_n$. Is A invertible?

Fact: Let A be an $n \times n$ matrix. If $E_p \cdots E_2 E_1 A = I_n$ then $A^{-1} = E_p \cdots E_2 E_1$, where E_1, \dots, E_p are elementary matrices.

Observation: Elementary row operations that transform A to I_n transform I_n to A^{-1} .

Note: If A is invertible, then the rref of $[A \mid I_n]$ is given by

$$\text{rref}([A \mid I_n]) = [E_k \cdots E_2 E_1 A \mid E_k \cdots E_2 E_1] = [I_n \mid A^{-1}].$$

Gauss-Jordan method for computing inverse

Question: Suppose $\text{rref}(A) = I_n$. Is A invertible?

Fact: Let A be an $n \times n$ matrix. If $E_p \cdots E_2 E_1 A = I_n$ then $A^{-1} = E_p \cdots E_2 E_1$, where E_1, \dots, E_p are elementary matrices.

Observation: Elementary row operations that transform A to I_n transform I_n to A^{-1} .

Note: If A is invertible, then the rref of $[A \mid I_n]$ is given by

$$\text{rref}([A \mid I_n]) = [E_k \cdots E_2 E_1 A \mid E_k \cdots E_2 E_1] = [I_n \mid A^{-1}].$$

So to find A^{-1} , use GJE to $[A \mid I_n]$.

Gauss-Jordan method:

$$[A \mid I_n] \longrightarrow [I_n \mid X] \Rightarrow A \text{ is invertible and } A^{-1} = X.$$

Example: Gauss-Jordan method

Let $A := \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ 3 & 7 & 9 \end{bmatrix}$. Then $[A \mid I] \rightarrow [I \mid A^{-1}]$ gives

Example: Gauss-Jordan method

Let $A := \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ 3 & 7 & 9 \end{bmatrix}$. Then $[A \mid I] \rightarrow [I \mid A^{-1}]$ gives

$$\left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 5 & 7 & 0 & 1 & 0 \\ 3 & 7 & 9 & 0 & 0 & 1 \end{array} \right] \rightarrow \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & 1 & -2 & 1 & 0 \\ 0 & 1 & 0 & -3 & 0 & 1 \end{array} \right] \rightarrow$$

Example: Gauss-Jordan method

Let $A := \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ 3 & 7 & 9 \end{bmatrix}$. Then $[A \mid I] \rightarrow [I \mid A^{-1}]$ gives

$$\begin{aligned} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 5 & 7 & 0 & 1 & 0 \\ 3 & 7 & 9 & 0 & 0 & 1 \end{array} \right] &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & 1 & -2 & 1 & 0 \\ 0 & 1 & 0 & -3 & 0 & 1 \end{array} \right] \rightarrow \\ \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 5 & -2 & 0 \\ 0 & 1 & 1 & -2 & 1 & 0 \\ 0 & 0 & -1 & -1 & -1 & 1 \end{array} \right] &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 4 & -3 & 1 \\ 0 & 1 & 0 & -3 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & -1 \end{array} \right] \end{aligned}$$

Example: Gauss-Jordan method

Let $A := \begin{bmatrix} 1 & 2 & 3 \\ 2 & 5 & 7 \\ 3 & 7 & 9 \end{bmatrix}$. Then $[A \mid I] \rightarrow [I \mid A^{-1}]$ gives

$$\begin{aligned} \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 2 & 5 & 7 & 0 & 1 & 0 \\ 3 & 7 & 9 & 0 & 0 & 1 \end{array} \right] &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 0 & 0 \\ 0 & 1 & 1 & -2 & 1 & 0 \\ 0 & 1 & 0 & -3 & 0 & 1 \end{array} \right] \rightarrow \\ \left[\begin{array}{ccc|ccc} 1 & 0 & 1 & 5 & -2 & 0 \\ 0 & 1 & 1 & -2 & 1 & 0 \\ 0 & 0 & -1 & -1 & -1 & 1 \end{array} \right] &\rightarrow \left[\begin{array}{ccc|ccc} 1 & 0 & 0 & 4 & -3 & 1 \\ 0 & 1 & 0 & -3 & 0 & 1 \\ 0 & 0 & 1 & 1 & 1 & -1 \end{array} \right] \\ \Rightarrow A^{-1} &= \begin{bmatrix} 4 & -3 & 1 \\ -3 & 0 & 1 \\ 1 & 1 & -1 \end{bmatrix}. \end{aligned}$$

Characterization of invertibility

Theorem: Let A be an $n \times n$ matrix. Then the following statements are equivalent.

- 1 A is invertible.
- 2 $Ax = b$ has a unique solution for every b in \mathbb{R}^n .
- 3 $Ax = 0$ has only the trivial solution.
- 4 The reduced row echelon form of A is I_n .
- 5 A is a product of elementary matrices.
- 6 $\text{rank}(A) = n$

LU Factorization

An $n \times n$ matrix A has an LU factorization if $A = LU$, where U is upper triangular and L is **unit lower triangular** (diagonals are 1).

Fact: If $A \rightarrow \text{ref}(A)$ without row interchange then A has an LU factorization.

LU Factorization

An $n \times n$ matrix A has an LU factorization if $A = LU$, where U is upper triangular and L is **unit lower triangular** (diagonals are 1).

Fact: If $A \rightarrow \text{ref}(A)$ without row interchange then A has an LU factorization.

- $E_p \dots E_2 E_1 A = \text{ref}(A) \Rightarrow A = LU$.
- $L := E_1^{-1} E_2^{-1} \dots E_p^{-1}$ and $U := \text{ref}(A)$.
- Each E_j is unit lower triangular and Type-II $\Rightarrow L$ is unit lower triangular.

LU Factorization

An $n \times n$ matrix A has an LU factorization if $A = LU$, where U is upper triangular and L is **unit lower triangular** (diagonals are 1).

Fact: If $A \rightarrow \text{ref}(A)$ without row interchange then A has an **LU factorization**.

- $E_p \dots E_2 E_1 A = \text{ref}(A) \Rightarrow A = LU.$
- $L := E_1^{-1} E_2^{-1} \dots E_p^{-1}$ and $U := \text{ref}(A).$
- Each E_j is unit lower triangular and Type-II $\Rightarrow L$ is unit lower triangular.

Solution of $Ax = b$ via LU factorization (if exists):

- Compute $A = LU.$
- Solve $Ly = b$ for y - forward substitution.
- Solve $Ux = y$ for x - back substitution.

LU factorization (cont.)

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$. Set $m_{21} := a_{21}/a_{11}$ and $m_{31} := a_{31}/a_{11}$

when $a_{11} \neq 0$ (pivot) and define

$$E_1 := \begin{bmatrix} 1 & 0 & 0 \\ -m_{21} & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } E_2 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -m_{31} & 0 & 1 \end{bmatrix}.$$

LU factorization (cont.)

Let $A = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix}$. Set $m_{21} := a_{21}/a_{11}$ and $m_{31} := a_{31}/a_{11}$

when $a_{11} \neq 0$ (pivot) and define

$$E_1 := \begin{bmatrix} 1 & 0 & 0 \\ -m_{21} & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \text{ and } E_2 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ -m_{31} & 0 & 1 \end{bmatrix}.$$

Then

$$E_2 E_1 A = E_2 \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} \\ 0 & a_{32}^{(1)} & a_{33}^{(1)} \end{bmatrix}.$$

Set $m_{32} := a_{32}^{(1)}/a_{22}^{(1)}$ if $a_{22}^{(1)} \neq 0$ (pivot) and define

LU factorization (cont.)

$$E_3 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -m_{32} & 1 \end{bmatrix}. \text{ Then we have}$$

$$E_3 E_2 E_1 A = E_3 \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} \\ 0 & a_{32}^{(1)} & a_{33}^{(1)} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} \\ 0 & 0 & a_{33}^{(2)} \end{bmatrix} = U.$$

LU factorization (cont.)

$$E_3 := \begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & -m_{32} & 1 \end{bmatrix}. \text{ Then we have}$$

$$E_3 E_2 E_1 A = E_3 \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} \\ 0 & a_{32}^{(1)} & a_{33}^{(1)} \end{bmatrix} = \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ 0 & a_{22}^{(1)} & a_{23}^{(1)} \\ 0 & 0 & a_{33}^{(2)} \end{bmatrix} = U.$$

$$\text{Hence } A = LU, \text{ where } L = E_1^{-1} E_2^{-1} E_3^{-1} = \begin{bmatrix} 1 & 0 & 0 \\ m_{21} & 1 & 0 \\ m_{31} & m_{32} & 1 \end{bmatrix}.$$

Examples: LU factorization

Let $A := \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix}$. Then $A = LU$, where

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} \text{ and } U = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Examples: LU factorization

Let $A := \begin{bmatrix} 1 & 1 & 1 \\ 1 & 2 & 2 \\ 1 & 2 & 3 \end{bmatrix}$. Then $A = LU$, where

$$L = \begin{bmatrix} 1 & 0 & 0 \\ 1 & 1 & 0 \\ 1 & 1 & 1 \end{bmatrix} \text{ and } U = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}.$$

Let $A := \begin{bmatrix} 2 & 4 & -1 \\ -4 & -5 & 3 \\ 2 & -5 & -4 \end{bmatrix}$. Then $A = LU$, where

$$L = \begin{bmatrix} 1 & 0 & 0 \\ -2 & 1 & 0 \\ 1 & -3 & 1 \end{bmatrix} \text{ and } U = \begin{bmatrix} 2 & 4 & -1 \\ 0 & 3 & 1 \\ 0 & 0 & 0 \end{bmatrix}.$$

Three theorems with similar proof structure

Theorem: rref of a matrix is unique. Equivalently, if R_1 and R_2 are in rref and are row equivalent, then $R_1 = R_2$.

Three theorems with similar proof structure

Theorem: rref of a matrix is unique. Equivalently, if R_1 and R_2 are in rref and are row equivalent, then $R_1 = R_2$.

Theorem: Let A and B be $m \times n$ matrices. Then the systems $A\mathbf{x} = \mathbf{0}$ and $B\mathbf{x} = \mathbf{0}$ are equivalent (i.e. have same solutions) **if and only if** A and B are row equivalent.

Three theorems with similar proof structure

Theorem: rref of a matrix is unique. Equivalently, if R_1 and R_2 are in rref and are row equivalent, then $R_1 = R_2$.

Theorem: Let A and B be $m \times n$ matrices. Then the systems $Ax = 0$ and $Bx = 0$ are equivalent (i.e. have same solutions) if and only if A and B are row equivalent.

Theorem: Let A and B be $m \times n$ matrices. Suppose that the systems $Ax = b$ and $Bx = c$ are consistent. Then the two systems are equivalent **if and only if** the matrices $[A \mid b]$ and $[B \mid c]$ are row equivalent.

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Proof: Note that R_1 and R_2 are row equivalent. Hence

$$R_1 \mathbf{x} = \mathbf{0} \iff R_2 \mathbf{x} = \mathbf{0}. \quad (1)$$

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Proof: Note that R_1 and R_2 are row equivalent. Hence

$$R_1 \mathbf{x} = \mathbf{0} \iff R_2 \mathbf{x} = \mathbf{0}. \quad (1)$$

Step-1: Claim $R_1 \mathbf{e}_1 = R_2 \mathbf{e}_1$.

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Proof: Note that R_1 and R_2 are row equivalent. Hence

$$R_1 \mathbf{x} = \mathbf{0} \iff R_2 \mathbf{x} = \mathbf{0}. \quad (1)$$

Step-1: Claim $R_1 \mathbf{e}_1 = R_2 \mathbf{e}_1$.

By (1), we have $R_1 \mathbf{e}_1 = \mathbf{0} \iff R_2 \mathbf{e}_1 = \mathbf{0}$.

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Proof: Note that R_1 and R_2 are row equivalent. Hence

$$R_1 \mathbf{x} = \mathbf{0} \iff R_2 \mathbf{x} = \mathbf{0}. \quad (1)$$

Step-1: Claim $R_1 \mathbf{e}_1 = R_2 \mathbf{e}_1$.

By (1), we have $R_1 \mathbf{e}_1 = \mathbf{0} \iff R_2 \mathbf{e}_1 = \mathbf{0}$.

If $R_1 \mathbf{e}_1 \neq \mathbf{0}$ then $R_1 \mathbf{e}_1$ and $R_2 \mathbf{e}_1$ are pivot columns. Hence

$$R_1 \mathbf{e}_1 = \mathbf{e}_1 = R_2 \mathbf{e}_1.$$

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Proof: Note that R_1 and R_2 are row equivalent. Hence

$$R_1 \mathbf{x} = \mathbf{0} \iff R_2 \mathbf{x} = \mathbf{0}. \quad (1)$$

Step-1: Claim $R_1 \mathbf{e}_1 = R_2 \mathbf{e}_1$.

By (1), we have $R_1 \mathbf{e}_1 = \mathbf{0} \iff R_2 \mathbf{e}_1 = \mathbf{0}$.

If $R_1 \mathbf{e}_1 \neq \mathbf{0}$ then $R_1 \mathbf{e}_1$ and $R_2 \mathbf{e}_1$ are pivot columns. Hence

$$R_1 \mathbf{e}_1 = \mathbf{e}_1 = R_2 \mathbf{e}_1.$$

Step-2: Induction: Suppose $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j$ for $j = 1 : k$. Claim:
 $R_1 \mathbf{e}_{k+1} = R_2 \mathbf{e}_{k+1}$.

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Proof: Note that R_1 and R_2 are row equivalent. Hence

$$R_1 \mathbf{x} = \mathbf{0} \iff R_2 \mathbf{x} = \mathbf{0}. \quad (1)$$

Step-1: Claim $R_1 \mathbf{e}_1 = R_2 \mathbf{e}_1$.

By (1), we have $R_1 \mathbf{e}_1 = \mathbf{0} \iff R_2 \mathbf{e}_1 = \mathbf{0}$.

If $R_1 \mathbf{e}_1 \neq \mathbf{0}$ then $R_1 \mathbf{e}_1$ and $R_2 \mathbf{e}_1$ are **pivot columns**. Hence

$$R_1 \mathbf{e}_1 = \mathbf{e}_1 = R_2 \mathbf{e}_1.$$

Step-2: Induction: Suppose $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j$ for $j = 1 : k$. Claim:
 $R_1 \mathbf{e}_{k+1} = R_2 \mathbf{e}_{k+1}$. If $R_1 \mathbf{e}_{k+1}$ is a **non-pivot column** then there exist $\alpha_1, \dots, \alpha_k$ such that

Uniqueness of rref

Theorem: Let R_1 and R_2 be rref of A . Then $R_1 = R_2$.

Proof: Note that R_1 and R_2 are row equivalent. Hence

$$R_1 \mathbf{x} = \mathbf{0} \iff R_2 \mathbf{x} = \mathbf{0}. \quad (1)$$

Step-1: Claim $R_1 \mathbf{e}_1 = R_2 \mathbf{e}_1$.

By (1), we have $R_1 \mathbf{e}_1 = \mathbf{0} \iff R_2 \mathbf{e}_1 = \mathbf{0}$.

If $R_1 \mathbf{e}_1 \neq \mathbf{0}$ then $R_1 \mathbf{e}_1$ and $R_2 \mathbf{e}_1$ are **pivot columns**. Hence

$$R_1 \mathbf{e}_1 = \mathbf{e}_1 = R_2 \mathbf{e}_1.$$

Step-2: Induction: Suppose $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j$ for $j = 1 : k$. Claim: $R_1 \mathbf{e}_{k+1} = R_2 \mathbf{e}_{k+1}$. If $R_1 \mathbf{e}_{k+1}$ is a **non-pivot column** then there exist $\alpha_1, \dots, \alpha_k$ such that

$$R_1 \mathbf{e}_{k+1} = \alpha_1 R_1 \mathbf{e}_1 + \dots + \alpha_k R_1 \mathbf{e}_k \implies R_1 \mathbf{x} = \mathbf{0}, \quad (2)$$

where $\mathbf{x} := [\alpha_1, \dots, \alpha_k, -1, 0, \dots, 0]^\top$.

Uniqueness of rref

Since $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j, j = 1, 2, \dots, k$, by (1) and (2), we have

Uniqueness of rref

Since $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j, j = 1, 2, \dots, k$, by (1) and (2), we have

$$R_2 \mathbf{x} = \mathbf{0} \Rightarrow R_2 \mathbf{e}_{k+1} = \alpha_1 R_2 \mathbf{e}_1 + \cdots + \alpha_k R_2 \mathbf{e}_k = R_1 \mathbf{e}_{k+1}.$$

Uniqueness of rref

Since $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j, j = 1, 2, \dots, k$, by (1) and (2), we have

$$R_2 \mathbf{x} = \mathbf{0} \Rightarrow R_2 \mathbf{e}_{k+1} = \alpha_1 R_2 \mathbf{e}_1 + \dots + \alpha_k R_2 \mathbf{e}_k = R_1 \mathbf{e}_{k+1}.$$

On the other hand, if $R_1 \mathbf{e}_{k+1}$ is a pivot column then $R_2 \mathbf{e}_{k+1}$ is also a pivot column.

Uniqueness of rref

Since $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j, j = 1, 2, \dots, k$, by (1) and (2), we have

$$R_2 \mathbf{x} = \mathbf{0} \Rightarrow R_2 \mathbf{e}_{k+1} = \alpha_1 R_2 \mathbf{e}_1 + \dots + \alpha_k R_2 \mathbf{e}_k = R_1 \mathbf{e}_{k+1}.$$

On the other hand, if $R_1 \mathbf{e}_{k+1}$ is a pivot column then $R_2 \mathbf{e}_{k+1}$ is also a pivot column.

If there are p pivot columns in the first k columns of R_1 then $R_1 \mathbf{e}_{k+1} = \mathbf{e}_{p+1}$ and $R_2 \mathbf{e}_{k+1} = \mathbf{e}_{p+1}$.

Uniqueness of rref

Since $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j, j = 1, 2, \dots, k$, by (1) and (2), we have

$$R_2 \mathbf{x} = \mathbf{0} \Rightarrow R_2 \mathbf{e}_{k+1} = \alpha_1 R_2 \mathbf{e}_1 + \dots + \alpha_k R_2 \mathbf{e}_k = R_1 \mathbf{e}_{k+1}.$$

On the other hand, if $R_1 \mathbf{e}_{k+1}$ is a pivot column then $R_2 \mathbf{e}_{k+1}$ is also a pivot column.

If there are p pivot columns in the first k columns of R_1 then $R_1 \mathbf{e}_{k+1} = \mathbf{e}_{p+1}$ and $R_2 \mathbf{e}_{k+1} = \mathbf{e}_{p+1}$. Hence $R_1 \mathbf{e}_{k+1} = R_2 \mathbf{e}_{k+1}$.

Uniqueness of rref

Since $R_1 \mathbf{e}_j = R_2 \mathbf{e}_j, j = 1, 2, \dots, k$, by (1) and (2), we have

$$R_2 \mathbf{x} = \mathbf{0} \Rightarrow R_2 \mathbf{e}_{k+1} = \alpha_1 R_2 \mathbf{e}_1 + \dots + \alpha_k R_2 \mathbf{e}_k = R_1 \mathbf{e}_{k+1}.$$

On the other hand, if $R_1 \mathbf{e}_{k+1}$ is a pivot column then $R_2 \mathbf{e}_{k+1}$ is also a pivot column.

If there are p pivot columns in the first k columns of R_1 then $R_1 \mathbf{e}_{k+1} = \mathbf{e}_{p+1}$ and $R_2 \mathbf{e}_{k+1} = \mathbf{e}_{p+1}$. Hence $R_1 \mathbf{e}_{k+1} = R_2 \mathbf{e}_{k+1}$.

Hence we have $R_1 = R_2$. ■

*** End ***