

Linear Algebra

Department of Mathematics
Indian Institute of Technology Guwahati

January – May 2019

MA 102 (RA, RKS, MGPP, KVK)

Linear Transformations

Topics:

- Linear Transformation
- Kernel and Range
- Matrix of a Linear Transformation

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}$ by $(F(p))(x) := p(3)$.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}$ by $(F(p))(x) := p(3)$.

What is common in all of these?

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}$ by $(F(p))(x) := p(3)$.

What is **common** in all of these? Well, they are maps (functions) with **domains** and **codomains** as **vector spaces**.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}$ by $(F(p))(x) := p(3)$.

What is **common** in all of these? Well, they are maps (functions) with **domains** and **codomains** as **vector spaces**. What else?

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}$ by $(F(p))(x) := p(3)$.

What is **common** in all of these? Well, they are maps (functions) with **domains** and **codomains** as **vector spaces**. What else? For all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha, \beta \in \mathbb{F}$, we have

$$F(\mathbf{u} + \mathbf{v}) = F(\mathbf{u}) + F(\mathbf{v}) \text{ and } F(\alpha\mathbf{v}) = \alpha F(\mathbf{v}).$$

Equivalently,

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}$ by $(F(p))(x) := p(3)$.

What is **common** in all of these? Well, they are maps (functions) with **domains** and **codomains** as **vector spaces**. What else? For all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha, \beta \in \mathbb{F}$, we have

$$F(\mathbf{u} + \mathbf{v}) = F(\mathbf{u}) + F(\mathbf{v}) \text{ and } F(\alpha\mathbf{v}) = \alpha F(\mathbf{v}).$$

Equivalently, $F(\alpha\mathbf{u} + \beta\mathbf{v}) = \alpha F(\mathbf{u}) + \beta F(\mathbf{v})$ for $\mathbf{u}, \mathbf{v} \in \mathbb{V}$, $\alpha, \beta \in \mathbb{F}$.

Linear Transformations

- Suppose $A \in \mathcal{M}_{m \times n}$. Let $\mathbf{v} \in \mathbb{R}^n$. Then $A\mathbf{v} \in \mathbb{R}^m$. Thus, we have a map (function) $F : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $F(\mathbf{v}) := A\mathbf{v}$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}[x]$ by $(F(p))(x) := \frac{dp(x)}{dx}$.
- Define $F : \mathcal{C}[0, 1] \rightarrow \mathcal{C}[0, 1]$ by $(F(f))(x) := \int_0^x f(t)dt$.
- Define $F : \mathbb{R}[x] \rightarrow \mathbb{R}$ by $(F(p))(x) := p(3)$.

What is common in all of these? Well, they are maps (functions) with domains and codomains as vector spaces. What else? For all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha, \beta \in \mathbb{F}$, we have

$$F(\mathbf{u} + \mathbf{v}) = F(\mathbf{u}) + F(\mathbf{v}) \text{ and } F(\alpha\mathbf{v}) = \alpha F(\mathbf{v}).$$

Equivalently, $F(\alpha\mathbf{u} + \beta\mathbf{v}) = \alpha F(\mathbf{u}) + \beta F(\mathbf{v})$ for $\mathbf{u}, \mathbf{v} \in \mathbb{V}$, $\alpha, \beta \in \mathbb{F}$. Such functions are called linear transformations (LT).

Linear Transformations

Definition: Let \mathbb{V}, \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a **linear transformation (LT)**

Linear Transformations

Definition: Let \mathbb{V}, \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a **linear transformation (LT)** if $\forall \mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v}) \text{ and } T(\alpha \mathbf{v}) = \alpha T(\mathbf{v}).$$

Linear Transformations

Definition: Let \mathbb{V}, \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a **linear transformation (LT)** if $\forall \mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v}) \text{ and } T(\alpha \mathbf{v}) = \alpha T(\mathbf{v}).$$

Equivalently,

$$T(\alpha \mathbf{u} + \beta \mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v}), \forall \mathbf{u}, \mathbf{v} \in \mathbb{V}, \alpha, \beta \in \mathbb{F}.$$

Linear Transformations

Definition: Let \mathbb{V}, \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a linear transformation (LT) if $\forall \mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v}) \text{ and } T(\alpha \mathbf{v}) = \alpha T(\mathbf{v}).$$

Equivalently,

$$T(\alpha \mathbf{u} + \beta \mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v}), \forall \mathbf{u}, \mathbf{v} \in \mathbb{V}, \alpha, \beta \in \mathbb{F}.$$

Obvious LT's:

Linear Transformations

Definition: Let \mathbb{V}, \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a **linear transformation (LT)** if $\forall \mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v}) \text{ and } T(\alpha \mathbf{v}) = \alpha T(\mathbf{v}).$$

Equivalently,

$$T(\alpha \mathbf{u} + \beta \mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v}), \forall \mathbf{u}, \mathbf{v} \in \mathbb{V}, \alpha, \beta \in \mathbb{F}.$$

Obvious LT's:

- $\mathbf{0} : \mathbb{V} \rightarrow \mathbb{W}, \mathbf{0}(\mathbf{v}) = \mathbf{0}, \forall \mathbf{v} \in \mathbb{V}$ (**zero transformation**).

Linear Transformations

Definition: Let \mathbb{V} , \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a **linear transformation (LT)** if $\forall \mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v}) \text{ and } T(\alpha \mathbf{v}) = \alpha T(\mathbf{v}).$$

Equivalently,

$$T(\alpha \mathbf{u} + \beta \mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v}), \forall \mathbf{u}, \mathbf{v} \in \mathbb{V}, \alpha, \beta \in \mathbb{F}.$$

Obvious LT's:

- $\mathbf{0} : \mathbb{V} \rightarrow \mathbb{W}$, $\mathbf{0}(\mathbf{v}) = \mathbf{0}$, $\forall \mathbf{v} \in \mathbb{V}$ (**zero transformation**).
- $I : \mathbb{V} \rightarrow \mathbb{V}$, $I(\mathbf{v}) = \mathbf{v}$, $\forall \mathbf{v} \in \mathbb{V}$ (**identity transformation**).

Linear Transformations

Definition: Let \mathbb{V}, \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a **linear transformation (LT)** if $\forall \mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v}) \text{ and } T(\alpha \mathbf{v}) = \alpha T(\mathbf{v}).$$

Equivalently,

$$T(\alpha \mathbf{u} + \beta \mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v}), \forall \mathbf{u}, \mathbf{v} \in \mathbb{V}, \alpha, \beta \in \mathbb{F}.$$

Obvious LT's:

- $\mathbf{0} : \mathbb{V} \rightarrow \mathbb{W}, \mathbf{0}(\mathbf{v}) = \mathbf{0}, \forall \mathbf{v} \in \mathbb{V}$ (zero transformation).
- $I : \mathbb{V} \rightarrow \mathbb{V}, I(\mathbf{v}) = \mathbf{v}, \forall \mathbf{v} \in \mathbb{V}$ (identity transformation).

Example: Is $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T([x, y]^T) = [2x, x + y]^T$ an

LT?

Linear Transformations

Definition: Let \mathbb{V}, \mathbb{W} be VS's over \mathbb{F} . A map $T : \mathbb{V} \rightarrow \mathbb{W}$ is called a **linear transformation (LT)** if $\forall \mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$

$$T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v}) \text{ and } T(\alpha \mathbf{v}) = \alpha T(\mathbf{v}).$$

Equivalently,

$$T(\alpha \mathbf{u} + \beta \mathbf{v}) = \alpha T(\mathbf{u}) + \beta T(\mathbf{v}), \forall \mathbf{u}, \mathbf{v} \in \mathbb{V}, \alpha, \beta \in \mathbb{F}.$$

Obvious LT's:

- $\mathbf{0} : \mathbb{V} \rightarrow \mathbb{W}, \mathbf{0}(\mathbf{v}) = \mathbf{0}, \forall \mathbf{v} \in \mathbb{V}$ (zero transformation).
- $I : \mathbb{V} \rightarrow \mathbb{V}, I(\mathbf{v}) = \mathbf{v}, \forall \mathbf{v} \in \mathbb{V}$ (identity transformation).

Example: Is $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ given by $T([x, y]^T) = [2x, x + y]^T$ an

LT? Yes. Note that $T\left(\begin{bmatrix} x \\ y \end{bmatrix}\right) = \begin{bmatrix} 2 & 0 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix}$.

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

① $T(\mathbf{0}) = \mathbf{0};$

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

① $T(\mathbf{0}) = \mathbf{0}$; $[T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).]$

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

① $T(\mathbf{0}) = \mathbf{0}$; $[T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).]$

② $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$;

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

① $T(\mathbf{0}) = \mathbf{0}$; $[T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).]$

② $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$; $[T(-\mathbf{v}) + T\mathbf{v} = \mathbf{0}.]$

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- 1 $T(\mathbf{0}) = \mathbf{0}$; $[T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).]$
- 2 $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$; $[T(-\mathbf{v}) + T\mathbf{v} = \mathbf{0}.]$
- 3 $T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$.

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- ① $T(\mathbf{0}) = \mathbf{0}$; [$T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).$]
- ② $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$; [$T(-\mathbf{v}) + T\mathbf{v} = \mathbf{0}.$]
- ③ $T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$.

Exercise: Is $T : \mathbb{R} \rightarrow \mathbb{R}$, where $T(x) = x + 1$ is an LT?

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- ❶ $T(\mathbf{0}) = \mathbf{0}$; [$T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).$]
- ❷ $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$; [$T(-\mathbf{v}) + T\mathbf{v} = \mathbf{0}.$]
- ❸ $T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$.

Exercise: Is $T : \mathbb{R} \rightarrow \mathbb{R}$, where $T(x) = x + 1$ is an LT? No.

$T(0) \neq 0.$

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- ❶ $T(\mathbf{0}) = \mathbf{0}$; [$T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).$]
- ❷ $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$; [$T(-\mathbf{v}) + T\mathbf{v} = \mathbf{0}.$]
- ❸ $T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$.

Exercise: Is $T : \mathbb{R} \rightarrow \mathbb{R}$, where $T(x) = x + 1$ is an LT? No.

$T(0) \neq 0$.

Exercise: Can there be an LT $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

$T([0, 1]^T) = [2, 3]^T$, $T([1, 0]^T) = [3, 2]^T$ and

$T([1, 1]^T) = [3, 3]^T$?

Basic properties of LT

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- ① $T(\mathbf{0}) = \mathbf{0}$; [$T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).$]
- ② $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$; [$T(-\mathbf{v}) + T\mathbf{v} = \mathbf{0}.$]
- ③ $T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$.

Exercise: Is $T : \mathbb{R} \rightarrow \mathbb{R}$, where $T(x) = x + 1$ is an LT? No.

$$T(0) \neq 0.$$

Exercise: Can there be an LT $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

$$T([0, 1]^T) = [2, 3]^T, \quad T([1, 0]^T) = [3, 2]^T \text{ and}$$

$$T([1, 1]^T) = [3, 3]^T?$$

No, since $[3, 3]^T = T([1, 1]^T) = T(\mathbf{e}_1) + T(\mathbf{e}_2) = [5, 5]^T$

Basic properties of LT

every linear transformation has to satisfy following properties

Fact: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

1 $T(\mathbf{0}) = \mathbf{0}$; $[T(\mathbf{0}) = T(\mathbf{0} + \mathbf{0}) = T(\mathbf{0}) + T(\mathbf{0}) \Rightarrow \mathbf{0} = T(\mathbf{0}).]$

2 $T(-\mathbf{v}) = -T(\mathbf{v})$ for all $\mathbf{v} \in \mathbb{V}$; $[T(-\mathbf{v}) + T\mathbf{v} = \mathbf{0}.]$

3 $T(\mathbf{u} - \mathbf{v}) = T(\mathbf{u}) - T(\mathbf{v})$ for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$.

Exercise: Is $T : \mathbb{R} \rightarrow \mathbb{R}$, where $T(x) = x + 1$ is an LT? No.

$$T(0) \neq 0.$$

Exercise: Can there be an LT $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ such that

$$T([0, 1]^T) = [2, 3]^T, \quad T([1, 0]^T) = [3, 2]^T \text{ and}$$

$$T([1, 1]^T) = [3, 3]^T?$$

No, since $[3, 3]^T = T([1, 1]^T) = T(\mathbf{e}_1) + T(\mathbf{e}_2) = [5, 5]^T$ is not possible.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$).

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} .

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as
$$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n.$$

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as $\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by $T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is **completely determined** by its action on any **basis** of the domain.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is **completely determined** by its action on any **basis** of the domain.

Example: Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$ is an LT.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is **completely determined** by its action on any **basis** of the domain.

Example: Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$ is an LT. Given that $T([1, 0]^\top) = 2 - 3x + x^2$ and $T([0, 1]^\top) = 1 - x^2$.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is **completely determined** by its action on any **basis** of the domain.

Example: Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$ is an LT. Given that $T([1, 0]^\top) = 2 - 3x + x^2$ and $T([0, 1]^\top) = 1 - x^2$. What is $T([2, 3]^\top)$?

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is **completely determined** by its action on any **basis** of the domain.

Example: Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$ is an LT. Given that $T([1, 0]^\top) = 2 - 3x + x^2$ and $T([0, 1]^\top) = 1 - x^2$. What is $T([2, 3]^\top)$? $T([2, 3]^\top) = T(2\mathbf{e}_1 + 3\mathbf{e}_2) =$

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is **completely determined** by its action on any **basis** of the domain.

Example: Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$ is an LT. Given that

$T([1, 0]^\top) = 2 - 3x + x^2$ and $T([0, 1]^\top) = 1 - x^2$. What is

$T([2, 3]^\top)$? $T([2, 3]^\top) = T(2\mathbf{e}_1 + 3\mathbf{e}_2) = 2T(\mathbf{e}_1) + 3T(\mathbf{e}_2) =$

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a basis of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be arbitrarily vectors in \mathbb{W} . Then there is a unique LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is completely determined by its action on any basis of the domain.

Example: Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$ is an LT. Given that

$T([1, 0]^\top) = 2 - 3x + x^2$ and $T([0, 1]^\top) = 1 - x^2$. What is

$T([2, 3]^\top)$? $T([2, 3]^\top) = T(2\mathbf{e}_1 + 3\mathbf{e}_2) = 2T(\mathbf{e}_1) + 3T(\mathbf{e}_2) = 2(2 - 3x + x^2) + 3(1 - x^2) = 7 - 6x - x^2$.

Action of an LT

Theorem: Let $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** of \mathbb{V} ($\dim(\mathbb{V}) = n$). Let $\mathbf{u}_1, \dots, \mathbf{u}_n$ be **arbitrarily vectors** in \mathbb{W} . Then there is a **unique** LT $T : \mathbb{V} \longrightarrow \mathbb{W}$ such that $T(\mathbf{v}_i) = \mathbf{u}_i$ for $i = 1 : n$.

Proof: Let $\mathbf{v} \in \mathbb{V}$. Then \mathbf{v} can be written uniquely as

$\mathbf{v} = \alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n$. Define $T : \mathbb{V} \longrightarrow \mathbb{W}$ by

$T\mathbf{v} := \alpha_1 \mathbf{u}_1 + \dots + \alpha_n \mathbf{u}_n \in \mathbb{W}$. Then T is the required LT. ■

REMARK: An LT is **completely determined** by its action on any **basis** of the domain.

Example: Suppose $T : \mathbb{R}^2 \rightarrow \mathbb{R}_2[x]$ is an LT. Given that

$T([1, 0]^\top) = 2 - 3x + x^2$ and $T([0, 1]^\top) = 1 - x^2$. What is

$T([2, 3]^\top)$? $T([2, 3]^\top) = T(2\mathbf{e}_1 + 3\mathbf{e}_2) = 2T(\mathbf{e}_1) + 3T(\mathbf{e}_2) =$

$2(2 - 3x + x^2) + 3(1 - x^2) = 7 - 6x - x^2$. What is $T([a, b]^\top)$?

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\};$

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$.

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Example: Consider $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T(\mathbf{x}) := A\mathbf{x}$, where $A \in \mathcal{M}_{m \times n}$. What is the range of T ?

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Example: Consider $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T(\mathbf{x}) := A\mathbf{x}$, where $A \in \mathcal{M}_{m \times n}$. What is the range of T ? It is $\{T(\mathbf{x}) \in \mathbb{R}^m \mid \mathbf{x} \in \mathbb{R}^n\}$

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Example: Consider $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T(\mathbf{x}) := A\mathbf{x}$, where $A \in \mathcal{M}_{m \times n}$. What is the range of T ? It is $\{T(\mathbf{x}) \in \mathbb{R}^m \mid \mathbf{x} \in \mathbb{R}^n\} = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} =$

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Example: Consider $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T(\mathbf{x}) := A\mathbf{x}$, where $A \in \mathcal{M}_{m \times n}$. What is the range of T ? It is $\{T(\mathbf{x}) \in \mathbb{R}^m \mid \mathbf{x} \in \mathbb{R}^n\} = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} = \text{col}(A) \subseteq \mathbb{R}^m$.

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Example: Consider $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T(\mathbf{x}) := A\mathbf{x}$, where $A \in \mathcal{M}_{m \times n}$. What is the range of T ? It is $\{T(\mathbf{x}) \in \mathbb{R}^m \mid \mathbf{x} \in \mathbb{R}^n\} = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} = \text{col}(A) \subseteq \mathbb{R}^m$. On the other hand, $\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\} =$

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Example: Consider $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T(\mathbf{x}) := A\mathbf{x}$, where $A \in \mathcal{M}_{m \times n}$. What is the range of T ? It is $\{T(\mathbf{x}) \in \mathbb{R}^m \mid \mathbf{x} \in \mathbb{R}^n\} = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} = \text{col}(A) \subseteq \mathbb{R}^m$. On the other hand, $\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\} = \{\mathbf{x} \in \mathbb{R}^n \mid T(\mathbf{x}) = \mathbf{0}\} = \ker(T)$ is a subspace of \mathbb{R}^n .

Kernel and range of an LT

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT.

- Kernel (or null space) of T : $\ker(T) := \{\mathbf{v} \in \mathbb{V} \mid T(\mathbf{v}) = \mathbf{0}\}$;
- range of T : $\text{range}(T) := \{T(\mathbf{v}) \in \mathbb{W} \mid \mathbf{v} \in \mathbb{V}\}$.

Fact: We have $\ker(T) \preceq \mathbb{V}$ and $\text{range}(T) \preceq \mathbb{W}$. Moreover,

If $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ spans V , then $T(B) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ spans $\text{range}(T)$.

Example: Consider $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ given by $T(\mathbf{x}) := A\mathbf{x}$, where $A \in \mathcal{M}_{m \times n}$. What is the range of T ? It is $\{T(\mathbf{x}) \in \mathbb{R}^m \mid \mathbf{x} \in \mathbb{R}^n\} = \{A\mathbf{x} \mid \mathbf{x} \in \mathbb{R}^n\} = \text{col}(A) \subseteq \mathbb{R}^m$. On the other hand, $\text{null}(A) = \{\mathbf{x} \in \mathbb{R}^n \mid A\mathbf{x} = \mathbf{0}\} = \{\mathbf{x} \in \mathbb{R}^n \mid T(\mathbf{x}) = \mathbf{0}\} = \ker(T)$ is a subspace of \mathbb{R}^n .

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of range}(T),$

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of } \text{range}(T),$
- $\text{nullity}(T) := \text{dimension of } \ker(T).$

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of } \text{range}(T),$
- $\text{nullity}(T) := \text{dimension of } \ker(T).$

Example: Let $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ be defined by $D(p(x)) = p'(x).$

Find $\ker(D)$, $\text{range}(D)$ and their dimensions.

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of } \text{range}(T),$
- $\text{nullity}(T) := \text{dimension of } \ker(T).$

Example: Let $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ be defined by $D(p(x)) = p'(x).$

Find $\ker(D)$, $\text{range}(D)$ and their dimensions. $\text{rank}(D) = 3$ and

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of range}(T)$,
- $\text{nullity}(T) := \text{dimension of ker}(T)$.

Example: Let $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ be defined by $D(p(x)) = p'(x)$.

Find $\text{ker}(D)$, $\text{range}(D)$ and their dimensions. $\text{rank}(D) = 3$ and $\text{nullity}(D) = 1$.

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of range}(T)$,
- $\text{nullity}(T) := \text{dimension of ker}(T)$.

Example: Let $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ be defined by $D(p(x)) = p'(x)$.

Find $\text{ker}(D)$, $\text{range}(D)$ and their dimensions. $\text{rank}(D) = 3$ and $\text{nullity}(D) = 1$.

The Rank-Nullity Theorem: Let \mathbb{V} be finite dimensional and $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

$$\text{rank}(T) + \text{nullity}(T) = \dim(V).$$

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of range}(T)$,
- $\text{nullity}(T) := \text{dimension of ker}(T)$.

Example: Let $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ be defined by $D(p(x)) = p'(x)$.

Find $\text{ker}(D)$, $\text{range}(D)$ and their dimensions. $\text{rank}(D) = 3$ and $\text{nullity}(D) = 1$.

The Rank-Nullity Theorem: Let \mathbb{V} be finite dimensional and $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

$$\text{rank}(T) + \text{nullity}(T) = \dim(V).$$

Proof: Take a basis $B = \{v_1, \dots, v_k\}$ of $\text{ker}(T)$.

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of range}(T)$,
- $\text{nullity}(T) := \text{dimension of ker}(T)$.

Example: Let $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ be defined by $D(p(x)) = p'(x)$.

Find $\text{ker}(D)$, $\text{range}(D)$ and their dimensions. $\text{rank}(D) = 3$ and $\text{nullity}(D) = 1$.

The Rank-Nullity Theorem: Let \mathbb{V} be finite dimensional and $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

$$\text{rank}(T) + \text{nullity}(T) = \dim(\mathbb{V}).$$

Proof: Take a basis $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ of $\text{ker}(T)$. Extend it to a basis $B \cup \{\mathbf{v}_{k+1}, \dots, \mathbf{v}_n\}$ of \mathbb{V} .

Rank-nullity theorem

Definition: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. We define

- $\text{rank}(T) := \text{dimension of range}(T)$,
- $\text{nullity}(T) := \text{dimension of ker}(T)$.

Example: Let $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ be defined by $D(p(x)) = p'(x)$. Find $\text{ker}(D)$, $\text{range}(D)$ and their dimensions. $\text{rank}(D) = 3$ and $\text{nullity}(D) = 1$.

The Rank-Nullity Theorem: Let \mathbb{V} be finite dimensional and $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

$$\text{rank}(T) + \text{nullity}(T) = \dim(\mathbb{V}).$$

Proof: Take a basis $B = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ of $\text{ker}(T)$. Extend it to a basis $B \cup \{\mathbf{v}_{k+1}, \dots, \mathbf{v}_n\}$ of \mathbb{V} . Then $\{T(\mathbf{v}_{k+1}), \dots, T(\mathbf{v}_n)\}$ spans $\text{range}(T)$. Moreover, $\{T(\mathbf{v}_{k+1}), \dots, T(\mathbf{v}_n)\}$ is LI. ■

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set

$\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set $\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Definition: For $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{F}$, define

$T + S : \mathbb{V} \rightarrow \mathbb{W}$, $\alpha T : \mathbb{V} \rightarrow \mathbb{W}$ by

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set $\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Definition: For $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{F}$, define

$T + S : \mathbb{V} \rightarrow \mathbb{W}$, $\alpha T : \mathbb{V} \rightarrow \mathbb{W}$ by

$$(T + S)(\mathbf{v}) := T(\mathbf{v}) + S(\mathbf{v}),$$

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set $\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Definition: For $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{F}$, define

$T + S : \mathbb{V} \rightarrow \mathbb{W}$, $\alpha T : \mathbb{V} \rightarrow \mathbb{W}$ by

$$(T + S)(\mathbf{v}) := T(\mathbf{v}) + S(\mathbf{v}), \quad (\alpha T)(\mathbf{v}) := \alpha(T(\mathbf{v})), \quad \mathbf{v} \in \mathbb{V}.$$

Exercise: Show that $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is a vector space.

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set $\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Definition: For $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{F}$, define

$T + S : \mathbb{V} \rightarrow \mathbb{W}$, $\alpha T : \mathbb{V} \rightarrow \mathbb{W}$ by

$$(T + S)(\mathbf{v}) := T(\mathbf{v}) + S(\mathbf{v}), \quad (\alpha T)(\mathbf{v}) := \alpha(T(\mathbf{v})), \quad \mathbf{v} \in \mathbb{V}.$$

Exercise: Show that $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is a vector space.

Composition:

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set $\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Definition: For $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{F}$, define

$T + S : \mathbb{V} \rightarrow \mathbb{W}$, $\alpha T : \mathbb{V} \rightarrow \mathbb{W}$ by

$$(T + S)(\mathbf{v}) := T(\mathbf{v}) + S(\mathbf{v}), \quad (\alpha T)(\mathbf{v}) := \alpha(T(\mathbf{v})), \quad \mathbf{v} \in \mathbb{V}.$$

Exercise: Show that $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is a vector space.

Composition: Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be LTs.

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is one-one iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is onto iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set $\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Definition: For $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{F}$, define

$T + S : \mathbb{V} \rightarrow \mathbb{W}$, $\alpha T : \mathbb{V} \rightarrow \mathbb{W}$ by

$$(T + S)(\mathbf{v}) := T(\mathbf{v}) + S(\mathbf{v}), \quad (\alpha T)(\mathbf{v}) := \alpha(T(\mathbf{v})), \quad \mathbf{v} \in \mathbb{V}.$$

Exercise: Show that $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is a vector space.

Composition: Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be LTs. Then the composition $S \circ T : \mathbb{U} \rightarrow \mathbb{W}$ is also an LT.

Composition of LTs

Corollary: Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT. Then

- T is **one-one** iff $\ker(T) = \{\mathbf{0}\}$.
- If $\dim(\mathbb{V}) = \dim(\mathbb{W}) = n$, then T is **onto** iff $\ker(T) = \{\mathbf{0}\}$.

Define $\mathcal{L}(\mathbb{V}, \mathbb{W}) := \{T : \mathbb{V} \rightarrow \mathbb{W} \mid T \text{ is an LT}\}$ and set $\mathcal{L}(\mathbb{V}) := \mathcal{L}(\mathbb{V}, \mathbb{V})$.

Definition: For $S, T \in \mathcal{L}(\mathbb{V}, \mathbb{W})$ and $\alpha \in \mathbb{F}$, define

$T + S : \mathbb{V} \rightarrow \mathbb{W}$, $\alpha T : \mathbb{V} \rightarrow \mathbb{W}$ by

$$(T + S)(\mathbf{v}) := T(\mathbf{v}) + S(\mathbf{v}), \quad (\alpha T)(\mathbf{v}) := \alpha(T(\mathbf{v})), \quad \mathbf{v} \in \mathbb{V}.$$

Exercise: Show that $\mathcal{L}(\mathbb{V}, \mathbb{W})$ is a vector space.

Composition: Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be LTs. Then the composition $S \circ T : \mathbb{U} \rightarrow \mathbb{W}$ is also an LT.

Ordered basis

Definition: Let \mathbb{V} be a VS and $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis .

Ordered basis

Definition: Let \mathbb{V} be a VS and $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis . An **ordered** basis of \mathbb{V} is a basis with a **specific order** of its elements.

Ordered basis

Definition: Let \mathbb{V} be a VS and $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis . An **ordered** basis of \mathbb{V} is a basis with a **specific order** of its elements.

We write an **ordered basis** as $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$. Note that $C := [\mathbf{v}_2, \mathbf{v}_1, \mathbf{v}_3, \dots, \mathbf{v}_n]$ is an ordered basis and $B \neq C$.

Ordered basis

Definition: Let \mathbb{V} be a VS and $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis. An **ordered basis** of \mathbb{V} is a basis with a **specific order** of its elements.

We write an **ordered basis** as $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$. Note that $C := [\mathbf{v}_2, \mathbf{v}_1, \mathbf{v}_3, \dots, \mathbf{v}_n]$ is an ordered basis and $B \neq C$.

Example: Let $A := [\mathbf{a}_1, \dots, \mathbf{a}_n] \in \mathcal{M}_n(\mathbb{F})$ be an invertible matrix. Then A is an **ordered basis** of \mathbb{F}^n . The identity matrix I_n is the **standard ordered basis** of \mathbb{F}^n .

Ordered basis

Definition: Let \mathbb{V} be a VS and $\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis. An **ordered** basis of \mathbb{V} is a basis with a **specific order** of its elements.

We write an **ordered basis** as $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$. Note that $C := [\mathbf{v}_2, \mathbf{v}_1, \mathbf{v}_3, \dots, \mathbf{v}_n]$ is an ordered basis and $B \neq C$.

Example: Let $A := [\mathbf{a}_1, \dots, \mathbf{a}_n] \in \mathcal{M}_n(\mathbb{F})$ be an invertible matrix. Then A is an **ordered basis** of \mathbb{F}^n . The identity matrix I_n is the **standard ordered basis** of \mathbb{F}^n .

Definition: Let $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$ be an ordered basis of \mathbb{V} . For $\mathbf{x} := [x_1, \dots, x_n]^\top \in \mathbb{F}^n$, we define

$$B\mathbf{x} := x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n.$$

Thus $B : \mathbb{F}^n \longrightarrow \mathbb{V}, \mathbf{x} \longmapsto B\mathbf{x}$, is an LT.

Coordinates

Theorem: Let $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$ be an ordered basis of \mathbb{V} . Then the LT $B : \mathbb{F}^n \longrightarrow \mathbb{V}, \mathbf{x} \longmapsto B\mathbf{x}$, is bijective.

Coordinates

Theorem: Let $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$ be an ordered basis of \mathbb{V} . Then the LT $B : \mathbb{F}^n \longrightarrow \mathbb{V}, \mathbf{x} \longmapsto B\mathbf{x}$, is bijective.

Proof: $B\mathbf{x} = \mathbf{0} \Rightarrow x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n = \mathbf{0} \Rightarrow \mathbf{x} = \mathbf{0}$. Hence B is injective. By rank-nullity theorem B is onto. ■

Coordinates

Theorem: Let $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$ be an ordered basis of \mathbb{V} . Then the LT $B : \mathbb{F}^n \longrightarrow \mathbb{V}, \mathbf{x} \longmapsto B\mathbf{x}$, is bijective.

Proof: $B\mathbf{x} = \mathbf{0} \Rightarrow x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n = \mathbf{0} \Rightarrow \mathbf{x} = \mathbf{0}$. Hence B is injective. By rank-nullity theorem B is onto. ■

Definition: Let $\mathbf{v} \in \mathbb{V}$. Then there is a unique $\mathbf{x} := [x_1, \dots, x_n]^T$ in \mathbb{F}^n such that $\mathbf{v} = B\mathbf{x} = x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n$. We say that \mathbf{x} is the **coordinate** of \mathbf{v} with respect to the **ordered basis** B and denote \mathbf{x} by $[\mathbf{v}]_B$. Thus $\mathbf{v} = B[\mathbf{v}]_B$ for all $\mathbf{v} \in \mathbb{V}$.

Coordinates

Theorem: Let $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$ be an ordered basis of \mathbb{V} . Then the LT $B : \mathbb{F}^n \longrightarrow \mathbb{V}, \mathbf{x} \longmapsto B\mathbf{x}$, is bijective.

Proof: $B\mathbf{x} = \mathbf{0} \Rightarrow x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n = \mathbf{0} \Rightarrow \mathbf{x} = \mathbf{0}$. Hence B is injective. By rank-nullity theorem B is onto. ■

Definition: Let $\mathbf{v} \in \mathbb{V}$. Then there is a unique $\mathbf{x} := [x_1, \dots, x_n]^\top$ in \mathbb{F}^n such that $\mathbf{v} = B\mathbf{x} = x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n$. We say that \mathbf{x} is the **coordinate** of \mathbf{v} with respect to the **ordered basis** B and denote \mathbf{x} by $[\mathbf{v}]_B$. Thus $\mathbf{v} = B[\mathbf{v}]_B$ for all $\mathbf{v} \in \mathbb{V}$.

Remark: Note that the coordinate $[\mathbf{v}]_B$ depends B as well as on the order of the basis in B .

Example: Consider the standard ordered basis $B := [1, x, x^2]$ of $\mathbb{R}_2[x]$.

Coordinates

Theorem: Let $B := [\mathbf{v}_1, \dots, \mathbf{v}_n]$ be an ordered basis of \mathbb{V} . Then the LT $B : \mathbb{F}^n \longrightarrow \mathbb{V}, \mathbf{x} \longmapsto B\mathbf{x}$, is bijective.

Proof: $B\mathbf{x} = \mathbf{0} \Rightarrow x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n = \mathbf{0} \Rightarrow \mathbf{x} = \mathbf{0}$. Hence B is injective. By rank-nullity theorem B is onto. ■

Definition: Let $\mathbf{v} \in \mathbb{V}$. Then there is a unique $\mathbf{x} := [x_1, \dots, x_n]^\top$ in \mathbb{F}^n such that $\mathbf{v} = B\mathbf{x} = x_1\mathbf{v}_1 + \dots + x_n\mathbf{v}_n$. We say that \mathbf{x} is the **coordinate** of \mathbf{v} with respect to the **ordered basis** B and denote \mathbf{x} by $[\mathbf{v}]_B$. Thus $\mathbf{v} = B[\mathbf{v}]_B$ for all $\mathbf{v} \in \mathbb{V}$.

Remark: Note that the coordinate $[\mathbf{v}]_B$ depends B as well as on the order of the basis in B .

Example: Consider the standard ordered basis $B := [1, x, x^2]$ of $\mathbb{R}_2[x]$. Let $p(x) := a + bx + cx^2$. Then, $[p]_B = [a, b, c]^\top$.

Isomorphism

Theorem: The coordinate map $S : \mathbb{V} \longrightarrow \mathbb{F}^n$ given by $S\mathbf{v} := [\mathbf{v}]_B$ is an LT. Thus, for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$, we have

$$[\mathbf{u} + \mathbf{v}]_B = [\mathbf{u}]_B + [\mathbf{v}]_B \quad \text{and} \quad [\alpha\mathbf{u}]_B = \alpha[\mathbf{u}]_B.$$

Further, S is bijective and is the inverse of B , that is, $SB = I_n$ and $BS = I_{\mathbb{V}}$.

Isomorphism

Theorem: The coordinate map $S : \mathbb{V} \longrightarrow \mathbb{F}^n$ given by $S\mathbf{v} := [\mathbf{v}]_B$ is an LT. Thus, for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$, we have

$$[\mathbf{u} + \mathbf{v}]_B = [\mathbf{u}]_B + [\mathbf{v}]_B \quad \text{and} \quad [\alpha\mathbf{u}]_B = \alpha[\mathbf{u}]_B.$$

Further, S is bijective and is the inverse of B , that is, $SB = I_n$ and $BS = I_{\mathbb{V}}$.

Proof: Use $\mathbf{v} = B[\mathbf{v}]_B$. ■

Isomorphism

Theorem: The coordinate map $S : \mathbb{V} \longrightarrow \mathbb{F}^n$ given by $S\mathbf{v} := [\mathbf{v}]_B$ is an LT. Thus, for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$, we have

$$[\mathbf{u} + \mathbf{v}]_B = [\mathbf{u}]_B + [\mathbf{v}]_B \quad \text{and} \quad [\alpha\mathbf{u}]_B = \alpha[\mathbf{u}]_B.$$

Further, S is bijective and is the inverse of B , that is, $SB = I_n$ and $BS = I_{\mathbb{V}}$.

Proof: Use $\mathbf{v} = B[\mathbf{v}]_B$. ■

Definition: A linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is called an isomorphism of \mathbb{V} onto \mathbb{W} , if it is one-one and onto.

Isomorphism

Theorem: The coordinate map $S : \mathbb{V} \longrightarrow \mathbb{F}^n$ given by $S\mathbf{v} := [\mathbf{v}]_B$ is an LT. Thus, for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$, we have

$$[\mathbf{u} + \mathbf{v}]_B = [\mathbf{u}]_B + [\mathbf{v}]_B \quad \text{and} \quad [\alpha\mathbf{u}]_B = \alpha[\mathbf{u}]_B.$$

Further, S is bijective and is the inverse of B , that is, $SB = I_n$ and $BS = I_{\mathbb{V}}$.

Proof: Use $\mathbf{v} = B[\mathbf{v}]_B$. ■

Definition: A linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is called an isomorphism of \mathbb{V} onto \mathbb{W} , if it is one-one and onto. In that case, we say that \mathbb{V} is isomorphic to \mathbb{W}

Isomorphism

Theorem: The coordinate map $S : \mathbb{V} \longrightarrow \mathbb{F}^n$ given by $S\mathbf{v} := [\mathbf{v}]_B$ is an LT. Thus, for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$, we have

$$[\mathbf{u} + \mathbf{v}]_B = [\mathbf{u}]_B + [\mathbf{v}]_B \quad \text{and} \quad [\alpha\mathbf{u}]_B = \alpha[\mathbf{u}]_B.$$

Further, S is bijective and is the inverse of B , that is, $SB = I_n$ and $BS = I_{\mathbb{V}}$.

Proof: Use $\mathbf{v} = B[\mathbf{v}]_B$. ■

Definition: A linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is called an isomorphism of \mathbb{V} onto \mathbb{W} , if it is one-one and onto. In that case, we say that \mathbb{V} is isomorphic to \mathbb{W} and we write $\mathbb{V} \cong \mathbb{W}$.

Example: The LT $S : \mathbb{V} \rightarrow \mathbb{F}^n, \mathbf{v} \longrightarrow [\mathbf{v}]_B$, is an isomorphism.

Isomorphism

Theorem: The coordinate map $S : \mathbb{V} \longrightarrow \mathbb{F}^n$ given by $S\mathbf{v} := [\mathbf{v}]_B$ is an LT. Thus, for all $\mathbf{u}, \mathbf{v} \in \mathbb{V}$ and $\alpha \in \mathbb{F}$, we have

$$[\mathbf{u} + \mathbf{v}]_B = [\mathbf{u}]_B + [\mathbf{v}]_B \quad \text{and} \quad [\alpha\mathbf{u}]_B = \alpha[\mathbf{u}]_B.$$

Further, S is bijective and is the inverse of B , that is, $SB = I_n$ and $BS = I_{\mathbb{V}}$.

Proof: Use $\mathbf{v} = B[\mathbf{v}]_B$. ■

Definition: A linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is called an isomorphism of \mathbb{V} onto \mathbb{W} , if it is one-one and onto. In that case, we say that \mathbb{V} is isomorphic to \mathbb{W} and we write $\mathbb{V} \cong \mathbb{W}$.

Example: The LT $S : \mathbb{V} \rightarrow \mathbb{F}^n, \mathbf{v} \longrightarrow [\mathbf{v}]_B$, is an isomorphism.

Thus, any VS over \mathbb{F} of dimension n is isomorphic to \mathbb{F}^n .

The matrix of an LT

Suppose $\dim(\mathbb{V}) = n$, $\dim(\mathbb{W}) = m$,

The matrix of an LT

Suppose $\dim(\mathbb{V}) = n$, $\dim(\mathbb{W}) = m$, $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ an ordered basis of \mathbb{V} ,

The matrix of an LT

Suppose $\dim(\mathbb{V}) = n$, $\dim(\mathbb{W}) = m$, $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ an ordered basis of \mathbb{V} , $C := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m]$ an ordered basis of \mathbb{W}

The matrix of an LT

Suppose $\dim(\mathbb{V}) = n$, $\dim(\mathbb{W}) = m$, $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ an ordered basis of \mathbb{V} , $C := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m]$ an ordered basis of \mathbb{W} and $T : V \rightarrow W$ an LT.

The matrix of an LT

Suppose $\dim(\mathbb{V}) = n$, $\dim(\mathbb{W}) = m$, $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ an ordered basis of \mathbb{V} , $C := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m]$ an ordered basis of \mathbb{W} and $T : V \rightarrow W$ an LT. Then the $m \times n$ matrix A defined by

$$A := \left[[T(\mathbf{v}_1)]_C, [T(\mathbf{v}_2)]_C, \dots, [T(\mathbf{v}_n)]_C \right]$$

is called the **matrix of T** with respect to the **ordered bases B and C** . The matrix A is written as $[T]_{C \leftarrow B}$.

The matrix of an LT

Suppose $\dim(\mathbb{V}) = n$, $\dim(\mathbb{W}) = m$, $B := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ an ordered basis of \mathbb{V} , $C := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_m]$ an ordered basis of \mathbb{W} and $T : V \rightarrow W$ an LT. Then the $m \times n$ matrix A defined by

$$A := [[T(\mathbf{v}_1)]_C, [T(\mathbf{v}_2)]_C, \dots, [T(\mathbf{v}_n)]_C]$$

is called the **matrix of T** with respect to the **ordered bases B and C** . The matrix A is written as $[T]_{C \leftarrow B}$.

Theorem: Let $\mathbf{v} \in \mathbb{V}$. Then $T\mathbf{v} = C[T\mathbf{v}]_C$ and $[T\mathbf{v}]_C = A[\mathbf{v}]_B$.

$$\left\{ \begin{array}{ccc} \mathbf{v} \in \mathbb{V} & \xrightarrow{T} & T(\mathbf{v}) \in \mathbb{W} \\ \downarrow & & \uparrow \\ [\mathbf{v}]_B \in \mathbb{F}^n & \xrightarrow{A} & A[\mathbf{v}]_B = [T(\mathbf{v})]_C \in \mathbb{F}^m \end{array} \right\}$$

The Matrix of an LT

Remark: If $\mathbb{V} = \mathbb{W}$ and $B = C$, then $[T]_{C \leftarrow B}$ is written as $[T]_B$.

The Matrix of an LT

Remark: If $\mathbb{V} = \mathbb{W}$ and $B = C$, then $[T]_{C \leftarrow B}$ is written as $[T]_B$.

The previous result shows that T is **completely determined** by its matrix.

The Matrix of an LT

Remark: If $\mathbb{V} = \mathbb{W}$ and $B = C$, then $[T]_{C \leftarrow B}$ is written as $[T]_B$.

The previous result shows that T is **completely determined** by its matrix. Indeed, if $A := [T]_{C \leftarrow B}$ is known then T is determined as follows.

The Matrix of an LT

Remark: If $\mathbb{V} = \mathbb{W}$ and $B = C$, then $[T]_{C \leftarrow B}$ is written as $[T]_B$.

The previous result shows that T is **completely determined** by its matrix. Indeed, if $A := [T]_{C \leftarrow B}$ is known then T is determined as follows. If $\mathbf{v} := \sum_{i=1}^n a_i \mathbf{v}_i = B[\mathbf{v}]_B$ and

$$A \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

then

The Matrix of an LT

Remark: If $\mathbb{V} = \mathbb{W}$ and $B = C$, then $[T]_{C \leftarrow B}$ is written as $[T]_B$.

The previous result shows that T is **completely determined** by its matrix. Indeed, if $A := [T]_{C \leftarrow B}$ is known then T is determined as follows. If $\mathbf{v} := \sum_{i=1}^n a_i \mathbf{v}_i = B[\mathbf{v}]_B$ and

$$A \begin{bmatrix} a_1 \\ a_2 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ \vdots \\ b_m \end{bmatrix}$$

then $T(\mathbf{v}) = \sum_{j=1}^m b_j \mathbf{u}_j = C[T\mathbf{v}]_C$. Thus $T\mathbf{v} = C[T]_{C \leftarrow B}[\mathbf{v}]_B$.

Example

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be defined by

$$T([x, y, z]^T) = [x - 2y, x + y - 3z]^T.$$

Example

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be defined by

$$T([x, y, z]^T) = [x - 2y, x + y - 3z]^T.$$

Consider the bases $B = [\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3]$ and $C = [\mathbf{e}_1, \mathbf{e}_2]$ for \mathbb{R}^3 and \mathbb{R}^2 , respectively.

Example

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be defined by

$$T([x, y, z]^T) = [x - 2y, x + y - 3z]^T.$$

Consider the bases $B = [\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3]$ and $C = [\mathbf{e}_1, \mathbf{e}_2]$ for \mathbb{R}^3 and \mathbb{R}^2 , respectively. Then

$$A = [T]_{C \leftarrow B} = [T(\mathbf{e}_1), T(\mathbf{e}_2), T(\mathbf{e}_3)] =$$

Example

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be defined by

$$T([x, y, z]^T) = [x - 2y, x + y - 3z]^T.$$

Consider the bases $B = [\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3]$ and $C = [\mathbf{e}_1, \mathbf{e}_2]$ for \mathbb{R}^3 and \mathbb{R}^2 , respectively. Then

$$A = [T]_{C \leftarrow B} = [T(\mathbf{e}_1), T(\mathbf{e}_2), T(\mathbf{e}_3)] = \begin{bmatrix} 1 & -2 & 0 \\ 1 & 1 & -3 \end{bmatrix}.$$

Example

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be defined by

$$T([x, y, z]^T) = [x - 2y, x + y - 3z]^T.$$

Consider the bases $B = [\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3]$ and $C = [\mathbf{e}_1, \mathbf{e}_2]$ for \mathbb{R}^3 and \mathbb{R}^2 , respectively. Then

$$A = [T]_{C \leftarrow B} = [T(\mathbf{e}_1), T(\mathbf{e}_2), T(\mathbf{e}_3)] = \begin{bmatrix} 1 & -2 & 0 \\ 1 & 1 & -3 \end{bmatrix}.$$

What is $T([1, 2, 3]^T)$?

Example

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^2$ be defined by

$$T([x, y, z]^T) = [x - 2y, x + y - 3z]^T.$$

Consider the bases $B = [\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3]$ and $C = [\mathbf{e}_1, \mathbf{e}_2]$ for \mathbb{R}^3 and \mathbb{R}^2 , respectively. Then

$$A = [T]_{C \leftarrow B} = [T(\mathbf{e}_1), T(\mathbf{e}_2), T(\mathbf{e}_3)] = \begin{bmatrix} 1 & -2 & 0 \\ 1 & 1 & -3 \end{bmatrix}.$$

What is $T([1, 2, 3]^T)$? In general $T([x, y, z]^T) = A[x, y, z]^T$.

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$.

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$.

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$. Since $D(1) = 0$, $D(x) = 1$, $D(x^2) = 2x$, $D(x^3) = 3x^2$,

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$. Since $D(1) = 0$, $D(x) = 1$, $D(x^2) = 2x$, $D(x^3) = 3x^2$, we get

$$[D]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$. Since $D(1) = 0$, $D(x) = 1$, $D(x^2) = 2x$, $D(x^3) = 3x^2$, we get

$$[D]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Consider $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3$.

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$. Since $D(1) = 0$, $D(x) = 1$, $D(x^2) = 2x$, $D(x^3) = 3x^2$, we get

$$[D]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Consider $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3$. Then $(Dp)(x) = a_1 + 2a_2x + 3a_3x^2$.

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$. Since $D(1) = 0$, $D(x) = 1$, $D(x^2) = 2x$, $D(x^3) = 3x^2$, we get

$$[D]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Consider $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3$. Then $(Dp)(x) = a_1 + 2a_2x + 3a_3x^2$. Note that

$$[p]_B = \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \end{bmatrix} \quad \text{and}$$

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$. Since $D(1) = 0$, $D(x) = 1$, $D(x^2) = 2x$, $D(x^3) = 3x^2$, we get

$$[D]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Consider $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3$. Then

$(Dp)(x) = a_1 + 2a_2x + 3a_3x^2$. Note that

$$[p]_B = \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \end{bmatrix} \quad \text{and} \quad [D]_B[p]_B = \begin{bmatrix} a_1 \\ 2a_2 \\ 3a_3 \\ 0 \end{bmatrix} =$$

Example

Consider $D : \mathbb{R}_3[x] \rightarrow \mathbb{R}_3[x]$ defined by $(Dp)(x) = p'(x)$. Take the standard (ordered) basis $B = [1, x, x^2, x^3]$ of $\mathbb{R}_3[x]$. Since $D(1) = 0$, $D(x) = 1$, $D(x^2) = 2x$, $D(x^3) = 3x^2$, we get

$$[D]_B = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \\ 0 & 0 & 0 & 0 \end{bmatrix}.$$

Consider $p(x) = a_0 + a_1x + a_2x^2 + a_3x^3$. Then

$(Dp)(x) = a_1 + 2a_2x + 3a_3x^2$. Note that

$$[p]_B = \begin{bmatrix} a_0 \\ a_1 \\ a_2 \\ a_3 \end{bmatrix} \quad \text{and} \quad [D]_B[p]_B = \begin{bmatrix} a_1 \\ 2a_2 \\ 3a_3 \\ 0 \end{bmatrix} = [Dp]_B.$$

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively.

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations.

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Now, the i -th column of $[S \circ T]_{D \leftarrow B}$ is

$$[(S \circ T)\mathbf{v}_i]_D$$

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Now, the i -th column of $[S \circ T]_{D \leftarrow B}$ is

$$[(S \circ T)\mathbf{v}_i]_D = [(S(T\mathbf{v}_i))]_D = [S]_{D \leftarrow C} [T\mathbf{v}_i]_C$$

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Now, the i -th column of $[S \circ T]_{D \leftarrow B}$ is

$$\begin{aligned} [(S \circ T)\mathbf{v}_i]_D &= [(S(T\mathbf{v}_i))]_D = [S]_{D \leftarrow C} [T\mathbf{v}_i]_C \\ &= [S]_{D \leftarrow C} [T]_{C \leftarrow B} [\mathbf{v}_i]_B = \end{aligned}$$

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Now, the i -th column of $[S \circ T]_{D \leftarrow B}$ is

$$\begin{aligned} [(S \circ T)\mathbf{v}_i]_D &= [(S(T\mathbf{v}_i))]_D = [S]_{D \leftarrow C} [T\mathbf{v}_i]_C \\ &= [S]_{D \leftarrow C} [T]_{C \leftarrow B} [\mathbf{v}_i]_B = [S]_{D \leftarrow C} [T]_{C \leftarrow B} \mathbf{e}_i, \end{aligned}$$

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Now, the i -th column of $[S \circ T]_{D \leftarrow B}$ is

$$\begin{aligned} [(S \circ T)\mathbf{v}_i]_D &= [(S(T\mathbf{v}_i))]_D = [S]_{D \leftarrow C} [T\mathbf{v}_i]_C \\ &= [S]_{D \leftarrow C} [T]_{C \leftarrow B} [\mathbf{v}_i]_B = [S]_{D \leftarrow C} [T]_{C \leftarrow B} \mathbf{e}_i, \end{aligned}$$

the i -th column of $[S]_{D \leftarrow C} [T]_{C \leftarrow B}$. ■

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Now, the i -th column of $[S \circ T]_{D \leftarrow B}$ is

$$\begin{aligned} [(S \circ T)\mathbf{v}_i]_D &= [(S(T\mathbf{v}_i))]_D = [S]_{D \leftarrow C} [T\mathbf{v}_i]_C \\ &= [S]_{D \leftarrow C} [T]_{C \leftarrow B} [\mathbf{v}_i]_B = [S]_{D \leftarrow C} [T]_{C \leftarrow B} \mathbf{e}_i, \end{aligned}$$

the i -th column of $[S]_{D \leftarrow C} [T]_{C \leftarrow B}$. ■

Corollary: Let \mathbb{V} be a VS with an ordered basis B and $T, S \in \mathcal{L}(V)$.

Matrix of composition of LTs

Theorem: Let \mathbb{U} , \mathbb{V} and \mathbb{W} be three vector spaces with ordered bases B , C and D , respectively. Let $T : \mathbb{U} \rightarrow \mathbb{V}$ and $S : \mathbb{V} \rightarrow \mathbb{W}$ be linear transformations. Then $[S \circ T]_{D \leftarrow B} = [S]_{D \leftarrow C} [T]_{C \leftarrow B}$.

Proof: We have

$$[S \circ T]_{D \leftarrow B} = \left[[(S \circ T)\mathbf{v}_1]_D, [(S \circ T)\mathbf{v}_2]_D, \dots, [(S \circ T)\mathbf{v}_n]_D \right].$$

Now, the i -th column of $[S \circ T]_{D \leftarrow B}$ is

$$\begin{aligned} [(S \circ T)\mathbf{v}_i]_D &= [(S(T\mathbf{v}_i))]_D = [S]_{D \leftarrow C} [T\mathbf{v}_i]_C \\ &= [S]_{D \leftarrow C} [T]_{C \leftarrow B} [\mathbf{v}_i]_B = [S]_{D \leftarrow C} [T]_{C \leftarrow B} \mathbf{e}_i, \end{aligned}$$

the i -th column of $[S]_{D \leftarrow C} [T]_{C \leftarrow B}$. ■

Corollary: Let \mathbb{V} be a VS with an ordered basis B and $T, S \in \mathcal{L}(V)$. Then $[S \circ T]_B = [S]_B [T]_B$.

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} .

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n .

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n . How are they related?

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n . How are they related?

Define $P_{C \leftarrow B} := [[\mathbf{u}_1]_C, [\mathbf{u}_2]_C, \dots, [\mathbf{u}_n]_C] = [I_{\mathbb{V}}]_{C \leftarrow B}$.

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n . How are they related?

Define $P_{C \leftarrow B} := [[\mathbf{u}_1]_C, [\mathbf{u}_2]_C, \dots, [\mathbf{u}_n]_C] = [I_{\mathbb{V}}]_{C \leftarrow B}$. The matrix $P_{C \leftarrow B}$ is called the **change of basis matrix** from B to C .

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n . How are they related?

Define $P_{C \leftarrow B} := [[\mathbf{u}_1]_C, [\mathbf{u}_2]_C, \dots, [\mathbf{u}_n]_C] = [I_{\mathbb{V}}]_{C \leftarrow B}$. The matrix $P_{C \leftarrow B}$ is called the **change of basis matrix** from B to C . Then

① $P_{C \leftarrow B}[\mathbf{v}]_B = [\mathbf{v}]_C$ for all $\mathbf{v} \in \mathbb{V}$;

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n . How are they related?

Define $P_{C \leftarrow B} := [[\mathbf{u}_1]_C, [\mathbf{u}_2]_C, \dots, [\mathbf{u}_n]_C] = [I_{\mathbb{V}}]_{C \leftarrow B}$. The matrix $P_{C \leftarrow B}$ is called the **change of basis matrix** from B to C . Then

- 1 $P_{C \leftarrow B}[\mathbf{v}]_B = [\mathbf{v}]_C$ for all $\mathbf{v} \in \mathbb{V}$;
- 2 $P_{C \leftarrow B}$ is invertible and $(P_{C \leftarrow B})^{-1} = P_{B \leftarrow C} = [I_{\mathbb{V}}]_{B \leftarrow C}$.

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n . How are they related?

Define $P_{C \leftarrow B} := [[\mathbf{u}_1]_C, [\mathbf{u}_2]_C, \dots, [\mathbf{u}_n]_C] = [I_{\mathbb{V}}]_{C \leftarrow B}$. The matrix $P_{C \leftarrow B}$ is called the **change of basis matrix** from B to C . Then

- 1 $P_{C \leftarrow B}[\mathbf{v}]_B = [\mathbf{v}]_C$ for all $\mathbf{v} \in \mathbb{V}$;
- 2 $P_{C \leftarrow B}$ is invertible and $(P_{C \leftarrow B})^{-1} = P_{B \leftarrow C} = [I_{\mathbb{V}}]_{B \leftarrow C}$.

Exercise: Find the change of basis matrix $P_{C \leftarrow B}$ and $P_{B \leftarrow C}$ for the bases $B = [1, x, x^2]$ and $C = [1 + x, x + x^2, 1 + x^2]$ of $\mathbb{R}_2[x]$.

Change of Basis

Let $B := [\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_n]$ and $C := [\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n]$ be ordered bases of \mathbb{V} . Let $\mathbf{v} \in \mathbb{V}$. Then $[\mathbf{v}]_B$ and $[\mathbf{v}]_C$ are in \mathbb{R}^n . How are they related?

Define $P_{C \leftarrow B} := [[\mathbf{u}_1]_C, [\mathbf{u}_2]_C, \dots, [\mathbf{u}_n]_C] = [I_{\mathbb{V}}]_{C \leftarrow B}$. The matrix $P_{C \leftarrow B}$ is called the **change of basis matrix** from B to C . Then

- 1 $P_{C \leftarrow B}[\mathbf{v}]_B = [\mathbf{v}]_C$ for all $\mathbf{v} \in \mathbb{V}$;
- 2 $P_{C \leftarrow B}$ is invertible and $(P_{C \leftarrow B})^{-1} = P_{B \leftarrow C} = [I_{\mathbb{V}}]_{B \leftarrow C}$.

Exercise: Find the change of basis matrix $P_{C \leftarrow B}$ and $P_{B \leftarrow C}$ for the bases $B = [1, x, x^2]$ and $C = [1 + x, x + x^2, 1 + x^2]$ of $\mathbb{R}_2[x]$. Then find the coordinate vector of $p(x) = 1 + 2x - x^2$ w.r.t. respect to the basis C .

Exercises

- Check whether the following LTs are one-one and onto.
 - $T : \mathbb{R} \rightarrow \mathbb{R}^2$ defined by $T(x) = [x, 0]^T$, $x \in \mathbb{R}$.
 - $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T[x, y]^T = x$, for $[x, y]^T \in \mathbb{R}^2$.
 - $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T[x, y]^T = [-x, -y]^T$, for $[x, y]^T \in \mathbb{R}^2$.

Exercises

- Check whether the following LTs are one-one and onto.
 - $T : \mathbb{R} \rightarrow \mathbb{R}^2$ defined by $T(x) = [x, 0]^T$, $x \in \mathbb{R}$.
 - $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T[x, y]^T = x$, for $[x, y]^T \in \mathbb{R}^2$.
 - $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T[x, y]^T = [-x, -y]^T$, for $[x, y]^T \in \mathbb{R}^2$.
- Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT and $\mathbf{v}_1, \dots, \mathbf{v}_k$ in \mathbb{V} be such that $T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)$ are linearly independent. Are $\mathbf{v}_1, \dots, \mathbf{v}_k$ linearly dependent? Justify?

Exercises

- Check whether the following LTs are one-one and onto.
 - $T : \mathbb{R} \rightarrow \mathbb{R}^2$ defined by $T(x) = [x, 0]^T$, $x \in \mathbb{R}$.
 - $T : \mathbb{R}^2 \rightarrow \mathbb{R}$ defined by $T[x, y]^T = x$, for $[x, y]^T \in \mathbb{R}^2$.
 - $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ defined by $T[x, y]^T = [-x, -y]^T$, for $[x, y]^T \in \mathbb{R}^2$.
- Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be an LT and $\mathbf{v}_1, \dots, \mathbf{v}_k$ in \mathbb{V} be such that $T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)$ are linearly independent. Are $\mathbf{v}_1, \dots, \mathbf{v}_k$ linearly dependent? Justify?
- Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be linear and one-one. If $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ is an LI subset of \mathbb{V} then show that $T(S) = \{T(\mathbf{v}_1), \dots, T(\mathbf{v}_k)\}$ is LI in \mathbb{W} .

Exercises

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by

$$T[x, y]^T = [x - y, -3x + 4y]^T \text{ and}$$

$$S[x, y]^T = [4x + y, 3x + y]^T$$

for $[x, y]^T \in \mathbb{R}^2$. Compute $T \circ S$ and $S \circ T$. What is your observation?

- Let \mathbb{V} and \mathbb{W} be vector spaces over \mathbb{F} .

Exercises

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by

$$T[x, y]^T = [x - y, -3x + 4y]^T \text{ and}$$

$$S[x, y]^T = [4x + y, 3x + y]^T$$

for $[x, y]^T \in \mathbb{R}^2$. Compute $T \circ S$ and $S \circ T$. What is your observation?

- Let \mathbb{V} and \mathbb{W} be vector spaces over \mathbb{F} .
 - If $T : \mathbb{V} \rightarrow \mathbb{W}$ is an one-one and onto (i.e., invertible). linear transformation, then show that $T^{-1} : \mathbb{W} \rightarrow \mathbb{V}$ is an LT.
 - Argue that if \mathbb{V} is isomorphic to \mathbb{W} , then \mathbb{W} is isomorphic to \mathbb{V} .

Exercises

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by
$$T[x, y]^T = [x - y, -3x + 4y]^T \text{ and}$$
$$S[x, y]^T = [4x + y, 3x + y]^T$$
for $[x, y]^T \in \mathbb{R}^2$. Compute $T \circ S$ and $S \circ T$. What is your observation?
- Let \mathbb{V} and \mathbb{W} be vector spaces over \mathbb{F} .
 - If $T : \mathbb{V} \rightarrow \mathbb{W}$ is an one-one and onto (i.e., invertible). linear transformation, then show that $T^{-1} : \mathbb{W} \rightarrow \mathbb{V}$ is an LT.
 - Argue that if \mathbb{V} is isomorphic to \mathbb{W} , then \mathbb{W} is isomorphic to \mathbb{V} .
- Let $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Then show that a linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is one-one iff T is onto.

Exercises

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ and $S : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by
$$T[x, y]^T = [x - y, -3x + 4y]^T \text{ and}$$
$$S[x, y]^T = [4x + y, 3x + y]^T$$
for $[x, y]^T \in \mathbb{R}^2$. Compute $T \circ S$ and $S \circ T$. What is your observation?
- Let \mathbb{V} and \mathbb{W} be vector spaces over \mathbb{F} .
 - If $T : \mathbb{V} \rightarrow \mathbb{W}$ is an one-one and onto (i.e., invertible). linear transformation, then show that $T^{-1} : \mathbb{W} \rightarrow \mathbb{V}$ is an LT.
 - Argue that if \mathbb{V} is isomorphic to \mathbb{W} , then \mathbb{W} is isomorphic to \mathbb{V} .
- Let $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Then show that a linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ is one-one iff T is onto.
- Let $\dim(\mathbb{V}) = \dim(\mathbb{W})$. Then a one-one linear transformation $T : \mathbb{V} \rightarrow \mathbb{W}$ maps a basis for \mathbb{V} onto a basis for \mathbb{W} .

Exercises

- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces. Then \mathbb{V} is isomorphic to \mathbb{W} iff $\dim(\mathbb{V}) = \dim(\mathbb{W})$.

Exercises

- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces. Then \mathbb{V} is **isomorphic** to \mathbb{W} iff **$\dim(\mathbb{V}) = \dim(\mathbb{W})$** .
- Show that
 - \mathbb{R}^3 and $\mathbb{R}_2[x]$ are **isomorphic**.
 - The vector spaces \mathbb{R}^n and $\mathbb{R}_n[x]$ are **not** isomorphic.

Exercises

- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces. Then \mathbb{V} is **isomorphic** to \mathbb{W} iff **$\dim(\mathbb{V}) = \dim(\mathbb{W})$** .
- Show that
 - \mathbb{R}^3 and $\mathbb{R}_2[x]$ are **isomorphic**.
 - The vector spaces \mathbb{R}^n and $\mathbb{R}_n[x]$ are **not** isomorphic.
- Let $B = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis for a vector space \mathbb{V} , and let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ be vectors in \mathbb{V} . Then **$\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$** is linearly independent in \mathbb{V} if and only if **$\{[\mathbf{u}_1]_B, [\mathbf{u}_2]_B, \dots, [\mathbf{u}_k]_B\}$** is linearly independent in \mathbb{R}^n .

Exercises

- Let \mathbb{V} and \mathbb{W} be two finite dimensional vector spaces. Then \mathbb{V} is **isomorphic** to \mathbb{W} iff **$\dim(\mathbb{V}) = \dim(\mathbb{W})$** .
- Show that
 - \mathbb{R}^3 and $\mathbb{R}_2[x]$ are **isomorphic**.
 - The vector spaces \mathbb{R}^n and $\mathbb{R}_n[x]$ are **not** isomorphic.
- Let $B = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_n\}$ be a basis for a vector space \mathbb{V} , and let $\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k$ be vectors in \mathbb{V} . Then **$\{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}$** is linearly independent in \mathbb{V} if and only if **$\{[\mathbf{u}_1]_B, [\mathbf{u}_2]_B, \dots, [\mathbf{u}_k]_B\}$** is linearly independent in \mathbb{R}^n .
- Suppose $T, S : \mathbb{V} \rightarrow \mathbb{W}$ are LT's, and B and C are ordered bases of \mathbb{V} and \mathbb{W} , resp. Show that

$$[T + S]_{C \leftarrow B} = [T]_{C \leftarrow B} + [S]_{C \leftarrow B},$$

$$[\alpha T]_{C \leftarrow B} = \alpha [T]_{C \leftarrow B}.$$

Exercises

- Let \mathbb{V}, \mathbb{W} be n dimensional with bases B and C , resp., and $T : V \rightarrow W$ an LT. Then T is invertible if and only if the matrix $[T]_{C \leftarrow B}$ is invertible. In that case,

$$([T]_{C \leftarrow B})^{-1} = [T^{-1}]_{B \leftarrow C}.$$

Exercises

- Let V, W be n dimensional with bases B and C , resp., and $T : V \rightarrow W$ an LT. Then T is invertible if and only if the matrix $[T]_{C \leftarrow B}$ is invertible. In that case,

$$([T]_{C \leftarrow B})^{-1} = [T^{-1}]_{B \leftarrow C}.$$

- Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}_1[x]$ be defined by $T([a, b]^T) = a + (a + b)x$ for $[a, b]^T \in \mathbb{R}^2$. Find $[T]_{C \leftarrow B}$ w.r.t. standard bases, show that T is invertible, and thus find T^{-1} .
