

Linear Algebra

Department of Mathematics
Indian Institute of Technology Guwahati

January – May 2019

MA 102 (RA, RKS, MGPP, KVK)

Eigenvalues and eigenvectors

Topics:

- Determinant of Matrices
- Eigenvalues and Eigenvectors

Determinant of a matrix

Determinant of a matrix

The determinant $\det(A)$ of 2×2 matrix A is defined by

$$\det(A) = \det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = a_{11}a_{22} - a_{12}a_{21}.$$

Determinant of a matrix

The determinant $\det(A)$ of 2×2 matrix A is defined by

$$\det(A) = \det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = a_{11}a_{22} - a_{12}a_{21}.$$

If $\det(A) \neq 0$ then $|\det(A)|$ is the **area** of the **parallelogram** formed by the vectors $A\mathbf{e}_1$ and $A\mathbf{e}_2$.

Determinant of a matrix

The determinant $\det(A)$ of 2×2 matrix A is defined by

$$\det(A) = \det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = a_{11}a_{22} - a_{12}a_{21}.$$

If $\det(A) \neq 0$ then $|\det(A)|$ is the **area** of the **parallelogram** formed by the vectors $A\mathbf{e}_1$ and $A\mathbf{e}_2$.

The determinant $\det(A)$ of 3×3 matrix A is defined by

$$\begin{aligned} \det(A) &= \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \\ &= a_{11} \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}. \end{aligned}$$

Determinant of a matrix

The determinant $\det(A)$ of 2×2 matrix A is defined by

$$\det(A) = \det \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix} = a_{11}a_{22} - a_{12}a_{21}.$$

If $\det(A) \neq 0$ then $|\det(A)|$ is the **area** of the **parallelogram** formed by the vectors $A\mathbf{e}_1$ and $A\mathbf{e}_2$.

The determinant $\det(A)$ of 3×3 matrix A is defined by

$$\begin{aligned} \det(A) &= \det \begin{bmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{bmatrix} \\ &= a_{11} \det \begin{bmatrix} a_{22} & a_{23} \\ a_{32} & a_{33} \end{bmatrix} - a_{12} \det \begin{bmatrix} a_{21} & a_{23} \\ a_{31} & a_{33} \end{bmatrix} + a_{13} \det \begin{bmatrix} a_{21} & a_{22} \\ a_{31} & a_{32} \end{bmatrix}. \end{aligned}$$

If $\det(A) \neq 0$ then $|\det(A)|$ is the **area** of the **parallelepiped** formed by the vectors $A\mathbf{e}_1$, $A\mathbf{e}_2$ and $A\mathbf{e}_3$.

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$.

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

where A_{ij} is the submatrix of A obtained by deleting the i -th row and the j -th column of A .

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

where A_{ij} is the submatrix of A obtained by deleting the i -th row and the j -th column of A . Note that $\det(A) \in \mathbb{F}$.

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

where A_{ij} is the submatrix of A obtained by deleting the i -th row and the j -th column of A . Note that $\det(A) \in \mathbb{F}$.

Example

$$\det \begin{bmatrix} 1 & 5 & 0 & 0 \\ 2 & 0 & 8 & 0 \\ 3 & 6 & 9 & 0 \\ 4 & 7 & 10 & 1 \end{bmatrix} =$$

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

where A_{ij} is the submatrix of A obtained by deleting the i -th row and the j -th column of A . Note that $\det(A) \in \mathbb{F}$.

Example

$$\det \begin{bmatrix} 1 & 5 & 0 & 0 \\ 2 & 0 & 8 & 0 \\ 3 & 6 & 9 & 0 \\ 4 & 7 & 10 & 1 \end{bmatrix} = 1 \cdot \det \begin{bmatrix} 0 & 8 & 0 \\ 6 & 9 & 0 \\ 7 & 10 & 1 \end{bmatrix} - 5 \cdot \det \begin{bmatrix} 2 & 8 & 0 \\ 3 & 9 & 0 \\ 4 & 10 & 1 \end{bmatrix} =$$

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

where A_{ij} is the submatrix of A obtained by deleting the i -th row and the j -th column of A . Note that $\det(A) \in \mathbb{F}$.

Example

$$\det \begin{bmatrix} 1 & 5 & 0 & 0 \\ 2 & 0 & 8 & 0 \\ 3 & 6 & 9 & 0 \\ 4 & 7 & 10 & 1 \end{bmatrix} = 1 \cdot \det \begin{bmatrix} 0 & 8 & 0 \\ 6 & 9 & 0 \\ 7 & 10 & 1 \end{bmatrix} - 5 \cdot \det \begin{bmatrix} 2 & 8 & 0 \\ 3 & 9 & 0 \\ 4 & 10 & 1 \end{bmatrix} = -18.$$

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

where A_{ij} is the submatrix of A obtained by deleting the i -th row and the j -th column of A . Note that $\det(A) \in \mathbb{F}$.

Example

$$\det \begin{bmatrix} 1 & 5 & 0 & 0 \\ 2 & 0 & 8 & 0 \\ 3 & 6 & 9 & 0 \\ 4 & 7 & 10 & 1 \end{bmatrix} = 1 \cdot \det \begin{bmatrix} 0 & 8 & 0 \\ 6 & 9 & 0 \\ 7 & 10 & 1 \end{bmatrix} - 5 \cdot \det \begin{bmatrix} 2 & 8 & 0 \\ 3 & 9 & 0 \\ 4 & 10 & 1 \end{bmatrix} = -18.$$

Theorem: The determinant of a **diagonal, upper or lower triangular matrix** is the **product of its diagonal entries**.

Determinant of an $n \times n$ matrix

Definition: Let $A = [a_{ij}] \in \mathcal{M}_n(\mathbb{F})$. The **determinant** of A is defined by

$\det(A) = a_{11}$, if $n = 1$, i.e., $A = [a_{11}]$. For $n \geq 2$:

$$\det(A) = a_{11}\det(A_{11}) - a_{12}\det(A_{12}) + \dots + (-1)^{1+n}a_{1n}\det(A_{1n}),$$

where A_{ij} is the submatrix of A obtained by deleting the i -th row and the j -th column of A . Note that $\det(A) \in \mathbb{F}$.

Example

$$\det \begin{bmatrix} 1 & 5 & 0 & 0 \\ 2 & 0 & 8 & 0 \\ 3 & 6 & 9 & 0 \\ 4 & 7 & 10 & 1 \end{bmatrix} = 1 \cdot \det \begin{bmatrix} 0 & 8 & 0 \\ 6 & 9 & 0 \\ 7 & 10 & 1 \end{bmatrix} - 5 \cdot \det \begin{bmatrix} 2 & 8 & 0 \\ 3 & 9 & 0 \\ 4 & 10 & 1 \end{bmatrix} = -18.$$

Theorem: The determinant of a **diagonal, upper or lower triangular matrix** is the **product of its diagonal entries**.

Proof: Use induction on the size of the matrix.

Properties of Determinant

Let A be a square matrix.

Properties of Determinant

Let A be a square matrix.

- 1 If B is obtained by **interchanging two rows** of A , then $\det(B) = -\det(A)$.

Properties of Determinant

Let A be a square matrix.

- 1 If B is obtained by **interchanging two rows** of A , then $\det(B) = -\det(A)$.
- 2 If A has a **zero row**, then $\det(A) = 0$.

Properties of Determinant

Let A be a square matrix.

- 1 If B is obtained by **interchanging two rows** of A , then $\det(B) = -\det(A)$.
- 2 If A has a **zero row**, then $\det(A) = 0$.
- 3 If A has **two identical rows**, then $\det(A) = 0$.

Properties of Determinant

Let A be a square matrix.

- 1 If B is obtained by **interchanging two rows** of A , then $\det(B) = -\det(A)$.
- 2 If A has a **zero row**, then $\det(A) = 0$.
- 3 If A has **two identical rows**, then $\det(A) = 0$.
- 4 If B is obtained by **multiplying a row** of A by a scalar α , then $\det(B) = \alpha \det(A)$.

Properties of Determinant

Let A be a square matrix.

- 1 If B is obtained by **interchanging two rows** of A , then $\det(B) = -\det(A)$.
- 2 If A has a **zero row**, then $\det(A) = 0$.
- 3 If A has **two identical rows**, then $\det(A) = 0$.
- 4 If B is obtained by **multiplying a row** of A by a scalar α , then $\det(B) = \alpha \det(A)$.
- 5 If the matrices A , B and C are identical except that one of the rows of C is the sum of the corresponding rows of A and B , then $\det(C) = \det(A) + \det(B)$.

Properties of Determinant

Let A be a square matrix.

- 1 If B is obtained by **interchanging two rows** of A , then $\det(B) = -\det(A)$.
- 2 If A has a **zero row**, then $\det(A) = 0$.
- 3 If A has **two identical rows**, then $\det(A) = 0$.
- 4 If B is obtained by **multiplying a row** of A by a scalar α , then $\det(B) = \alpha \det(A)$.
- 5 If the matrices A , B and C are identical except that one of the rows of C is the sum of the corresponding rows of A and B , then $\det(C) = \det(A) + \det(B)$.
- 6 If B is obtained by adding a **multiple of one row** of A to another row, then $\det(B) = \det(A)$.

Properties of Determinant

Let A be a square matrix.

- 1 If B is obtained by **interchanging two rows** of A , then $\det(B) = -\det(A)$.
- 2 If A has a **zero row**, then $\det(A) = 0$.
- 3 If A has **two identical rows**, then $\det(A) = 0$.
- 4 If B is obtained by **multiplying a row** of A by a scalar α , then $\det(B) = \alpha \det(A)$.
- 5 If the matrices A , B and C are identical except that one of the rows of C is the sum of the corresponding rows of A and B , then $\det(C) = \det(A) + \det(B)$.
- 6 If B is obtained by adding a **multiple of one row** of A to another row, then $\det(B) = \det(A)$.
- 7 $\det(\alpha A) = \alpha^n \det A$.

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$.

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\det(A) = (-1)^{i+1} a_{i1} \det(A_{i1}) + \dots + (-1)^{i+n} a_{in} \det(A_{in})$$

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\det(A) = (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in})$$

(Expansion along the i -th row)

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj})\end{aligned}$$

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj}) \\ &\quad \text{(Expansion along the } j\text{-th column).}\end{aligned}$$

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj}) \\ &\quad \text{(Expansion along the } j\text{-th column).}\end{aligned}$$

Theorem:

- Determinants of elementary matrices:

$$\det E_{ij} =$$

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj}) \\ &\quad \text{(Expansion along the } j\text{-th column).}\end{aligned}$$

Theorem:

- Determinants of elementary matrices:

$$\det E_{jj} = -1, \quad \det E_i(\alpha) =$$

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj}) \\ &\quad \text{(Expansion along the } j\text{-th column).}\end{aligned}$$

Theorem:

- Determinants of elementary matrices:

$$\det E_{jj} = -1, \quad \det E_i(\alpha) = \alpha, \quad \det E_{ij}(\alpha) =$$

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj}) \\ &\quad \text{(Expansion along the } j\text{-th column).}\end{aligned}$$

Theorem:

- Determinants of elementary matrices:

$$\det E_{jj} = -1, \quad \det E_i(\alpha) = \alpha, \quad \det E_{ij}(\alpha) = 1.$$

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj}) \\ &\quad \text{(Expansion along the } j\text{-th column).}\end{aligned}$$

Theorem:

- Determinants of elementary matrices:
 $\det E_{jj} = -1, \quad \det E_i(\alpha) = \alpha, \quad \det E_{ij}(\alpha) = 1.$
- For any elementary matrix E , $\det(E) = \det(E^T)$.
- For any elementary matrix E , $\det(EA) = \det(E)\det(A)$.

Expansion of det along rows and columns

Theorem: Let $A \in \mathcal{M}_n(\mathbb{F})$. Then for any $i, j \in \{1, \dots, n\}$

$$\begin{aligned}\det(A) &= (-1)^{i+1} a_{i1} \det(A_{i1}) + \cdots + (-1)^{i+n} a_{in} \det(A_{in}) \\ &\quad \text{(Expansion along the } i\text{-th row)} \\ &= (-1)^{1+j} a_{1j} \det(A_{1j}) + \cdots + (-1)^{n+j} a_{nj} \det(A_{nj}) \\ &\quad \text{(Expansion along the } j\text{-th column).}\end{aligned}$$

Theorem:

- Determinants of elementary matrices:
 $\det E_{jj} = -1, \quad \det E_i(\alpha) = \alpha, \quad \det E_{ij}(\alpha) = 1.$
- For any elementary matrix E , $\det(E) = \det(E^T)$.
- For any elementary matrix E , $\det(EA) = \det(E) \det(A)$.

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$.

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j .

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$$\det(A) \neq 0 \text{ iff}$$

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$$\det(A) \neq 0 \text{ iff } \det(\text{rref}(A)) \neq 0 \text{ iff}$$

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$$\det(A) \neq 0 \text{ iff } \det(\text{rref}(A)) \neq 0 \text{ iff } \text{rref}(A) = I_n \text{ iff}$$

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$\det(A) \neq 0$ iff $\det(\text{rref}(A)) \neq 0$ iff $\text{rref}(A) = I_n$ iff A is invertible.

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$\det(A) \neq 0$ iff $\det(\text{rref}(A)) \neq 0$ iff $\text{rref}(A) = I_n$ iff A is invertible.

Definition A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is said to be **singular** or **non-singular** according as

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$\det(A) \neq 0$ iff $\det(\text{rref}(A)) \neq 0$ iff $\text{rref}(A) = I_n$ iff A is invertible.

Definition A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is said to be **singular** or **non-singular** according as $\det(A) = 0$ or $\det(A) \neq 0$.

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$\det(A) \neq 0$ iff $\det(\text{rref}(A)) \neq 0$ iff $\text{rref}(A) = I_n$ iff A is invertible.

Definition A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is said to be **singular** or **non-singular** according as $\det(A) = 0$ or $\det(A) \neq 0$.

Theorem: A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is **invertible** if and only if it is **non-singular**.

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$\det(A) \neq 0$ iff $\det(\text{rref}(A)) \neq 0$ iff $\text{rref}(A) = I_n$ iff A is invertible.

Definition A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is said to be **singular** or **non-singular** according as $\det(A) = 0$ or $\det(A) \neq 0$.

Theorem: A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is **invertible** if and only if it is **non-singular**.

Theorem:

- (a) For any matrix $A \in \mathcal{M}_n(\mathbb{F})$, $\det(A) = \det(A^T)$.
- (b) For any $A, B \in \mathcal{M}_n(\mathbb{F})$, $\det(AB) = \det(A)\det(B)$.

Two important properties

Suppose $A \in \mathcal{M}_n(\mathbb{F})$. Then $A = E_k \cdots E_1 \cdot \text{rref}(A)$ for some elementary E_j . We have

$$\det(A) = \det(E_k) \cdots \det(E_1) \det(\text{rref}(A)).$$

Thus,

$\det(A) \neq 0$ iff $\det(\text{rref}(A)) \neq 0$ iff $\text{rref}(A) = I_n$ iff A is invertible.

Definition A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is said to be **singular** or **non-singular** according as $\det(A) = 0$ or $\det(A) \neq 0$.

Theorem: A matrix $A \in \mathcal{M}_n(\mathbb{F})$ is **invertible** if and only if it is **non-singular**.

Theorem:

- (a) For any matrix $A \in \mathcal{M}_n(\mathbb{F})$, $\det(A) = \det(A^T)$.
- (b) For any $A, B \in \mathcal{M}_n(\mathbb{F})$, $\det(AB) = \det(A)\det(B)$.

Eigenvalues and Eigenvectors

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation.

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$.

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is nicely described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is nicely described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Remark

- $[T\mathbf{v}_i]_B = \lambda_i [\mathbf{v}_i]_B$.

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is nicely described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Remark

- $[T\mathbf{v}_i]_B = \lambda_i [\mathbf{v}_i]_B$.
- $[T]_B =$

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is nicely described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Remark

- $[T\mathbf{v}_i]_B = \lambda_i [\mathbf{v}_i]_B$.
- $[T]_B = [[T(\mathbf{v}_1)]_B, \dots, [T(\mathbf{v}_n)]_B] =$

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is *nicely* described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Remark

- $[T\mathbf{v}_i]_B = \lambda_i [\mathbf{v}_i]_B$.
- $[T]_B = [[T(\mathbf{v}_1)]_B, \dots, [T(\mathbf{v}_n)]_B] = \text{diag}\{\lambda_1, \dots, \lambda_n\}$.

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is nicely described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Remark

- $[T\mathbf{v}_i]_B = \lambda_i [\mathbf{v}_i]_B$.
- $[T]_B = [[T(\mathbf{v}_1)]_B, \dots, [T(\mathbf{v}_n)]_B] = \text{diag}\{\lambda_1, \dots, \lambda_n\}$.

Definition: Let $T \in \mathcal{L}(\mathbb{V})$. If there is a scalar $\lambda \in \mathbb{F}$ and a nonzero vector $\mathbf{v} \in \mathbb{V}$ such that $T(\mathbf{v}) = \lambda \mathbf{v}$

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is nicely described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Remark

- $[T\mathbf{v}_i]_B = \lambda_i [\mathbf{v}_i]_B$.
- $[T]_B = [[T(\mathbf{v}_1)]_B, \dots, [T(\mathbf{v}_n)]_B] = \text{diag}\{\lambda_1, \dots, \lambda_n\}$.

Definition: Let $T \in \mathcal{L}(\mathbb{V})$. If there is a scalar $\lambda \in \mathbb{F}$ and a nonzero vector $\mathbf{v} \in \mathbb{V}$ such that $T(\mathbf{v}) = \lambda \mathbf{v}$ then λ is called an **eigenvalue** of T and

Eigenvalues and Eigenvectors

Suppose \mathbb{V} is a vector space and $T : \mathbb{V} \rightarrow \mathbb{V}$ is a linear transformation. Suppose $B = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a basis of \mathbb{V} and $\lambda_i \in \mathbb{F}$ are such that $T(\mathbf{v}_i) = \lambda_i \mathbf{v}_i$. Then, T is nicely described by

$$T(\alpha_1 \mathbf{v}_1 + \dots + \alpha_n \mathbf{v}_n) = \lambda_1 \alpha_1 \mathbf{v}_1 + \dots + \lambda_n \alpha_n \mathbf{v}_n.$$

Remark

- $[T\mathbf{v}_i]_B = \lambda_i [\mathbf{v}_i]_B$.
- $[T]_B = [[T(\mathbf{v}_1)]_B, \dots, [T(\mathbf{v}_n)]_B] = \text{diag}\{\lambda_1, \dots, \lambda_n\}$.

Definition: Let $T \in \mathcal{L}(\mathbb{V})$. If there is a scalar $\lambda \in \mathbb{F}$ and a nonzero vector $\mathbf{v} \in \mathbb{V}$ such that $T(\mathbf{v}) = \lambda \mathbf{v}$ then λ is called an **eigenvalue** of T and \mathbf{v} an **eigenvector** of T corresponding to λ .

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$.

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$,

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT.

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff $[T(\mathbf{v})]_B = \lambda[\mathbf{v}]_B$ for any ordered basis B of \mathbb{V} .

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff $[T(\mathbf{v})]_B = \lambda[\mathbf{v}]_B$ for any ordered basis B of \mathbb{V} .

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff $[T(\mathbf{v})]_B = \lambda[\mathbf{v}]_B$ for any ordered basis B of \mathbb{V} .

Thus it is enough to study eigenvalues/eigenvectors of square matrices.

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff $[T(\mathbf{v})]_B = \lambda[\mathbf{v}]_B$ for any ordered basis B of \mathbb{V} .

Thus it is enough to study eigenvalues/eigenvectors of square matrices.

Example: Consider the matrix $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$.

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff $[T(\mathbf{v})]_B = \lambda[\mathbf{v}]_B$ for any ordered basis B of \mathbb{V} .

Thus it is enough to study eigenvalues/eigenvectors of square matrices.

Example: Consider the matrix $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$. Then 4 is an eigenvalue of A with $[1, 1]^T$ as a corresponding eigenvector.

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff $[T(\mathbf{v})]_B = \lambda[\mathbf{v}]_B$ for any ordered basis B of \mathbb{V} .

Thus it is enough to study eigenvalues/eigenvectors of square matrices.

Example: Consider the matrix $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$. Then 4 is an eigenvalue of A with $[1, 1]^T$ as a corresponding eigenvector.

Next, consider the rotation matrix $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$. Then $A\mathbf{v} \neq \lambda\mathbf{v}$ for any $\lambda \in \mathbb{R}$ and $\mathbf{0} \neq \mathbf{v} \in \mathbb{R}^2$.

Eigenvalues and eigenvectors

Definition: Let $A \in \mathcal{M}_n(\mathbb{F})$. If $A\mathbf{x} = \lambda\mathbf{x}$ for some $\lambda \in \mathbb{F}$ and $\mathbf{0} \neq \mathbf{x} \in \mathbb{F}^n$, then λ is called an **eigenvalue** of A and \mathbf{x} an **eigenvector** of A corresponding to λ .

Fact: Suppose $\dim(\mathbb{V}) = n$, and $T : \mathbb{V} \rightarrow \mathbb{V}$ is an LT. Then $T(\mathbf{v}) = \lambda\mathbf{v}$ iff $[T(\mathbf{v})]_B = \lambda[\mathbf{v}]_B$ for any ordered basis B of \mathbb{V} .

Thus it is enough to study eigenvalues/eigenvectors of square matrices.

Example: Consider the matrix $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$. Then 4 is an eigenvalue of A with $[1, 1]^T$ as a corresponding eigenvector.

Next, consider the rotation matrix $A = \begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix}$. Then $A\mathbf{v} \neq \lambda\mathbf{v}$ for any $\lambda \in \mathbb{R}$ and $\mathbf{0} \neq \mathbf{v} \in \mathbb{R}^2$.

Henceforth, we consider $\mathbb{F} = \mathbb{C}$.

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$.

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** .

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Theorem: Let $A \in \mathcal{M}_n(\mathbb{C})$. Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$ is an eigenvalue of A .

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Theorem: Let $A \in \mathcal{M}_n(\mathbb{C})$. Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$ is an eigenvalue of A .
- $E_\lambda = \text{null}(A - \lambda I)$ is a **nontrivial subspace** of \mathbb{C}^n .

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Theorem: Let $A \in \mathcal{M}_n(\mathbb{C})$. Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$ is an eigenvalue of A .
- $E_\lambda = \text{null}(A - \lambda I)$ is a **nontrivial subspace** of \mathbb{C}^n .
- $\text{rank}(A - \lambda I) < n$.

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Theorem: Let $A \in \mathcal{M}_n(\mathbb{C})$. Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$ is an eigenvalue of A .
- $E_\lambda = \text{null}(A - \lambda I)$ is a **nontrivial subspace** of \mathbb{C}^n .
- $\text{rank}(A - \lambda I) < n$.
- $\det(A - \lambda I) = 0$.

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Theorem: Let $A \in \mathcal{M}_n(\mathbb{C})$. Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$ is an eigenvalue of A .
- $E_\lambda = \text{null}(A - \lambda I)$ is a **nontrivial subspace** of \mathbb{C}^n .
- $\text{rank}(A - \lambda I) < n$.
- $\det(A - \lambda I) = 0$.

Definition: Let A be an $n \times n$ matrix.

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Theorem: Let $A \in \mathcal{M}_n(\mathbb{C})$. Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$ is an eigenvalue of A .
- $E_\lambda = \text{null}(A - \lambda I)$ is a **nontrivial subspace** of \mathbb{C}^n .
- $\text{rank}(A - \lambda I) < n$.
- $\det(A - \lambda I) = 0$.

Definition: Let A be an $n \times n$ matrix. Then

- $\det(A - \lambda I)$ is called the **characteristic polynomial** of A .

Eigenspace

Definition: Let λ be an eigenvalue of a matrix $A \in \mathcal{M}_n(\mathbb{C})$. Then

$$E_\lambda = \{\mathbf{x} \mid A\mathbf{x} = \lambda\mathbf{x}\}$$

is called the **eigenspace of λ** . E_λ consists of all eigenvectors of A corresponding to λ , together with the **zero vector**.

Theorem: Let $A \in \mathcal{M}_n(\mathbb{C})$. Then the following conditions are equivalent.

- $\lambda \in \mathbb{C}$ is an eigenvalue of A .
- $E_\lambda = \text{null}(A - \lambda I)$ is a **nontrivial subspace** of \mathbb{C}^n .
- $\text{rank}(A - \lambda I) < n$.
- $\det(A - \lambda I) = 0$.

Definition: Let A be an $n \times n$ matrix. Then

- $\det(A - \lambda I)$ is called the **characteristic polynomial** of A .
- $\det(A - \lambda I) = 0$ is called the **characteristic equation** of A .

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces?

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) =$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \Leftrightarrow \lambda = 4, -2,$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \Leftrightarrow \lambda = 4, -2,$$

$$E_4 = \text{null}(A - 4I) =$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \Leftrightarrow \lambda = 4, -2,$$

$$E_4 = \text{null}(A - 4I) = \left\{ \mathbf{x} : \begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} =$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \Leftrightarrow \lambda = 4, -2,$$

$$E_4 = \text{null}(A - 4I) = \left\{ \mathbf{x} : \begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, 1]^T),$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \Leftrightarrow \lambda = 4, -2,$$

$$E_4 = \text{null}(A - 4I) = \left\{ \mathbf{x} : \begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, 1]^T),$$

$$E_{(-2)} = \text{null}(A + 2I) =$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \Leftrightarrow \lambda = 4, -2,$$

$$E_4 = \text{null}(A - 4I) = \left\{ \mathbf{x} : \begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, 1]^T),$$

$$E_{(-2)} = \text{null}(A + 2I) = \left\{ \mathbf{x} : \begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} =$$

Facts:

- The eigenvalues of A are the **zeros** of the characteristic polynomial $\det(A - \lambda I)$ of A . Hence A has at most n eigenvalues.
- The eigenvalues of a **triangular** matrix are its **diagonal** entries.
- A is **invertible** iff 0 is not an eigenvalue of A .

Example: What are the eigenvalues of $A = \begin{bmatrix} 1 & 3 \\ 3 & 1 \end{bmatrix}$ and their eigenspaces? We have

$$\det(A - \lambda I) = \det \begin{bmatrix} 1 - \lambda & 3 \\ 3 & 1 - \lambda \end{bmatrix} = 0 \Leftrightarrow \lambda = 4, -2,$$

$$E_4 = \text{null}(A - 4I) = \left\{ \mathbf{x} : \begin{bmatrix} -3 & 3 \\ 3 & -3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, 1]^T),$$

$$E_{(-2)} = \text{null}(A + 2I) = \left\{ \mathbf{x} : \begin{bmatrix} 3 & 3 \\ 3 & 3 \end{bmatrix} \mathbf{x} = \mathbf{0} \right\} = \text{span}([1, -1]^T),$$

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.
- 2 Solve the char. eqn. $\det(A - \lambda I) = 0$ for eigenvalues λ .

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.
- 2 Solve the char. eqn. $\det(A - \lambda I) = 0$ for eigenvalues λ .
- 3 For each eigenvalue λ , find a basis of $\text{null}(A - \lambda I) = E_\lambda$.

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.
- 2 Solve the char. eqn. $\det(A - \lambda I) = 0$ for eigenvalues λ .
- 3 For each eigenvalue λ , find a basis of $\text{null}(A - \lambda I) = E_\lambda$.

Example

A real matrix may have complex eigenvalues and complex eigenvectors.

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.
- 2 Solve the char. eqn. $\det(A - \lambda I) = 0$ for eigenvalues λ .
- 3 For each eigenvalue λ , find a basis of $\text{null}(A - \lambda I) = E_\lambda$.

Example

A real matrix may have complex eigenvalues and complex eigenvectors. Consider $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$.

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.
- 2 Solve the char. eqn. $\det(A - \lambda I) = 0$ for eigenvalues λ .
- 3 For each eigenvalue λ , find a basis of $\text{null}(A - \lambda I) = E_\lambda$.

Example

A real matrix may have complex eigenvalues and complex eigenvectors. Consider $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. The char. poly. of A is $\lambda^2 + 1$ and so the eigenvalues are $\pm i$.

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.
- 2 Solve the char. eqn. $\det(A - \lambda I) = 0$ for eigenvalues λ .
- 3 For each eigenvalue λ , find a basis of $\text{null}(A - \lambda I) = E_\lambda$.

Example

A real matrix may have complex eigenvalues and complex eigenvectors. Consider $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. The char. poly. of A is $\lambda^2 + 1$ and so the eigenvalues are $\pm i$. Corresponding to the eigenvalues $\pm i$, it has eigenvectors $[1, i]^\top$ and $[i, 1]^\top$, respectively.

Computation of eigenvalues and bases for eigenspaces

- 1 Compute the char. poly. $\det(A - \lambda I)$.
- 2 Solve the char. eqn. $\det(A - \lambda I) = 0$ for eigenvalues λ .
- 3 For each eigenvalue λ , find a basis of $\text{null}(A - \lambda I) = E_\lambda$.

Example

A real matrix may have complex eigenvalues and complex eigenvectors. Consider $A = \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}$. The char. poly. of A is $\lambda^2 + 1$ and so the eigenvalues are $\pm i$. Corresponding to the eigenvalues $\pm i$, it has eigenvectors $[1, i]^\top$ and $[i, 1]^\top$, respectively. The eigenspaces E_i and $E_{(-i)}$ are one dimensional.

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

- The **algebraic multiplicity** of λ is the multiplicity of λ as a root of the characteristic polynomial of A .

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

- The **algebraic multiplicity** of λ is the multiplicity of λ as a root of the characteristic polynomial of A .
- The **geometric multiplicity** of λ is the dimension of E_λ .

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

- The **algebraic multiplicity** of λ is the multiplicity of λ as a root of the characteristic polynomial of A .
- The **geometric multiplicity** of λ is the dimension of E_λ .

Theorem: Let λ be an eigenvalue of A . Then the **algebraic multiplicity** of $\lambda \geq$ the **geometric multiplicity** of λ .

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

- The **algebraic multiplicity** of λ is the multiplicity of λ as a root of the characteristic polynomial of A .
- The **geometric multiplicity** of λ is the dimension of E_λ .

Theorem: Let λ be an eigenvalue of A . Then the **algebraic multiplicity** of $\lambda \geq$ the **geometric multiplicity** of λ .

Example: What are the eigenvalues and the corresponding

eigenspaces of $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$?

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

- The **algebraic multiplicity** of λ is the multiplicity of λ as a root of the characteristic polynomial of A .
- The **geometric multiplicity** of λ is the dimension of E_λ .

Theorem: Let λ be an eigenvalue of A . Then the **algebraic multiplicity** of $\lambda \geq$ the **geometric multiplicity** of λ .

Example: What are the eigenvalues and the corresponding eigenspaces of $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$? Eigenvalues are 1, 1, 1 (1 appearing three times, **multiplicity** three).

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

- The **algebraic multiplicity** of λ is the multiplicity of λ as a root of the characteristic polynomial of A .
- The **geometric multiplicity** of λ is the dimension of E_λ .

Theorem: Let λ be an eigenvalue of A . Then the **algebraic multiplicity** of $\lambda \geq$ the **geometric multiplicity** of λ .

Example: What are the eigenvalues and the corresponding eigenspaces of $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$? Eigenvalues are 1, 1, 1 (1 appearing three times, **multiplicity** three). Moreover,

$$E_1 = \text{null} \left(\begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \right) = \text{span} \left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right).$$

Algebraic and geometric multiplicities

Definition: Let λ be an eigenvalue of $A \in \mathcal{M}_n(\mathbb{C})$.

- The **algebraic multiplicity** of λ is the multiplicity of λ as a root of the characteristic polynomial of A .
- The **geometric multiplicity** of λ is the dimension of E_λ .

Theorem: Let λ be an eigenvalue of A . Then the **algebraic multiplicity** of $\lambda \geq$ the **geometric multiplicity** of λ .

Example: What are the eigenvalues and the corresponding

eigenspaces of $A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 1 \\ 0 & 0 & 1 \end{bmatrix}$? Eigenvalues are 1, 1, 1 (1

appearing three times, **multiplicity** three). Moreover,

$$E_1 = \text{null} \left(\begin{bmatrix} 0 & 1 & 1 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{bmatrix} \right) = \text{span} \left(\begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \right).$$

We have $\dim(E_1) = 1$.

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are linearly independent.

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A ,

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i .

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI.

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not.

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$.

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$. Then,

$$A\mathbf{x}_k = \lambda_1 \alpha_1 \mathbf{x}_1 + \dots + \lambda_{k-1} \alpha_{k-1} \mathbf{x}_{k-1}.$$

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$. Then,

$$A\mathbf{x}_k = \lambda_1 \alpha_1 \mathbf{x}_1 + \dots + \lambda_{k-1} \alpha_{k-1} \mathbf{x}_{k-1}.$$

Therefore, $\mathbf{0} = \lambda_k \mathbf{x}_k - A\mathbf{x}_k$

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$. Then,

$$A\mathbf{x}_k = \lambda_1 \alpha_1 \mathbf{x}_1 + \dots + \lambda_{k-1} \alpha_{k-1} \mathbf{x}_{k-1}.$$

Therefore,
$$\begin{aligned} \mathbf{0} &= \lambda_k \mathbf{x}_k - A\mathbf{x}_k \\ &= (\lambda_k - \lambda_1) \alpha_1 \mathbf{x}_1 + \dots + (\lambda_k - \lambda_{k-1}) \alpha_{k-1} \mathbf{x}_{k-1}. \end{aligned}$$

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$. Then,

$$A\mathbf{x}_k = \lambda_1 \alpha_1 \mathbf{x}_1 + \dots + \lambda_{k-1} \alpha_{k-1} \mathbf{x}_{k-1}.$$

Therefore,
$$\begin{aligned} \mathbf{0} &= \lambda_k \mathbf{x}_k - A\mathbf{x}_k \\ &= (\lambda_k - \lambda_1) \alpha_1 \mathbf{x}_1 + \dots + (\lambda_k - \lambda_{k-1}) \alpha_{k-1} \mathbf{x}_{k-1}. \end{aligned}$$

Since $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI,

$$(\lambda_k - \lambda_1) \alpha_1 = \dots = (\lambda_k - \lambda_{k-1}) \alpha_{k-1} = 0,$$

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$. Then,

$$A\mathbf{x}_k = \lambda_1 \alpha_1 \mathbf{x}_1 + \dots + \lambda_{k-1} \alpha_{k-1} \mathbf{x}_{k-1}.$$

$$\begin{aligned} \text{Therefore, } \mathbf{0} &= \lambda_k \mathbf{x}_k - A\mathbf{x}_k \\ &= (\lambda_k - \lambda_1) \alpha_1 \mathbf{x}_1 + \dots + (\lambda_k - \lambda_{k-1}) \alpha_{k-1} \mathbf{x}_{k-1}. \end{aligned}$$

Since $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI,

$$(\lambda_k - \lambda_1) \alpha_1 = \dots = (\lambda_k - \lambda_{k-1}) \alpha_{k-1} = 0,$$

i.e., $\alpha_1 = \dots = \alpha_{k-1} = 0$

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$. Then,

$$A\mathbf{x}_k = \lambda_1 \alpha_1 \mathbf{x}_1 + \dots + \lambda_{k-1} \alpha_{k-1} \mathbf{x}_{k-1}.$$

$$\begin{aligned} \text{Therefore, } \mathbf{0} &= \lambda_k \mathbf{x}_k - A\mathbf{x}_k \\ &= (\lambda_k - \lambda_1) \alpha_1 \mathbf{x}_1 + \dots + (\lambda_k - \lambda_{k-1}) \alpha_{k-1} \mathbf{x}_{k-1}. \end{aligned}$$

Since $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI,

$$(\lambda_k - \lambda_1) \alpha_1 = \dots = (\lambda_k - \lambda_{k-1}) \alpha_{k-1} = 0,$$

i.e., $\alpha_1 = \dots = \alpha_{k-1} = 0$ (as λ_i are distinct),

Linearly independent eigenvectors

Theorem: Eigenvectors corresponding to **distinct** eigenvalues are **linearly independent**.

PROOF. Suppose that $\lambda_1, \dots, \lambda_m$ are distinct eigenvalues of A , \mathbf{x}_i is an eigenvector of A corresponding to λ_i . Suppose, if possible, \mathbf{x}_i are not LI. Then there is $k > 1$ such that $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI, but $\{\mathbf{x}_1, \dots, \mathbf{x}_k\}$ is not. Let $\mathbf{x}_k = \alpha_1 \mathbf{x}_1 + \dots + \alpha_{k-1} \mathbf{x}_{k-1}$. Then,

$$A\mathbf{x}_k = \lambda_1 \alpha_1 \mathbf{x}_1 + \dots + \lambda_{k-1} \alpha_{k-1} \mathbf{x}_{k-1}.$$

$$\begin{aligned} \text{Therefore, } \mathbf{0} &= \lambda_k \mathbf{x}_k - A\mathbf{x}_k \\ &= (\lambda_k - \lambda_1) \alpha_1 \mathbf{x}_1 + \dots + (\lambda_k - \lambda_{k-1}) \alpha_{k-1} \mathbf{x}_{k-1}. \end{aligned}$$

Since $\{\mathbf{x}_1, \dots, \mathbf{x}_{k-1}\}$ is LI,

$$(\lambda_k - \lambda_1) \alpha_1 = \dots = (\lambda_k - \lambda_{k-1}) \alpha_{k-1} = 0,$$

i.e., $\alpha_1 = \dots = \alpha_{k-1} = 0$ (as λ_i are distinct), i.e. $\mathbf{x}_k = \mathbf{0}$, a contradiction.

CAYLEY-HAMILTON THEOREM

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A .

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem.]

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

Exercise

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$.

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

Exercise

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$.

- Verify that char. poly. of A is $\lambda^2 - 2\lambda - 3$.

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

Exercise

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$.

- Verify that char. poly. of A is $\lambda^2 - 2\lambda - 3$.
- Verify that $A^2 - 2A - 3I = \mathbf{0}$.

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

Exercise

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$.

- Verify that char. poly. of A is $\lambda^2 - 2\lambda - 3$.
- Verify that $A^2 - 2A - 3I = \mathbf{0}$.
- Use the fact $A^2 = 2A + 3I$ to compute A^5 without computing any matrix multiplication.

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

Exercise

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$.

- Verify that char. poly. of A is $\lambda^2 - 2\lambda - 3$.
- Verify that $A^2 - 2A - 3I = \mathbf{0}$.
- Use the fact $A^2 = 2A + 3I$ to compute A^5 without computing any matrix multiplication.
- Argue that A is invertible using its char. poly.

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

Exercise

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$.

- Verify that char. poly. of A is $\lambda^2 - 2\lambda - 3$.
- Verify that $A^2 - 2A - 3I = \mathbf{0}$.
- Use the fact $A^2 = 2A + 3I$ to compute A^5 without computing any matrix multiplication.
- Argue that A is invertible using its char. poly. Since $3I = A^2 - 2A$, we should have $A^{-1} = \frac{1}{3}[A - 2I]$.

CAYLEY-HAMILTON THEOREM

Let $p(\lambda)$ be the characteristic polynomial of a matrix A . Then $p(A) = \mathbf{0}$, the zero matrix.

[This is a beautiful and useful theorem. However, we omit the proof.]

Exercise

Let $A = \begin{bmatrix} 1 & 2 \\ 2 & 1 \end{bmatrix}$.

- Verify that char. poly. of A is $\lambda^2 - 2\lambda - 3$.
- Verify that $A^2 - 2A - 3I = \mathbf{0}$.
- Use the fact $A^2 = 2A + 3I$ to compute A^5 without computing any matrix multiplication.
- Argue that A is invertible using its char. poly. Since $3I = A^2 - 2A$, we should have $A^{-1} = \frac{1}{3}[A - 2I]$. Verify.

Exercise

- Let A be a matrix with eigenvalue λ and corresponding eigenvector \mathbf{x} .
 - For any positive integer n , show that λ^n is an eigenvalue of A^n with corresponding eigenvector \mathbf{x} .
 - If A is invertible, then show that $\frac{1}{\lambda}$ is an eigenvalue of A^{-1} with corresponding eigenvector \mathbf{x} .
 - If A is invertible then show that for any integer n , λ^{-n} is an eigenvalue of A^{-n} with corresponding eigenvector \mathbf{x} .
- Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$ be eigenvectors of a matrix A with corresponding eigenvalues $\lambda_1, \lambda_2, \dots, \lambda_m$, respectively. Let $\mathbf{x} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_m\mathbf{v}_m$. Show that for any positive integer k ,

$$A^k \mathbf{x} = c_1 \lambda_1^k \mathbf{v}_1 + c_2 \lambda_2^k \mathbf{v}_2 + \dots + c_m \lambda_m^k \mathbf{v}_m.$$

Google Search Engine

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

Google Search Engine

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

Google search engine does three basic things:

- **Crawl the web** and locate all web pages with **public access**.

Google Search Engine

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

Google search engine does three basic things:

- **Crawl the web** and locate all web pages with **public access**.
- **Index the data** (create term-document matrix) from step-1 so that they can be searched efficiently for relevant **key words or phrases**.

Google Search Engine

The task of **extracting information from all the web pages** available on the Internet is performed by **search engines**.

Google search engine does three basic things:

- **Crawl the web** and locate all web pages with **public access**.
- **Index the data** (create term-document matrix) from step-1 so that they can be searched efficiently for relevant **key words or phrases**.
- **Rank each page in the database**, so that when a user does a search and the subset of pages in the database with the desired information has been found, the **more important pages** can be presented first.

PageRank Algorithm

Google PageRank algorithm assigns ranks to all the web pages and is formulated as a matrix eigenvalue problem.

PageRank Algorithm

Google PageRank algorithm assigns ranks to all the web pages and is formulated as a matrix eigenvalue problem.

The web is an example of a directed graph. Let all the web pages be ordered as P_1, \dots, P_n . Link from P_i to P_j represents an arrow. Google assigns rank to a page based on its in-links and out-links.

PageRank Algorithm

Google PageRank algorithm assigns ranks to all the web pages and is formulated as a matrix eigenvalue problem.

The web is an example of a directed graph. Let all the web pages be ordered as P_1, \dots, P_n . Link from P_i to P_j represents an arrow. Google assigns rank to a page based on its in-links and out-links.

In-links of P_j : All the incoming links (pages) to P_j .

Out-links of P_j : All the outgoing links (pages) from P_j .

PageRank Algorithm

Google PageRank algorithm assigns ranks to all the web pages and is formulated as a matrix eigenvalue problem.

The web is an example of a directed graph. Let all the web pages be ordered as P_1, \dots, P_n . Link from P_i to P_j represents an arrow. Google assigns rank to a page based on its in-links and out-links.

In-links of P_j : All the incoming links (pages) to P_j .

Out-links of P_j : All the outgoing links (pages) from P_j .

Google assigns a high rank to a web page if it has backlinks from other pages that have a high rank. This self-referencing statement can be formulated mathematically as an eigenvalue problem.

PageRank Algorithm

Google PageRank algorithm assigns ranks to all the web pages and is formulated as a matrix eigenvalue problem.

The web is an example of a directed graph. Let all the web pages be ordered as P_1, \dots, P_n . Link from P_i to P_j represents an arrow. Google assigns rank to a page based on its in-links and out-links.

In-links of P_j : All the incoming links (pages) to P_j .

Out-links of P_j : All the outgoing links (pages) from P_j .

Google assigns a high rank to a web page if it has backlinks from other pages that have a high rank. This self-referencing statement can be formulated mathematically as an eigenvalue problem.

However, a ranking system based only on the number of backlinks is easy to manipulate.

PageRank Algorithm

Google PageRank algorithm assigns ranks to all the web pages and is formulated as a matrix eigenvalue problem.

The web is an example of a directed graph. Let all the web pages be ordered as P_1, \dots, P_n . Link from P_i to P_j represents an arrow. Google assigns rank to a page based on its in-links and out-links.

In-links of P_j : All the incoming links (pages) to P_j .

Out-links of P_j : All the outgoing links (pages) from P_j .

Google assigns a high rank to a web page if it has backlinks from other pages that have a high rank. This self-referencing statement can be formulated mathematically as an eigenvalue problem.

However, a ranking system based only on the number of backlinks is easy to manipulate.

PageRank Algorithm (eigenvalue problem)

Let $x_j \geq 0$ be the rank of page P_j . Then $x_j > x_k \implies P_j$ is **more important** than page P_k .

PageRank Algorithm (eigenvalue problem)

Let $x_j \geq 0$ be the rank of page P_j . Then $x_j > x_k \implies P_j$ is **more important** than page P_k .

If page P_j contains n_j **out-links**, one of which links to page P_k , then **page P_k 's score is boosted by x_j/n_j** .

PageRank Algorithm (eigenvalue problem)

Let $x_j \geq 0$ be the rank of page P_j . Then $x_j > x_k \implies P_j$ is **more important** than page P_k .

If page P_j contains n_j **out-links**, one of which links to page P_k , then **page P_k 's score is boosted by x_j/n_j** .

Let L_k denote the **set of in-links P_k** . Then the rank x_k of P_k is defined by

$$x_k = \sum_{j \in L_k} \frac{x_j}{n_j} \quad (1)$$

where n_j is the **number of outgoing links (outlinks)** from page P_j .

Setting $\mathbf{v} := [x_1, \dots, x_n]^T$, equation (1) is rewritten as

PageRank Algorithm (eigenvalue problem)

Let $x_j \geq 0$ be the rank of page P_j . Then $x_j > x_k \implies P_j$ is **more important** than page P_k .

If page P_j contains n_j **out-links**, one of which links to page P_k , then **page P_k 's score is boosted by x_j/n_j** .

Let L_k denote the **set of in-links P_k** . Then the rank x_k of P_k is defined by

$$x_k = \sum_{j \in L_k} \frac{x_j}{n_j} \quad (1)$$

where n_j is the **number of outgoing links (outlinks)** from page P_j .

Setting $\mathbf{v} := [x_1, \dots, x_n]^T$, equation (1) is rewritten as

$$\text{Eigenvalue problem} \quad \mathbf{H}\mathbf{v} = \mathbf{v} ,$$

where \mathbf{H} is the **hyperlink matrix** and \mathbf{v} is the **page rank vector**.

The Google matrix

The hyperlink matrix is modified to obtain the Google matrix

$$\mathbf{G} := (1 - m)\mathbf{H} + m\mathbf{S},$$

The Google matrix

The hyperlink matrix is modified to obtain the Google matrix

$$\mathbf{G} := (1 - m)\mathbf{H} + m\mathbf{S},$$

where \mathbf{S} is an $n \times n$ matrix with all entries $1/n$ and $0 \leq m \leq 1$.

The Google matrix

The hyperlink matrix is modified to obtain the **Google matrix**

$$\mathbf{G} := (1 - m)\mathbf{H} + m\mathbf{S},$$

where \mathbf{S} is an $n \times n$ matrix with all entries $1/n$ and $0 \leq m \leq 1$. Both \mathbf{H} and \mathbf{G} have 1 as the **dominant eigenvalue**.

The Google matrix

The hyperlink matrix is modified to obtain the **Google matrix**

$$\mathbf{G} := (1 - m)\mathbf{H} + m\mathbf{S},$$

where \mathbf{S} is an $n \times n$ matrix with all entries $1/n$ and $0 \leq m \leq 1$. Both \mathbf{H} and \mathbf{G} have 1 as the **dominant eigenvalue**.

By Perron-Frobenius theorem, there is a unique vector $\mathbf{v} = [x_1, \dots, x_n]^\top$ with $x_j > 0$ for $j = 1 : n$ such that

$$\mathbf{G}\mathbf{v} = \mathbf{v} \text{ and } \sum_{j=1}^n x_j = 1.$$

The Google matrix

The hyperlink matrix is modified to obtain the **Google matrix**

$$\mathbf{G} := (1 - m)\mathbf{H} + m\mathbf{S},$$

where \mathbf{S} is an $n \times n$ matrix with all entries $1/n$ and $0 \leq m \leq 1$. Both \mathbf{H} and \mathbf{G} have 1 as the **dominant eigenvalue**.

By Perron-Frobenius theorem, there is a unique vector $\mathbf{v} = [x_1, \dots, x_n]^\top$ with $x_j > 0$ for $j = 1 : n$ such that

$$\mathbf{G}\mathbf{v} = \mathbf{v} \text{ and } \sum_{j=1}^n x_j = 1.$$

The **rank of the web page** P_j is given by the j -th component x_j of the **eigenvector** \mathbf{v} of the **Google matrix** \mathbf{G} . Google sets $m = 0.15$.

References

For more on Google search engine, read the following paper/book.

References

For more on Google search engine, read the following paper/book.

- K. Bryan and T. Leise, [The \\$25,000,000,000 Eigenvector: The Linear Algebra behind Google](#), SIAM Review, 48(2006), pp.569-581.

References

For more on Google search engine, read the following paper/book.

- K. Bryan and T. Leise, [The \\$25,000,000,000 Eigenvector: The Linear Algebra behind Google](#), SIAM Review, 48(2006), pp.569-581.
- Amy N. Langville and Carl D. Meyer, [Googles PageRank and Beyond: The Science of Search Engine Rankings](#), Princeton University Press, 2006
