

MA 102 (Mathematics II)
IIT Guwahati

Tutorial Sheet No. 6

Linear Algebra

March 14, 2019

1. (a) If A is an $n \times n$ matrix with $\text{nullity}(A) = k$, then show that there exists an invertible matrix P such that $P^{-1}AP = \begin{bmatrix} \mathbf{0} & B \\ \mathbf{0} & D \end{bmatrix}$, where D is an $(n - k) \times (n - k)$ matrix.

(b) Hence show that for any eigenvalue λ of A , the algebraic multiplicity of $\lambda \geq$ the geometric multiplicity of λ .

(c) Deduce that $\text{rank}(A) \geq$ the number of nonzero eigenvalues of A .
2. Find the eigenvalues of the $n \times n$ matrix A which has all diagonal entries equal to 3 and all other entries equal to 2. Find two eigenvectors \mathbf{x}, \mathbf{y} corresponding to two distinct eigenvalues of A .
Hint: $A = B + I$, where B is the matrix having all entries equal to 2. Check that \mathbf{x} is an eigenvector of B corresponding to an eigenvalue λ if and only if \mathbf{x} is an eigenvector of A corresponding to eigenvalue $\lambda + 1$.
3. If $A = \mathbf{u}\mathbf{u}^T$ where $\mathbf{0} \neq \mathbf{u} \in \mathbb{R}^n$, then find the eigenvalues of A and show that A is diagonalizable.
Hint: Check that $\text{rank}(A) = 1$ and $\mathbf{u}^T \mathbf{u}$ is the nonzero eigenvalue of A .
4. True or False? Give justifications.
 - (a) If both A and A^{-1} has only integer entries, then $\det(A) = +1$ or -1 .
 - (b) If A and B are non square matrices such that both AB and BA are defined then either AB or BA has a zero eigenvalue.
 - (c) If A is a 3×3 matrix with eigenvalues 0, 3, 4 and D is a 2×2 matrix with eigenvalues 0, 3 then the matrix $C = \begin{bmatrix} A & B \\ \mathbf{0} & D \end{bmatrix}$ has eigenvalues 0, 3, 4 with algebraic multiplicities 2, 2, 1, respectively.
Hint: $\det(C - \lambda I) = \det(A - \lambda I)\det(D - \lambda I)$.
 - (d) An upper triangular matrix with all diagonal entries equal to a is diagonalizable only if A is a diagonal matrix.
Hint: Look at $\text{null}(A - aI)$.
 - (e) Eigenvalues of real matrices occur in conjugate pairs (that is if $a + ib$ is an eigenvalue of A then $a - ib$ is also an eigenvalue of A).
Hint: $\det(A - cI) = \det(\overline{A - cI})$.
5. Let $A \in \mathcal{M}_n(\mathbb{C})$. If $\langle A\mathbf{x}, \mathbf{x} \rangle$ is real for all $\mathbf{x} \in \mathbb{C}^n$ then show that A is Hermitian.
6. Use Gram Schmidt procedure to transform the basis $\mathcal{B} = \left\{ \begin{bmatrix} 2 \\ 3 \\ -1 \end{bmatrix}, \begin{bmatrix} 3 \\ 1 \\ -0 \end{bmatrix}, \begin{bmatrix} 4 \\ -1 \\ 2 \end{bmatrix} \right\}$ to an orthogonal basis. What happens if the order in which the vectors are taken changes, does the elements of the orthogonal basis remain the same?
7. Let \mathbb{V} be a finite dimensional inner product space and $B := \{\mathbf{u}_1, \dots, \mathbf{u}_m\}$ be an orthonormal subset of \mathbb{V} .

- (a) Let $\mathbf{x}, \mathbf{y} \in \mathbb{V}$. Show that $\mathbf{x} = \mathbf{y} \Leftrightarrow \langle \mathbf{x}, \mathbf{w} \rangle = \langle \mathbf{y}, \mathbf{w} \rangle$ for all $\mathbf{w} \in \mathbb{V}$.
- (b) Show that $|\langle \mathbf{v}, \mathbf{u}_1 \rangle|^2 + \cdots + |\langle \mathbf{v}, \mathbf{u}_m \rangle|^2 \leq \|\mathbf{v}\|^2$ for all $\mathbf{v} \in \mathbb{V}$.
- (c) Show that $|\langle \mathbf{v}, \mathbf{u}_1 \rangle|^2 + \cdots + |\langle \mathbf{v}, \mathbf{u}_m \rangle|^2 = \|\mathbf{v}\|^2$ for all $\mathbf{v} \in \mathbb{V} \Leftrightarrow B$ is an orthonormal basis of \mathbb{V} .
- (d) Suppose that B is an orthonormal basis of \mathbb{V} . For any $\mathbf{x}, \mathbf{y} \in \mathbb{V}$ prove that

$$\langle \mathbf{x}, \mathbf{y} \rangle = \langle \mathbf{x}, \mathbf{u}_1 \rangle \overline{\langle \mathbf{y}, \mathbf{u}_1 \rangle} + \cdots + \langle \mathbf{x}, \mathbf{u}_m \rangle \overline{\langle \mathbf{y}, \mathbf{u}_m \rangle}.$$

- 8. Let $\mathcal{M}_2(\mathbb{R})$ be the inner product space of all real 2×2 matrices with respect to the inner product $\langle A, B \rangle = \text{trace}(B^T A)$ for all $A, B \in \mathcal{M}_2(\mathbb{R})$. Find the orthogonal complement of the set $\mathcal{S} = \left\{ \begin{bmatrix} 1 & 0 \\ 2 & 0 \end{bmatrix}, \begin{bmatrix} 1 & 0 \\ 1 & 0 \end{bmatrix} \right\}$ in $\mathcal{M}_2(\mathbb{R})$.
- 9. Let \mathbb{V} be a finite dimensional complex inner product space. Fix a vector $\mathbf{v} \in \mathbb{V}$ and define $T : \mathbb{V} \rightarrow \mathbb{C}$ by $T(\mathbf{u}) = \langle \mathbf{u}, \mathbf{v} \rangle$ for all $\mathbf{u} \in \mathbb{V}$. Determine T^* , that is, determine $T^* \mathbf{w}$ for all $\mathbf{w} \in \mathbb{C}$.
- 10. Let \mathbb{V} and \mathbb{W} be finite dimensional inner product spaces. Let $T : \mathbb{V} \rightarrow \mathbb{W}$ be a linear transformation. Prove that:
 - (i) T is injective if and only if T^* is surjective.
 - (ii) T is surjective if and only if T^* is injective.
 - (iii) $\dim(\text{null}(T^*)) = \dim(\text{null}(T)) + \dim(\mathbb{W}) - \dim(V)$.
 - (iv) $\dim(\text{range}(T^*)) = \dim(\text{range}(T))$.
