

# List of Tables

5.1	Run range and luminosity of the proton-proton collisions data collected at the center-of-mass energy of 8 TeV by the CMS experiment in the year of 2012 in different run periods. . . . .	75
5.2	The official Monte Carlo samples produced in phase space slices in $H_T$ with the generator MADGRAPH5 and interfaced to PYTHIA6. . .	76
5.3	The single jet HLT trigger paths used in the analysis. . . . .	77
5.4	The jet identification criteria (ID) removes noise and non-physical jets based on the properties of the reconstructed jets and the clustered particle candidates. . . . .	82
5.5	The jet energy resolution in the data is actually worse than in simulation. To match the resolution in the data, the reconstructed jet transverse momentum in simulated events need to be smeared by applying the scale factors. . . . .	90
5.6	The parameters obtained by fitting the relative resolution as a function of $H_{T,2}/2$ . . . . .	94
5.7	An overview of all experimental uncertainties affecting the measurement of cross-sections and the cross-section ratio. . . . .	112
6.1	NLO PDF sets available via LHAPDF6 with various assumptions on the value of $\alpha_s(M_Z)$ . . . . .	115

6.2	Overview of all systematic theoretical uncertainties affecting the measurement of cross-sections and the cross-section ratio. . . . .	124
7.1	Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross-sections using five PDF sets. . . . .	138
7.2	Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross-sections simultaneously and from their ratio $R_{32}$ using five PDF sets. . . . .	138
7.3	Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet event cross-section using five PDF sets at NLO without (left) and with (right) electroweak (EW) corrections. . . . .	139
7.4	Determination of $\alpha_s(M_Z)$ from the ratio $R_{32}$ using the two most compatible PDF sets MSTW2008 and MMHT2014. . . . .	141
7.5	Determination of $\alpha_s(M_Z)$ from the ratio $R_{32}$ in the $H_{T,2}/2$ range from 300 up to 1680 GeV at the central scale and for the six scale factor combinations for the two PDF sets MSTW2008 and MMHT2014. . .	141
7.6	Uncertainty decomposition for $\alpha_s(M_Z)$ from the determination of $\alpha_s$ from the jet event rate $R_{32}$ in bins of $H_{T,2}/2$ . . . . .	141
7.7	Evolution of the strong coupling constant. . . . .	143
A.1	Differential cross-sections ( $\times 10^{-3}(\text{pb/GeV})$ ) and the cross-section ratio $R_{32}$ at detector level in each bin of $H_{T,2}/2$ , along with statistical uncertainty (in %). . . . .	162
A.2	Experimental uncertainties (in %) affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 2-jet events. . . . .	167
A.3	Experimental uncertainties (in %) affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 3-jet events. . . . .	168

---

A.4	Experimental uncertainties (in %) affecting the cross-section measurement in each bin of $H_{T,2}/2$ for cross-section ratio. . . . .	169
A.5	Theoretical uncertainties (in %), calculated using CT10-NLO PDF set, affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 2-jet events. . . . .	170
A.6	Theoretical uncertainties (in %), calculated using CT10-NLO PDF set, affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 3-jet events. . . . .	171
A.7	Theoretical uncertainties (in %), calculated using CT10-NLO PDF set, affecting the measurement of cross-section ratio in each bin of $H_{T,2}/2$ . . . . .	172