

**MEASUREMENT OF MULTIJET CROSS-SECTION RATIOS
IN PROTON-PROTON COLLISIONS WITH THE CMS
DETECTOR AT THE LHC**

A THESIS

**Submitted to the
FACULTY OF SCIENCE
PANJAB UNIVERSITY, CHANDIGARH
for the degree of**

DOCTOR OF PHILOSOPHY

2018

ANTERPREET KAUR

**DEPARTMENT OF PHYSICS
CENTRE OF ADVANCED STUDY IN PHYSICS
PANJAB UNIVERSITY
CHANDIGARH, INDIA**

*Dedicated to
my Parents*

Acknowledgement

I would like to take this opportunity to thank the many people who helped make this dissertation possible.

In the first and foremost place, I gratefully express my sincere gratitude to my supervisor Prof.(Mrs.) Manjit Kaur for her invaluable assistance, support, guidance and blessings. She inspired me greatly to work in this project. Her willingness to motivate me contributed tremendously to this project.

I am highly obliged to Prof. C.S. Aulakh Chairperson of the Department of Physics, Panjab University, Chandigarh, for his helping hand in providing a good environment and adequate facilities to complete this project successfully.

I feel deeply indebted to my parents for their unconditional love, understanding, support, strength, blessings and for everything.

I am greatly thankful to my well wishers, my friends Anu, Sohail, Junaid, Mandeep and my classmates for all that they meant to me during the crucial times of the completion of my project.

I owe my very special thanks to Mr. Inderpal Singh, Mrs. Prabhdeep Kaur, Mrs. Manuk Z. Mehta, Ms. Ritu Aggarwal, Ms. Ruchi Gupta, Ms. Monika Mittal and Ms. Genius, the members of high energy physics Laboratory, Dept. of Physics, P.U. for helping me out from all kinds of problems with their constant support, suggestions and encouragement.

Finally and the most importantly, I would like to pay my greatest regards to the Almighty God for His blessings and kindness, a boundless and an endless source of inspiration for me always.

Abstract

The hadrons colliding at very high centre-of-mass energies provide a direct probe to the nature of the underlying parton-parton scattering physics. The scattering of the elementary quarks and gluons, constituents of the incoming hadron beams, produces a high momentum partons which then fragment and hadronize producing a spray of particles. These particles get clustered in the form of jets. The jets being the final structures observed in the detector, preserve the energy and direction of the initial partons. Hence jets can serve as a direct test of theory of strong interactions called Quantum Chromodynamics. The inclusive multijet production cross-section is an important observable which provides the details of parton distribution functions of the colliding hadrons and the precise measurement of the strong coupling constant α_S . Instead of individual cross-sections, the ratio of cross-sections is a better tool to determine the value of α_S as many theoretical and experimental uncertainties cancel in the ratio.

A measurement of inclusive multijet event cross-sections and the cross-section ratio is presented using data from proton-proton collisions collected with the CMS detector at a centre-of-mass energy of 8 TeV corresponding to an integrated luminosity of 19.7 fb^{-1} . Jets are reconstructed with the anti- k_t clustering algorithm for a jet size parameter $R = 0.7$. The inclusive 2-jet and 3-jet event cross-sections as well as the ratio of the 3-jet over 2-jet event cross-section (R_{32}) are measured as a function of the average transverse momenta p_T of the two leading jets in a phase space region ranging up to jet p_T of 2.0 TeV and an absolute rapidity of $|y| = 2.5$. The measurements after correcting for detector effects are well described by predictions at next-to-leading order in perturbative quantum chromodynamics and additionally are compared to several Monte Carlo event generators. The strong coupling constant at the scale of the Z boson mass is extracted from a fit of the measured R_{32} which gives $\alpha_s(M_Z) = 0.1150 \pm 0.0010 \text{ (exp)} \pm 0.0013 \text{ (PDF)} \pm 0.0015 \text{ (NP)} {}^{+0.0050}_{-0.0000} \text{ (scale)}$ using MSTW2008 PDF set. The current measurement agrees well with the world

average value of $\alpha_s(M_Z) = 0.1181 \pm 0.0011$ as well as previous measurements.

Contents

List of Figures	xv
List of Tables	xxi
<hr/>	
1 Introduction	1
2 Theoretical Background	5
2.1 Standard Model	5
2.2 Quantum Chromodynamics	8
2.2.1 Perturbative Quantum Chromodynamics	12
2.2.2 Renormalization and Running of the Strong Coupling	14
2.3 Hadronic Collisions	16
2.3.1 Parton Shower and Hadronization	19
2.3.2 Underlying Event	21
2.4 Jets	23
2.4.1 Jet Algorithms	24
3 Experimental Setup	29
3.1 The Large Hadron Collider	30
3.1.1 Luminosity Measurement	32

3.1.2	Pileup Interactions	34
3.2	The Compact Muon Solenoid	35
3.2.1	Coordinate System	37
3.2.2	Inner Tracker System	39
3.2.3	Electromagnetic Calorimeter	42
3.2.4	Hadronic Calorimeter	43
3.2.5	Superconducting Magnet	47
3.2.6	Muon System	47
3.2.7	Trigger and Data Acquisition System	49
3.2.7.1	Jet Triggers	51
3.2.7.2	Data Acquisition System	52
3.2.8	Data Management	52
3.3	Software Tools	53
3.3.1	CMSSW Framework	54
3.3.2	ROOT	54
3.3.3	NLOJET++ and FASTNLO	55
4	Event Generation, Simulation and Reconstruction	57
4.1	Event Generation and Simulation Software	57
4.1.1	PYTHIA	59
4.1.2	MADGRAPH5	59
4.1.3	HERWIG	60
4.1.4	POWHEG	60
4.2	Detector Simulation	60
4.3	Event Reconstruction	62
4.3.1	Particle Flow Algorithm	64

4.3.2	Jet Energy Corrections	66
5	Measurement of the Differential Inclusive Multijet Cross-sections and their Ratio	69
5.1	Data Samples	70
5.1.1	Monte Carlo Samples	71
5.2	Event Selection	72
5.2.1	Trigger Selection	72
5.2.2	Primary Vertex Selection	74
5.2.3	Missing Transverse Energy	76
5.2.4	Jet Identification	77
5.2.4.1	Jet ID Efficiency	78
5.2.5	Jet Selection	81
5.3	Comparison with Simulation	82
5.3.1	Pileup Reweighting	82
5.3.2	Comparison of Cross-sections and their Ratio	82
5.4	Jet Energy Resolution (JER)	85
5.5	Unfolding	90
5.5.1	Response Matrices	92
5.5.1.1	Inclusive Cross-sections	92
5.5.1.2	Cross-section Ratio, R_{32}	95
5.5.2	Closure Test	96
5.5.3	Unfolding of the Measurement	98
5.6	Experimental Uncertainties	99
5.6.1	Statistical Uncertainty	100
5.6.2	Jet Energy Corrections Uncertainty	101

5.6.3	Unfolding Uncertainty	104
5.6.4	Luminosity Measurement Uncertainty	105
5.6.5	Residual Uncertainty	106
5.6.6	Total Experimental Uncertainty	106
6	Theoretical Calculations	109
6.1	Fixed Order NLO Calculations	109
6.1.1	NLO Correction Factors	111
6.1.2	Non-perturbative Corrections	112
6.1.3	Electroweak Corrections	114
6.2	Theoretical Uncertainties	116
6.2.1	Scale Uncertainty	116
6.2.2	PDF Uncertainty	117
6.2.3	Non-perturbative Uncertainty	118
6.2.4	Total Theoretical Uncertainty	118
6.3	Comparison of Theory to Data	120
7	Determination of the Strong Coupling Constant	125
7.1	Sensitivity of Measurements to $\alpha_s(M_Z)$	126
7.2	Determination of $\alpha_s(M_Z)$	127
7.2.1	Fitting Procedure	127
7.2.2	Fit Results	133
7.3	Running of the Strong Coupling Constant	136
8	Summary	141
Appendix A		145

A.1 Cross-section Ratio, R_{32}	146
A.2 Individual Sources of Jet Energy Correction Uncertainties	147
A.3 Experimental Uncertainties	150
A.4 Theoretical Uncertainties	153
Bibliography	157
List of Publications	173
Reprints	200

List of Figures

2.1	The Standard Model summarizing the properties of elementary particles and their forces of interaction.	6
2.2	The fundamental Feynman rules of QCD.	11
2.3	The Feynman diagrams of leading-order (LO), next-to-leading order (NLO) and next-to-next-to-leading order (NNLO) processes in quantum chromodynamics.	13
2.4	Running of the strong coupling constant.	16
2.5	Schematic illustration of factorization theorem in a collision of two protons.	18
2.6	Illustration of the hadronization process in Lund string model. . . .	21
2.7	Proton-proton collision.	22
2.8	Formation of a jet in a proton-proton collision.	24
2.9	Effects of emission of infrared radiations and collinear splitting in jet algorithms.	25
2.10	The clustering of particles into jets using different jet algorithms. . .	28
3.1	Overview of the different experiments of the Large Hadron Collider (LHC), a complex particle accelerator and collider located at CERN. .	31
3.2	The integrated luminosity, delivered to CMS during stable beams for proton-proton collisions.	34

3.3	Pileup Interactions	35
3.4	The three dimensional view of the CMS detector along with its sub-detector components.	36
3.5	Front view of the CMS detector along with its components.	37
3.6	The right-handed coordinate system used by the CMS detector.	38
3.7	Longitudinal section of the CMS detector in the y - z plane.	40
3.8	The longitudinal section of the inner tracking system.	41
3.9	A geometric view of one quarter of the electromagnetic calorimeter (ECAL) in y - z plane.	43
3.10	Longitudinal view of one quarter of the hadronic calorimeter (HCAL) detector in r - η plane.	45
3.11	A longitudinal view presenting the location of the three muon sub-systems.	48
3.12	Work flow of the L1 trigger system	50
3.13	Architecture of the CMS DAQ system.	52
3.14	Schematic overview of the CMS computing grid.	53
4.1	The comparison of Monte Carlo (MC) simulations generated by event generators with the real data produced by the collisions of observed in the detectors.	58
4.2	Formation of jets in a proton-proton collision at different levels.	63
4.3	Schematic association of sub-detector measurements to physical particle objects using the Particle Flow (PF) event reconstruction technique used by CMS.	65
4.4	A schematic diagram of the factorized jet energy corrections (JEC).	66
5.1	Trigger efficiencies turn-on curves for the single jet HLT trigger paths. .	75

5.2	Missing transverse energy distribution.	77
5.3	The fractions of jet constituents for different types of PF candidates for inclusive 2-jet events.	79
5.4	The fractions of jet constituents for different types of PF candidates for inclusive 3-jet events.	80
5.5	The jet ID efficiency is studied as a function of $H_{T,2}/2$	81
5.6	Number of reconstructed vertices before (left) and after (right) the pileup reweighting.	83
5.7	Comparison of differential cross-sections for data with simulated events and CT10-NLO theory predictions.	84
5.8	Comparison of the cross-section ratio for data with simulated events and CT10-NLO theory predictions.	84
5.9	Fitting of the jet energy resolution distribution as a function of $H_{T,2}/2$. .	87
5.10	Comparison of jet energy resolution calculated using Crystal-Ball fit function and Gaussian fit function	88
5.11	Jet energy resolution (JER) is shown as a function of Gen $H_{T,2}/2$. . .	88
5.12	Additional unfolding uncertainty.	90
5.13	Fitted CT10-NLO spectrum of differential cross-section as a function of $H_{T,2}/2$	94
5.14	The response matrices are derived using the Toy Monte Carlo and forward smearing method.	95
5.15	Left : The ratio of cross-sections for inclusive 3-jet to that of 2-jet events as a function of $H_{T,2}/2$. Right : The response matrix is derived using the Toy Monte Carlo and forward smearing method, for the cross-section ratio R_{32}	96
5.16	Closure test of the unfolding technique	97

5.17 Reco differential cross-section distributions unfolded with the response matrices.	98
5.18 The measured differential cross-sections as well as the cross-section ratio R_{32} are unfolded as a function of $H_{T,2}/2$ using the response matrices derived using the Toy Monte Carlo and forward smearing method.	99
5.19 The fractional statistical uncertainties of the unfolded data are compared with those of the measured one.	101
5.20 The unfolding procedure introduces the correlations of the statistical uncertainty through bin migrations.	102
5.21 Experimental uncertainties from different sources affecting the measurement of cross-sections and cross-section ratio.	107
6.1 The k-factors using five different PDF sets.	112
6.2 The nonperturbative (NP) corrections are presented as a function of $H_{T,2}/2$	114
6.3 The electroweak (EW) corrections as a function of $H_{T,2}/2$	115
6.4 Ratio of data over theory obtained using the CT10-NLO PDF set.	116
6.5 The systematic theoretical uncertainties affecting the cross-section measurement and the cross-section ratio.	119
6.6 Comparison of the measured differential inclusive 2-jet and 3-jet event cross-sections as a function of $H_{T,2}/2$ to theoretical predictions.	121
6.7 Cross-section ratio as a function of $H_{T,2}/2$ calculated from data in comparison to that from NLO pQCD predictions obtained using the CT10-NLO PDF set.	121
6.8 Ratio of data over theory using the CT10-NLO PDF set.	122
6.9 Ratio of data over the predictions from Monte Carlo simulations.	123

7.1	Ratio of the measured inclusive 2-jet differential cross-section to theory predictions using different PDF sets.	128
7.2	Ratio of the measured inclusive 3-jet differential cross-section to theory predictions using different PDF sets.	129
7.3	Ratio of the measured cross-section ratio to theory predictions using different PDF sets.	130
7.4	The running $\alpha_S(Q)$ as a function of the scale Q	139
A.1	The fractional jet energy correction (JEC) uncertainties from individual sources (Part I).	147
A.2	The fractional jet energy correction (JEC) uncertainties from individual sources (Part II).	148
A.3	The fractional jet energy correction (JEC) uncertainties from individual sources (Part III).	149

List of Tables

5.1	Data sets of the 2012 LHC run period.	71
5.2	The official Monte Carlo samples produced in phase space slices in H_T with the generator MADGRAPH5 and interfaced to PYTHIA6.	72
5.3	The single jet HLT trigger paths used in the analysis.	73
5.4	The jet ID criteria.	78
5.5	Jet energy resolution scale factors.	86
5.6	The parameters obtained by fitting the relative resolution as a function of $H_{T,2}/2$	89
5.7	Overview of all experimental uncertainties affecting the measurement of cross-sections and the cross-section ratio.	108
6.1	NLO PDF sets available via LHAPDF6 with various assumptions on the value of $\alpha_s(M_Z)$	111
6.2	Overview of all systematic theoretical uncertainties affecting the measurement of cross-sections and the cross-section ratio.	120
7.1	Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross-sections using five PDF sets.	134
7.2	Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross-sections simultaneously and from their ratio R_{32} using five PDF sets.	134

7.3	Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet event cross-section using five PDF sets at NLO without (left) and with (right) electroweak (EW) corrections.	135
7.4	Determination of $\alpha_s(M_Z)$ from the ratio R_{32} using the two most compatible PDF sets MSTW2008 and MMHT2014.	137
7.5	Determination of $\alpha_s(M_Z)$ from the ratio R_{32} in the $H_{T,2}/2$ range from 0.3 up to 1.68 TeV at the central scale and for the six scale factor combinations for the two PDF sets MSTW2008 and MMHT2014. . .	137
7.6	Uncertainty decomposition for $\alpha_s(M_Z)$ from the determination of α_S from the jet event rate R_{32} in bins of $H_{T,2}/2$	137
7.7	Evolution of the strong coupling constant.	139
A.1	Differential cross-sections ($\times 10^{-3}(\text{pb}/\text{GeV})$) and cross-section ratio R_{32} at detector level in each bin of $H_{T,2}/2$, along with statistical uncertainty (in %).	146
A.2	Experimental uncertainties (in %) affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 2-jet events.	150
A.3	Experimental uncertainties (in %) affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 3-jet events.	151
A.4	Experimental uncertainties (in %) affecting the cross-section measurement in each bin of $H_{T,2}/2$ for cross-section ratio.	152
A.5	Theoretical uncertainties (in %), calculated using CT10-NLO PDF set, affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 2-jet events.	153

A.6 Theoretical uncertainties (in %), calculated using CT10-NLO PDF set, affecting the cross-section measurement in each bin of $H_{\mathrm{T},2}/2$ for inclusive 3-jet events	154
A.7 Theoretical uncertainties (in %), calculated using CT10-NLO PDF set, affecting the measurement of cross-section ratio in each bin of $H_{\mathrm{T},2}/2$	155

Chapter 1

Introduction

Particle physics deals with the study of the basic constituents of matter and the forces governing the interactions among them. The Standard Model (SM) is the most accepted theory which describes the nature and properties of the fundamental particles and their interactions. The elementary particles leptons and quarks, known as fermions, interact through the exchange of gauge bosons and acquire mass through a scalar boson called the Higgs. The four fundamental forces of interaction existing in nature are : the electromagnetic force, the strong force, the weak force and the gravitational force. Quantum Chromodynamics (QCD) is the theory of the strong interactions between the quarks mediated by the massless gluons. The partons (quarks and gluons) have a peculiar property of “color” charge. The quarks strongly binds into colorless particles called hadrons such as protons and neutrons together known as nucleons, pions etc. The structure and the properties of sub-atomic particles can be explored by first accelerating them using particle accelerators and then colliding at very high energies. The end products of these collisions get detected in the particle detectors constituting the real data. The data sets analyzed in details to reveal the structure and characteristic properties of the fundamental particles.

To investigate the very rare particles or to search for physics beyond SM, the

particle accelerators have become ever bigger and more complex. The Large Hadron Collider (LHC) is one of the today's biggest and most powerful collider where the protons are accelerated and collided at extremely high center-of-mass energies to probe their internal structure described by parton distribution functions (PDFs). The PDF sets give the probability for finding a parton at an energy scale Q with a fractional momentum x of the proton. Since the proton is not elementary and is made up of partons, the proton-proton (pp) collisions are viewed as interactions between their constituent partons. The final products of the scattering are observed by Compact Muon Solenoid (CMS), one of the detectors located around the interaction points. The scattering cross-section can be defined as a sum of terms with increasing powers of the strong coupling constant α_S convoluted with PDFs. The lowest-order α_S^2 term represents the production of two partons in final states whereas terms of higher-order α_S^3 , α_S^4 etc. signify the existence of multi partons in final states. The final state partons give a parton shower (PS) due to decrease in energy through emission of other quarks and gluons. The colored products of parton shower hadronize to a spray of colorless hadrons known as jets. The jets are the final structures observed in the detector. So they carry the significant information about the energy and direction of the initial partons and hence are important to study. The final partons have the probability to radiate more gluons resulting in multijets in the final state. Such events are produced in large number and are an important source for testing the predictions given by QCD. They also serve as an important background in the searches for new particles and physics beyond SM.

The inclusive multijet event cross-section σ_{i-jet} given by $pp \rightarrow i \text{ jets} + X$, where every jet counts, is proportional to α_s^i . The inclusive jet cross-section studied in terms of jet p_T and rapidity y is one of the important observables as it provides the essential information about the PDFs and the precise measurement of α_S . Also the ratio of cross-sections given by Eq. 1.1 is proportional to the QCD coupling α_S

and hence can be used to determine the value of α_S .

$$R_{mn} = \frac{\sigma_{m-jet}}{\sigma_{n-jet}} \propto \alpha_S^{m-n} \quad (1.1)$$

Instead of studying inclusive cross-sections, the cross-section ratio is more useful because of the partial or complete cancellation of many theoretical and experimental uncertainties between numerator and denominator. The CMS Collaboration has previously measured the ratio of the inclusive 3-jet cross-section to the inclusive 2-jet cross-section as a function of the average transverse momentum, $\langle p_{T1,2} \rangle$, of the two leading jets in the event at 7 TeV [1] and lead to an extraction of $\alpha_s(M_Z) = 0.1148 \pm 0.0055$, where the dominant uncertainty stems from the estimation of higher-order corrections to the NLO prediction. In this analysis, a measurement of inclusive 2-jet and 3-jet event cross-sections as well as ratio of 3-jet event cross-section over 2-jet R_{32} , is presented using an event sample collected by the CMS experiment during 2012 at the LHC and corresponding to an integrated luminosity of 19.7 fb^{-1} of pp collisions at a centre-of-mass energy of 8 TeV. The event scale is chosen as before to be the average transverse momentum of the two leading jets, but will be referred to as $H_{T,2}/2$ in this thesis. The measurements are used to determine the value of the strong coupling constant at the scale of the Z boson mass $\alpha_s(M_Z)$ and the running of α_S with energy scale Q is studied.

This thesis is organized as :

Chapter 2 gives a brief overview of the Standard Model of particle physics and the theory of strong interactions QCD with main emphasis on the jets and jet algorithms.

Chapter 3 deals with experimental apparatus which covers the details of the CMS detector and its various sub-detectors.

Chapter 4 describes the methods of event generation using different Monte-Carlo event generators, detector geometry simulation and reconstruction of the par-

ticles in the detector. This chapter also gives an overview of the different approaches of jet reconstruction at CMS and application of jet-energy corrections along with the description of the software framework used in the analysis presented in the current thesis.

Chapter 5 presents the measurement of inclusive differential multijet cross-sections and the cross-section ratio. The measurements are corrected for detector effects by unfolding procedure which is discussed in details in this chapter. The sources of the experimental uncertainties are studied in details.

Chapter 6 contains a detailed description of the NLO pQCD theory predictions compared to data and the extraction of α_s . The NLO calculations are corrected with the non-perturbative and electroweak corrections. The theoretical uncertainties are calculated from various sources. The unfolded measurements are compared to the predictions at NLO in pQCD and additionally as well as to predictions from several Monte Carlo event generators.

Chapter 7 describes the method to extract $\alpha_s(M_Z)$ from the current measurement and the running of α_s with energy scale Q is presented along with the previous measurements from different experiments.

Chapter 8 summarizes the results and conclusions of the work done in this thesis.

The common unit convention based on International System of Units (SI), followed in particle physics, will be used throughout the thesis. It is supplemented with the units electron volt (eV) for energy and barn (b) for the interaction cross section. The speed of light (c) and the reduced Planck constant (\hbar) are set to unity, i.e. $\hbar = c = 1$.

Chapter 2

Theoretical Background

Since 1930s, many theories and discoveries in particle physics have revealed the fundamental structure of matter. The matter is made up of fundamental particles and their interactions are mediated by four fundamental forces [2]. The theoretical models describe all the phenomena of particle physics as well as predict the properties of particles. These models must be either confirmed experimentally or totally excluded giving hints of new physics. This interplay between experimental discoveries and the corresponding theoretical predictions leads to a theoretical model called Standard Model, which describes the fundamental particles and their interactions. The world's most powerful particle accelerators and detectors are used by physicists to test the predictions and limits of the Standard Model where it has successfully explained almost all experimental results. This chapter describes the Standard Model with main focus on the theory of strong interactions called Quantum Chromodynamics and its features which serve as the theoretical base of this thesis.

2.1 Standard Model

The Standard Model (SM) of particle physics [3–5] was developed in 1970s. It is a mathematical framework which describes the nature and properties of the fun-

damental particles and the three of the four known forces of interactions between them, as summarized in Fig. 2.1. According to the SM, there are 12 elementary particles i.e. without any internal structure, known as fermions. The fermions have half integral spin and obey Fermi-Dirac statistics. They follow the Pauli exclusion principle according to which two or identical fermions cannot occupy the same quantum state. Each fermion has an associated anti-particle having the same properties but opposite-sign quantum numbers. Depending on how the fermions interact, these




Figure 2.1: The Standard Model¹ summarizing the properties of elementary particles known as fermions (leptons and quarks) grouped into three generations, gauge bosons as mediators for the interactions, the scalar Higgs boson and not incorporated graviton for the gravitational force.

are classified into two categories - leptons (ℓ) and quarks (q). The leptons are of

¹Source : <http://www.texample.net/tikz/examples/model-physics>

six types : electron (e), muon (μ) and tau (τ) with electric charge $Q = -1^2$ and the corresponding neutrinos : electron neutrino (ν_e), muon neutrino (ν_μ) and tau neutrino (ν_τ) having electric charge $Q = 0$. The quarks exist in six “flavors” : up (u), down (d), strange (s), charm (c), bottom (b) and top (t). u , c and t carry electric charge $Q = \pm \frac{2}{3}$ whereas d , s and b carry $Q = \pm \frac{1}{3}$. The quarks and leptons are categorized into three generations. The first generation has the lightest and the most stable particles whereas the heavier and less stable particles belong to the second and third generations.

The four fundamental forces exist in nature : electromagnetic, strong, weak and gravitational force. Every interaction involves the exchange of a gauge boson : the photon (γ) for the electromagnetic force, gluons (g) for the strong force, two W ’s and a Z for the weak force and the graviton (not yet found) for the gravitational force. However, the gravitational force has not been incorporated into SM yet. Along with this, the existence of dark matter or dark energy and the matter-antimatter asymmetry are still missing pieces in the SM. The interaction between fundamental particles acts because of some peculiar property of the particles - charge for the electromagnetic force, color for the strong force and flavor for the weak force.

In the SM, the first three forces are unified into one quantum field theory [6], known as Grand Unified Theory (GUT) [7–9]. The SM framework based on quantum field theories is described by $SU(3)_C \otimes SU(2)_L \otimes U(1)_Y$ gauge symmetry where C stands for the color charge, L for weak isospin and Y for hypercharge. Here $SU(3)_C$, $SU(2)_L$ and $U(1)_Y$ terms give rise to strong, weak and electromagnetic forces, respectively. $U(n)$ are the unitary and $SU(n)$ are the special unitary groups of degree n . The electromagnetic interaction of particles is explained by a well established modern theory called Quantum Electrodynamics (QED). In SM, the weak and electromagnetic interactions are combined by an electroweak theory. The

²all charges are expressed in units of elementary charge e

spontaneous symmetry is broken due to the coupling to the scalar Higgs field which results in the massive W^\pm and Z bosons and the massless photon (γ). The Higgs boson, named after Peter Higgs, is the field quantum of the Higgs field responsible for electroweak symmetry breaking. Its existence was confirmed by the CMS [10] and ATLAS [11] collaborations in 2012, with the properties consistent with the SM. The $SU(3)_C$ term defines the strong interaction between quarks and gluons mediated by gluons, with the three degrees of freedom of the color charge (C). In contrast to the electroweak symmetry, the $SU(3)_C$ of the strong interaction is an exact symmetry and hence the gluons are massless. The strong interaction between quarks and gluons is described by theory called Quantum Chromodynamics (QCD), explained in details in the next section of this thesis.

2.2 Quantum Chromodynamics

The strong interactions between the quarks and gluons are described by a non-abelian gauge theory called Quantum Chromodynamics (QCD) [12, 13]. The gauge group of QCD is the special unitary group $SU(3)_C$ with color charges C as the generators of the gauge group. Color charge is the peculiar property of QCD and has a same role as the electric charge in electromagnetic interactions. However, the mediator of electromagnetic interactions i.e. photon, itself does not carry any electric charge whereas the gluon itself carry color charge. This allows the self coupling of gluons and hence make the theory non-abelian. Both the quarks and gluons carry three types of color charges : red (r), green (g) and blue (b), and three types of anti-color charges : anti-red (\bar{r}), anti-green (\bar{g}) and anti-blue (\bar{b}). The quarks carry a single color charge whereas gluons carry a combination of color charges. There are nine eigen states of gluons but one of them $\frac{1}{\sqrt{3}}(r\bar{r} + g\bar{g} + b\bar{b})$ is totally symmetric color singlet which has no net color charge and does not take part in interaction.

The remaining eight eigen states of the gluons are :

$$r\bar{b}, \ r\bar{g}, \ g\bar{r}, \ g\bar{b}, \ b\bar{g}, \ b\bar{r}, \ \frac{1}{\sqrt{2}}(r\bar{r} - b\bar{b}), \ \frac{1}{\sqrt{6}}(r\bar{r} + b\bar{b} - 2g\bar{g}) \quad (2.1)$$

The dynamics of the quarks and gluons are controlled by the gauge invariant QCD Lagrangian \mathcal{L}_{QCD} which is composed of four terms as :

$$\mathcal{L}_{QCD} = \underbrace{-\frac{1}{4}F_{\mu\nu}^A F_A^{\mu\nu}}_{\mathcal{L}_{gluons}} + \underbrace{\sum_{flavors} \bar{q}_a (i\gamma^\mu (D_\mu)_{ab} - m_q) q_b}_{\mathcal{L}_{quarks}} + \mathcal{L}_{gauge} + \mathcal{L}_{ghost} \quad (2.2)$$

where \mathcal{L}_{gluons} describes the kinetic term of the gluon fields \mathcal{A}_μ^A ; \mathcal{L}_{quarks} defines the interaction between spin- $\frac{1}{2}$ quark fields q_a of mass m_q and spin-1 gluon fields \mathcal{A}_μ^A summing over all presently known six flavors of quarks; \mathcal{L}_{gauge} describes the chosen gauge and \mathcal{L}_{ghost} is the so-called ghost term required to treat the degeneracy of equivalent gauge field configurations in non-abelian gauge theories. In Eq. 2.2, the Greek letters $\mu, \nu, \dots \in \{0,1,2,3\}$ are the space-time indices; $a,b,c \in \{1,2,3\}$ and $A,B,C \in \{1,\dots,8\}$ are the indices of the triplet and octet representations, respectively, of the gauge symmetry group $SU(3)_C$. The field tensor $F_{\mu\nu}^A$ is defined as

$$F_{\mu\nu}^A = \partial_\mu \mathcal{A}_\nu^A - \partial_\nu \mathcal{A}_\mu^A - g_s f_{ABC} \mathcal{A}_\mu^B \mathcal{A}_\nu^C \quad (2.3)$$

where g_s is the coupling constant determining the strength of the interaction between colored partons and f_{ABC} are the structure constants of the $SU(3)_C$ group. The last term in Eq. 2.3 is a non-abelian term which distinguishes QCD from QED and gives rise to a three- and a four-gluon vertex. In the term \mathcal{L}_{quarks} , $(D_\mu)_{ab}$ is the covariant

derivative given by Eq. 2.4 and γ_μ are the Dirac γ -matrices.

$$(D_\mu)_{ab} = \partial_\mu \delta_{ab} + ig_s T_{ab}^A \mathcal{A}_\mu^A \quad (2.4)$$

\mathcal{A}_μ^A are the gluon fields with factors T_{ab}^A factors corresponding to the generators of the $SU(3)_C$ gauge group. The generators are represented via $T^A = \lambda^A/2$ by the Hermitian and traceless Gell-Mann matrices λ^A [14]. The generator matrices T^A follow the commutation relations :

$$\left[T^A, T^B \right] = if_{ABC} T^C \quad (2.5)$$

In \mathcal{L}_{QCD} , the classical contribution comes from \mathcal{L}_{gluons} and \mathcal{L}_{quarks} terms which give rise to the free quark- and gluon-field terms, and the quark-gluon interaction terms presented in Fig. 2.2. The cubic and quartic gluon self-interaction vertices proportional to g_s and g_s^2 , respectively, come into play due to the non-abelian property of QCD.

It is impossible to use perturbation theory on a gauge invariant Lagrangian without choosing a specific gauge in which to calculate. The usual gauge-fixing term is given by

$$\mathcal{L}_{gauge} = -\frac{1}{2\xi} (\partial^\mu \mathcal{A}_\mu^A)^2 \quad (2.6)$$

where ξ may be any finite constant. This choice fixes the class of covariant gauges with ξ as the gauge parameter. As QCD is non-abelian, the gauge fixing term must be supplemented by a ghost Lagrangian as

$$\mathcal{L}_{ghost} = \partial_\alpha \eta^{A\dagger} (D_{AB}^\mu \eta^B) \quad (2.7)$$

where η^A is a complex scalar field which obeys Fermi-Dirac statistics. The ghost




Figure 2.2: The fundamental Feynman rules of a free quark-field term (top left), quark-gluon interaction term (top right), free gluon-field term (bottom left), cubic gluon self-interaction term (bottom middle) and quartic gluon self-interaction term (bottom right). Taken from [15].

fields cancel unphysical degrees of freedom arising due to use of covariant gauges. This completes the QCD Lagrangian shown in Eq. 2.2.

The strength of an interaction is given by a fundamental parameter called the coupling constant α . In QED, the coupling constant $\alpha_e = e^2/4\pi = 1/137$ is constant. In contrast to this, in QCD, the coupling constant $\alpha_S(Q) = g_s^2/4\pi$ is not constant and depends on the separation between the interacting particles. It increases with the increase in the distance or decrease in the energy scale Q . At large distances or low energies, the quarks can never be found as free particles but exist in color neutral bound states known as hadrons. Hadrons are of two types : baryons and mesons. According to the quark model [2] every (anti-)baryon is made up of three (anti-)quarks and every meson is made up of a quark-antiquark pair. When the colored partons within a hadron are pulled farther and farther apart, there is an increase in the strength of force between them. This results in creation of new quark-antiquark pair making difficult to liberate a free quark or gluon. This property of QCD is

known as confinement according to which at low energy, the partons are forever bound into hadrons such as protons (uud), neutrons (udd). Although the gluons are massless but the confinement leads to the finite range of the strong interactions. On the other hand, at small distances, the strength of coupling decreases. The quarks and gluons interact very weakly and are treated as free particles. This property is known as asymptotic freedom. This indicates that perturbative theory is only applicable at high energies or small distances.

2.2.1 Perturbative Quantum Chromodynamics

At high energies, the property of asymptotic freedom allows a perturbative treatment in QCD calculations. In perturbative Quantum Chromodynamics (pQCD), any physical observable X such as cross-section of a scattering process, can be expanded as a perturbative series in terms of coupling constant α_s as :

$$X = \sum_{i=0}^N \alpha_s^n c_i = c_0 + \alpha_s^1 c_1 + \alpha_s^2 c_2 + \dots \quad (2.8)$$

where c_i are the perturbative coefficients. In a process, the pQCD calculation of X is determined by summing over the amplitudes of all Feynman diagrams contributing to that process. For a given Feynman diagram, the power of α_s is determined by the number of vertices associated with quark-gluon or gluon-gluon interactions. A leading order (LO) prediction sums over only the lowest-order contribution whereas next-to-leading order (NLO) includes terms with the additional powers of α_s . The next-to-next-to-leading order (NNLO) includes emission of another gluon or a virtual gluon loop. The different order of the QCD processes are shown in Fig. 2.3. The calculations become complex with the loop diagrams where the momenta of the virtual particles in a loop are not fully constrained by four-momentum conservation and the associated integrals are divergent. Such ultraviolet (UV) divergences enter




Figure 2.3: The Feynman diagrams³ of leading-order (LO), next-to-leading order (NLO) and next-to-next-to-leading order (NNLO) processes in quantum chromodynamics along with the perturbative expansion of any observable X in powers of the strong coupling constant α_S . At each successive step in perturbation series, the emission of an additional gluon take place.

the calculations beyond LO either due to loop or vertex corrections. These are overcome by a procedure known as renormalization, described in next section. Apart from the UV divergences, the QCD also suffers from infrared and collinear divergences (IRC) due to the presence of massless gluons and neglected quark masses. These need to be handled in pQCD calculations. The observable to be studied must be IRC safe.

³Drawn using ROOT

2.2.2 Renormalization and Running of the Strong Coupling

The renormalization is a mathematical procedure which allows the finite calculation of momenta integrals of virtual loop by removing UV divergences. It introduces a regulator for the infinities, the renormalization scale μ_r . At first, the divergences are regularized temporarily by introducing a cut-off to the loop momenta at μ_r scale. Then the free parameters of the Lagrangian, i.e. the coupling constant are redefined or renormalized to absorb the UV divergences. Due to this, both $\alpha_s(Q)$ and observable X become a function of μ_r . The exact dependence of $\alpha_s(\mu_r^2)$ on μ_r is described by the renormalization group equation (RGE) which determines the running of $\alpha_s(\mu_r^2)$. According to RGE, the dependence of X on μ_r must cancel. Mathematically this can be expressed as :

$$\mu_r^2 \frac{d}{d\mu_r^2} X \left(\frac{Q^2}{\mu_r^2}, \alpha_s(\mu_r^2) \right) = \left(\mu_r^2 \frac{\partial}{\partial \mu_r^2} + \mu_r^2 \frac{\partial \alpha_s(\mu_r^2)}{\partial \mu_r^2} \frac{\partial}{\partial \alpha_s(\mu_r^2)} \right) X = 0 \quad (2.9)$$

Using beta function $\beta(\alpha_s) = \mu_r^2 \frac{\partial \alpha_s(\mu_r^2)}{\partial \mu_r^2}$, Eq. 2.9 can be re-written as

$$\left(\mu_r^2 \frac{\partial}{\partial \mu_r^2} + \beta(\alpha_s) \frac{\partial}{\partial \alpha_s(\mu_r^2)} \right) X = 0 \quad (2.10)$$

By setting the renormalization scale equal to the physical scale i.e. $\mu^2 = Q^2$, $X(1, \alpha_s(Q))$ is a solution to above equation. Q -dependence of the X is only from the renormalization of the theory which is present in the classical theory. Hence measuring the Q -dependence of X will directly probe the quantum structure of the theory. The β function in QCD has a perturbative expansion as :

$$\beta(\alpha_s) = -\alpha_s^2 \left(b_0 + b_1 \alpha_s + b_2 \alpha_s^2 + \mathcal{O}(\alpha_s^3) \right) \quad (2.11)$$

where b_n is the $n+$ 1-loop β -function coefficients giving the dependence of the coupling on the energy scale Q . In the modified minimal subtraction ($\overline{\text{MS}}$) scheme [16, 17], the beta coefficient functions have following values :

$$b_0 = \frac{33 - 2n_f}{12\pi}, \quad b_1 = \frac{153 - 19n_f}{24\pi^2}, \quad b_2 = \frac{77139 - 15099n_f + 325n_f^2}{3456\pi^3} \quad (2.12)$$

where n_f is the number of quark flavours with masses $m_q < \mu_r$. On integration of Eq. 2.11, the energy dependence of α_S is yielded. Neglecting the higher orders, the first order solution of RGE is :

$$\alpha_s(\mu_r^2) = \frac{1}{b_0 \ln(\mu_r^2/\Lambda_{QCD}^2)} \quad (2.13)$$

where Λ_{QCD} is the constant of integration. The perturbative coupling becomes large at the scale Λ_{QCD} and the perturbative series diverge. With $b_0 > 0$, the coupling becomes weaker at higher scales Q , i.e. the effective color charge is small at small distances or large energies. This allows the quarks to behave as free particles within the hadron, leading to the property called asymptotic freedom. It is always convenient to express α_S at some fixed scale. Since some of the best measurements come from Z^0 decays, it is common practise to determine the strong coupling at the scale of the Z boson mass $\alpha_s(M_Z)$. So, Eq. 2.13 can be expressed as :

$$\alpha_S\left(\mu_r, \alpha_s(M_Z)\right) = \frac{\alpha_s(M_Z)}{1 + \alpha_s(M_Z)b_0 \ln(\mu_r^2/M_z^2)} \quad (2.14)$$

Since α_S is the free parameter of QCD theory, it is always extracted from the experimental measurements and evolved to the scale of the Z boson. According to Particle Data Group (PDG) [18], the current world average value of the strong

coupling constant at the scale of mass of Z boson is

$$\alpha_s(M_Z) = 0.1181 \pm 0.0011 \quad (2.15)$$

This value is derived using data from deep inelastic scattering process, electron-positron annihilation processes, hadronic τ lepton decays, lattice QCD calculations and electroweak precision fits. The different experimental determinations of the strong coupling constant evolved at the scale Q are shown as a function of Q in Fig. 2.4 which describe the running of the α_S up to the 1 TeV scale.




Figure 2.4: The different experimental measurements of the strong coupling constant α_S evolved at the energy scale Q are shown as a function of Q . These describe the running of the α_S up to the 1 TeV scale. Taken from [18].

2.3 Hadronic Collisions

At a large momentum transfer, the collision between two hadrons can be visualized as an interaction between their constituents - quarks and gluons. In this thesis, we

are studying the proton-proton collisions taking place at the Large Hadron Collider (LHC). A proton is a complex composite particle consisting of three valence quarks (uud), gluons for the exchange of the strong force and the sea quarks. The sea quarks consist of quark-antiquark pairs coming into and out of existence rapidly and continuously due to gluon colour field splitting. In any collision, one of the most important quantities to evaluate is the cross-section (σ) of a certain process which gives the probability that the two hadrons interact and give rise to that specific process. In a hadronic collision, the perturbation theory is only valid at the parton-level but due to property of confinement at low energies, free partons cannot exist in nature. Only hadrons with a complex internal structure are available for the high energy collisions. Here, a factorization theorem of QCD [19] comes into play which allows the calculation of σ by separating into two parts : a short-distance partonic cross-section calculable with pQCD, and a non-perturbative long-distance part described by parton distribution functions $f_i(x, \mu_f)$ (PDFs). The PDFs describe the partonic content of the colliding hadrons and give the probability to find a parton i with momentum fraction x within a hadron. μ_f is a factorization scale which corresponds to the resolution with which the hadron is being probed. The particles which are emitted with transverse momenta $p_T > \mu_f$ are considered in the calculation of hard scattering perturbative coefficients. The particles emitted with $p_T < \mu_f$ are accounted for within the PDFs. Applying the factorization theorem in a proton-proton collision, the cross-section of a hard scattering process can be written as :

$$\begin{aligned} \sigma_{P_1 P_2 \rightarrow X} = & \sum_{i,j} \int dx_1 dx_2 f_{i,P_1}(x_1, \mu_f) f_{j,P_2}(x_2, \mu_f) \\ & \times \hat{\sigma}_{ij \rightarrow X} \left(x_1 p_1, x_2 p_2, \alpha(\mu_r^2), \frac{Q^2}{\mu_f^2} \right) \end{aligned} \quad (2.16)$$

where f_i and f_j are the proton PDFs which depend on momentum fractions x_1 and x_2 of parent protons P_1 and P_2 respectively as well as on the factorization scale μ_f .

The sum extends over all contributing initial-state parton flavours i, j . The cross-section for the production of final state X at parton level ($\hat{\sigma}_{ij}$) depends on the final state phase, the factorization scale μ_f and the renormalization scale μ_r . Figure. 2.5 illustrates the factorization into the PDFs and the hard scattering cross-section in a proton-proton collision. The PDFs of the proton are a necessary input to almost




Figure 2.5: Schematic illustration⁴ of the factorization theorem in a collision of two protons P_1 and P_2 having momenta p_1 and p_2 , respectively. In a hard-scattering process at a scale Q^2 , the two partons x_1 and x_2 participate with momenta $x_1 p_1$ and $x_2 p_2$. The total cross-section is factorized into the hard scattering cross-section $\hat{\sigma}_{ij}$ calculable using perturbative Quantum Chromodynamics and the PDFs $f_i(x_1, \mu_f)$ and $f_j(x_2, \mu_f)$ with factorization scale μ_f .

all theory predictions of a proton-proton collision. The QCD does not predict the parton content of the proton. So the shapes of PDFs are determined in fits to experimental measurements of different experiments. The dependence of PDFs on μ_f is given by the Dokshitzer-Gribov-Lipatov-Altarelli-Parisi (DGLAP) [20–22] equations which use α_S and the RGE as inputs. The knowledge of proton PDFs mainly

⁴Drawn using ROOT

comes from the Deep Inelastic Scattering (DIS) HERA, fixed target and Tevatron data. The LHC data has a potential to improve constraints of the PDFs further as done in one of the recent CMS measurements [23]. There are several groups which determine the PDFs using the different minimization methods, phenomenological approaches and the methods to estimate the uncertainties. The global PDFs are the CTEQ [24], MMHT [25], NNPDF [26], ABM [27] and HERAPDF [28] at LO, NLO and NNLO.

2.3.1 Parton Shower and Hadronization

The partons involved in a hard scattering process get accelerated due to large momentum transfers. These accelerated partons emit QCD radiation in the form of gluons with successively lower energy. Unlike the uncharged photons in QED, the gluons themselves carry color charge and hence also emit further gluons. The emitted gluons in turn splits into $q\bar{q}$ pairs. This successive emission of partons lead to a parton shower. In a parton shower, the main contribution is by the collinear parton splitting and the soft gluon emissions. The parton shower mimics the effect of higher-order corrections to the hard process. These cannot be calculated exactly and are taken into account using the parton shower approximation. The two incoming partons which are constituents of two colliding hadrons and taking part in hard scattering process can also develop parton showers, commonly known as Initial-State Radiation (ISR). The initial partons produce showers till they collide to initiate the hard scattering process. The final outgoing partons produced from a hard scattering process can also undergo parton showering giving rise to Final-State Radiation (FSR). A parton shower terminates when the scale Q is below the hadronization scale ~ 1 GeV for QCD.

At the end of the shower, there is a decrease in the energy of partons due to successive emission of gluons. Due to this the coupling constant of QCD α_s evolves and become large. This leads to the confinement of colored quarks and

gluons into the color-neutral composite particles called hadrons and this process is known as hadronization. The hadronization takes place at low momentum transfer and hence non-perturbative in nature. Although no exact theory for hadronization is known, the different phenomenological models have been developed to simulate the hadronization process. The two main models implemented in Monte Carlo event generators to simulate the hadronization process are :

Lund String Model - In the Lund string model of hadronization [29], the highly energetic gluons are treated as field lines. Due to the gluon self-interaction, the gluons are attracted to each other forming a narrow tube or string of strong color field between a $q\bar{q}$ pair. This model is based on an observation that at distances greater than about a femtometer (fm), the potential energy $V(r)$ of colored quarks grows linearly with the increase in distance between them (r) as :

$$V(r) = \kappa r \quad (2.17)$$

where $\kappa \sim \text{GeV}/\text{fm}^2$ is the tension of the string connecting the quarks. When the q and \bar{q} are pulled apart from each other move apart, the gluonic string stretches. Due to this, the potential energy of the string grows at the expense of the kinetic energy of the quarks. As the potential energy becomes of the order of hadron masses, the string breaks at some point along its length, creating a new $q\bar{q}$ pair. The newly formed two string segments again stretch and break producing further $q\bar{q}$ pairs. This process of stretching and breaking continues until all the potential energy gets converted to $q\bar{q}$ pairs. This whole process is illustrated in Fig 2.6. The $q\bar{q}$ pairs then undergo hadronization due to confinement property. PYTHIA Monte Carlo generator uses the Lund string model.

Cluster Model - The cluster model of hadronization [30, 31] is based on pre-confinement property of QCD [32]. According to this property, at evolution scales Q_0 much less than the hard process scale Q , the partons produced in a shower are

clustered in colourless groups with an invariant mass distribution, depending on nature of hard process and Q_0 , not on Q . This model contains two steps : firstly all gluons split into $q\bar{q}$ pairs at the end of the parton shower and in the second step, a new set of low-mass color-singlet clusters are obtained which decay into either secondary clusters or directly into hadrons. The generator HERWIG is based on the cluster fragmentation model. However, this model has problems in dealing with the decay of very massive clusters.




Figure 2.6: Illustration of the hadronization process in Lund string model⁵. When the quark q and anti-quark \bar{q} are pulled apart from each other, the potential energy of the gluonic string connecting the quarks increases. As it becomes of the order of hadron masses, the string breaks and a new $q\bar{q}$ pair is created. The breaking of string and creation of $q\bar{q}$ continues till all the potential energy gets converted to $q\bar{q}$ pairs which then get hadronized.

2.3.2 Underlying Event

Due to the composite nature of the protons, their collisions are not clean events. The event structure is significantly more complex than that of the lepton collisions. The final states of the collisions involves the multiparticle calculations. In a high energy proton-proton collisions, the underlying event (UE) includes the effects which are not coming from the primary hard scattering process. The UE include the contributions from relatively small momentum transfer processes : initial and final-state radiations (ISR, FSR), leftover partons in the collisions called beam remnants

⁵Source : <http://inspirehep.net/record/806744>

and multiple parton interactions (MPI). Due to composite nature of proton, the remaining two partons which do not participate in a hard collision may also interact giving rise to multiple parton interactions. The UE induces an additional energy in an event which is not related to the main hard interaction. This acts an unavoidable background which needs to be removed. Hence, it is very crucial to study and understand the UE. The UE activity increases with Q and the center-of-mass energy \sqrt{s} . Figure 2.7 shows the complex variety of processes taking place in a single proton-proton collision.




Figure 2.7: A proton-proton collision⁶ involving the main hard scattering process along with the low momentum transfer underlying event (UE) contributions coming from initial- and final-state radiations (ISR and FSR) complemented with multiple parton interactions (MPI) and collisions from leftover partons called beam remnants.

The bunch of hadrons, produced from hadronization of quarks and gluons, gets collimated in the form of “jets” with the direction towards the direction of the initial parton that originated them. The jets are the final structures observed experimentally in the detectors. These act as a bridge between the elementary quarks and gluons of QCD and the final hadrons produced in high energy collisions. Therefore, at large momentum transfer of the interacting partons, the jets and their observables are the best tools to test the predictions of perturbative QCD. Also,

⁶Source : [The Energy Dependence of Min-Bias and the Underlying Event at CDF](#)

the jet production is sensitive to the strong coupling constant α_S . The precise knowledge of the jet production cross-section can help to extract the value of α_S and also to reduce the uncertainties of the PDFs of proton. In LHC, the simplest jet production process is $2 \rightarrow 2$ scattering process at leading-order giving dijet events. But the partons originating from ISR, FSR or MPI can also hadronize to produce jets greater than 2 in a single proton-proton collision. This results in the production of multijet events. The investigation of inclusive multijet event cross-sections permits more elaborate tests of QCD. Also, a precise study of jet variables helps to understand the signal and background modelling for the new physics search in hadronic final states. In this thesis, the inclusive multijet event cross-sections as well as the ratio of cross-sections are exploited to extract the value of strong coupling constant α_S . In the next section, we focus on the definition of a jet.

2.4 Jets

Jets [33] are the conical structures which group the hadrons into a single physics entity. Figure 2.8 shows the the outgoing partons of the hard scattering process in a proton-proton collision, undergoing fragmentation and hadronization processes and forming a conical jet with radius R. The jet structure was observed for the first time hadron production of e^+e^- annihilation process at SLAC in 1975 [34]. The partons can not be measured directly by the experiments because they can not exist freely in nature. The information about the dynamics of the partons can be obtained indirectly from jets. The configurations of high-energy quarks and gluons at short distances are truly reflected in the energy and angular distributions of the jets. Hence the jets are important to study. To perform the clustering of particles, a certain set of rules are followed in the form of jet algorithms.




Figure 2.8: In a proton-proton collision, the outgoing partons of the hard scattering process undergo fragmentation and hadronization processes producing a shower of partons which get collimated into a conical jet with radius R .

2.4.1 Jet Algorithms

Jet algorithms [35] provide a set of rules which determine how the particles can be clustered into a jet. In a jet algorithm, usually one or more parameters are involved that indicate how close two particles must be for them to belong to the same jet. These parameters can either measure closeness in coordinate space (cone algorithms) or in momentum space (sequential algorithms). The jet algorithms are applicable on parton, particle and calorimeter levels. A recombination scheme is always associated with a jet algorithm which calculates the momentum assigned to the combined particles. A jet algorithm along with its parameters and a recombination scheme forms a “jet definition”. A jet definition [36] must be simple to implement in an experimental analysis as well as in the theoretical calculation. It should be defined at any order of perturbation theory. It must yield finite cross-section at any order of perturbation theory that is relatively intensive to hadronization. In addition to these requirements, a jet algorithm must be infrared and collinear (IRC) safe. Infrared safety is the property by which the addition of a soft emission i.e. addition of a soft gluon should not change or modify the number of hard jets found in an event. In an infrared unsafe algorithm, a soft gluon emission in the middle of two cone

jets can lead to overlap of the two initial cones, as shown in Fig. 2.9 (top). This produces a single jet instead of initial two jets resulting in the change of number of jets. Collinear safety is the property by virtue of which the collinear splitting




Figure 2.9: Top : Infrared unsafe behavior of jet algorithm is illustrated where the presence of soft radiation between two jets may cause a merging of the jets that would not occur in the absence of the soft radiation. Bottom : Collinear unsafe behavior of jet algorithm is shown in which the number of jets change due to a collinear splitting⁷.

i.e. replacement of one parton by two at the same place should not modify the number of jets formed in a jet. This implies that the output of the jet algorithm should remain the same if the energy of a particle is distributed among two collinear particles. According to the collinear safety property, the two cases shown in Fig. 2.9 (bottom) should always produce a single jet. If an algorithm produces zero or two jets after collinear splitting, then it is not collinear safe.

The jet algorithms can be classified mainly into two types :

Cone algorithms - In the iterative cone (IC) algorithm [37], the jet is defined as a cone with fixed radius R in $\eta\phi$ space drawn around the highest energy seed. The relative distance (d) of all the particles is iteratively calculated and compared

⁷Source : <http://inspirehep.net/record/1251416>

with R . If the calculated $d < R$, the considered particles are clustered together in a jet and the directions of the clustered particles give the direction of the jet. On the other side i.e. if $d > R$, the considered particles initiate two different jets. The algorithm iterates until the cone is stable which means that the direction of sum of momentum of all the particles is same as that of the center of cone. But IC algorithm is not IRC safe. There is an another cone algorithm, Seedless Infrared-Safe cone (SIS-Cone) [38], which is an exact seedless i.e. does not rely on seed threshold and is IRC safe. This is a complex approach which tests the stability of all subsets of particles and has a complexity of $\mathcal{O}(N2^N)$ for N particles. But this algorithm is much slower and hence not preferred.

Sequential algorithms - The sequential algorithms [39] cluster the particles by defining a distance between pairs of particles and recombine the pair of closest particles successively. This is collinear and infrared safe algorithm. It is possible for the jet cones to overlap such that one particle is contained in more than one jet but the sequential algorithm never assigns a particle to more than one jet. The sequential algorithm is based on transverse momentum p_T of the particles and follows the below procedure :

1. First the distance d_{ij} between two particles i and j and distance d_{iB} of the particle to the beam are calculated.

$$d_{ij} = \min(p_{Ti}^{2p}, p_{Tj}^{2p}) \frac{\Delta R_{ij}^2}{R^2}, \quad d_{iB} = p_{Ti}^{2p} \quad (2.18)$$

$$\text{where } \Delta R_{ij}^2 = (\eta_i - \eta_j)^2 + (\phi_i - \phi_j)^2$$

2. If $d_{ij} < d_{iB}$, then the particles i and j are merged into a new single particle k , summing four-momenta of two initial particles by recombination scheme and step 1 is repeated.

3. If $d_{iB} < d_{ij}$, particle i is declared as a final-state jet and the particle gets removed from the list.

This procedure continues until all particles get clustered into jets. The value of the parameter p defines the three different sequential algorithms having distinct properties. For $p = 1$, we have k_t algorithm [40, 41], $p = 0$ gives the Cambridge/Aachen (C/A) algorithm [42] whereas $p = -1$ defines the anti- k_T algorithm [43]. The k_t algorithm involves clustering of soft particles first resulting in an area that fluctuates considerably. This algorithm is susceptible to the underlying and pile-up events. The C/A algorithm involves energy independent clusterings. Both k_t and C/A produce jets of irregular shapes. Instead of jet analysis, these are widely considered for studying the jet substructure. The anti- k_T algorithm tends to cluster hard particles first and produce jets with circular regular shapes. It is less sensitive to underlying and pile-up events. It is the most preferred algorithm for jet studies at the LHC. Figure 2.10 shows the clustering of same particles but using the different jet algorithms.

A jet algorithm must specify how to combine the momenta of different partons or particles going to be clustered into a jet. This is given by the recombination scheme. The most widely used recombination scheme is the E -scheme [37] which corresponds to vector addition of four-momenta where the four-momenta of the jet is obtained by simply adding the four-momenta vector of merging particles.

The sequential clustering algorithms have always been favoured by theorists but not by experimentalists because of slow computational performance. However, the introduction of the FASTJET program [44] enhanced the speed of clustering algorithms and hence are preferred by experimentalists as well. This thesis studies the particles produced in proton-proton collisions by clustering them in to jets using anti- k_t algorithm with distance parameter $R = 0.7$. These jets are observed in the Compact Muon Solenoid detector of the Large Hadron Collider, whose details are discussed in the coming chapter.




Figure 2.10: The clustering of particles, in y - ϕ space at the parton level, into jets clustered with the k_t (top left), Cambridge/Aachen (top right), SIScone (bottom left) and anti- k_t (bottom right) algorithms with $R = 1$. The towers represent the jet p_T . The anti- k_t algorithm gives circular jets while the jets produced with other three algorithms have irregular shapes. Taken from [35].

Chapter 3

Experimental Setup

The hadron colliders aim at search for elementary particles and their interactions as predicted in the Standard Model or theories beyond the Standard Model. For the same beam energy, higher center-of-mass energy can be achieved by the hadron colliders as compared to fixed target experiments. Due to the availability of very high center-of-mass energy for the collisions, it becomes possible for the researchers to understand the fundamental structure of the universe deeply and to look back in its history. The precise measurement of mass of the Z and W boson discovered by the UA1 and UA2 experiments was done at the Large Electron-Positron (LEP) collider. The discovery of top quark and an acute measurement of its mass was performed by the proton-antiproton collider Tevatron at FNAL. The search for the long awaited Higgs boson was carried out by the currently running most powerful accelerator, the Large Hadron Collider (LHC). Still many questions related to the the nature of dark matter, the existence of super-symmetry (SUSY) or the extra dimensions are yet to be answered.

The European Organization for Nuclear Research (CERN) is a world-class fundamental physics research organization founded in 1954. In the beginning, it concentrated on pure physics research to understand the inside of the atom, justifying the word “nuclear” in name. At present, the main area of research at CERN

is particle physics - the study of the fundamental constituents of matter and the forces of interactions acting between them. To complete this task, several particle accelerators have been built by CERN to explore the physics at the TeV energy scale.

3.1 The Large Hadron Collider

The Large Hadron Collider (LHC) [45] is the world's biggest and the most powerful particle accelerator and collider built by CERN. It occupies the 27 km circumference circular tunnel (between the border of France and Switzerland), previously used by LEP collider [46], at a depth ranging from 50 to 175 metres (164 to 574 ft) underground. Two beams of particles of the same kind, either protons or lead or xenon ions, are accelerated in direction opposite to each other. 1,232 dipole magnets maintain the beams in their circular path and 392 quadrupole magnets keep the beams focused to increase the probabilities of interaction between the particles. Since this thesis is based on the proton-proton (pp) collisions data, the main focus is on protons.

The protons pass through a series of accelerators which increase their energy successively before their injection into the main ring of LHC. Figure 3.1 gives an overview of the various accelerators and detectors comprising the complex structure of the LHC. The protons are obtained by stripping of electrons from hydrogen gas atoms using an electric field. The protons are accelerated up to 100 keV through a radiofrequency quadrupole which provides the first focusing and a further acceleration to 750 keV energy. The linear particle accelerator (LINAC2) increases the energy of protons to 50 MeV. Then these protons are injected into the Proton Synchrotron Booster (PSB) in the form of bunches where they get accelerated to 1.4 GeV energy. The energy of protons is further enhanced to 25 GeV by Proton Synchrotron (PS) and then to 450 GeV by the Super Proton Synchrotron (SPS). Finally

the protons are injected into two beam pipelines of the main LHC ring where their energy increases to the collision energy. In head on collisions for colliding beams of the same mass particles, the center of mass system and the laboratory system coincide and the total center of mass energy is twice the energy of the beams.




Figure 3.1: Overview of the different experiments of the Large Hadron Collider (LHC), a complex particle accelerator and collider located at CERN⁸.

The accelerated beams are made to collide at four interaction points around which six detectors : ALICE (A Large Ion Collider Experiment) [47], ATLAS (A Toroidal LHC Apparatus) [48], CMS (Compact Muon Solenoid) [49–51], LHCb (Large Hadron Collider for Beauty) [52], LHCf (Large Hadron Collider forward) [53] and TOTEM (Total, elastic and diffractive cross-section measurement) [54], are located. The CMS and ATLAS are the two general purpose detectors dedicated to the search of the Higgs boson, existence of super-symmetry (SUSY) and also looking

⁸Source : https://en.wikipedia.org/wiki/Large_Hadron_Collider

for extra dimensions. The ALICE is a heavy-ion detector which collides lead ions to study quark-gluon plasma, a state of matter believed to be present just after the Big Bang. The LHCb experiment will explore the differences between matter and anti-matter and new physics through b-quark (beauty) studies. TOTEM experiment is dedicated to cross-section measurements whereas LHCf focuses on forward physics.

The LHC successfully injected the first protons on September 10, 2008 but after few days magnetic quench occurred in about 100 bending magnets leading to a loss of ~ 6 tonnes of liquid helium. The low-energy beams circulated in the tunnel for the first time on November 20, 2009 and after three days, the first particle collisions took place in all four detectors at $\sqrt{s} = 450$ GeV. The LHC achieved 1.18 TeV energy per beam on November 30, 2009 and become the world's highest energy particle accelerator leaving behind the Tevatron with record of 0.98 TeV per beam for eight years. The pp collisions at $\sqrt{s} = 2.36$ TeV were recorded around December 15, 2009. On March 19, 2010, the beam energy was ramped up to 3.5 TeV, resulting in the first pp collisions at $\sqrt{s} = 7$ TeV on March 30, 2010. The beam energy was kept at 3.5 TeV throughout 2011, and increased to 4 TeV in 2012. After a long shutdown for two years, the LHC restarted in 2015 and collided the proton beams at a much higher centre-of-mass energy of 13 TeV and is running successfully till now. In the coming years, protons will be made to collide at a designed $\sqrt{s} = 14$ TeV with luminosity up to 10^{34} cm $^{-2}$ s $^{-1}$. In this thesis, work has been carried out using the pp collisions data collected by the CMS detector at $\sqrt{s} = 8$ TeV in the year of 2012.

3.1.1 Luminosity Measurement

Luminosity (\mathcal{L}) is one of the most important parameters of an accelerator which characterizes its performance. It gives the rate at which collisions occur and given by the number of collisions produced in a detector per cm 2 and per second. Cross-

section (σ) is a measurement of the probability that an event will occur. It is related to total number of events N of a process over a time period T and \mathcal{L} as :

$$N = \int_0^T \mathcal{L} \sigma dt = \mathcal{L}_{int} \sigma \quad (3.1)$$

where $\int_0^T \mathcal{L} dt = \mathcal{L}_{int}$ is the total integrated luminosity. It is expressed in units of area, usually in barn $^{-1}$ and gives a direct indication of the number of produced events for a process. For example, an integrated luminosity of 10 fb $^{-1}$ means that 10 events are produced in a process with cross-section equal to 1 fb.

The luminosity depends on the particle beam parameters and is given by :

$$\mathcal{L} = \frac{N_p^2 N_b f_{rev} \gamma F}{4\pi \epsilon_n \beta^*} \quad (3.2)$$

where N_p is the number of particles per bunch, N_b is the number of bunches per beam, f_{rev} is the revolution frequency of the beam, γ is the relativistic gamma factor and F gives the geometric luminosity reduction factor. The effective collision area of the two beams is related to the normalized transverse beam emittance ϵ_n and the value of the betatron function β^* at the interaction point.

The CMS experiment constantly monitors the instantaneous luminosity delivered by LHC which is shown versus time in Fig. 3.2 for proton-proton collisions at nominal center-of-mass energy for the years 2010-2017. The relative instantaneous luminosity is calculated by using two methods [55] : Hadron Forward (HF) method by measuring the particle flux in the hadron forward calorimeter and by counting the number of reconstructed vertices in the pixel tracker. The absolute luminosity measurement relies on van-der-Meer scans done in special runs of the LHC [56]. The uncertainty on the luminosity measured for 2012 data set is 2.5% (syst.) and 0.5% (stat.).




Figure 3.2: The integrated luminosity, delivered to CMS during stable beams for proton-proton collisions at nominal center-of-mass energy, is shown versus time for data-taking in 2010 (green), 2011 (red), 2012 (blue), 2015 (purple), 2016 (orange) and 2017 (light blue) run periods of the LHC⁹.

3.1.2 Pileup Interactions

To observe the extremely rare events, the event rate in a collider should be very high. This demands delivered luminosity to be high which is achieved by increasing the number of bunches or increasing the number of protons per bunch. However, this comes at the cost of multiple proton-proton interactions coming from independent hadron-hadron collisions occurring in the same bunch crossing, called pileup (PU) interactions. The hard interaction in every event is accompanied by a large amount of PU interactions which give rise to low p_T jets. The vertex of pile-up interaction is reconstructed from tracks pointing to it as shown in Fig. 3.3. The pileup due to additional collisions within a single bunch crossing is called in-time pileup whereas pileup coming from collisions other than hard scattering in other bunch crossings is

⁹Source : <https://twiki.cern.ch/twiki/bin/view/CMSPublic/LumiPublicResults>

known as out-of-time pileup. The pileup itself cannot be directly measured, it can be correlated to various other directly measurable quantities. Since pileup comes from additional proton-proton interactions, the number of primary vertices (N_{PV}) is directly correlated to the amount of pileup. The greater the N_{PV} , the more pileup energy is added to the jets which needs to be subtracted.




Figure 3.3: In a proton-proton collision, the particles produced from the hard interaction are clustered into a jet. The hard interaction corresponds to the main vertex. The particles produced in the interactions other than the hard one, form a pileup jet¹⁰.

3.2 The Compact Muon Solenoid

The Compact Muon Solenoid (CMS) detector is a general purpose detector located at the interaction point 5 (P5) of the main LHC ring, near the village of Cessy in France. The name of CMS comes from its compact size with main emphasis on the detection of muons and enclosed within high solenoidal magnetic field. The CMS detector aims at identifying the different types of particles produced in proton-

¹⁰Source : <http://cds.cern.ch/record/1747055>

proton and heavy ion collisions and measuring their energies and momenta. This is achieved by concentric layers of different sub-detectors arranged in a cylindrical complex structure with 21.6 m length and 15 m diameter. The silicon-based tracker surrounds the the interaction point and forms the innermost layer. It is surrounded by a scintillating crystal electromagnetic calorimeter (ECAL) and a sampling hadron calorimeter (HCAL) which are enclosed inside the superconducting solenoid. Outside the magnet lies the large muon detectors embedded inside an iron yoke. The




Figure 3.4: The three dimensional view of the CMS detector along with its sub-detector components¹¹.

three dimensional view of the CMS detector along with its components is presented in Fig. 3.4. The CMS was constructed in parts at ground and assembled later on in the cavern. The components are easily accessible for upgrades or repairs as the

¹¹Source : <https://orbiterchspacenews.blogspot.in/2013/04/cern-cms-prepares-for-future.html>

detector can be opened up into movable slices. Figure 3.5 shows the front view of the CMS detector differentiating individual components which contribute to event reconstruction. The path of reconstructed particles is represented by dashed (invisible track) and solid (visible track) lines for different particle classes : photons (γ), muons (μ^\pm), electrons (e^-), neutrons (n) and charged hadrons (pions π^\pm).




Figure 3.5: Front view of the CMS detector along with its components : inner tracker, outer tracker, electromagnetic calorimeter, hadronic calorimeter, solenoid and muon system. The path of different particles detected by dedicated sub-detectors are shown by dashed (invisible track) and solid (visible track) lines. \otimes and \odot gives the direction of magnetic field inside the solenoid and in the return yoke, respectively. Taken from [51].

A brief overview of the CMS detector has been presented and the details of the its design as well as physics performance are available in Ref. [50, 51]. Before going into the details of each sub-detector, first the CMS coordinate system is described in the next section.

3.2.1 Coordinate System

CMS uses right-handed coordinate system, illustrated in Fig. 3.6, having origin at the nominal interaction point (IP) of the collision inside the detector. The x -

axis points horizontally from the IP, towards the center of the LHC ring, the y -axis vertically upwards and the z -axis along the beam direction towards the Jura mountains. The radial coordinate in x - y plane is denoted by r . Following customary




Figure 3.6: The right-handed coordinate system used by the CMS detector¹² with origin at the interaction point (IP), x -axis points horizontally from the IP towards the center of the LHC ring, the y -axis vertically upwards and the z -axis along the beam direction towards the Jura mountains, ϕ as the azimuthal angle measured from the x -axis in the x - y plane and θ as the polar angle calculated from the z -axis in the z - y plane.

polar coordinate conventions : the azimuthal angle ϕ is measured from the x -axis in the x - y plane as $\phi = \tan^{-1}(\frac{y}{x})$ where $\phi = 0$ points to the $+x$ axis and $\phi = \pi/2$ points to the $+y$ axis. The polar angle θ , is calculated from the z -axis in the z - y plane as $\theta = \tan^{-1}(\frac{x^2 + y^2}{z})$ with $\theta = 0$ corresponding to the $+z$ direction and $\theta = \pi$ to the $-z$ direction. The quantities pseudorapidity η and the rapidity y are preferred over the angles θ and ϕ . The pseudorapidity and rapidity are given by Eq. 3.3. Both the

¹²Source : https://wiki.physik.uzh.ch/cms/latex:example_spherical_coordinates

quantities are equal for massless particles.

$$\begin{aligned}\eta &= -\ln\left(\tan\left(\frac{\theta}{2}\right)\right) \\ y &= \frac{1}{2} \ln\left(\frac{E + p_z}{E - p_z}\right)\end{aligned}\tag{3.3}$$

The difference between rapidities Δy is invariant under longitudinal Lorentz boost whereas it does not hold for η . Hence y is considered in this thesis. The angular distance between the two particles is defined by $\Delta R = \sqrt{(\Delta\eta)^2 + (\Delta\phi)^2}$. The momentum component transverse to the direction of beam p_T , is computed from the x - and y -components as $p_T = \sqrt{p_x^2 + p_y^2}$ and the transverse energy is given by $E_T = E \sin\theta$. After introducing the CMS coordinate system, further the detector subsystems are described briefly in the following sections. A longitudinal section of the CMS detector in the y - z plane, shown in Fig. 3.7, locates the different subsystems along with the superconducting solenoid.

3.2.2 Inner Tracker System

The charged particles produced from the LHC collisions leave their trajectories as they move outward from the interaction point. The particle flux within the detector decreases as $1/r^2$. So the tracks of the particles need to be measured as close to the collision point as possible and in a precise manner. The innermost tracking system of the CMS consisting of silicon detectors measures the hits of charged particles. It surrounds the interaction point and has a cylindrical volume of length of 5.8 m and a diameter of 2.5 m and covers a pseudorapidity range up to $|\eta| < 2.5$. When charged particles pass through the silicon detector material, small ionization currents are produced which are detected as a hit. Such multiple hits when combined, reconstruct the track which gives the information about the direction and trans-




Figure 3.7: Longitudinal section of the CMS detector in the y - z plane¹³. It shows the tracking detector (TIB, TID, TOB, TEC) close to the nominal interaction point at $(0,0)$, the electromagnetic (EB, EE) and hadronic (HB, HE, HO, HF) calorimeters. The coil of the solenoid magnet (CB) surrounds the inner barrel region. The iron return yoke (YB, YE) is interleaved with the muon chambers (MB, ME).

verse momentum p_T of the charged particle. Silicon detectors have a much higher resolution in tracking charged particles as compared to the older ones such as cloud chambers or wire chambers. CMS inner tracking system shown in Fig. 3.8 consists of two sub-systems :

Pixel Detector - A pixel detector lying close to the beam pipe have three concentric barrel layers at radii of 4.4, 7.3 and 10.2 cm from the beam pipe. It has two disks of pixel modules on each side of barrel. At the LHC design luminosity of $10^{34} \text{ cm}^{-2}\text{s}^{-1}$, about 1000 particles from more than 20 overlapping proton-proton interactions traverse through the tracker for each bunch crossing, i.e. every 25 ns.

¹³Source : <http://cds.cern.ch/record/1747055>

The size of each pixel is $100 \mu\text{m} \times 150 \mu\text{m}$ which gives an average occupancy of 10^{-4} per bunch crossing. By taking an advantage of the large Lorentz effect, the pixel tracker has a resolution of $10 \mu\text{m} \times 20 \mu\text{m}$ needed for a precise determination of the primary and secondary vertices and the required momentum resolution.

Strip Detector - After coming out of the pixel detector the charged particles pass through ten layers of silicon strip detectors, reaching out to a radius of 130 cm. The silicon strip detector consists of four inner barrel (TIB) layers assembled in shells with two inner endcaps (TID), each composed of three small discs. The outer barrel (TOB) consists of six concentric layers. Finally two endcaps (TEC) close off the tracker. Each part has silicon modules designed differently for its place within the detector. The strip detector measures the particle tracks with a reduced resolution of $23 \mu\text{m}$ reflecting the smaller particle flux at larger distances from the interaction point. The active silicon area of CMS tracker is about 200m^2 which




Figure 3.8: The longitudinal section of the inner tracking system consisting of the silicon pixel detector and the silicon strip detector is shown in r - z plane. The silicon strip detector has four components : The Tracker Inner Barrel (TIB) complemented by the Tracker Inner Disks (TID) which are further surrounded by the Tracker Outer Barrel (TOB) in barrel region. Tracker End Cap (TEC) covers high η ranges up to $\eta = 2.5$. Taken from [49].

makes it the largest silicon tracker. Along with the measurement of tracks, the en-

ergy also needs to be measured for which the calorimeters are present outside the tracker. The tracker should interfere with the particles to a minimum extent so that their momentum can be measured precisely but to measure their energy, they are required to interact with the calorimeters fully.

3.2.3 Electromagnetic Calorimeter

The electromagnetic calorimeter (ECAL) is a homogeneous and hermetic calorimeter used to slow down the produced photons and electrons/positrons and measure their energy by absorb them into the detector material. ECAL consists of 61200 lead tungstate (PbWO_4) crystals in the barrel part and 7324 crystals in each of the two endcaps. PbWO_4 is a very dense material with a short radiation length of $X_0 = 0.89$ cm and covers the pseudorapidity up to $|\eta| < 3.0$. The particles interact with matter and produce electromagnetic shower through the subsequent processes of bremsstrahlung and electron-positron pair production. The energy of the particles deposited by Compton scattering and the photoelectric effect causes excitation of the material atomic state and the emission of photons. These photons are detected by silicon avalanche photo diodes (APDs) in the barrel region and vacuum phototriodes (VPT) in the end-cap region. The number of created photons gives the direct measure of energy of the incident particle. The incorporation of oxygen makes it highly transparent and enables to emit scintillation light. The small Molière radius of 2.19 cm gives a fine granularity. These properties leads to compact size of ECAL and can be placed easily within the solenoid magnet.

Figure 3.9 presents a geometric view of ECAL in the y - z plane showing the arrangement of different parts of ECAL : the ECAL barrel (EB) extending up to $|\eta| < 1.479$ using more than 60000 crystals and ECAL endcaps (EE) covering the region $1.479 < |\eta| < 3.0$ with an additional 15000 crystals. The preshower detectors (ES) made of lead absorbers and silicon detectors are put in front of the endcaps to

distinguish high energetic single photons from low energetic photon pairs originating from neutral pions decays.




Figure 3.9: A geometric view of one quarter of the electromagnetic calorimeter (ECAL) in y - z plane showing the arrangement of sub-modules covering the barrel region (EB) and the endcaps (EE). ECAL is complemented with preshower detector (ES) mounted in front of the endcaps. Taken from [50].

The relative energy resolution of the ECAL has been measured to be [57] :

$$\left(\frac{\sigma(E)}{E}\right)^2 = \left(\frac{2.8\%}{\sqrt{E}}\right)^2 + \left(\frac{12\%}{E}\right)^2 + \left(0.30\%\right)^2 \quad (3.4)$$

where E is the energy in GeV. The first term is the stochastic component caused by fluctuations in lateral shower containment and in the energy deposited in the preshower absorber. The second term is the contribution by noise and the last is the constant term comes from the non-uniformity of the longitudinal light collection, inter-calibration errors, and leakage of energy from the back of the crystal.

3.2.4 Hadronic Calorimeter

At CMS, the major fraction of the produced particles in proton-proton collisions is hadrons. The combined CMS calorimeter system measures directions and energies

of quarks, gluons and neutrinos by measuring the energy and direction of particle jets. The neutral hadrons do not leave track and hence their energy is measured by taking into account the missing transverse energy (E_T^{miss}). The determination of E_T^{miss} is a crucial tool in searching the new particles and phenomena. The hadron calorimeter (HCAL) are particularly important in the identification of hadron jets, neutrinos as well as electrons, photons and muons in conjunction with the ECAL and the muon system. Hence HCAL is an essential sub-system of the CMS detector and contributes to most of CMS's physics studies.

HCAL is a sampling calorimeter installed inside the solenoid coil. It is made up of non-magnetic brass absorber having a short interaction length of $\lambda_I = 16$ cm interleaved with plastic scintillators having wavelength-shifting (WLS) fibres as readout. The hadrons produced with high energy showers a large number of pions and nucleons by inelastic interactions. The hadronic shower spreads more than the electromagnetic showers because of large transverse momentum of the secondary particles. As the energy of the particles is lower than a certain threshold, the ionization and low-energy hadronic processes come into play. The active scintillation material gets excited and emits blue-violet light. All scintillators are connected to photodiodes using wavelength shifters which read out the signals and pass them to the data acquisition system. The longitudinal view of one quarter of the HCAL presented in Fig. 3.10 shows the different parts :

Hadron Barrel - The hadron barrel (HB) is divided into two identical half barrel sections on either side of the interaction point. Each half barrel is made of 18 azimuthal wedges which are further divided into four azimuthal sectors each giving a granularity of $\Delta\phi = 0.087$. In z direction, the plastic scintillators are divided into 16 intervals of granularity $\Delta\eta = 0.087$. HB covers the region up to $|\eta| < 1.305$ and overlaps with endcaps for $1.305 \leq |\eta| \leq 1.392$. Since HB has the highest resolution ($\Delta\eta \times \Delta\phi = 0.087 \times 0.087$), it constitute the optimal region for a data-driven calibration of the jet energy scale. The thickness of the HCAL amounts to 7-11




Figure 3.10: Longitudinal view of one quarter of the hadronic calorimeter (HCAL) detector in r - η plane showing its different parts : hadron barrel (HB), hadron outer (HO), hadron endcap (HE) and hadron forward (HF). Taken from [49].

interaction lengths and which should be sufficient enough to stop nearly all hadrons in the calorimeter.

Hadron Outer - The total amount of material in barrel region to absorb the hadronic shower is not sufficient. This requirement is fulfilled by placing an outer hadron (HO) calorimeter as a tail catcher on top of the coil of the magnet. The HO utilizes the solenoid coil as an additional absorber equal to $1.4/\sin\theta$ interaction lengths and measures the tails of hadron showers penetrating the HB and the coil. Since the HO is physically located inside the muon system, it is strongly constrained by its geometry. The muon system is subdivided into 5 rings along the z -axis. Each of these rings is 2.536 m wide in z -direction and the HO is placed as first sensitive layer in these rings, with a scintillator thickness of 10 mm. The central ring ($\eta = 0$) has two scintillator layers placed on each side of 19.5 cm thick iron layer.

Hadron Endcap - The hadron endcaps (HE) extends the pseudorapidity range

up to $|\eta| < 3.0$, a region containing about 34% of the particles produced in the final state. The granularity in $\Delta\eta \times \Delta\phi$ is 0.087×0.087 up to $|\eta| < 1.6$ and 0.17×0.17 for $|\eta| > 1.6$. The usage of non-magnetic material in order to not disturb the magnetic field and the close distance to the beam line were the main challenges faced during the construction of the HE. The continuous radiation damages decrease the detector response which need to be corrected and monitored at regular intervals.

Hadron Forward - The hadron forward (HF) calorimeter lies $2.8 < |\eta| < 5.2$ region and is placed more closer to the beam pipe at distance of $z = \pm 11.2$ m from the interaction point. It is essentially a cylindrical steel structure with an outer radius of 130.0 cm which is azimuthally subdivided into 36 20° modular wedges. The HF is made of 5 mm thick grooved steel plates which have quartz fibers inserted into the grooves. The fibres run parallel to the beam line which are bundled to form 0.175×0.175 ($\Delta\eta \times \Delta\phi$) towers. HF detects the jets with very high η and the hadronization products of the beam remnants. It is built using iron absorbers and quartz fibers as active material, which measures the emitted Cerenkov light and produces the signal in the photomultipliers (PMT).

The relative hadronic energy resolution of the barrel HCAL and ECAL combination can be parametrized as :

$$\left(\frac{\sigma(E)}{E}\right)^2 = \left(\frac{a}{\sqrt{E}}\right)^2 + b^2 \quad (3.5)$$

with a stochastic term a and a constant term b . These values have been measured [58] as $a = (0.847 \pm 0.016) \sqrt{\text{GeV}}$ and $b = 0.074 \pm 0.008$ whereas for HF the measured values are $a = 1.98 \sqrt{\text{GeV}}$ and $b = 0.09$.

3.2.5 Superconducting Magnet

The superconducting magnet is the key feature of the CMS detector which is 13m long and 6m in diameter. Its refrigerated superconducting high-purity aluminium-stabilized niobium-titanium coils cooled at 4 Kelvin produces a magnetic field of 4 Teslas (T). The magnet will run at 3.8 T in order to maximize its lifetime. This intense solenoidal field makes the compactness and cylindrical symmetry of the detector possible. The magnet is placed between the calorimeters and the muon system. The solenoidal magnetic field parallel to the beam bends the tracks of the high momentum charged particles in the transverse plane. The curvature of the trajectory increases with the strength of the magnetic field which make possible to determine the transverse momentum more precisely. The magnet is complemented by a \sim 10000 tonnes iron yoke which returns the magnetic field at 2 T.

3.2.6 Muon System

As the name suggests, the detection of muons is of central importance to CMS. Only the muons and neutrinos, out of all the known stable particles, pass through the calorimeter without depositing most or all of their energy. They interact very little with matter and can travel long distances through the dense matter. The charged muons can be detected by having an additional tracking system outside the calorimeters whereas the neutrinos are practically undetectable as they escape completely without being tracked in any of the layers. Their presence can be detected from the missing energy carried by them. The CMS muon system is installed outside calorimeters in the iron return yoke of the magnet. The muon system perform the muon identification, momentum measurement and triggering. Good muon momentum resolution and trigger capability are enabled by the high-field solenoidal magnet and its flux-return yoke. The latter also serves as a hadron absorber for the identification of muons. The CMS muon system is designed to have the capability of reconstructing the momentum and charge of muons over the entire kinematic range

of the LHC. The muon system shown in Fig. 3.11 consists of three types of gaseous particle detectors :

Drift Tube - The muon barrel (MB) detector has four concentric layers of drift




Figure 3.11: A longitudinal view presenting the location of the three muon subsystems : four Drift Tube (DT) stations in the barrel (MB1-MB4, green), four stations of Cathode Strip Chambers (CSC) in the endcap (ME1-ME4, blue), and the Resistive Plate Chambers (RPC) stations (red)¹⁴.

tube (DT) chambers inside the iron yoke which covers the region up to $|\eta| < 1.2$. DT stations are distributed into 5 wheels along the z direction. Each wheel is divided into 12 sectors, each covering a 30° azimuthal angle. The DTs are aluminium tubes of 2.5 m of length and $4.2 \times 1.3 \text{ cm}^2$ of area, filled with gas mixture (58% Ar + 15 % CO₂).

Cathode Strip Chambers - In the forward region, the muon and background

¹⁴Source : <https://arxiv.org/abs/1209.2646>

flux is higher. So cathode strip chambers (CSC) are preferred because of their fast response time, radiation tolerance and fine segmentation. The four stations of CSCs are installed in each end cap covering in total the region of $0.9 < |\eta| < 2.4$. Each CSC is trapezoidal in shape and consists of 6 gas gaps, each gap having a plane of radial cathode strips and a plane of anode wires running almost perpendicularly to the strips.

Resistive Plate Chambers - Both DT and CSC are accompanied by resistive plate chambers (RPC) which are double-gap chambers, operated in avalanche mode to ensure good operation at high rates. They help to resolve ambiguities in attempting to make tracks from multiple hits in a chamber and provide additional points for determination of a muon trajectory. They provide fast response to the trigger system which is described in the following section.

3.2.7 Trigger and Data Acquisition System

The proton-proton collisions (events) take place at high interaction rates at LHC. In the 2012 run period, at beam crossing frequencies of 25 ns there are 40 million bunch crossings per second with an average of around 20 collisions per bunch crossing. Both the collision and the overall data rates are much higher than the rate at which the information can be stored. So a dramatic rate reduction has to be achieved which is possible with an efficient trigger system by retaining interesting signal events and rejecting background events. The decision of accepting or rejecting an event has to be performed very quickly and it is based on signals of certain physics objects inside the detector. CMS has a two-level complex trigger system :

Level-1 Trigger - The Level-1 (L1) trigger system is based on custom electronics which stores the events at maximum rate of 100 kHz and then forward them to the next level triggers. The L1 system uses only coarsely segmented data from calorimeter and muon detectors, while holding all the high-resolution data in pipeline

memories in the front-end electronics. Figure 3.12 shows the work flow of the L1 trigger system. L1 trigger consists of local, regional and global components. The




Figure 3.12: Work flow of the L1 trigger system. Taken from [49].

local triggers known as Trigger Primitive Generators (TPG), are based on energy deposits in calorimeter trigger towers and tracks in muon chambers. Regional Triggers combine their information and use pattern logic to determine ranked and sorted trigger objects such as electron or muon candidates in limited spatial regions. The rank is determined as a function of energy or momentum and quality, which reflects the level of confidence attributed to the L1 parameter measurements, based on detailed knowledge of the detectors and trigger electronics and on the amount of information available. The Global Calorimeter and Global Muon Triggers determine the highest-rank calorimeter and muon objects across the entire experiment and transfer them to the Global Trigger, the top entity of the Level-1 hierarchy.

The latter takes the decision to reject an event or to accept it for further evaluation by the HLT.

High Level Trigger - At the second step, a software-based High-Level Trigger (HLT) is designed to reduce the maximum L1 accept rate of 100 kHz to a final output rate of 100 Hz. The HLT system filter events by performing physics selections using faster versions of the offline reconstruction software. The HLT is provided by a subset of the on-line processor farm which, in turn, passes a fraction of the accepted events to the Data Acquisition (DAQ) system for more complete processing.

3.2.7.1 Jet Triggers

Triggers based on jet and missing transverse energy (E_T^{miss}) triggers are important for search for new physics whereas the single-jet triggers are mainly designed to study quantum chromodynamics (QCD). In this thesis, the single-jet triggers are used to select the events for analysis. At L1, they use mainly information from the calorimeter by looking for an energy cluster and a high energy deposit. The sums of transverse energy from ECAL and HCAL are computed in 4×4 trigger towers, except in the HF region where this quantity is measured in the whole trigger tower itself. If this deposit is greater than a certain threshold, the event is selected at L1 and it is passed to the HLT. Jets are reconstructed in the HLT using the anti- k_t jet clustering algorithm. The inputs for the jet algorithm are either calorimeter towers giving “CaloJet” objects, or the reconstructed particle flow objects giving “PFJet” objects. The processing time of reconstruction algorithm is high and hence the jet trigger paths are divided into multiple selection steps. At first, jets are reconstructed from calorimeter towers. Only for events in which at least one calorimeter jet passes a certain p_T threshold, the particle flow algorithm is run and the jets are clustered again from the particle flow candidates. In 2012, most of the jet trigger paths use PFJets as their inputs. The rate of jet events is quite high, so PFJet trigger paths have a pre-selection based on CaloJets. The matching between CaloJets and PFJets

is required in single PFJet paths. Due to the flexibility of the HLT, it is possible to apply the jet energy corrections during the HLT selection.

3.2.7.2 Data Acquisition System

As the L1 trigger accepts events at a rate of 100 kHz, the Data Acquisition (DAQ) system has to process the events at the same speed. It reads out the data of all detector sub-components and assembles the complete events, see Fig. 3.13. The data is subsequently passed to the HLT which further reduces the rates to a few hundred events per second. Finally, the events are merged and saved to a local storage system, from which they are continuously transferred to the Tier-0 computing center at CERN.




Figure 3.13: Architecture of the CMS DAQ system. Taken from [49].

3.2.8 Data Management

Although the trigger system reduces the collision rate enough to be stored in tape, still there is a huge amount of data need to be analyzed. An efficient computing infrastructure and the software is required for storing and distributing the data. To meet this need, the LHC has a data storage infrastructure called the Worldwide LHC Computing Grid (WLCG) [59]. WLCG provides a hierarchical structure, as

shown in Fig. 3.14, in a series of four levels or Tiers. Each Tier is made up of several computer centres. All the raw collision data collected by CMS is converted into a format suitable for offline analysis and sorted in the form of data sets at the Tier-0 site at CERN. This processed data is then transferred to Tier-1 centers all over the world where reconstruction algorithms are run. Further reconstructed and simulated data is distributed to Tier-2 sites, where it is available for physics analysis mainly performed on Tier-3 sites.




Figure 3.14: Schematic overview of the CMS computing grid. All data collected by CMS is stored at the Tier-0 site at CERN which is then transferred to Tier-1 centers all over the world. Further reconstructed and simulated data is distributed to Tier-2 sites, where it is available for physics analysis mainly performed on Tier-3 sites. Taken from [59].

3.3 Software Tools

Every year, the CMS is recording a huge amount of data, from collisions as well as simulations. This data is analyzed iteratively to improve the understanding of the detector and the measured physics. So a dedicated data structure and software

tools are required for data analysis. These are included in the software framework referred to as CMSSW. This section describes the CMSSW framework and software tools used in the current study.

3.3.1 CMSSW Framework

The CMS software framework (CMSSW) [60] provides all necessary tools for a physics analysis. The CMSSW framework is built on top of an event data model (EDM). It is a container for arbitrary C++ objects, e.g. recorded raw data and reconstructed physical objects (e.g particles, missing transverse energy) or derived quantities of an event. The reconstruction and distribution algorithms in CMSSW are divided into modules, which can be dynamically loaded and run. The CMSSW event processing model consists of one executable, called cmsRun, and many plug-in modules which are managed by the framework. SCRAM (Source Configuration, Release, And Management) is a configuration and management tool in the framework. It builds a runtime environment and make available all the necessary shared libraries. The shared libraries reduces memory consumption by only loading required modules during runtime. The CMSSW framework performs calibration, event generation, detector simulation, event reconstruction as well as data analysis by implementing the codes either in C++ or Python languages. To reduce the event content, a process called skimming is performed where only necessary data is preserved.

3.3.2 ROOT

ROOT [61] is an object-oriented data analysis framework, developed by CERN. ROOT consists of a huge C++ library provided with all the functionalities to store and analyze large amounts of data. It provides histogramming methods in 1, 2 and 3 dimensions, curve fitting, function evaluation, minimization, graphics and visualization classes. The command language of ROOT is command line interpreter (CINT), with several extensions to C++ which makes ROOT a versatile package.

ROOT is an open-source system which can be dynamically extended by linking external libraries. The events generated or analyzed in CMSSW framework are stored in a tree structure in files using ROOT libraries. In this thesis, ROOT has been used extensively for storing information of events or objects, for fitting as well as plotting purposes.

3.3.3 NLOJET++ and FASTNLO

The leading order (LO) as well as next-to-leading order (NLO) cross-sections for jet production are evaluated using a C++ program called NLOJET++ [62, 63]. It uses the dipole subtraction method for the separation of the divergences. NLOJET++ can calculate up to three-jet observables at NLO precision. The perturbative QCD cross-section calculations in NLOJET++ are determined in Monte Carlo integration and are very time consuming. So it is not feasible to repeatedly calculate the cross-sections which is required for PDF fits or uncertainty estimations. The NLOJET++ is interfaced to the FASTNLO project [64, 65] which performs fast re-evaluations of cross-sections. It stores the perturbative coefficients obtained with NLOJET++ in a way that the strong coupling constant and the PDFs can be changed afterwards without a recalculation of the perturbative coefficients.

All the event generators and cross-section calculation tools takes the PDFs as an input. They are either hard coded in the generators or accessed via a standardized interface with the LHAPDF library [66, 67]. LHAPDF provides a unified and easy way to use the PDF sets by storing them in data files. It provides interpolation routines to read the PDFs and interpolate the PDFs at all scales. It also allows access to single PDF members without needing to load whole sets. LHAPDF is supported by many MC event generators and other physics programs.

Chapter 4

Event Generation, Simulation and Reconstruction

Event simulation plays a significant role in the operation of any experiment. Before the real data taking, the reconstruction algorithms, efficiency of trigger paths, analysis strategies and other operational details of the experiment need to be studied and optimized well. This is achieved by simulation of the apparatus and the expected processes using the Monte Carlo (MC) method [68]. In high energy physics, the simulation of experimental data is done in two steps : event generation and detector simulation. Event generators simulate a collision starting from the proton-proton interaction up to the production of the final decay products, to be observed with the detector. The output of an event generator is taken as input in a detector simulation program which models the interactions between the generated final-state particles and the detector. This requires a sophisticated and complex simulation of the detector material and the behaviour of the interacting particles.

4.1 Event Generation and Simulation Software

In real world, the machine or collider produces interactions which are observed by detectors. The interesting events are stored and reconstructed afterwards for a

physics analysis. In the MC world, the role of machines is played by the event generators. The event generators generate simulated events as detailed as observed by a detector. The output produced of an event generator is in the form of “events” with the same behaviour and fluctuations as the real data. The detector simulation takes the output of the generator as an input and allows a precise prediction and verification of the entire experimental setup. The comparison of real and Monte Carlo world is presented in Fig. 4.1.




Figure 4.1: The comparison of Monte Carlo (MC) simulations generated by event generators with the real data produced by the collisions of observed in the detectors.

There are a variety of MC event generators which are commonly used in high energy physics. In this thesis, the three leading order (LO) generators : PYTHIA, MADGRAPH5 and HERWIG as well as the next-to-leading order (NLO) generator POWHEG are used. These generators are described one by one in the following sections.

4.1.1 PYTHIA

PYTHIA is the most widely used program to generate the collisions at high energies for p-p, e-e and e-p colliders. It contains theoretical models for a number of physics aspects which includes hard and soft processes, parton distribution functions, initial-state and final-state parton showers, multiparton interactions, fragmentation and decay. It also has a set of utilities and interfaces to external programs. It uses the Lund string hadronization model [69] to describe the hadronization process. PYTHIA was originally coded in FORTRAN language under the version 6 i.e. PYTHIA6 [70]. In 2004, it was rewritten in C++ and was released as PYTHIA8 [71] in 2007. The two versions differ in the description of multi-parton interactions. Both the versions use leading order (LO) calculations to derive the colored partons from the hard interaction. From these partons, colorless objects like hadrons, leptons and photons are produced. For the studies in this thesis, PYTHIA6 with tune Z2* [72] and PYTHIA8 with tunes CUETS1 and CUETM1 [73] have been used.

4.1.2 MADGRAPH5

MADGRAPH5 [74] generates matrix elements for high energy physics processes, such as decays and $2 \rightarrow n$ scatterings. The event information of the hard process such as particle ID, momenta, spin etc. is stored in the Les Houches format [75] and can be interfaced to other generators. In this thesis, MADGRAPH5 has been interfaced to PYTHIA6 with tune Z2* to handle the rest of the generation steps which involves parton showering and hadronization. Matching algorithms make sure that there is no double-counting between the tree-level and the PS-model-generated partons. MADGRAPH5+PYTHIA6 (MG5+P6) samples are used mainly for general comparisons to data and calculating the detector resolution.

4.1.3 HERWIG

HERWIG (Hadron Emission Reactions With Interfering Gluons) [76] is a multi-purpose event generator. It includes the simulation of hard lepton-lepton, lepton-hadron and hadron-hadron scattering and soft hadron-hadron collisions. It uses angular ordering for parton showers and cluster model for hadronization. The HERWIG generator includes a number of hard scattering processes and has the possibility to interface external matrix element generators. It uses angular ordering for parton showers and cluster model for hadronization. HERWIG was FORTRAN based and has a version in C++, named HERWIG++ [77]. The samples generated using HERWIG++ generator with the default tune of version 2.3 [78] have been used to study non-perturbative effects.

4.1.4 POWHEG

POWHEG generator performs the fixed next-to-leading order (NLO) calculations merged with parton showers [79–81]. A computer framework known as POWHEG BOX [82], implements NLO calculations in shower Monte Carlo programs according to the POWHEG method. It can be interfaced with all modern shower Monte Carlo programs that support the Les Houches Interface for User Generated Processes. It contains the hard matrix elements for NLO dijet production. For the parton shower and hadronization, POWHEG is interfaced to PYTHIA8 with tunes CUETS1 and CUETM1.

4.2 Detector Simulation

The particles generated by Monte Carlo event generators are passed through the detector simulation. It is a computer program which defines the detector system. The detector definition includes the representation of its geometrical elements, their materials and electronics properties. The geometrical representation of detector el-

ements focuses on the definition of solid models and their spatial position. The detector simulation describes the interactions of the passing particles with the material of the detector. While propagating through the detector material, these particles are allowed to decay according to their known branching fractions and decay kinematics. The interactions of the particles with the detector material take place through several physical processes, including electron bremsstrahlung, energy loss by ionization, multiple scattering, hadron showering etc., which are simulated or parametrized in the corresponding parts of the detector.

In CMS, the detector response is simulated by two approaches [50] : Full Simulation and Fast Simulation. Full Simulation is based on a C++ simulation toolkit GEANT4 (GEometry AND Tracking) [83]. It is a successor of a FORTRAN based GEANT3 and handles the interactions of particles with matter over a wide range of energy. In GEANT4, the magnetic fields, electric fields and electromagnetic uniform and non-uniform fields can be specified. The equation of motion of the particle in the field gives the track of the particle. A physical interaction of a track in the sensitive region of a detector is called a hit. The secondary particles produced are stored in a stack with the information of their kinematic properties as well as the vertex position where the interaction has occurred. A large number of Monte Carlo events may have to be produced for a feasible physics analysis. The complete detector simulation of CMS using GEANT4 is rather time consuming. For the fast simulation of the detector response, a Fast Simulation framework [84] has been developed in the general software framework of the CMS. In Fast Simulation, detector effects are parametrized instead of simulating these from first principles as done in Full Simulation. Fast Simulation package produces events at rates of the order of 100 times faster than the corresponding Full Simulation ones, while maintaining almost the same level of accuracy for physics studies. The format of the Fast Simulation data output is fully compatible with the standard Full Simulation one.

After simulating the detector response, it is then transformed into a digital

signal with the help of electronics and this step is called digitization. The simulated output of the detector response needs to be as close as possible to the real data coming from the CMS detector. After this, event reconstruction algorithms are applied to both simulated and real events.

4.3 Event Reconstruction

The aim of the event reconstruction is to identify the particles passing through the detector by interpreting the electrical signals produced in digitization. These particles are produced either directly from the interaction point of pp collisions or from the hadronization process. In event reconstruction, analysis-level objects are created by combining recorded signals from the tracker, calorimeters and muon detectors. At initial level, the reconstructed hits are collected which are combined to form tracks and calorimetric towers. Then higher level objects such as electrons, photons, muons and jets are reconstructed by combining the tracks and energy deposits. This thesis presents the study of jets. The hadrons and other particles produced by the hadronization of a quark or gluon are clustered into a narrow cone called “jets”. The detailed description of jets is given in Sec. 2.4. The jet algorithms, discussed in Sec. 2.4.1, are used for clustering by taking partons, stable particles or reconstructed particle candidates as inputs. The different levels at which the jets are formed are parton level, particle level and detector or reconstructed level, as illustrated in Fig. 4.2. In the CMS detector, jets are the localized deposits of energy in the calorimeter cells along with the large number of tracks in the direction of the deposited energy. Depending on the type of input to the jet algorithm, jets can be categorized into different types :

Generator Jets - The stable particles generated by the Monte Carlo event generators are clustered into generator jets (GenJets). At this particle level, the passage through the detector simulation has not been carried out. The objects at this level




Figure 4.2: In a proton-proton collision, the hard scattered quarks and gluons fragment and hadronize to produce the showers of partons, hadrons, or detector measurements which are clustered into parton jets, particle jets and reconstructed jets, respectively. Taken from [85].

are charged hadrons, photons and neutral hadrons. Since the energy of GenJets is independent of the detector response, these are considered as reference objects for jet energy corrections.

Calorimetric Jets - The jets reconstructed using the energy clusters deposited in the ECAL and HCAL calorimeter towers are called Calorimetric jets (CaloJets). One calorimetric tower consists of one HCAL cell surrounded by an array of $5 \times$

5 ECAL cells. The tower's four-momenta are computed taking the direction from the interaction point to the tower center and assuming zero mass. All towers with a transverse-energy measurement above 300 MeV are considered in the clustering process. CaloJets are relatively simple objects because only calorimeter information is deployed, but they are strongly affected by the non-linearity of the calorimeter response. Since the readout of calorimeter measurements is fast, CaloJets are commonly used by the trigger system.

Particle Flow Jets - The clustering of particle flow candidates give detector level jets called Particle Flow jets (PFJets). The four-momenta of the particles is taken as the input. The use of the tracking detectors and high granularity of the ECAL improves the energy resolution through the independent measurements of charged hadrons and photons inside a jet. Hence PFJets perform better than CaloJets and are the standard jets used at CMS.

The study presented in this thesis uses the jets clustered using the anti- k_t algorithm with a jet size parameter of $R = 0.7$ and particle flow candidates. In CMS, all the particles are reconstructed and identified using a Particle Flow (PF) algorithm, discussed in details in next section.

4.3.1 Particle Flow Algorithm

To identify and reconstruct the particles, the CMS employs the event reconstruction technique called Particle Flow (PF) algorithm [86,87]. Basically, the detector signals are converted back to physical objects by using PF event reconstruction algorithm, as shown in Fig. 4.3. It combines the information from the individual sub-detectors. The additional identification of the tracks, using the Combinatorial Track Finder (CTF) algorithm [88], enhances the reconstruction performance. Based on these tracks, the primary vertices in an event are identified. The transverse momenta of final state stable particles or energies of the calorimeter towers are the inputs to PF

algorithm. The PF algorithm first collects reconstructed hits in each sub-detector independently and creates a list of reconstructed elements (referred as blocks) : charged tracks in tracker, energy clusters in calorimeters and muon tracks in muon system. Then a link algorithm connects topologically compatible blocks producing PF objects. The PF objects consists of all stable particles : electrons, muons, photons, charged and neutral hadrons. The combination of the track momentum at the main interaction vertex, the corresponding ECAL energy deposit and the energy sum of all bremsstrahlung photons associated with the track is used to determine the energy of electrons. The curvature of the tracks in tracker and muon chamber




Figure 4.3: Schematic association of sub-detector measurements to physical particle objects using the Particle Flow (PF) event reconstruction technique used by CMS. Taken from [15].

is used to estimate the energy of muons. The energy of photons is obtained directly from the ECAL measurement, corrected for zero-suppression effects¹⁵. The energy of charged hadrons is calculated by combining the track momentum and corresponding energy clusters in ECAL and HCAL, corrected for zero-suppression effects as well as calibrated for the nonlinear response of the calorimeters. The energy of neutral hadrons is obtained from the corresponding calibrated ECAL and HCAL energies only. Along with the reconstruction of these objects, missing transverse energy

¹⁵To suppress noise in the calorimeters, only cells with energies above a given threshold are considered and this procedure is known as zero-suppression.

(E_T^{miss}) is also determined using PF algorithm. E_T^{miss} is defined as the negative vector sum of transverse momenta p_T of all the isolated stable particles reconstructed in an event i.e. $E_T^{\text{miss}} = -\sum_i \vec{p}_{T,i}$. To avoid any kind of double-counting of energy, blocks of all PF reconstructed particle objects are removed and the energy of the calorimeter clusters is recalculated. Finally, the collection of PF objects is used to reconstruct the jets by using jet clustering algorithms.

4.3.2 Jet Energy Corrections

The measured energy of jets cannot be directly translated to the true particle or parton level. This is because of the nonlinear and nonuniform response of the calorimeters, effects of pileup and small residual effects in the data after corrections based on MC simulation. Hence to relate the measured jet energy to the corresponding true particle jet energy, jet energy corrections (JEC) [89, 90] are used to correct the measured jet energy. CMS follows a factorized approach, as presented in Fig. 4.4, where JEC are applied in a sequential manner with fixed order, i.e. the output of one step serves as the input for the next one. Each level of correction takes care of a different effect and is independent of each other. At each step, the jet four momentum is scaled with a correction factor which depends on jet p_T , η , flavor etc.




Figure 4.4: A schematic diagram of the factorized jet energy corrections (JEC) for data (upper half) and simulation (lower half) [90]. The reconstructed jets are corrected for pileup effects, non-uniform p_T and η response and residual differences between data and Monte Carlo simulations along with optional flavor corrections. All corrections marked with MC are derived from simulation studies, RC stands for random cone, and MJB refers to the analysis of multijet events.

The corrected jet transverse momentum p_T^{corr} is obtained by applying all correction factors subsequently on raw or uncorrected jet transverse momentum p_T^{raw} as below :

$$p_T^{\text{corr}} = c_{\text{res}}(\eta, p_T'') \cdot c_{\text{mc}}(\eta, p_T') \cdot c_{\text{pileup}}(\eta, \rho, A_j, p_T^{\text{raw}}) \cdot p_T^{\text{raw}} \quad (4.1)$$

where p_T' is the transverse momentum after applying the pileup correction factor c_{pileup} on p_T^{raw} , p_T'' is the transverse momentum after applying the additional correction factor c_{mc} because of relative and absolute effects derived from MC. At last a correction factor c_{res} for residual effects derived from data is applied. The corrections applied at each step are discussed as :

Pileup Corrections - The additional proton-proton collisions occurring within the same bunch-crossing produce particles which got clustered into the jets coming from the hard interaction. This extra energy must be subtracted from the reconstructed jet energy. This is done by applying the pileup corrections to raw jet p_T^{raw} . The pileup corrections are determined from the simulation of a sample of QCD dijet events processed with and without pileup effects. The pileup correction factor, c_{pileup} is calculated using jet area method from the pileup density ρ in the event and the jet area A_j and is parametrized as a function of ρ , A_j , jet p_T and η . The corrections for residual differences between data and detector simulation as a function of eta are determined using the random cone (RC) method in zero-bias events. Hence the different pileup corrections are applied to data and MC.

MC Corrections - The next correction applied to the pile up corrected jets is based on simulated QCD events. There are some differences between the reconstructed and generated jet p_T because of inefficiencies introduced by the detector simulation. The correction factor, c_{mc} is derived by comparing the measured jet p_T to the particle level jet p_T . The corrections are determined as a function of jet p_T and η which make the detector response uniform over these two variables.

Residual Data Corrections - The jets corrected with above mentioned corrections are further corrected for remaining small differences between data and Monte Carlo simulations. This correction is only applied to data. The correction factor c_{res} is derived using data-driven methods. The relative residual corrections are based on well-balanced dijet events in which a forward probe jet is calibrated using a tag jet in the well understood barrel region. The last correction step is the absolute residual correction in which reconstructed Z bosons balanced to a jet are used to calibrate the jet energy using the very precisely reconstructed Z boson.

Flavor Corrections - These corrections correct the jets for flavor dependence (b , τ etc.) and are optional. These are extracted using Z +jet and photon+jets simulated events. The flavor corrections have not been applied for 8 TeV data.

The process of correction of jets by using JEC introduces uncertainties in the final corrected jet energy which are discussed in Sec. 5.6.2.

Chapter 5

Measurement of the Differential Inclusive Multijet Cross-sections and their Ratio

In a proton-proton collision, the inclusive jet cross-section studied as a function of jet properties, provides essential information about the parton distribution functions of proton and the strong coupling constant. This chapter describes the measurement of inclusive differential jet event cross-sections and the cross-section ratio in details which includes event and jet selections, trigger studies, spectrum construction, corrections applied and calculation of experimental uncertainties.

The inclusive differential jet event cross-sections are studied as a function of the average transverse momentum, $H_{T,2}/2 = \frac{1}{2}(p_{T,1} + p_{T,2})$, where $p_{T,1}$ and $p_{T,2}$ denote the transverse momenta of the two leading jets, and are defined by :

$$\frac{d\sigma}{d(H_{T,2}/2)} = \frac{1}{\epsilon \mathcal{L}_{int,eff}} \frac{N_{event}}{\Delta(H_{T,2}/2)} \quad (5.1)$$

where N_{event} is the number of inclusive n-jet events counted in an $H_{T,2}/2$ bin, ϵ is the product of the trigger and jet selection efficiencies, which are greater than 99%,

$\mathcal{L}_{\text{int,eff}}$ is the effective integrated luminosity, and $\Delta(H_{\text{T},2}/2)$ are the bin widths. The measurements are reported in units of (pb/GeV). The inclusive n-jet event samples include the events with number of jets $\geq n$, where $n = 2$ and 3 in the current study.

The cross-section ratio R_{32} , defined in Eq. 5.2 is obtained by dividing the differential cross-sections of inclusive 3-jet events to that of inclusive 2-jet one, for each bin in $H_{\text{T},2}/2$.

$$R_{32} = \frac{\frac{d\sigma_{3\text{-jet}}}{d(H_{\text{T},2}/2)}}{\frac{d\sigma_{2\text{-jet}}}{d(H_{\text{T},2}/2)}} \quad (5.2)$$

For inclusive 2-jet events ($n_j \geq 2$) sufficient data are available up to $H_{\text{T},2}/2 = 2000$ GeV, while for inclusive 3-jet events ($n_j \geq 3$) and the ratio R_{32} , the accessible range in $H_{\text{T},2}/2$ is limited to $H_{\text{T},2}/2 = 1680$ GeV.

5.1 Data Samples

This measurement uses the data which was collected at the centre-of-mass energy of 8 TeV by CMS experiment in the 2012 run period of the LHC. The 2012 data is taken in four periods A, B, C, D and the data sets are divided into samples according to the run period. Further each sample is grouped into subsets based on the trigger decision. For run B-D, the **JetMon** stream data sets contain prescaled low trigger threshold paths (HLTPFJet40, 80, 140, 200 and 260) while the **JetHT** stream data sets contain unprescaled high threshold trigger paths (HLT PFJet320 and 400). For run A, the **Jet** stream contains all the above mentioned trigger paths. The data to be used in physics analysis must satisfy a certain criteria according to which it should fulfill the validation requirements of data quality monitoring procedure. CMS uses JSON (Java Script Object Notation) format files to store the range of good lumi

sections within a run. In the current analysis, the applied certification file¹⁶ is based on the final event reconstruction of the 2012 CMS data sets. The data sets used in the current study are mentioned in the Table 5.1 along with the luminosity of each data set which increases with period. Full 2012 data sample corresponds to an integrated luminosity of 19.71 fb^{-1} .

Table 5.1: Run range and luminosity of the proton-proton collisions data collected at the centre-of-mass energy of 8 TeV by CMS experiment in the year of 2012 in four different run periods A, B, C and D.

Run	Run range	Data set	Luminosity fb^{-1}
A	190456-193621	/Jet/Run2012A-22Jan2013-v1/AOD	0.88
B	193834-196531	/Jet[Mon,HT]/Run2012B-22Jan2013-v1/AOD	4.41
C	198022-203742	/Jet[Mon,HT]/Run2012C-22Jan2013-v1/AOD	7.06
D	203777-208686	/Jet[Mon,HT]/Run2012D-22Jan2013-v1/AOD	7.37

5.1.1 Monte Carlo Samples

To have a comparison of data results with the simulated events, the MADGRAPH5 Monte-Carlo (MC) event generator has been used. It has been interfaced to PYTHIA6 by the LHE event record, to generate the rest of the higher-order effects using the Parton Showering (PS) model, with tune Z2* to model the underlying event. The MC samples are processed through the complete CMS detector simulation to allow studies of the detector response and compare to measured data on detector level.

The cross-section measured as a function of the transverse momentum p_T or the scalar sum of the transverse momentum of all jets H_T falls steeply with the increasing p_T . So in the reasonable time, it is not possible to generate a large number of high p_T events. Hence, the events are generated in the different phase-space region binned in H_T or the leading jet p_T . Later on, the different phase-space regions are added together in the data analyses by taking into account the cross-section of the

¹⁶Cert_190456-208686_8TeV_22Jan2013ReReco_Collisions12_JSON

different phase-space regions. The official CMS MADGRAPH5+PYTHIA6 (MG5+P6) MC samples used in this analysis are generated as slices in the H_T phase-space are tabulated in Table 5.2 along with their cross-sections and number of events generated.

Table 5.2: The official Monte Carlo samples are produced in phase space slices in H_T with the generator MADGRAPH5 and interfaced to PYTHIA6 for the parton shower and motorization of the events. The cross-section and number of events generated are mentioned for each sample.

Generator	Sample	Events	Cross-section pb
MADGRAPH5 + PYTHIA6	/QCD_HT-100To250_TuneZ2star_8TeV-madgraph-pythia6/ Summer12_DR53X-PU_S10_START53_V7A-v1/AODSIM	50129518	1.036×10^7
	/QCD_HT-250To500_TuneZ2star_8TeV-madgraph-pythia6/ Summer12_DR53X-PU_S10_START53_V7A-v1/AODSIM	27062078	2.760×10^5
	/QCD_HT-500To1000_TuneZ2star_8TeV-madgraph-pythia6/ Summer12_DR53X-PU_S10_START53_V7A-v1/AODSIM	30599292	8.426×10^3
	/QCD_HT-1000ToInf_TuneZ2star_8TeV-madgraph-pythia6/ Summer12_DR53X-PU_S10_START53_V7A-v1/AODSIM	13843863	2.040×10^2

5.2 Event Selection

The events are selected according to several quality criteria which ensures the high purity and high selection efficiency of the sample to be studied. This event selection also reduces beam induced background, detector-level noise and jets arising from fake calorimeter energy deposits.

5.2.1 Trigger Selection

CMS implements a two-level trigger system to reduce the amount of recorded events to a sustainable rate. In this analysis the jets are the final objects to study. So single jet trigger paths with varying thresholds are used to select events in data. It consists of one L1 trigger seed and multiple HLT filters. The L1 jet trigger uses transverse energy sums computed using both HCAL and ECAL in the central region ($|\eta| < 3.0$) or HF in the forward region ($|\eta| > 3.0$). The single jet triggers (HLT_PFJetX), same

as the ones used for other CMS 8 TeV measurements [23, 91], are used in the current study and are tabulated in Table 5.3. A single jet trigger selects an event in which at least one jet has the transverse momentum above the threshold. HLT_PFJetX implies that there is at-least one jet in the event, whose $p_T > X$ (GeV). The L1 trigger has a lower threshold to ensure full efficiency versus p_T of the HLT trigger. The p_T spectrum is steeply falling and hence the rates for low- p_T jets are very high. So it is not feasible to use a single unprescaled trigger for the selection of all required events. To collect sufficient data in the lower part of the p_T spectrum, different five prescaled low- p_T trigger paths, each with different prescale value, are used. Also, one unprescaled trigger i.e. HLT_Jet320 is used in the high p_T region, in which the rate is sufficiently small to collect and store all events. During the reconstruction of the spectrum, the prescales have been taken into the account.

Table 5.3: The single jet HLT trigger paths used in the analysis are listed here. The column $H_{T,2}/2$, 99% indicates the value of $H_{T,2}/2$ at which each trigger exhibits an efficiency larger than 99%. The last column gives the effective luminosity seen by each trigger which divided by the total integrated luminosity of 19.71 fb^{-1} , gives the effective prescale applied on a trigger over the whole run period.

Trigger Path	L1 threshold GeV	HLT threshold GeV	$H_{T,2}/2$, 99% GeV	Eff. Lumi fb^{-1}
HLT_PFJet80	36	80	120.0	0.0021
HLT_PFJet140	68	140	187.5	0.056
HLT_PFJet200	92	200	262.5	0.26
HLT_PFJet260	128	260	345.0	1.06
HLT_PFJet320	128	320	405.0	19.71

The efficiency of each trigger path as a function of $H_{T,2}/2$ is described by the turn-on curves with a rising part where the trigger is partly inefficient, until a plateau region where the trigger is fully efficient. Hence it is important to determine the threshold above which a trigger becomes fully efficient. The threshold is the value at which the trigger efficiency exceeds 99%. The trigger efficiency for HLT_PFJetY is given by Eq. 5.3 where HLT_PFJetX is the reference trigger and is assumed to be fully efficient in the considered phase space region. The value of X is chosen

previous to that of Y in p_T ordering from the trigger list so that the higher trigger condition can be emulated from the lower trigger path.

$$\epsilon_{\text{HLT_PFJetY}} = \frac{H_{T,2}/2 \left(\text{HLT_PFJetX} + (\text{L1Object_}p_T > Z) + (\text{HLTOBJECT_}p_T > Y) \right)}{H_{T,2}/2(\text{HLT_PFJetX})} \quad (5.3)$$

where Y is the p_T threshold of HLT_PFJetY and Z is the L1 seed value corresponding to the trigger path HLT_PFJetY. The denominator represents the number of events for which the reference trigger path HLT_PFJetX has been fired. The numerator is the number of events for which HLT_PFJetX has been fired along the p_T of L1Object \geq Z and the p_T of HLTOBJECT \geq Y. For example, to obtain turn-on curve for HLT_PFJet260, HLT_PFJet200 is the reference HLT path. The p_T cut on L1Object is 128 GeV and p_T cut on HLTOBJECT is 260 GeV. The threshold point at which the trigger efficiency is larger than 99% is determined by fitting the turn-on distribution with a sigmoid function described in Eq. 5.4. The trigger turn-on curves as a function of $H_{T,2}/2$ can be seen in Fig. 5.1 which are described by a sigmoid function (blue line). The error bars give the uncertainty on the efficiency which is calculated using Clopper-Pearson confidence intervals [92].

$$f_{fit}(x) = \frac{1}{2} \left(1 + \text{erf} \left(\frac{x - \mu}{\sqrt{2}\sigma} \right) \right) \quad (5.4)$$

5.2.2 Primary Vertex Selection

The reconstructed tracks, number of strip and pixel hits and the normalized track χ^2 , identify the primary vertex (PV). The tracks are clustered according to the z-coordinate of their point of closest approach to the beam axis. A selection criteria for primary vertex should be followed which helps to identify and reject the beam




Figure 5.1: Trigger efficiencies turn-on curves for the single jet HLT trigger paths are fitted with a sigmoid function (blue line) to obtain the 99% efficiency threshold. The error bars give the uncertainty on the efficiency which is calculated using Clopper-Pearson confidence intervals [92].

background events. At-least one good primary vertex reconstructed from at least four tracks within a distance of $|z(PV)| < 24$ cm to the nominal interaction point in a collision, is required in each event. The radial distance in x-y plane, $\rho(PV)$ should be not be greater than 2 cm. The number of degrees of freedom in fitting for the position of each vertex using its associated tracks should be at-least four in number.

5.2.3 Missing Transverse Energy

In an ideal detector where all particles could be identified and perfectly measured, the transverse momentum of all particles would sum up to zero. But the neutral weakly interacting particles, such as neutrinos, escape from typical collider detectors and do not produce any direct response in the detector elements. The imbalance of total momentum of all visible particles can give the hints of the presence of such particles. The vector momentum imbalance in the plane perpendicular to the beam direction is known as missing transverse momentum or energy (E_T^{miss}). It is one of the most important observables for discriminating leptonic decays of W bosons and top quarks from background events which do not contain neutrinos, such as multijet and Drell–Yan events or searches for physics beyond the Standard Model.

The ratio of missing transverse energy to the total transverse energy $E_T^{\text{miss}}/\sum E_T$, shown in Fig. 5.2 for $n_j \geq 2$ (left) and $n_j \geq 3$ events (right), shows a discrepancy between data (black solid circles) and simulated MC (blue histogram), at the tail part of the distribution. This is because of a finite contribution from $Z(\rightarrow \nu\bar{\nu}) + \text{jet}$ events which gives rise to non-zero E_T in the events in data. Such events are absent in QCD simulated events in MC. Hence $E_T^{\text{miss}}/\sum E_T$ is required to be less than 0.3 to reject events with high E_T^{miss} .




Figure 5.2: Missing transverse energy fraction of the total transverse energy per event in data (black solid circles) and simulated Monte Carlo events (blue histogram) in inclusive 2-jet (left) and 3-jet events (right). To remove background and noise, events with a fraction exceeding a certain threshold, here indicated with the red dashed line, are rejected.

5.2.4 Jet Identification

In order to suppress fake jets, arising from detector noise or misreconstructed particles, jet identification criteria (ID) has been applied. Instead of applying it event-wise, it is applied it on each jet. The algorithm works on reconstructed jets using information of the clustered particle candidates. The official tight jet ID [93], recommended by JETMET group [94] is used. Due to pileup and electronic noise the jet constituent fractions may vary from event to event. In order to reject the noisy jets, some jet selection criteria are optimized to select only good quality jets. The selection criteria are implemented as selection cut on jet fractions. Table 5.4 summarizes the properties of the reconstructed jets and their respective cuts. Each jet should contain at least two particles, one of which should be a charged hadron. The cut on the fraction of neutral hadrons and photons removes HCAL noise and ECAL noise, respectively. Muons that are falsely identified and clustered as jets are removed by the muon fraction criterion. Based on information of the tracker, additional selection cuts are enforced in the region $|\eta| < 2.4$. The charged electromagnetic fraction cut removes the jets clustered from misidentified electrons. Furthermore, the fraction of charged hadrons in the jet must be larger than zero and jets without any charged

hadrons are very likely to be pileup jets. The Figs. 5.3 and 5.4 show the distributions of the jet constituents observed in data (black solid circles) and simulated MC events (blue histogram) for $n_j \geq 2$ and $n_j \geq 3$, respectively.

Table 5.4: The jet ID removes noise and fake jets based on the properties of the reconstructed jets and the clustered particle candidates. All the selection cuts which are recommended by the JETMET group are applied [94].

	Property	Loose ID	Tight ID
Whole η region	neutral hadron fraction	< 0.99	< 0.90
	neutral EM fraction	< 0.99	< 0.90
	number of constituents	> 1	> 1
	muon fraction	< 0.80	< 0.80
only $ \eta < 2.4$	charged hadron fraction	> 0	> 0
	charged multiplicity	> 0	> 0
	charged EM fraction	< 0.99	< 0.90

5.2.4.1 Jet ID Efficiency

The efficiency of the jet ID as a function of $H_{T,2}/2$ is studied using a tag-and-probe technique with dijet events. The two leading jets are required to be back-to-back in the azimuthal plane such that $|\Delta\phi - \pi| < 0.3$. One of the dijets is selected randomly as a “tag” jet which is required to fulfill the tight jet ID criteria. The other jet is called “probe” jet for which it is examined, whether it also passes the tight jet ID. The ID efficiency is defined as the ratio of events where the probe jet passes the ID requirements, over the total number of dijet events. It is shown as function of $H_{T,2}/2$ in Fig. 5.5 and as expected, it is always greater than 99%. The QCD cross-section decreases as a function of $H_{T,2}/2$ and hence the number of events decrease on moving to higher $H_{T,2}/2$. Consequently the statistical fluctuations for ID efficiency are larger at higher $H_{T,2}/2$.




Figure 5.3: The fractions of jet constituents as observed in data (black solid circles) and simulated Monte Carlo events (blue histogram) for different types of PF candidates for inclusive 2-jet events. Data and simulations are normalized to the same number of events. The distributions are shown after the application of the jet ID.




Figure 5.4: The fractions of jet constituents as observed in data (black solid circles) and simulated Monte Carlo events (blue histogram) for different types of PF candidates for inclusive 3-jet events. Data and simulations are normalized to the same number of events. The distributions are shown after the application of the jet ID.




Figure 5.5: The jet ID efficiency is studied as a function of $H_{T,2}/2$ with tag-and-probe technique using dijet event topologies and it always exceeds 99%.

5.2.5 Jet Selection

The measurement of differential cross-sections and their ratio uses jets clustered from particle flow candidates using the anti- k_t jet algorithm with a size parameter, $R = 0.7$. The energy scale of the jets is corrected with the CMS recommended jet energy corrections, described in Sec. 4.3.2. These corrections are applied to jets in both data¹⁷ as well those in simulated events¹⁸. As a convention, the jets in one event are in decreasing order of p_T , with the first (leading) jet being the jet with highest p_T . The jet selection, based on phase space cuts on transverse momentum and rapidity of jets in an event, is as follows :

- All jets having $p_T > 150$ GeV and $|y| < 5.0$ are selected.
- Events with at least two jets are selected.
- The two leading jets should have $|y| < 2.5$ and further jets are counted only, if they lie within the same central rapidity range of $|y| < 2.5$.

These cuts assure high detector acceptance and exactly same selection is ap-

¹⁷Winter14_V8 jet energy corrections

¹⁸START53_V27 jet energy corrections

plied in the measurement, simulated events as well in theoretical calculations for a consistent comparison.

5.3 Comparison with Simulation

5.3.1 Pileup Reweighting

While generating the official Monte-Carlo samples, the number of pileup interactions describing the conditions expected for each data-taking period are taken care of. But the number of pileup events implemented in the simulation $N_{\text{MC}}(N_{\text{PU,truth}})$, does not match exactly with the one measured actually in data $N_{\text{data}}(N_{\text{PU,est.}})$. To match the pileup distributions in data, a reweighting factor w_{PU} , as given by Eq. 5.5 is applied to the simulated events. In Fig. 5.6 the number of reconstructed vertices are shown before (left) and after pileup reweighting (right). It is observed that before pileup reweighting there was a significant mismatch of the pileup distributions in data (black solid circles) and simulated MC events (blue histogram), which completely vanishes after reweighting.

$$w_{\text{PU}} = \frac{N_{\text{data}}(N_{\text{PU,est.}}) / \sum N_{\text{data}}}{N_{\text{MC}}(N_{\text{PU,truth}}) / \sum N_{\text{MC}}} \quad (5.5)$$

5.3.2 Comparison of Cross-sections and their Ratio

The measured data distribution of differential cross-section at detector level is compared to the predictions of Monte Carlo simulation using **MADGRAPH5** generator interfaced with **PYTHIA6** (MG5+P6) including the detector simulation as well as to a fixed-order theory prediction obtained using CT10-NLO PDF set. Figure 5.7 shows the comparison of differential cross-section as a function of $H_{\text{T},2}/2$ for $n_j \geq 2$ (left) and $n_j \geq 3$ events (right), for data (black solid circles), MG5+P6 MC (red empty




Figure 5.6: Number of reconstructed vertices in data (black solid circles) and simulated Monte Carlo events (blue histogram) before (left) and after (right) the pileup reweighting.

circles) and CT10-NLO (blue histogram). The bottom panel in each plot shows the ratio of data to the MC predictions (red line) as well as to the CT10-NLO theory predictions (blue line). The NLO predictions on parton level are not corrected for non-perturbative effects. Still the NLO predictions describe the data better as compared to the LO MC simulations which roughly describes the spectrum on detector level. The sufficient data for $n_j \geq 2$ and $n_j \geq 3$ events are available up to $H_{T,2}/2 = 2000$ GeV and 1680 GeV, respectively. Due to some kinematical constraints, the minimum cut on $H_{T,2}/2$ is 300 GeV (explained in Sec. 6.1.1). Hence the differential cross-sections are studied in the range $300 \text{ GeV} \leq H_{T,2}/2 < 2000 \text{ GeV}$ for $n_j \geq 2$ and $300 \text{ GeV} \leq H_{T,2}/2 < 1680 \text{ GeV}$ for $n_j \geq 3$ events.

The ratio of differential cross-sections, R_{32} as a function of $H_{T,2}/2$, is extracted by dividing the cross-section of selected inclusive 3-jet events to that of inclusive 2-jet events at any given bin size of $H_{T,2}/2$. In the cross-section ratios, the numerator and denominator are not independent samples. So to calculate the statistical uncertainty for the cross-section ratios at reconstructed level, the Wilson score interval method is used which takes into account the correlation between the




Figure 5.7: The reconstructed level differential cross-sections are compared for data (black solid circles) and LO MADGRAPH5+PYTHIA6 (MG5+P6) Monte Carlo (red empty circles) simulations with CT10-NLO theory predictions (blue histogram), as a function of $H_{T,2}/2$ for inclusive 2-jet (left) and 3-jet events (right). Ratios of data to the Monte Carlo predictions (red line) as well as to the CT10-NLO predictions (blue line) are shown in bottom panel of each plot.




Figure 5.8: Comparison of the reconstructed level cross-section ratio R_{32} as a function of $H_{T,2}/2$, for data (black solid circles) and LO MADGRAPH5+PYTHIA6 (MG5+P6) Monte Carlo (red empty circles) with CT10-NLO theory predictions (blue line). The error bars give the asymmetrical statistical uncertainty, calculated by the Wilson score interval method which takes into the account the correlation between the numerator and denominator.

numerator and the denominator and give asymmetric errors. Figure 5.8 shows the comparison of the cross-section ratio R_{32} as a function of $H_{T,2}/2$, for data (black solid circles) and LO MADGRAPH5+PYTHIA6 (MG5+P6) Monte Carlo (red empty

circles), at reconstructed level with CT10-NLO theory predictions (blue line). Since in $n_j \geq 3$ events, the enough statistics for differential cross-section is available up to 1680 GeV of $H_{T,2}/2$ only, R_{32} is also studied in the range $300 \text{ GeV} \leq H_{T,2}/2 < 1680 \text{ GeV}$. The bin-wise inclusive 2-jet and 3-jet events differential cross-sections as well as their ratio R_{32} , calculated at detector level, along with statistical uncertainty (in %) are tabulated in Table A.1.

5.4 Jet Energy Resolution (JER)

In an ideal experiment, the value of a physical quantity would be determined exactly with an infinite precision. For e.g. whenever a particle with energy E passes an ideal calorimeter having infinite resolution, the measured energy should always be equal to E . But in real world, the measured energy of the above mentioned particle might differ from the value E . This difference of the measured quantity from its true value may be due to detector noise, uncertainties in the calibration, non-linearity of the response etc. Hence this results in the finite value of the resolution of the detector known as jet energy resolution (JER). In such case, the measured values of energy of different particles, passing through the same detector with same energy E , will be different. Such measurements are described by a Gaussian distribution which is centered around the true value of the measured quantity and its width is generally interpreted as detector resolution. Hence the importance of the detector resolution lies in the fact that it indicates how much the measured value of the observable differs from the true one i.e. how precisely a physical observable can be measured. The narrower the distribution, the higher the resolution is and hence the more efficient is the detector.

Due to finite resolution of the CMS detector, the measured transverse momentum of jets gets smeared. Since the observable in this study i.e. $H_{T,2}/2$ is the average sum of transverse momentum of leading and sub-leading jets, the resolution

of the detector has to be studied in terms of the observable. CMS detector simulation based on MG5+P6 MC event generators is used to determine the resolution as both the particle and reconstructed level information is available. The jets clustered from stable generator particles called Gen jets as well as from particle flow candidates reconstructed from the simulated detector output called Reco jets, are used. The studies of the JETMET working group at CMS has shown that the jet energy resolution in data is actually worse than in simulation [95]. So the reconstructed jet transverse momentum needs to be smeared additionally to match the resolution in data. Table 5.5 shows the scaling factors (c) which need to be applied on the transverse momentum of simulated reconstructed jets. The scaling factors depend on the absolute η of the jet. The uncertainty on these measured scaling factors ($c_{central}$) needs to be taken into account in a physics analysis. This is done by smearing the reconstructed jets with two additional sets of scaling factors, c_{up} and c_{down} , that correspond to varying the factors up and down respectively, by one sigma and evaluating the impact of these new sets.

Table 5.5: JETMET working group at CMS has shown that the jet energy resolution in data is actually worse than in simulation [95]. To match the resolution in data, the reconstructed jet transverse momentum in simulated events need to be smeared by applying the scale factors. The uncertainty on the resolution is given by an upwards and downwards variation c_{up} and c_{down} of the measured scaling factor $c_{central}$.

η	0.0 - 0.5	0.5 - 1.1	1.1 - 1.7	1.7 - 2.3	2.3 - 2.8
$c_{central}$	1.079	1.099	1.121	1.208	1.254
c_{down}	1.053	1.071	1.092	1.162	1.192
c_{up}	1.105	1.127	1.150	1.254	1.316

The reconstructed jet p_T is smeared randomly using a Gaussian width widened by the scaling factor ($c_{central}$)

$$p_T \rightarrow Gauss \left(\mu = p_T, \sigma = \sqrt{c_{central}^2 - 1} \cdot JER(p_T) \right) \quad (5.6)$$

where $\text{JER}(p_T)$ is the resolution determined as a function of jet p_T using MG5+P6 MC simulated events. After smearing transverse momentum of each reco jet, $H_{T,2}/2$ is calculated from both generator particle jets (Gen $H_{T,2}/2$) as well as the particle flow or reconstructed jets (Reco $H_{T,2}/2$). Then the response is calculated as defined in the Eq. 5.7.

$$R = \frac{\text{Reco } H_{T,2}/2}{\text{Gen } H_{T,2}/2} \quad (5.7)$$




Figure 5.9: Fitting of the jet energy resolution distribution, obtained using LO MADGRAPH5+PYTHIA6 (MG5+P6) Monte Carlo simulated events, as a function of $H_{T,2}/2$ for inclusive 2-jet (left) and 3-jet events (right). The blue line shows the double-sided Crystal Ball function fit of $\frac{\text{Reco } H_{T,2}/2}{\text{Gen } H_{T,2}/2}$ in each Gen $H_{T,2}/2$ bin, overlayed by Gaussian fitting the core of the resolution (red line).

The width of the response distribution in a given Gen $H_{T,2}/2$ bin is interpreted as the resolution which in good approximation can be described by the σ of a Gaussian fit of the response distribution. A double-sided Crystal-Ball function takes into account the non-Gaussian tails of the jet response distribution. The resolution as a function of $H_{T,2}/2$ is calculated separately for both $n_j \geq 2$ and $n_j \geq 3$ events. A fit example for one Gen $H_{T,2}/2$ bin is shown in Fig. 5.9 for $n_j \geq 2$ (left) and 3-jet events (right). Here the black dots represent the jet response distribution and the

double-sided Crystal-Ball fit (blue line) is overlayed by the Gaussian fit (red line).

The resolution in each Gen $H_{T,2}/2$ bin is then plotted as a function of Gen $H_{T,2}/2$.




Figure 5.10: Comparison of jet energy resolution calculated using Crystal-Ball fit function (blue solid circles) and Gaussian fit function (red solid circles) for inclusive 2-jet (left) and 3-jet events (right).




Figure 5.11: Jet energy resolution (JER) is shown as a function of Gen $H_{T,2}/2$ for inclusive 2-jet (left) and 3-jet events (right). JER (black solid circles) is fitted by using the modified NSC-formula (blue solid line) which is extrapolated to 80 GeV and up to 2000 GeV (red dashed line) to consider the migration into lower as well as higher bins.

As expected, it has been observed from Fig. 5.10 that the Crystal Ball function (blue solid circles) describes the measured distributions better as compared to Gaussian function fit (red solid circles), especially in the low- $H_{T,2}/2$ region where the non-Gaussian tails are more pronounced. Hence JER is determined using Crystal Ball function fit. Figure 5.11 shows the final relative jet energy resolution (JER)

which is described by a modified version of the NSC formula (blue solid line) [96], as mentioned in Equation 5.8. To consider the migration to lower as well higher bins and to obtain the resolution with reasonable statistics over the full range of Gen $H_{T,2}/2$, the fit function is extrapolated to 80 GeV and up to 2000 GeV which is shown by red dashed line. The fit formula used here is basically the usual NSC formula which describes the resolution in terms of noise N originating due to electronic and pileup noise and is independent of $H_{T,2}/2$; a stochastic component S due to sampling fluctuation and EM fraction fluctuation per hadrons; and a constant term C because of dead material, magnetic field and calorimeter cell to cell fluctuation. In the low $H_{T,2}/2$ region the tracking has a non-negligible influence on the resolution due to the particle flow algorithm, so the additional parameter s is introduced to obtain slightly better fits. The parameters obtained after fitting the relative resolution using the above mentioned NSC formula are tabulated in Table 5.6 for $n_j \geq 2$ and $n_j \geq 3$ events. This calculated JER is used in unfolding procedure to smear the generated truth spectrum which is used as input in getting the response matrices and is explained in details in Sec. 5.5.1. Since JER in $n_j \geq 2$ events is similar to that one in $n_j \geq 3$ events, so N , S and C fit parameters obtained for $n_j \geq 3$ events are used for unfolding R_{32} .

$$\frac{\sigma(x)}{x} = \sqrt{\operatorname{sgn}(N) \cdot \frac{N^2}{x^2} + S^2 \cdot x^{s-1} + C^2} \quad (5.8)$$

Table 5.6: The parameters obtained by fitting the relative resolution as a function of $H_{T,2}/2$, using the modified NSC formula, for inclusive 2-jet and 3-jet events.

	N	S	C	s
Inclusive 2-jet	3.32	1.62	0.0333	-0.318
Inclusive 3-jet	-6.03	3.32	0.0333	-0.515

Since the JER is calculated using MG5+P6 Reco and Gen $H_{T,2}/2$ distributions, so it is expected that if Gen $H_{T,2}/2$ is smeared using this JER, it should match the

Reco $H_{T,2}/2$. But this extracted JER in one large rapidity bin, smears the Gen $H_{T,2}/2$ too much because Smeared Gen/Gen ratio (red line) shows a discrepancy from simulated Reco/Gen ratio (blue line), as observed in Fig. 5.12 for $n_j \geq 2$ (left) and $n_j \geq 3$ events (right). Some shortcomings in the detector simulation of the theory spectra leads to these small nonclosures. When the 30% reduced JER is used to smear Gen, then the ratio Smeared Gen/Gen (pink line) matches with simulated Reco/Gen ratio (blue line) within the statistical fluctuations. Hence an additional unfolding uncertainty is attributed by comparison to 30% reduced JER for both $n_j \geq 2$ and $n_j \geq 3$ events. Due to high statistical fluctuations at high $H_{T,2}/2$, range is presented up to 1680 GeV only.




Figure 5.12: MADGRAPH5+PYTHIA6 (MG5+P6) Gen smeared using extracted jet energy resolution (JER) shows a discrepancy from simulated Reco as Smeared Gen/Gen ratio (red line) does not match with Reco/Gen ratio (blue line), for both inclusive 2-jet (left) and 3-jet events (right). Smeared Gen/Gen ratio (pink line) where Gen is smeared using 30% reduced JER matches with simulated Reco/Gen ratio (blue line) within the statistical fluctuations. Hence an additional unfolding uncertainty is attributed by comparison to 30% reduced JER.

5.5 Unfolding

One of the main goals in an experimental measurement is to do the comparison of data with theory predictions or with the results obtained from other experiments. But the finite resolution of a detector and the steeply falling jet p_T spectrum distorts

the physical quantities. As a result, the measured observables are different from their corresponding true values. Each p_T bin content contains the migrated events from neighbouring bins along with the original events. So an unfolding process of the data should be followed in order to remove detector effects. In this analysis, the measurements are corrected for detector smearing effects and unfolded to stable particle level by using the iterative D'Agostini Bayesian algorithm [97, 98] as implemented in RooUnfold software package [99]. In this algorithm, the number of iterations regularize the unfolding process. The obtained distribution in one iteration is taken as the input in the next one. χ^2 between two successive iterations is given by Eq. 5.9. The number of iterations stop when χ^2/N_{bins} is < 1 . A reduced χ^2 is obtained by a higher number of iterations but this will also increase the uncertainty and there are larger bin-by-bin fluctuations and correlations. So the optimization of number of iterations is very important. In the current analysis, unfolding done with “four” iterations gives the best results with low χ^2 and low bin-by-bin correlations.

$$\chi^2 = \sum_{i=1}^{N_{bins}} \left(\frac{n_i^{j+} - n_i^j}{\sqrt{n_i^j}} \right)^2 \quad (5.9)$$

where n_i^j number of events in i -th bin for j -th iteration.

The measured differential cross-sections as a function of $H_{T,2}/2$, are unfolded separately for $n_j \geq 2$ and $n_j \geq 3$ events. The measured cross-section ratio R_{32} is also corrected for detector smearing effects and unfolded to particle level. There can be two ways to obtain unfolded cross-section ratio :

- **Method I :** First unfold separately the inclusive 2-jet and 3-jet measured cross-sections and then construct the ratio R_{32}
- **Method II :** Unfold directly the cross-section ratio R_{32}

In further analysis, unfolded cross-section ratio R_{32} and its systematic uncer-

tainties are calculated using Method I, whereas Method II is used only to propagate the statistical uncertainties including bin-by-bin correlations and statistical correlations between the inclusive 3-jet and 2-jet events cross-sections. Unfolding takes the response matrix as an input which are explained in the next section.

5.5.1 Response Matrices

The response matrix is a two dimensional mapping between the true and measured distributions. It is usually derived from simulated Monte Carlo (MC) samples, which takes the true distribution from MC as an input and smears it by taking into account the detector resolution. Then this response matrix is used to unfold the measured data spectrum. But there are several drawbacks of constructing response matrix using this method. In some phase space regions, the shape of the distribution is not well described by the LO predictions. Also, the limited number of events in the MC samples at high transverse momenta introduces high statistical fluctuations in the response matrix.

However, there is an indirect way of constructing the response matrix which uses a custom Toy Monte Carlo method. In this method, the particle level or true $H_{T,2}/2$ spectrum is obtained by fitting the theoretically predicted NLO spectrum. Then this distribution is smeared with forward smearing technique, using the extracted jet energy resolution (JER) to obtain the reconstructed level or measured $H_{T,2}/2$ spectrum. After that, the response matrix is constructed from these two distributions is used for the unfolding procedure.

5.5.1.1 Inclusive Cross-sections

The NLO spectrum of the differential cross-sections for $n_j \geq 2$ and $n_j \geq 3$ events obtained using CT10-NLO PDF set are fitted with the following two different functions defined in Eq. 5.10 and 5.13. These functions describes the shape as well as

normalization of the distribution.

- **Function I :**

$$f(H_{T,2}/2) = N[x_T]^{-a}[1 - x_T]^b \times \exp[-c/x_T] \quad (5.10)$$

where N is normalization factor and a, b, c are fit parameters.

This function is derived from the below function [100] :

$$f(p_T; \alpha, \beta, \gamma) = N_0[p_T]^{-\alpha} \left[1 - \frac{2 p_T \cosh(y_{min})}{\sqrt{s}} \right]^\beta \times \exp[-\gamma/p_T] \quad (5.11)$$

using

$$\alpha = a, \quad \beta = b, \quad \gamma = c * \sqrt{s}/2, \quad x_T = \frac{2 * H_{T,2}/2 * \cosh(y_{min})}{\sqrt{s}} = \frac{2 * H_{T,2}/2}{\sqrt{s}} \quad (5.12)$$

where transverse scaling variable x_T corresponds to the proton fractional momentum x for dijets with rapidity $y = 0$, $\sqrt{s} = 8000$ GeV and y_{min} is low-edge of the rapidity bin y under consideration (here y_{min} is taken equal to 0)

- **Function II :**

$$f(H_{T,2}/2) = A_0 \left(1 - \frac{H_{T,2}/2}{A_6} \right)^{A_7} \times 10^{F(H_{T,2}/2)}, \text{ where } F(x) = \sum_{i=1}^5 A_i \left(\log \left(\frac{x}{A_6} \right) \right)^i \quad (5.13)$$

where the parameter A_6 is fixed to $\frac{\sqrt{s}}{2 \cosh(y_{min})}$, where $\sqrt{s} = 8000$ GeV and y_{min} is the minimum rapidity. The other parameters are derived from the fitting.

Figure 5.13 shows the fitted CT10-NLO spectrum of differential cross-section as a function of $H_{T,2}/2$ (green solid circles) using Function I (top) and using Function




Figure 5.13: Fitted CT10-NLO spectrum of differential cross-section as a function of $H_{T,2}/2$ (green solid circles) using Function I (top) defined in Eq. 5.10 and using Function II (bottom) given by Eq. 5.13, for inclusive 2-jet (left) and 3-jet events (right). To consider the migration to lower $H_{T,2}/2$ bins, the fit functions described by red lines are extrapolated to 80 GeV (blue dashed lines).

II (bottom) : for inclusive 2-jet (left) and 3-jet events (right). Function I is used primarily to generate response matrices and perform the closure tests and Function II is used as an alternative function to calculate unfolding uncertainty, described in Sec. 5.6.3. To include the migration to lower bins, the fit functions described by red lines are extrapolated to 80 GeV (blue dashed lines).

A flat $H_{T,2}/2$ spectrum is generated by using toy Monte Carlo events and the fit parameters obtained from the NLO spectrum using function I (as shown in Fig. 5.13) provides weights to the flat spectrum. A total of ten million events are generated randomly (in $H_{T,2}/2$ range 80-2000). These generated values are then smeared with a

Gaussian function, where σ of the Gaussian is determined from the relative resolution parametrization as a function of $H_{T,2}/2$ calculated from NSC formula mentioned in equation 5.8. The parameters N, S, C used for smearing are taken from Table 5.6. These randomly generated (Gen_{Toy}) and smeared ($\text{Measured}_{\text{Toy}}$) values are used to fill the response matrices. Figure 5.14 shows the response matrices derived using the Toy MC for $n_j \geq 2$ (left) and $n_j \geq 3$ events (right). The matrices are normalized to the number of events in each column. The response matrices are diagonal as the migrations in off-diagonal bins are much smaller than the bins along the diagonal.




Figure 5.14: The response matrices are derived using the Toy Monte Carlo and forward smearing method, for inclusive 2-jet (left) and 3-jet events (right). The matrices are normalized to the number of events in each column and are diagonal with small off-diagonal migrations between close-by $H_{T,2}/2$ bins.

5.5.1.2 Cross-section Ratio, R_{32}

To obtain the statistical uncertainty on the unfolded cross-section ratio R_{32} , Method II is used. In this method, the response matrix is constructed using Toy MC method as done in Sec. 5.5.1.1 for differential cross-sections. To obtain the true spectrum for R_{32} , the ratio of cross-section spectrum described by Eq. 5.10 for inclusive 3-jet to that of 2-jet events is taken. This ratio is shown by green solid circles in Fig. 5.15 (left) which is fitted using a polynomial function of degree 8 (red line). Then as explained in above section, response matrix is derived for R_{32} using the Toy Monte

Carlo and forward smearing method which is shown in Fig. 5.15 (right). The matrix is normalized to the number of events in each column and is diagonal with small off-diagonal migrations between close-by $H_{T,2}/2$ bins.




Figure 5.15: Left : The ratio of cross-sections described by Eq. 5.10 for inclusive 3-jet to that of 2-jet events is shown as a function of $H_{T,2}/2$ (green solid circles). It is fit using a polynomial function of degree 8 (red line). Right : The response matrix is derived using the Toy Monte Carlo and forward smearing method, for the cross-section ratio R_{32} . The matrix is normalized to the number of events in each column and is diagonal with small off-diagonal migrations between close-by $H_{T,2}/2$ bins.

5.5.2 Closure Test

A closure test has been performed to confirm the working of the unfolding procedure. In this test, $\text{Measured}_{\text{Toy}}$ spectrum is unfolded using the constructed response matrices shown in Figure 5.14. It is expected that the same Gen_{Toy} spectrum should be re-obtained after unfolding. Figure 5.16 confirms that the unfolded $\text{Measured}_{\text{Toy}}$ spectrum matches exactly with Gen_{Toy} spectrum as the ratio of these distributions is perfectly flat at one for both $n_j \geq 2$ (top left) and $n_j \geq 3$ events (top right) cross-sections as well as the cross-section ratio R_{32} (bottom).

For another closure test, Reco MG5+P6 MC differential cross-section distribution is unfolded using the above constructed response matrices using JER for forward smearing the randomly generated spectrum. While taking ratio of the un-




Figure 5.16: Closure test of the unfolding technique where the smeared spectrum obtained from Toy Monte Carlo method (Measured_{Toy}), is unfolded using the constructed response matrices (obtained by forward smearing the randomly generated spectrum (Gen_{Toy}) using extracted jet energy resolution (JER)). As expected, the unfolded measured_{Toy} spectrum matches exactly with Gen_{Toy} spectrum as the ratio of these distributions is perfectly flat at one for both inclusive 2-jet (top left) and 3-jet events (top right) cross-sections as well as the cross-section ratio R_{32} (bottom).

folded distribution to that of Gen MG5+P6 MC, it is observed that a well closure is not obtained. This is represented by blue line in Fig. 5.17 for $n_j \geq 2$ (top left) and $n_j \geq 3$ events (top right). As observed in Fig. 5.12 in Sec. 5.4, if Reco MG5+P6 MC is unfolded using the response matrices obtained using 30% reduced JER, then the good closure is obtained as shown by red line in Fig. 5.17. Since unfolded cross-section ratio R_{32} is the ratio of unfolded differential cross-sections (Method I), same behaviour is observed for R_{32} (bottom).




Figure 5.17: Reco MADGRAPH5+PYTHIA6 Monte Carlo (MG5+P6 MC) differential cross-section distributions unfolded with the response matrices (obtained by forward smearing the randomly generated spectrum (Gen) using extracted jet energy resolution (JER)), does not give a good closure with Gen MG5+P6 MC (blue line), for inclusive 2-jet (top left) and 3-jet events (top right). After performing the unfolding using 30% reduced JER, a good closure is obtained (red line). Since unfolded the cross-section ratio R_{32} is the ratio of unfolded differential cross-sections, same behaviour is observed for R_{32} (bottom).

5.5.3 Unfolding of the Measurement

After validity the unfolding method, the measured differential cross-sections as well as R_{32} are unfolded using the above reconstructed response matrices. The unfolded data spectrum is compared to that of measured one in Fig. 5.18 for $n_j \geq 2$ (top left) and $n_j \geq 3$ events (top right) cross-sections and for the cross-section ratio R_{32} (bottom). As already discussed that 30% reduced JER gives better closures than JER, so the unfolding of data is done with response matrices using JER (blue solid circles) as well as 30% reduced JER (red solid circles) for smearing. The difference

between both is taken as an additional uncertainty on the unfolded measurement.




Figure 5.18: The measured differential cross-sections as well as the cross-section ratio R_{32} are unfolded as a function of $H_{T,2}/2$ using the response matrices derived using the Toy Monte Carlo and forward smearing method. The unfolded spectrum are compared with that of the measured one for inclusive 2-jet (top left) and 3-jet events cross-sections (right) as well as for R_{32} (bottom). The unfolding is done with response matrices using JER (blue solid circles) as well as 30% reduced JER (red solid circles) for smearing. The difference between both is taken as an additional uncertainty on the unfolded measurement.

5.6 Experimental Uncertainties

In an experimental measurement of any physical observable, the uncertainties play a key role and hence are important to study in a physics analysis. The uncertainties can be categorized into two types : statistical and systematic. The statistical uncertainties arise due to random fluctuations depending on the number of events. The more the number of events, less is the statistical uncertainty. The systematic

uncertainties may be due to known detector effects, model dependence, assumptions made or various corrections applied. In general, if the statistical and systematic uncertainties are uncorrelated, these can be added in quadrature to obtain the total uncertainty on the measurement. In this section, all the experimental uncertainties affecting the measurement of cross-sections and the cross-section ratio R_{32} are described. The systematic experimental uncertainties for R_{32} are propagated from the cross-sections to the ratio taking into account correlations. Due to this, the systematic uncertainties may cancel for R_{32} completely or partially as compared to those for the individual cross-sections.

5.6.1 Statistical Uncertainty

Statistical uncertainty on the measurement is obtained through the unfolding procedure using a toy MC method. The measured data points are smeared within their statistical uncertainties to get the smeared spectrum. Such smeared spectrums are produced million in number and the unfolding is performed multiple times for each smeared spectra. The differences between the unfolded spectrums and the measured one give the statistical uncertainty. The unfolding process introduces more statistical fluctuations which can be observed in Fig. 5.19. Here the fractional statistical uncertainties of the unfolded data (red line) are compared with those of the measured one (blue line) for $n_j \geq 2$ (top left) and $n_j \geq 3$ events cross-sections (top right) as well as for the cross-section ratio R_{32} (bottom).

After the unfolding, the final statistical uncertainties become correlated among the bins such that the size of these correlations varies between 10 and 20%. The correlation (anti-) is more significant for neighbouring bins in $H_{T,2}/2$ as compared to the far off ones. In Fig. 5.20, the correlations of the statistical uncertainty after the unfolding can be seen for $n_j \geq 2$ (top left) and $n_j \geq 3$ events cross-sections (top right) and for the cross-section ratio R_{32} (bottom). These correlations must be considered while performing the fits to extract the value of the strong coupling

constant, α_S .




Figure 5.19: The fractional statistical uncertainties of the unfolded data (red line) are compared with those of the measured one (blue line) for inclusive 2-jet (top left) and 3-jet events cross-sections (top right) as well as for the cross-section ratio R_{32} (bottom). After unfolding, the statistical uncertainty increases slightly.

5.6.2 Jet Energy Corrections Uncertainty

As explained in Sec. 4.3.2, the measured jet energy is corrected for a variety of detector effects by using jet energy corrections (JEC). This process introduces uncertainties in the final corrected jet energy. There are 25 mutually independent sources which contribute to JEC. Each source presents a 1σ shift and is fully correlated in p_T and η but uncorrelated to all other sources. The observable is studied with the nominal values of the jet energy which gives nominal distributions as well as by varying up and down the energy of all jets by the uncertainty. The differences




Figure 5.20: The unfolding procedure introduces the correlations of the statistical uncertainty through bin migrations which are shown here for inclusive 2-jet (top left) and and 3-jet events cross-sections (top right) as well as for the cross-section ratio R_{32} (bottom). The correlation (anti-) is more significant between neighbouring bins than far-ff ones.

between the nominal distributions and the ones obtained by varying the jet energy gives the uncertainties from each source. The JEC uncertainties can be asymmetric in nature which leads to separate treatment of upwards and downwards variation of each source. The sum in quadrature of uncertainties from all sources gives the total JEC uncertainty. In the current analysis, JEC uncertainties are a dominant source of experimental uncertainty at low $H_{T,2}/2$. The JEC uncertainty ranges from 3% to 10% for $n_j \geq 2$ and from 3% to 8% for $n_j \geq 3$ events cross-sections measurement. To calculate JEC uncertainty for ratio R_{32} , the inclusive 2-jet and 3-jet events cross-sections are measured as a function of $H_{T,2}/2$ by shifting the jet p_T according to the JEC uncertainty for each source of JEC separately. Then the ratio of these cross-

sections is taken and the difference of these from the central ratio R_{32} , gives the JEC uncertainty for R_{32} . As expected, JEC uncertainty for R_{32} is small as compared to that for individual cross-sections and is about 1 to 2% over all $H_{T,2}/2$ bins.

The sources of JEC considered in the current measurements are : AbsoluteStat, AbsoluteScale, AbsoluteFlavMap, AbsoluteMPFBias, Fragmentation, SinglePionECAL, SinglePionHCAL, FlavorQCD, RelativeJEREC1, RelativeJEREC2, RelativeJERHF, RelativePtBB, RelativePtEC1, RelativePtEC2, RelativePtHF, RelativeFSR, RelativeStatFSR, RelativeStatEC2, RelativeStatHF, PileUpDataMC, PileUpPtRef, PileUpPtBB, PileUpPtEC1, PileUpPtEC2 and PileUpPtHF. The AbsoluteFlavMap uncertainty is exactly zero for the 8 TeV and can be ignored. For the four sources : RelativeJERHF, RelativePtHF, RelativeStatHF, PileUpPtHF, the JEC uncertainty is exactly zero because of $|y| < 2.5$ cut used in the analysis. So only 20 sources contribute to the total JEC uncertainty. The Figs. A.1-A.3 show the JEC uncertainty from each source separately for inclusive 2-jet (top) and 3-jet events cross-sections (middle) as for cross-section R_{32} (bottom). Depending on the origin of sources, they are categorized into four groups which are described below in brief :

1. **Pileup** - This uncertainty originates from the differences in the transverse momentum between the true offset and the Random Cone method (i.e. essentially difference of pileup inside and outside of jets), in simulated events. This uncertainty is derived from $Z/\gamma+jet$, dijet and multijet data using fit procedure to estimate the residual pileup uncertainty after the calibration.
2. **Relative** - The forward jets are calibrated by the relative η -dependent corrections using dijet events. The main contribution to the uncertainty comes from jet energy resolution (JER), derived by varying JER scale factors up and down by quoted uncertainties and the initial and final state radiation bias corrections.

3. **Absolute** - A global fit to Z/γ +jet and multijet events gives the absolute calibration of the jet energy scale. The uncertainties are related to the lepton momentum scale for muons in $Z (\rightarrow \mu\mu)$ +jet and the single pion response in the HCAL.
4. **Flavor** - Flavor response differences are studied from simulation by cross-checking the results with quark- and gluon-tagged γ +jet and Z +jet events. These uncertainties are based on PYTHIA6.4 and HERWIG++2.3 differences propagated through the data-based calibration method.

The details of the jet energy corrections and uncertainties can be found in [101].

5.6.3 Unfolding Uncertainty

The unfolding uncertainty is comprised of three uncertainties which are explained as follows :

1. **Jet Energy Resolution** - The calculation of the jet energy resolution (JER) using simulated MG5+P6 Monte Carlo events is already explained in Sec. 5.4. As mentioned before, the measured jet transverse momentum (p_T) in simulated MC events needs to be smeared additionally to match the resolution in data. This smearing is done by using measured scale factors ($c_{central}$) mentioned in Table 5.5. It is recommended by JETMET group that the uncertainty on these measured scaling factors must be taken into account in a physics analysis. Since JER is used in constructing the response matrix which is an input in unfolding procedure, so the uncertainty on scale factors accounts for the unfolding uncertainty. To calculate JER uncertainty, p_T is smeared with two additional sets of scale factors corresponding to varying the factors up and down by one sigma, and corresponding $H_{T,2}/2$ is calculated. Then again JER is calculated as a function of $H_{T,2}/2$ using these upwards (c_{up}) and downwards (c_{down}) variations of the scaling factors. Alternative response matrices are built

using the JER with above variations and the unfolding is performed again. The differences of the obtained unfolded spectrums to the nominal ones accounts for a systematic JER uncertainty.

2. **Model Dependence** - It is explained in Sec. 5.5.1 that to obtain the true $H_{T,2}/2$ spectrum to be used in constructing response matrix using Toy MC method, the fitting of the CT10-NLO predictions is performed with the Function I described in Eq. 5.10. Using the alternative function, Function II given by Eq. 5.13, for this fitting and then constructing different response matrix, gives the model dependence of the true $H_{T,2}/2$ spectrum. The differences in unfolded distributions using the above mentioned two different response matrices gives the model dependence uncertainty.
3. **Additional Uncertainty** - Small nonclosures observed in Fig. 5.12 introduces a supplementary uncertainty which is attributed by comparison of distributions unfolded using response matrices constructed using JER from simulation with that obtained with a 30% reduced JER.

All the three above mentioned uncertainties are added in quadrature to get the total unfolding uncertainty which increases from about 1% at low $H_{T,2}/2$ up to 2% at the high $H_{T,2}/2$ ends of the cross-sections for both $n_j \geq 2$ and $n_j \geq 3$ events. This uncertainty account for about less than 1% for R_{32} .

5.6.4 Luminosity Measurement Uncertainty

As discussed in Sec. 3.1.1, the luminosity delivered to CMS detector by LHC in the proton-proton collisions in the year of 2012 is measured by using the silicon pixel cluster counting method [55]. The uncertainty related to the integrated luminosity measurement is estimated to be 2.5% (syst.) and 0.5% (stat.). This uncertainty propagates directly to any absolute cross-section measurement. Hence, a total systematic uncertainty of 2.6% is considered across all the $H_{T,2}/2$ bins. At low $H_{T,2}/2$,

it is similar in size as the one from JEC. This uncertainty cancels completely for R_{32} .

5.6.5 Residual Uncertainty

The small trigger and jet identification inefficiencies account for smaller than 1% uncertainties on the cross-section measurements [91, 102]. Hence, an uncorrelated residual uncertainty of 1% is assumed across all $H_{T,2}/2$ bins for both $n_j \geq 2$ and $n_j \geq 3$ events cross-sections whereas for R_{32} , it gets cancel completely.

5.6.6 Total Experimental Uncertainty

After calculating the uncertainties from all the above mentioned sources, the total experimental uncertainty on measurement of cross-sections as well as cross-section ratio R_{32} , is obtained by adding in quadrature the uncertainties from individual sources. Figure 5.21 shows the experimental uncertainties, from different sources as well as the total uncertainty, affecting the measurement of $n_j \geq 2$ (top left) and $n_j \geq 3$ events cross-sections (top right) and cross-section ratio R_{32} (bottom). The error bars represent the statistical uncertainty obtained after unfolding. The systematic uncertainties due to jet energy corrections (JEC by blue line), luminosity (red dashed line), unfolding (green dashed line) and residual effects (light purple line) are also presented. The uncertainties due to luminosity and residual effects cancel completely in R_{32} . The total uncertainty (black dashed line) on the measurements is asymmetric in nature and dominated by the uncertainty due to the jet energy corrections (JEC) at lower $H_{T,2}/2$ values and by statistical uncertainty at higher $H_{T,2}/2$ values.

The experimental uncertainties from each source as well as total uncertainty are also quoted in Table 5.7. The values of uncertainties (in %) from each source as well as total uncertainty, for each $H_{T,2}/2$ bin, are tabulated in Tables A.2, A.3




Figure 5.21: Experimental uncertainties from different sources affecting the measurement of cross-sections for inclusive 2-jet (top left) and 3-jet events (top right) and cross-section ratio R_{32} (bottom). The error bars represent the statistical uncertainty after unfolding. The systematic uncertainties due to jet energy corrections (JEC by blue line), luminosity (red dashed line), unfolding (green dashed line) and residual effects (light purple line) are also presented. The uncertainties due to luminosity and residual effects cancel completely in R_{32} . The total uncertainty (black dashed line) is the quadrature sum of the individual sources of uncertainty.

and A.4 for $n_j \geq 2$ and $n_j \geq 3$ events cross-sections and cross-section ratio R_{32} , respectively.

The complete data analysis of the differential inclusive 2-jet and 3-jet events cross-sections as well as their ratio R_{32} has been presented as a function of $H_{T,2}/2$. The measured spectrums after correcting for detector effects through the unfolding procedure, are compared with the next-to-leading order (NLO) pQCD calculations in the next chapter.

Table 5.7: Overview of all experimental uncertainties affecting the measurement of cross-sections for inclusive 2-jet (left) and 3-jet events (middle) and cross-section ratio R_{32} (right). The uncertainties due to luminosity and residual effects cancel completely in R_{32} . The total uncertainty is the quadrature sum of the individual sources of uncertainty.

Uncertainty Source	Inclusive 2-jet	Inclusive 3-jet	R_{32}
Statistical	< 1 to 30%	< 1 to 27%	< 1 to 28%
Jet energy corrections (JEC)	3 to 10%	3 to 8%	1 to 2%
Unfolding	1 to 2%	1 to 2%	< 1%
Luminosity	2.6%	2.6%	cancels
Residual	1%	1%	cancels
Total	4 to 32%	4 to 28%	1 to 28%

Chapter 6

Theoretical Calculations

In an experiment, the measurements are validated by doing the comparison with the perturbative QCD (pQCD) theoretical calculations. The lowest order (LO) calculations describe well the shapes of the measured distributions but not the normalization due to the dependence on the unphysical renormalization (μ_r) and factorization (μ_f) scales. The next-to-leading order calculations (NLO) improves the precision by reducing the dependence on μ_r and μ_f scales and become an essential feature in the determination of fundamental parameters such as α_S and the parton distribution functions (PDF). This chapter describes the next-to-leading order pQCD calculations used for comparison with the measurements. NLO pQCD calculations need to be corrected for the multiparton interactions (MPI) and hadronization effects by applying non-perturbative (NP) corrections and also for the electroweak interactions (EW).

6.1 Fixed Order NLO Calculations

The predictions of the inclusive differential jet event cross-section at NLO accuracy in pQCD are computed with the NLOJET++ program version 4.1.3 [62, 63]. The results are provided within the framework of FASTNLO version 2.3 [64, 65]. The

PDFs are accessed through the LHAPDF6 library [66, 67]. The FASTNLO is preferred over the direct calculation with NLOJET++ because with FASTNLO the calculations of the cross-sections can be repeated several times with different PDFs and scale choices required for calculating the PDF and scale uncertainties. Here the factorization and renormalization scales are chosen equal to $H_{T,2}/2$, i.e. $\mu_f = \mu_r = H_{T,2}/2$.

In the current study, different PDF sets available for a series of different assumptions on the strong coupling constant at the scale of the Z boson mass $\alpha_s(M_Z)$ are used for NLO calculations. Table 6.1 summarizes the already existing PDF sets in LHC Run 1 (upper rows) and the newer PDF sets for Run 2 (lower rows). The different columns list the number of flavours N_f , the assumed masses M_t and M_Z of the top quark and the Z boson, respectively, the default values of $\alpha_s(M_Z)$, and the range in $\alpha_s(M_Z)$ variation available for fits for different PDF sets. All sets uses a variable-flavour number scheme with at most five or six flavours apart from the ABM11 PDFs, which employ a fixed-flavour number scheme with $N_F = 5$. Out of these eight PDF sets the following three are not considered further because of the below mentioned reasons :

- At NLO, predictions based on ABM11 do not describe LHC jet data at small jet rapidity [103–106].
- The HERAPDF2.0 set exclusively fits HERA DIS data with only weak constraints on the gluon PDF.
- The range in values available for $\alpha_s(M_Z)$ is too limited for the NNPDF3.0 set.

Mainly CT10 PDF set is considered for comparison between data and theory predictions as well as for calculating theoretical uncertainties.

Table 6.1: NLO PDF sets are available via LHAPDF6 with various assumptions on the value of $\alpha_s(M_Z)$. The upper rows list the already existing sets in LHC Run 1 and newer ones for Run 2 are listed in lower rows, along with the corresponding number of flavours N_f , the assumed masses M_t and M_Z of the top quark and the Z boson, respectively, the default values of $\alpha_s(M_Z)$, and the range in $\alpha_s(M_Z)$ variation available for fits.

Base set	N_F	M_t (GeV)	M_Z (GeV)	$\alpha_s(M_Z)$	$\alpha_s(M_Z)$ range
ABM11 [27]	5	180	91.174	0.1180	0.110 - 0.130
CT10 [107]	≤ 5	172	91.188	0.1180	0.112 - 0.127
MSTW2008 [108, 109]	≤ 5	10^{10}	91.1876	0.1202	0.110 - 0.130
NNPDF2.3 [110]	≤ 6	175	91.1876	0.1180	0.114–0.124
CT14 [24]	≤ 5	172	91.1876	0.1180	0.111–0.123
HERAPDF2.0 [28]	≤ 5	173	91.1876	0.1180	0.110–0.130
MMHT2014 [25]	≤ 5	10^{10}	91.1876	0.1200	0.108–0.128
NNPDF3.0 [26]	≤ 5	173	91.2	0.1180	0.115–0.121

6.1.1 NLO Correction Factors

The differences between LO predictions and NLO predictions give the effect of the higher-order contributions to the pQCD predictions. These are described by a NLO correction factor, k-factor, which is derived as the ratio of cross-sections at NLO accuracy to that at LO i.e.

$$\text{k-factor} = \frac{\sigma_{\text{NLO}}}{\sigma_{\text{LO}}} \quad (6.1)$$

The impact of the higher-order corrections is determined by the size of k-factor. The small size of k-factor indicates that the cross-section predictions are precisely described at the LO whereas the larger size hints the contributions from NLO. Figure 6.1 shows the k-factors of the NLOJET++ calculations, for inclusive 2-jet and 3-jet event cross-sections and their ratio R_{32} , using five different PDF sets. k-factor for R_{32} is obtained by taking the ratio of k-factors for inclusive 3-jet event cross-sections to that of inclusive 2-jet. The k-factors are similar for all the PDF sets in the lower region, but the differences increase in regions with larger $H_{T,2}/2$. It is observed that for inclusive 3-jet event cross-sections, k-factor jumps at lowest

$H_{T,2}/2$. This is because some jet configurations are kinematically forbidden near the p_T cut bin i.e. 150 GeV. Since the first few bins in $H_{T,2}/2$ (below 225 GeV) still suffer from these kinematical effects, the minimum value of $H_{T,2}/2$ studied is 300 GeV.




Figure 6.1: The k-factors of the NLOJET++ calculations, for inclusive 2-jet and 3-jet event cross-sections and their ratio R_{32} , using five different PDF sets.

6.1.2 Non-perturbative Corrections

The fixed-order pQCD NLO calculations predict the parton-level cross-section but lacks accuracy due to several effects. The partons which are emitted close to each other in phase space are not handled well in lower order perturbation theories and hence requires a parton shower (PS) correction. The scattering phenomena between partons within a colliding proton, other than the hard scattering, give rise to multi-parton interactions (MPI). The partons of the hard scattering forms colorless bound states called hadrons through a process of hadronization (HAD). The MPI and hadronization cannot be modelled well within the perturbative framework. Since the fixed-order NLO calculations do not include these additional soft QCD effects, these calculations cannot be compared directly to unfolded data. So the corrections for non-perturbative effects (NP) should be taken into account in NLO calculations. The ratio of cross-section predicted with a nominal event generation interfaced to

the simulation of UE contributions and to the one without hadronization and MPI effects gives the NP correction factors which are defined as :

$$C^{\text{NP}} = \frac{\sigma^{\text{PS+HAD+MPI}}}{\sigma^{\text{PS}}} \quad (6.2)$$

In the current study, the NP effects are estimated by using samples obtained from various MC event generators with a simulation of parton shower and underlying-event (UE) contributions. The leading order (LO), HERWIG++ with the default tune of version 2.3 and PYTHIA6 with tune Z2*, and the NLO POWHEG MC event generators are considered. The matrix-element calculation is performed with POWHEG interfaced to PYTHIA8 with tune CUETS1 for the UE simulation. The ratio, defined in Eq. 6.2, is obtained for each MC generator and is fitted by a power-law function defined in Eq. 6.3. Since this ratio obtained from different MC generators have large differences, so the average of the envelope, which covers all the differences, is taken as the correction factor which is then applied as bin-by-bin multiplicative factor to the parton-level NLO cross-section. The half of the envelope it is taken as the uncertainty on the NP correction factor.

$$f(H_{\text{T},2}/2) = a \cdot (H_{\text{T},2}/2)^b + c \quad (6.3)$$

The NP correction factors, $C_{3\text{-jet}}^{\text{NP}}$ and $C_{2\text{-jet}}^{\text{NP}}$ are calculated for $n_j \geq 2$ and $n_j \geq 3$ event cross-sections respectively and then their ratio gives the correction factor for R_{32} . The correction factors are shown in Fig. 6.2 for the inclusive 2-jet (top left) and 3-jet event cross-sections (top right), and for ratio R_{32} (bottom). At $H_{\text{T},2}/2 \sim 300$ GeV, the NP corrections amount to $\sim 4\text{-}5\%$ for inclusive 2-jet and 3-jet event cross-sections and $\sim 1\%$ for R_{32} , and decrease rapidly for increasing $H_{\text{T},2}/2$. On comparing the NP correction factors of R_{32} with that for individual cross-sections, it has been observed that the non-perturbative effects get reduced in R_{32} .




Figure 6.2: The nonperturbative (NP) corrections are presented as a function of $H_{T,2}/2$ for inclusive 2-jet (top left) and 3-jet (top right) event cross-sections, as well as their ratio R_{32} . These corrections are calculated from the leading order HERWIG++ with the default tune of version 2.3 (red line) and PYTHIA6 with tune Z2* (blue line); and the next-to-leading order POWHEG interfaced to PYTHIA8 with tune CUETS1 (green line) Monte Carlo event generators. The black solid circles give the average NP correction factor along with the uncertainty shown by the error bars.

6.1.3 Electroweak Corrections

In LHC, the centre-of-mass energy of proton-proton collisions is well beyond the electroweak (EW) scale $\sim \mathcal{O}(100 \text{ GeV})$. At such a high energy, the impact of higher order EW corrections is much more with respect to QCD effects [111] and affect jet cross-sections at large $H_{T,2}/2$. The quark-quark scattering processes involving virtual exchanges of massive W and Z bosons contribute to electroweak (EW) corrections. The fixed-order QCD calculations do not include EW corrections and hence the NLO theory calculations are corrected for EW effects. The EW corrections have




Figure 6.3: The electroweak (EW) corrections [112] in the phase space of the measurement are shown as a function of $H_{T,2}/2$ for inclusive 2-jet event cross-sections. These corrections are applied as a bin-by-bin correction factor to the fixed-order calculation of NLOJET++ as well as the MC predictions of MADGRAPH5+PYTHIA6. The EW correction factor increases up to 13% at high ends of $H_{T,2}/2$ and significantly improves the agreement between data and prediction.

been calculated for inclusive 1-jet and 2-jet case, in Ref. [112]. The EW correction factors in the phase space of the measurement are shown as a function of $H_{T,2}/2$ in Fig. 6.3 for inclusive 2-jet event cross-sections. These correction factor increases up to 13% at high ends of $H_{T,2}/2$ which are applied as a bin-by-bin correction factor to the fixed-order NLOJET++ calculations. To see the effects of EW corrections, a ratio of data to theory predictions obtained using CT10-NLO PDF set and corrected with NP effects without including EW corrections (left) and including EW corrections (right) is plotted for inclusive 2-jet event cross-sections in Fig. 6.4. On comparing both the figures, it is observed that the EW corrections significantly improve the agreement between data and prediction in the high $H_{T,2}/2$ region. EW corrections are not available yet for inclusive 3-jet production and hence not applied for inclusive 3-jet event cross-sections. The guess from theory side is that EW for inclusive 2-jet and 3-jet will be similar, so for R_{32} , it is assumed to be equal to the factor of 1. Since the EW effects are not taken care of in MC simulations so these corrections are applied to MC predictions also.




Figure 6.4: Ratio of data over theory obtained using the CT10-NLO PDF set and corrected with non-perturbative effects (NP) without including electroweak (EW) corrections (left) and including EW corrections (right) is shown for inclusive 2-jet event cross-sections. The error bars represents the statistical uncertainty of the data and the shaded rectangles represents the total experimental systematic uncertainty. The shaded band around unity indicate the total uncertainty of the theory. The EW corrections significantly improve the agreement between data and prediction in the high $H_{T,2}/2$ region.

6.2 Theoretical Uncertainties

The measurements are not only sensitive to experimental uncertainties but also to the theoretical uncertainties. The renormalization and factorization scale variations, PDF uncertainties and the non-perturbative corrections contribute to theoretical uncertainties which are described below :

6.2.1 Scale Uncertainty

In perturbative QCD calculations of cross-sections, one has to choose a renormalization (μ_r) and factorization (μ_f) scale. The dependence on scales is negligible if these calculations are performed for all orders of the perturbative series. But the perturbative series is truncated at NLO, so there is a scale dependence of the measurement which is covered by systematic uncertainty known as scale uncertainty. The scale uncertainty is evaluated with the conventional recipe of varying the default scale $H_{T,2}/2$ chosen for μ_r and μ_f independently in the following six combinations: $(\mu_r/H_{T,2}/2, \mu_f/H_{T,2}/2) = (1/2, 1/2), (1/2, 1), (1, 1/2), (1, 2), (2, 1)$ and $(2, 2)$. The

maximal upwards and downwards deviations in cross-section from the central prediction give the scale uncertainty. To calculate the scale uncertainty for cross-section ratio R_{32} , first R_{32} is obtained for each above mentioned scale choice and then its difference from central R_{32} is taken. The scale uncertainty calculated using CT10-NLO PDF set ranges from 5% to 13% and 11% to 17% for inclusive 2-jet and 3-jet events cross-sections respectively, and from 6% to 8% for R_{32} .

6.2.2 PDF Uncertainty

The calculation of jet cross-sections in proton-proton collisions relies upon the knowledge of PDFs. These PDF sets are determined by global fits to all the available deep inelastic scattering (DIS) and related hard scattering data from different experiments. The various sources affect the PDFs such as theory model, input parameters like the strong coupling constant α_S , the quark masses and the statistical and systematic uncertainty sources of the data included in the PDF fit. These sources contribute to PDF uncertainty which is evaluated according to the prescriptions given for each PDF set. The CT10-NLO PDF set [107, 113] employ the eigenvector method to evaluate the PDF uncertainties. The CT10-PDF set consists of $N_{\text{ev}} = 26$ eigenvectors with two PDF members per eigenvector k , which are varied upwards and downwards to generate a set of eigenvector pairs. The asymmetric uncertainties, ΔX^+ and ΔX^- , of a quantity X are given by Eq. 6.4 where X_0 is the central prediction, X_k^+ and X_k^- are the predictions using the upwards and downwards variation of each eigenvector k .

$$\Delta X^+ = \sqrt{\sum_{k=1}^{N_{\text{ev}}} [\max(X_k^+ - X^0, X_k^- - X^0, 0)]^2} \quad (6.4)$$

$$\Delta X^- = \sqrt{\sum_{k=1}^{N_{\text{ev}}} [\min(X_k^+ - X^0, X_k^- - X^0, 0)]^2}$$

The symmetric uncertainty (ΔX^\pm) is given by half the difference of the upwards and downwards variations :

$$\Delta X^\pm = \sqrt{\sum_{k=1}^{N_{\text{ev}}} \left[\frac{X_k^+ - X_k^-}{2} \right]^2} \quad (6.5)$$

The CT10-NLO PDF set uncertainties are downscaled by a factor of 1.64 in order to have the uncertainties at the 68.3% confidence level $\text{CL}(1\sigma)$ instead of 90% $\text{CL}(2\sigma)$ such that to have a uniform treatment with respect to other PDF sets. The PDF uncertainty as derived with the CT10-NLO PDF set is the dominant source of uncertainty and ranges from 3% to 30% for inclusive 2-jet and from 4% to 32% for 3-jet cross-sections. For R_{32} , the ratio of predictions for inclusive 3-jet to that of 2-jet is taken for each eigen vector with upwards and downwards variations separately and then PDF uncertainty is calculated as done for individual cross-sections. The PDF uncertainty ranges and from 2% to 10% for cross-section ratio R_{32} .

6.2.3 Non-perturbative Uncertainty

As discussed in [6.1.2](#), the differences in the non-perturbative (NP) corrections calculated from various Monte Carlo event generators introduce the NP uncertainty which is of the order of 1% and 1 to 2% for inclusive 2-jet and 3-jet event cross-sections respectively, and < 1% for cross-section ratio R_{32} .

6.2.4 Total Theoretical Uncertainty

The total systematic theoretical uncertainties are obtained as the quadratic sum of the scale, PDF and NP uncertainties. Figure [6.5](#) presents the systematic theoretical uncertainties affecting the cross-section measurement for inclusive 2-jet (top left) and 3-jet events (top right) and the cross-section ratio R_{32} (bottom), using CT10-




Figure 6.5: The systematic theoretical uncertainties affecting the cross-section measurement for inclusive 2-jet (top left) and 3-jet events (top right) and their ratio R_{32} (bottom). The scale (red dashed line), PDF (green line) and NP (blue dashed line) uncertainties as well as total uncertainty (black dashed line) obtained using CT10-NLO PDF set are shown. The total theoretical uncertainty is asymmetric and is dominated by PDF uncertainty.

NLO PDF set. The scale (red dashed line), PDF (green line) and NP (blue dashed line) uncertainties as well as total theoretical uncertainty (black dashed line) are shown. The total theoretical uncertainty is asymmetric and is dominated by PDF uncertainty which grows in magnitude with increasing value of $H_{T,2}/2$. Table 6.2 quotes the values of the theoretical uncertainty from each source as well as total uncertainty affecting the measurements. The bin-wise values of uncertainties (in %) from each source as well as total uncertainty are shown in Tables A.5, A.6 and A.7 for $n_j \geq 2$ and $n_j \geq 3$ event cross-sections and cross-section ratio R_{32} , respectively. The computation of the NLO predictions with NLOJET++ is also

subject to statistical fluctuations from the complex numerical integrations. For the inclusive 2-jet event cross-sections this uncertainty is smaller than about a per mille, while for the inclusive 3-jet event cross-section it amounts to 1-9 per mille. Hence the statistical uncertainty is not considered in the total theoretical uncertainty. The small dips at ~ 700 and 1000 GeV in the PDF uncertainty for inclusive 3-jet events cross-sections and cross-section ratio R_{32} is a feature of the CT10-NLO PDF set.

Table 6.2: Overview of all systematic theoretical uncertainties, obtained using CT10-NLO PDF set, affecting the measurement of cross-sections for inclusive 2-jet (left) and 3-jet events (middle) and cross-section ratio R_{32} (right).

Uncertainty Source	Inclusive 2-jet	Inclusive 3-jet	R_{32}
Scale	5 to 13%	11 to 17%	6 to 8%
PDF	3 to 30%	4 to 32%	2 to 10%
Non-perturbative (NP)	1%	1 to 2%	< 1%
Total	3 to 30%	5 to 34%	3 to 11%

6.3 Comparison of Theory to Data

After correcting the measurement for detector effects as well as NLO pQCD calculations for non-perturbative (NP) and electroweak (EW) effects, it is now feasible to compare the measured cross-sections with the theory predictions. Figure 6.6 shows the measured differential inclusive 2-jet and 3-jet event cross-sections as a function of $H_{T,2}/2$ after unfolding for detector effects. On the left, the measurements (points) are compared to the NLOJET++ predictions using the CT10-NLO PDF set (line), corrected for NP effects and in addition for EW effects in the 2-jet case. On the right, the comparison is made to the predictions from MADGRAPH5+PYTHIA6 (MG+P6) with tune Z2* (line), corrected for EW effects in the 2-jet case. The error bars give the total experimental uncertainty, given by the quadrature sum of the statistical and systematic uncertainties. On a logarithmic scale, the data are in well agreement with the NLO predictions over the whole range of $H_{T,2}/2$ from 300 GeV up to 2000

(2-jet) and 1680 GeV (3-jet) respectively.




Figure 6.6: Comparison of the measured differential inclusive 2-jet and 3-jet event cross-sections as a function of $H_{T,2}/2$ to theoretical predictions. On the left, the data (points) are shown together with NLO JET++ predictions (line) using the CT10-NLO PDF set, corrected for non-perturbative (NP) and electroweak (EW) effects (2-jet) or only NP effects (3-jet). On the (right), the data (points) are compared to predictions from MADGRAPH5+PYTHIA6 (MG+P6) with tune Z2* (line), corrected for EW effects in the 2-jet case. The error bars give the total experimental uncertainty, given by the quadratic sum of the statistical and systematic uncertainties.




Figure 6.7: Cross-section ratio R_{32} as a function of $H_{T,2}/2$ calculated from data (solid circles) in comparison to that from NLO pQCD predictions obtained using the CT10-NLO PDF set corrected with non-perturbative (NP) corrections (line). The error bars correspond to the total experimental uncertainty derived as quadratic sum from all uncertainty sources.

Figure 6.7 shows the cross-section ratio R_{32} obtained from unfolded data (solid circles) in comparison to that from NLO pQCD predictions obtained using the CT10-NLO PDF set corrected with NP corrections (line). The error bars here

represents the total experimental uncertainty derived as quadratic sum from all uncertainty sources. The deviations of measured R_{32} from the theoretical predicted value can be explained by the electroweak effects which are not considered yet because of their unavailability for inclusive 3-jet event cross-sections.




Figure 6.8: Ratio of data over theory using the CT10-NLO PDF set for inclusive 2-jet (top left) and 3-jet event cross-sections (top right) and their ratio R_{32} (bottom). The theory predictions are corrected for non-perturbative effects (NP) and also for electroweak effects (EW) for inclusive 2-jet only. For comparison predictions employing two other PDF sets, MSTW2008 and NNPDF2.3, are also shown. The error bars represents the statistical uncertainty of the data and the shaded rectangles represents the total experimental systematic uncertainty. The shaded band around unity indicate the total uncertainty of the theory.

For better visibility, the ratios of data over the theory at NLO are also studied in details. In Fig. 6.8, the ratios of data over NLOJET++ predictions using the CT10-NLO PDF set are shown for inclusive 2-jet (top left) and 3-jet event cross-sections (top right) as well as their ratio R_{32} (bottom). The data are well described by the predictions within their uncertainty, which is dominated at large $H_{T,2}/2$ by

PDF effects in the upwards and by scale variations in the downwards direction. A trend towards an increasing systematic excess of the 2-jet data with respect to theory, starting at about 1 TeV in $H_{T,2}/2$, is remedied by the inclusion of EWK corrections. In the 3-jet case the statistical precision of the data and the reach in $H_{T,2}/2$ is insufficient to observe any effect. The alternative PDF sets MSTW2008 and NNPDF2.3 exhibit a small underestimation of the cross-sections at high $H_{T,2}/2$.




Figure 6.9: Ratio of data over the predictions from POWHEG+PYTHIA8 with tune CUETS1 are presented for inclusive 2-jet (top left) and 3-jet event cross-sections (top right) as well as their ratio R_{32} (bottom). For comparison the alternative tune CUETM1 of POWHEG+PYTHIA8, the tree-level multi-leg improved prediction by MADGRAPH5+PYTHIA6 with tune $Z2^*$, and the the LO MC predictions from PYTHIA6 tune $Z2^*$ are shown as well. The error bars correspond to the statistical uncertainty of the data and the shaded rectangles to the total experimental systematic uncertainty. EW corrections have been accounted for in this comparison in the 2-jet case only.

The POWHEG framework providing a NLO dijet calculation matched to the parton showers of PYTHIA8 employed with the CUETS1 and CUETM1 tunes [73] is also

used for a comparison. The ratios of data over theory from POWHEG+PYTHIA8 with tune CUETS1 are shown for inclusive 2-jet (top left) and 3-jet event cross-sections (top right) as well as their ratio R_{32} (bottom) in Fig. 6.9. For comparison, the LO prediction from PYTHIA6 with tune Z2*, the tree-level multi-leg improved prediction by MADGRAPH5+PYTHIA6 with tune Z2*, and the matched NLO prediction from POWHEG+PYTHIA8 with tune CUETM1 are shown as well. EW corrections have been accounted for in this comparison in the 2-jet case only. Significant discrepancies, which are cancelled to a large extent in the ratio R_{32} , are visible in the comparison with the LO prediction from MADGRAPH5+PYTHIA6 with tune Z2*, in particular for small $H_{T,2}/2$. In contrast, the employed dijet MC POWHEG+PYTHIA8 better describe the 2-jet event cross-section, but fail for the 3-jet case.

The jet measurements at hadron colliders can be used to extract the strong coupling constant α_S , which is discussed in the next chapter.

Chapter 7

Determination of the Strong Coupling Constant

The inclusive jet production cross-section at hadron colliders mainly depends on the strong coupling constant α_s for a given centre-of-mass energy. Hence the measurements of the inclusive jet cross-section and jet properties provide a direct probe to measure the strong coupling constant. The measurement of α_s has been already done by various experiments such as CMS [1, 91, 105, 114, 115], ATLAS [116], D0 [117, 118], H1 [119, 120], and ZEUS [121]. In this thesis, the measurements of differential inclusive 2-jet and 3-jet event cross-sections as well as the cross-section ratio R_{32} , as a function of $H_{\text{T},2}/2$ are used to extract the value of the strong coupling constant at the scale of the Z boson mass $\alpha_s(M_Z)$. The differential inclusive jet production cross-section up to at NLO is given by [122] :

$$\frac{d\sigma}{d(H_{\text{T},2}/2)} = \alpha_s^2(\mu_r) \hat{X}^{(0)}(\mu_f, H_{\text{T},2}/2) [1 + \alpha_s(\mu_r) K_1(\mu_r, \mu_f, H_{\text{T},2}/2)] \quad (7.1)$$

where $\frac{d\sigma}{d(H_{\text{T},2}/2)}$ is the differential inclusive jet production cross-section as a function of $H_{\text{T},2}/2$, μ_r and μ_f are the renormalization and factorization scales set equal to $H_{\text{T},2}/2$, $\alpha_s^2(\mu_r) \hat{X}^{(0)}(\mu_f, H_{\text{T},2}/2)$ is the leading order

(LO) contribution to the differential inclusive jet production cross-section and $\alpha_S^3(\mu_r)\hat{X}^{(0)}(\mu_f, H_{T,2}/2)K1(\mu_r, \mu_f, H_{T,2}/2)$ is the next-to-leading order (NLO) contribution. Equation 7.1 shows how the inclusive jet production cross-section varies with $\alpha_S(\mu_r)$.

7.1 Sensitivity of Measurements to $\alpha_s(M_Z)$

For a fixed choice of μ_r and μ_f , different input values of $\alpha_s(M_Z)$ to a PDF set will lead to different theory predictions of the differential cross-section distribution. This will give an estimate of the sensitivity of the theory predictions to the varying input value of $\alpha_s(M_Z)$. A comparison of the measured spectrum with the theory predictions obtained using all $\alpha_s(M_Z)$ inputs will give a hint of the input value of $\alpha_s(M_Z)$ for which the theory distribution has the closest matching with data. In this section, the sensitivity of the inclusive differential jet event cross-sections and cross-section ratio, R_{32} to varying input values of $\alpha_s(M_Z)$ for different PDF sets is demonstrated by plotting the ratios of data over theory predictions with central value of $\alpha_s(M_Z)$.

Figures 7.1, 7.2 and 7.3 present the ratio of data to the theory predictions, corrected for NP effects, for all variations in $\alpha_s(M_Z)$ available for the PDF sets CT10, CT14, MSTW2008, MMHT2014 and NNPDF2.3 at NLO evolution order as specified in Table 6.1, for inclusive 2-jet event cross-sections, inclusive 3-jet events cross-sections and ratio R_{32} respectively. The $\alpha_s(M_Z)$ value is varied in the range 0.112-0.127, 0.111-123, 0.110-0.130, 0.108-0.128 and 0.114-0.124 in steps of 0.001 for the CT10, CT14, MSTW2008, MMHT2014 and NNPDF2.3PDF sets, respectively. The error bars correspond to the total experimental uncertainty derived as quadratic sum from all uncertainty sources. The theory predictions are also corrected for EW effects for inclusive 2-jet events cross-section. A small slope increasing with $H_{T,2}/2$ is visible for most PDFs in both cross-sections. This effect is largely cancelled in

the cross-section ratio. R_{32} exhibits a flat behaviour with respect to the predictions for all five PDF sets in the whole range of $H_{T,2}/2$ up to 1680 GeV. Therefore, these data can be used to determine the strong coupling constant, although only up to 1 TeV for the cross-sections as long as electroweak corrections are not taken into account.

Moreover, the different sensitivity to $\alpha_s(M_Z)$ caused by the leading power in α_S in the expansion of the 2-jet inclusive ($\propto \alpha_S^2$) and the 3-jet inclusive cross-section ($\propto \alpha_S^3$), and their ratio ($\propto \alpha_S^1$) is clearly visible from the spread between the calculations for the smallest and largest value of $\alpha_s(M_Z)$ within the same PDF set when passing through Figures 7.1–7.3. This also demonstrates the potential of ratios R_{mn} with $m-n > 1$.

7.2 Determination of $\alpha_s(M_Z)$

As discussed in the previous section, the measured inclusive 2-jet and 3-jet event cross-sections and their ratio R_{32} can be used for a determination of the strong coupling constant $\alpha_s(M_Z)$. To extract the value of $\alpha_s(M_Z)$, a general fit procedure [1, 91, 115] has been followed and is described in the following section.

7.2.1 Fitting Procedure

The value of $\alpha_s(M_Z)$ is determined by minimizing the chi-square (χ^2) between the experimental measurements and the theoretical predictions. The χ^2 is given by the following equation :

$$\chi^2 = M^T C^{-1} M \quad (7.2)$$




Figure 7.1: Ratio of the measured inclusive 2-jet differential cross-section to theory predictions using the CT10 (top left), the CT14 (top right), the MSTW2008 (middle left), the MMHT2014 (middle right) and NNPDF2.3 (bottom) NLO PDF sets for a series of values of $\alpha_s(M_Z)$. The $\alpha_s(M_Z)$ value is varied in the range 0.112-0.127, 0.111-0.123, 0.110-0.130, 0.108-0.128 and 0.114-0.124 in steps of 0.001 for the CT10, CT14, MSTW2008, MMHT2014 and NNPDF2.3 NLO PDF sets, respectively. The error bars correspond to the total experimental uncertainty. The theory predictions are corrected for non-perturbative (NP) and electroweak (EW) effects.




Figure 7.2: Ratio of the measured inclusive 3-jet differential cross-section to theory predictions using the CT10 (top left), the CT14 (top right), the MSTW2008 (middle left), the MMHT2014 (middle right) and NNPDF2.3 (bottom) NLO PDF sets for a series of values of $\alpha_s(M_Z)$. The $\alpha_s(M_Z)$ value is varied in the range 0.112-0.127, 0.111-0.123, 0.110-0.130, 0.108-0.128 and 0.114-0.124 in steps of 0.001 for the CT10, CT14, MSTW2008, MMHT2014 and NNPDF2.3 NLO PDF sets, respectively. The error bars correspond to the total experimental uncertainty. The theory predictions are corrected for non-perturbative (NP) effects.




Figure 7.3: Ratio of the measured cross-section ratio, R_{32} to theory predictions using the CT10 (top left), the CT14 (top right), the MSTW2008 (middle left), the MMHT2014 (middle right) and NNPDF2.3 (bottom) NLO PDF sets for a series of values of $\alpha_s(M_Z)$. The $\alpha_s(M_Z)$ value is varied in the range 0.112-0.127, 0.111-123, 0.110-0.130, 0.108-0.128 and 0.114-0.124 in steps of 0.001 for the CT10, CT14, MSTW2008, MMHT2014 and NNPDF2.3 NLO PDF sets, respectively. The error bars correspond to the total experimental uncertainty. The theory predictions are corrected for non-perturbative (NP) effects.

where M is the vector of the differences between the data (D^i) and the theoretical values (T^i) in each bin i ,

$$M^i = D^i - T^i \quad (7.3)$$

and C is the covariance matrix including all experimental uncertainties as described in Sec. 5.6 and some theoretical uncertainties discussed in Sec. 6.2. The covariance matrix $C = C_{\text{exp}} + C_{\text{theo}}$ is defined as the sum of covariances of experimental and theoretical sources of uncertainty as follows :

$$C_{\text{exp}} = \text{Cov}^{\text{ExpStat}} + \sum \text{Cov}^{\text{JEC}} + \text{Cov}^{\text{Unfolding}} + \text{Cov}^{\text{Lumi}} + \text{Cov}^{\text{Residual}} \quad (7.4)$$

$$C_{\text{theo}} = \text{Cov}^{\text{TheoStat}} + \text{Cov}^{\text{NP}} + \text{Cov}^{\text{PDF}} \quad (7.5)$$

where the labelled covariance matrices account for the following effects:

- $\text{Cov}^{\text{ExpStat}}$: statistical uncertainty of the data including correlations introduced by the unfolding
- Cov^{JEC} : the jet energy corrections (JEC) systematic uncertainty
- $\text{Cov}^{\text{Unfolding}}$: the unfolding systematic uncertainty including the resolution (JER) and model dependence
- Cov^{Lumi} : the luminosity uncertainty
- $\text{Cov}^{\text{Residual}}$: a residual uncorrelated systematic uncertainty summarizing individual causes such as small trigger and identification inefficiencies, time dependence of the jet p_T resolution, and uncertainty on the trigger prescale factors
- $\text{Cov}^{\text{TheoStat}}$: statistical uncertainty caused by numerical integrations in the cross-section computations

- Cov^{NP} : the systematic uncertainty of the non-perturbative (NP) corrections
- Cov^{PDF} : the PDF uncertainties

While taking the differences between theory and data, the treatment of experimental and theoretical systematic uncertainties is crucial. The Unfolding, JEC, Lumi and PDF and NP systematic uncertainties are treated as 100% correlated among $H_{T,2}/2$ bins. If δ_i is the total uncertainty on the differential cross-section, for the i -th $H_{T,2}/2$ bin, for any of these fully correlated sources, then the i, j -th element of the corresponding covariance matrix is given by $\text{COV}_{ij} = \delta_i \times \delta_j$. The JEC, unfolding, and luminosity uncertainties are treated as multiplicative to avoid the statistical bias that arises when estimating uncertainties from data. In fits of the ratio R_{32} , the luminosity and residual uncorrelated uncertainties cancel completely. Partial cancellations between the other sources of uncertainty are taken into account in the fit.

The evaluation of PDF uncertainty depends on the individual PDF set as already discussed in Sec. 6.2.2. The PDF covariance matrix construction varies among different PDF sets. The CT10, CT14, MMHT2014 and MSTW2008 NLO PDF sets employ the eigenvector method to evaluate the PDF uncertainties as explained in Sec. 6.2.2. The number of eigenvectors (N_{ev}) with two PDF members per eigenvector for CT10, CT14, MMHT2014 and MSTW2008 NLO PDF sets are 26, 28, 25 and 20, respectively. The NNPDF2.3 PDF set comes with hundred different replicas (N_{rep}) instead of different eigenvectors, as for CT10 or CT14 PDF sets. The mean uncertainty is calculated as average uncertainty from 100 different replicas. Following the prescription given in [123], the PDF uncertainty is calculated as :

$$(\Delta X)^2 = \frac{1}{N_{\text{rep}} - 1} \sum_{k=1}^{N_{\text{rep}}} [X_k - \langle X_k \rangle]^2 \quad (7.6)$$

where ΔX is the uncertainty on predicted differential cross-section, X_k is the differ-

ential cross-section for k -th replica and $\langle X_k \rangle$ is the average differential cross-section from all the replicas.

Scale uncertainties of the pQCD predictions are taken into account by employing the offset method, i.e. by performing separate fits with varying scale factors as described in the Sec. 6.2.1. The largest upwards and downwards deviations from the default factors are defined as the uncertainty. At NLO such scale variations predominantly lead to smaller cross-sections and also a smaller ratio R_{32} as visible in Fig. 6.5. As a consequence the scale uncertainty in fits is equally asymmetric, where smaller cross-sections or ratios are compensated by an increase in the fitted value for $\alpha_s(M_Z)$.

7.2.2 Fit Results

To determine the value of the strong coupling constant at the scale of the Z boson mass $\alpha_s(M_Z)$, fits to the differential inclusive 2-jet and 3-jet events cross-sections are performed using five different NLO PDF sets : CT10, CT14, MSTW2008, MMHT2014 and NNPDF2.3. The range in $H_{\mathrm{T},2}/2$ is restricted to be between 300 GeV and 1 TeV to avoid the region close to the minimal p_{T} threshold of 150 GeV for each jet at low p_{T} and the onset of electroweak effects at high $H_{\mathrm{T},2}/2$, which are available for the dijet case only. The $\alpha_s(M_Z)$ results obtained from a simultaneous fit to all 19 $H_{\mathrm{T},2}/2$ bins in the above mentioned range are reported in Table 7.1. For comparison, a simultaneous fit to both cross-sections ignoring any correlations, and a fit to the cross-section ratio R_{32} , fully accounting for correlations is also performed and the results are tabulated in Table 7.2. The electroweak effects are assumed to cancel in the ratio as do the luminosity and the uncorrelated uncertainty.

All cross-section fits give compatible values for $\alpha_s(M_Z)$ in the range of 0.115–0.118 whereas for the ratio R_{32} somewhat smaller values are obtained. But for individual cross-sections, χ^2/n_{dof} values are small as compared to the cross-section

ratio R_{32} . A possible explanation is an overestimation of the residual uncorrelated uncertainty of 1% that is cancelled for R_{32} . If the fits are repeated with an assumed uncertainty of 0.25% instead, the χ^2/n_{dof} values lie around unity while the $\alpha_s(M_Z)$ values are still compatible with the previous results but with slightly reduced uncertainties.

Table 7.1: Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross-sections using five PDF sets at NLO. Only total uncertainties without scale variations are quoted. The results are obtained from a simultaneous fit to all 19 $H_{\text{T},2}/2$ bins in the restricted range of $0.3 < H_{\text{T},2}/2 < 1.0$ TeV.

PDF set	Inclusive 2-jets			Inclusive 3-jets		
	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}
CT10	0.1174	0.0032	3.0/18	0.1169	0.0027	5.4/18
CT14	0.1160	0.0035	3.5/18	0.1159	0.0031	6.1/18
MSTW2008	0.1159	0.0025	5.3/18	0.1161	0.0021	6.7/18
MMHT2014	0.1165	0.0034	5.9/18	0.1166	0.0025	7.1/18
NNPDF2.3	0.1183	0.0025	9.7/18	0.1179	0.0021	9.1/18

Table 7.2: Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross-sections simultaneously and from their ratio R_{32} using five PDF sets at NLO. Only total uncertainties without scale variations are quoted. The results are obtained from a simultaneous fit to all 38 (left) and 19 (right) $H_{\text{T},2}/2$ bins in the restricted range of $0.3 < H_{\text{T},2}/2 < 1.0$ TeV. For comparison, correlations between the two cross-sections are neglected in the simultaneous fit on the left, but fully taken into account in the ratio fit on the right.

PDF set	Inclusive 2- and 3-jets			R_{32}		
	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}
CT10	0.1170	0.0026	8.2/37	0.1141	0.0028	19./18
CT14	0.1161	0.0029	9.1/37	0.1139	0.0032	15./18
MSTW2008	0.1161	0.0021	11./37	0.1150	0.0023	21./18
MMHT2014	0.1168	0.0025	11./37	0.1142	0.0022	19./18
NNPDF2.3	0.1188	0.0019	15./37	0.1184	0.0021	12./18

To investigate how the electroweak (EW) corrections affect the fit results for $\alpha_s(M_Z)$, the range in $H_{\text{T},2}/2$ is extended to $0.3 < H_{\text{T},2}/2 < 1.68$ TeV. $\alpha_s(M_Z)$ values are obtained from fits to the inclusive 2-jet event cross-section in this range with or without EW correction factors and the results are presented in Table 7.3. The largest impact is a reduction in χ^2/n_{dof} , which indicates a better agreement when

EW effects are included. In addition, a tendency to slightly smaller $\alpha_s(M_Z)$ values is observed without the EW corrections. For the ratio R_{32} , it is expected that these effects are much reduced.

Table 7.3: Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet event cross-section using five PDF sets at NLO without (left) and with (right) electroweak (EW) corrections. Only total uncertainties without scale variations are quoted. The results are obtained from a simultaneous fit to all 29 $H_{T,2}/2$ bins in the range of $0.3 < H_{T,2}/2 < 1.68$ TeV.

PDF set	without EW			with EW		
	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}
CT10	0.1163	0.0034	15./28	0.1165	0.0032	14./28
CT14	0.1137	0.0033	24./28	0.1144	0.0033	17./28
MSTW2008	0.1093	0.0028	27./28	0.1133	0.0023	19./28
MMHT2014	0.1127	0.0032	32./28	0.1141	0.0032	21./28
NNPDF2.3	0.1162	0.0024	31./28	0.1168	0.0024	23./28

From Fig. 7.3 follows that only the PDF sets MSTW2008 and MMHT2014 provide a large enough range in $\alpha_s(M_Z)$ values to ensure fits without extrapolation. The other three PDF sets are at the limit such that reliable fits cannot be performed for all scale settings and/or bins in scale $Q = H_{T,2}/2$. Since many systematic uncertainties cancel completely or partially in the cross-section ratio R_{32} as compared to the individual cross-sections, R_{32} is used mainly to determine the value of $\alpha_s(M_Z)$. Table 7.4 give the complete results for MSTW2008 and MMHT2014 for the full range in $H_{T,2}/2$ of 300 GeV up to 1.68 TeV along with the corresponding components of PDF, scale, NP and total experimental except scale uncertainties are shown. In contrast to fits at NLO using cross-sections where the scale uncertainty recipe usually leads to a very asymmetric behaviour with larger downward uncertainties in the case, this is inverted for the fits to the cross-section ratio R_{32} . The scale uncertainty is the most dominant source of total uncertainty on $\alpha_s(M_Z)$. These values are determined with the central renormalization and factorization scales i.e. $\mu_r = \mu_f = H_{T,2}/2$. The values are also determined for the six scale factor combinations for the two PDF sets MSTW2008 and MMHT2014 and results are shown in Table 7.5.

The uncertainty decomposition for $\alpha_s(M_Z)$ determined from cross-section ratio R_{32} is performed in four sub-ranges of $H_{T,2}/2$ and the results are shown in Table 7.6. The statistical uncertainty of the NLO computation is negligible in comparison to any of the other sources of uncertainty. Electroweak corrections, significant only at high $H_{T,2}/2$, are assumed to cancel between the numerator and denominator.

Using the MSTW2008 PDF set, which dates from before the LHC start, the strong coupling constant finally is determined to

$$\begin{aligned} \alpha_s(M_Z) &= 0.1150 \pm 0.0010 \text{ (exp)} \pm 0.0013 \text{ (PDF)} \pm 0.0015 \text{ (NP)} {}^{+0.0050}_{-0.0000} \text{ (scale)} \\ &= 0.1150 \pm 0.0023 \text{ (all except scale)} {}^{+0.0050}_{-0.0000} \text{ (scale)} \end{aligned} \tag{7.7}$$

The MMHT2014 PDF set, although using LHC jet data to determine the PDF parameters, leads to a very similar result of

$$\begin{aligned} \alpha_s(M_Z) &= 0.1142 \pm 0.0010 \text{ (exp)} \pm 0.0013 \text{ (PDF)} \pm 0.0014 \text{ (NP)} {}^{+0.0049}_{-0.0006} \text{ (scale)} \\ &= 0.1142 \pm 0.0022 \text{ (all except scale)} {}^{+0.0049}_{-0.0006} \text{ (scale)} \end{aligned} \tag{7.8}$$

7.3 Running of the Strong Coupling Constant

The value of the strong coupling constant α_S depends on the energy scale Q and it decreases with the increase of scale Q . To study this dependence, the determination of α_S is carried out at different energies. The procedure to extract $\alpha_S(Q)$ is same as the one followed for the $\alpha_s(M_Z)$. To have different energy scales, the fitted $H_{T,2}/2$ range 300 - 1680 GeV is divided into four different sub-ranges as shown by the first column in Table 7.7. Each of the $H_{T,2}/2$ range is associated with a scale Q , which

Table 7.4: Determination of $\alpha_s(M_Z)$ from the ratio R_{32} using the two most compatible PDF sets MSTW2008 and MMHT2014 at NLO along with the corresponding components of PDF, scale, NP and total (except scale) experimental uncertainties. The results are obtained from a simultaneous fit to all 29 $H_{T,2}/2$ bins in the full range of $0.3 < H_{T,2}/2 < 1.68$ TeV.

PDF set	$\alpha_s(M_Z)$	exp	PDF	NP	all exc.	scale	χ^2/n_{dof}
MSTW2008	0.1150	± 0.0010	± 0.0013	± 0.0015	± 0.0023	$^{+0.0050}_{-0.0000}$	26./28
MMHT2014	0.1142	± 0.0010	± 0.0013	± 0.0014	± 0.0022	$^{+0.0049}_{-0.0006}$	24./28

Table 7.5: Determination of $\alpha_s(M_Z)$ from the ratio R_{32} in the $H_{T,2}/2$ range from 0.3 up to 1.68 TeV at the central scale and for the six scale factor combinations for the two PDF sets MSTW2008 and MMHT2014.

$\mu_r/H_{T,2}/2$	$\mu_f/H_{T,2}/2$	MSTW2008		MMHT2014	
		$\alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	χ^2/n_{dof}
1	1	0.1150	26./28	0.1142	24./28
1/2	1/2	0.1165	77./28	0.1160	73./28
2	2	0.1200	18./28	0.1191	18./28
1/2	1	0.1150	53./28	0.1136	48./28
1	1/2	0.1150	30./28	0.1142	28./28
1	2	0.1155	23./28	0.1147	22./28
2	1	0.1180	19./28	0.1175	19./28

Table 7.6: Uncertainty decomposition for $\alpha_s(M_Z)$ from the determination of α_s from the jet event rate R_{32} in bins of $H_{T,2}/2$. The statistical uncertainty of the NLO computation is negligible in comparison to any of the other sources of uncertainty. Electroweak corrections, significant only at high $H_{T,2}/2$, are assumed to cancel between the numerator and denominator.

$H_{T,2}/2$ (GeV)	MSTW2008					MMHT2014				
	$\alpha_s(M_Z)$	exp	PDF	NP	scale	$\alpha_s(M_Z)$	exp	PDF	NP	scale
300-420	0.1157	± 0.0015	± 0.0014	± 0.0019	$^{+0.0053}_{-0.0000}$	0.1158	± 0.0014	± 0.0010	± 0.0019	$^{+0.0052}_{-0.0000}$
420-600	0.1153	± 0.0011	± 0.0014	± 0.0018	$^{+0.0057}_{-0.0000}$	0.1154	± 0.0011	± 0.0012	± 0.0017	$^{+0.0056}_{-0.0000}$
600-1000	0.1134	± 0.0013	± 0.0016	± 0.0019	$^{+0.0052}_{-0.0000}$	0.1140	± 0.0012	± 0.0012	± 0.0018	$^{+0.0045}_{-0.0000}$
1000-1680	0.1147	± 0.0029	± 0.0017	± 0.0018	$^{+0.0063}_{-0.0011}$	0.1154	± 0.0025	± 0.0014	± 0.0015	$^{+0.0056}_{-0.0011}$
300-1680	0.1150	± 0.0010	± 0.0013	± 0.0015	$^{+0.0050}_{-0.0000}$	0.1142	± 0.0010	± 0.0013	± 0.0014	$^{+0.0049}_{-0.0006}$

is the differential cross-section weighted average $H_{T,2}/2$ scale from the inclusive 2-jet calculations and integrated over all the measured $H_{T,2}/2$ bins contributing to that given $H_{T,2}/2$ range. Let N_{bin}^j be the total number of measured $H_{T,2}/2$ bins contributing to the j -th $H_{T,2}/2$ range, then the corresponding scale Q_j , shown in second column of Table 7.7, is calculated as :

$$Q_j = \frac{\sum_{i=1}^{N_{bin}^j} H_{T,2}^i \left[\frac{d\sigma}{d(H_{T,2}/2)} \right]^i}{\sum_{i=1}^{N_{bin}^j} \left[\frac{d\sigma}{d(H_{T,2}/2)} \right]^i} \quad (7.9)$$

The value of $\alpha_s(M_Z)$ is extracted in each $H_{T,2}/2$ range. These extracted $\alpha_s(M_Z)$ values are evolved to the corresponding values $\alpha_s(Q)$ and are quoted in Table 7.7 along with the extracted $\alpha_s(M_Z)$ values and the total uncertainty. The evolution is performed for five flavours at 2-loop order with the RUNDEC program [124, 125]. The obtained $\alpha_s(Q)$ points (black solid circles) are shown as a function of scale Q in Fig. 7.4. The black solid line and the yellow uncertainty band are evolved using $\alpha_s(M_Z) = 0.1150 \pm 0.0023$ (all except scale) $^{+0.0050}_{-0.0000}$ (scale) obtained using MSTW2008 NLO PDF set. The world average [18] (dashed line) and results from other measurements of the CMS [1, 91, 105, 114, 115], ATLAS [116], D0 [117, 118], H1 [119, 120], and ZEUS [121] experiments are also imposed. The current measurement is in very good agreement within the uncertainty with other results obtained by previous experiments as well as with the world average value of $\alpha_s(M_Z) = 0.1181 \pm 0.0011$ derived in Ref. [18].

Table 7.7: Evolution of the strong coupling constant between the scale of the Z boson mass and the cross-section averaged $H_{T,2}/2$ scale $\langle Q \rangle$ for the separate determinations in each respective fit range. The evolution is performed for five flavours at 2-loop order with the RUNDEC program [124, 125].

$H_{T,2}/2$ (GeV)	$\langle Q \rangle$ (GeV)	$\alpha_s(M_Z)$	$\alpha_S(Q)$	No. of data points	χ^2/n_{dof}
300-420	340	$0.1157^{+0.0060}_{-0.0030}$	$0.0969^{+0.0041}_{-0.0021}$	4	2.8/3
420-600	476	$0.1153^{+0.0062}_{-0.0025}$	$0.0928^{+0.0039}_{-0.0016}$	6	6.1/5
600-1000	685	$0.1134^{+0.0059}_{-0.0028}$	$0.0879^{+0.0035}_{-0.0017}$	9	7.1/8
1000-1680	1114	$0.1147^{+0.0074}_{-0.0040}$	$0.0841^{+0.0039}_{-0.0021}$	10	5.4/9




Figure 7.4: The running $\alpha_S(Q)$ as a function of the scale Q is shown as obtained by using the MSTW2008 NLO PDF set. The solid line and the uncertainty band are drawn by evolving the extracted $\alpha_s(M_Z)$ values using the 2-loop 5-flavour renormalization group equations as implemented in RUNDEC [124, 125]. The dashed line represents the evolution of the world average [18] and the black circles correspond to the $\alpha_S(Q)$ determinations presented in Table 7.7. Results from other measurements of CMS [1, 91, 105, 114, 115], ATLAS [116], D0 [117, 118], H1 [119, 120], and ZEUS [121] are superimposed.

Chapter 8

Summary

Inclusive multijet production cross-section measured precisely in terms of jet transverse momentum is one of the important observables in understanding physics at hadron colliders. It provides the essential information about the structure of parton through parton distribution functions (PDFs) and the precise measurement of the strong coupling constant α_S . The value of the strong coupling constant at the scale of the Z boson mass $\alpha_s(M_Z)$ can be determined using cross-section ratio instead of individual cross-sections because many uncertainties of theoretical and experimental origin cancel between numerator and denominator which reduces the dependence on PDFs, renormalization and factorization scales, luminosity etc.

In this thesis, a measurement of the inclusive 2-jet and 3-jet event cross-sections as well as the cross-section ratio R_{32} has been presented. The data sample has been collected from proton-proton collisions recorded with the CMS detector at a centre-of-mass energy of 8 TeV and corresponds to an integrated luminosity of 19.7 fb^{-1} . The jets are reconstructed with the anti- k_t clustering algorithm for a jet size parameter $R = 0.7$. The inclusive 2-jet and 3-jet event cross-sections are measured differentially as a function of the average transverse momentum of the two leading jets, referred as $H_{\text{T},2}/2$. The ratio R_{32} is obtained by dividing the differential cross-sections of inclusive 3-jet events to that of inclusive 2-jet one in

each bin of $H_{\mathrm{T},2}/2$. An appropriate selection criteria has been designed for choosing the best events for analysis. The measurements are performed at a central rapidity of $|y| < 2.5$ in a range of $0.3 < H_{\mathrm{T},2}/2 < 2.0 \text{ TeV}$ for inclusive 2-jet event cross-sections and $0.3 < H_{\mathrm{T},2}/2 < 1.68 \text{ TeV}$ for inclusive 3-jet event cross-sections and ratio R_{32} .

The measured cross-sections after correcting for detector effects by using an iterative unfolding procedure are compared to the perturbative QCD predictions computed, using NLOJET++ program, at next-to-leading order (NLO) accuracy and complemented with non-perturbative (NP) corrections that are important at low $H_{\mathrm{T},2}/2$. The data are found to be well described by NLO calculations. The upwards trend observed in the inclusive 2-jet and 3-jet data at high $H_{\mathrm{T},2}/2$ in comparison to the prediction at NLO QCD, is explained by the onset of electroweak (EW) corrections in the 2-jet case. For the 3-jet event cross-sections these corrections have not yet been computed yet. In the 3-jet to 2-jet cross-section ratio R_{32} , the EW corrections are assumed to cancel. In fact, NLO QCD provides an adequate description of R_{32} in the accessible range of $H_{\mathrm{T},2}/2$. In contrast, leading order (LO) tree-level Monte Carlo (MC) predictions obtained using MADGRAPH5 event generator interfaced to PYTHIA6 exhibit significant deviations. The sources of experimental and theoretical uncertainties are studied in details. The experimental uncertainty ranges from 4 to 32% for inclusive 2-jet event cross-sections, from 4 to 28% for 3-jet event cross-sections and from 1 to 28% for cross-section ratio R_{32} . It is dominated by the uncertainty due to the jet energy corrections (JEC) at lower $H_{\mathrm{T},2}/2$ values and by statistical uncertainty at higher $H_{\mathrm{T},2}/2$ values. The theoretical uncertainty ranges from 3 to 30% and 5 to 34% for inclusive 2-jet and 3-jet event cross-sections respectively and from 3 to 11% for ratio R_{32} . The PDF uncertainty derived with the CT10-NLO PDF set is the dominant source of theoretical uncertainty.

The inclusive multijet cross-sections being proportional to the powers of the strong coupling constant α_S ($\sigma_{n\text{-jet}} \propto \alpha_S^n$) are used to extract the value of the strong coupling constant at the scale of the Z boson mass $\alpha_s(M_Z)$. In cross-section ratio R_{32} which proportional to α_S , many uncertainties and PDF dependencies largely cancel and hence becomes the better tool to extract the value of $\alpha_s(M_Z)$. In this thesis, a fit of the ratio of the inclusive 3-jet event cross-section to that of 2-jet, R_{32} in the range $0.3 < H_{T,2}/2 < 1.68$ TeV using the MSTW2008 PDF set gives :

$$\begin{aligned}\alpha_s(M_Z) &= 0.1150 \pm 0.0010 \text{ (exp)} \pm 0.0013 \text{ (PDF)} \pm 0.0015 \text{ (NP)} {}^{+0.0050}_{-0.0000} \text{ (scale)} \\ &= 0.1150 \pm 0.0023 \text{ (all except scale)} {}^{+0.0050}_{-0.0000} \text{ (scale)}\end{aligned}$$

Very similar results are obtained using the MMHT2014 PDF set which gives :

$$\begin{aligned}\alpha_s(M_Z) &= 0.1142 \pm 0.0010 \text{ (exp)} \pm 0.0013 \text{ (PDF)} \pm 0.0014 \text{ (NP)} {}^{+0.0049}_{-0.0006} \text{ (scale)} \\ &= 0.1142 \pm 0.0022 \text{ (all except scale)} {}^{+0.0049}_{-0.0006} \text{ (scale)}\end{aligned}$$

The equally compatible values of $\alpha_s(M_Z)$ are determined with separate fits to the inclusive 2-jet and 3-jet event cross-sections provided the range in $H_{T,2}/2$ is restricted to $0.3 < H_{T,2}/2 < 1.0$ TeV. The extracted $\alpha_s(M_Z)$ values in sub-ranges of $H_{T,2}/2$ are evolved to corresponding $\alpha_S(Q)$ along with the error bars at different scales Q . The current measurement of $\alpha_s(M_Z)$ and the running of $\alpha_S(Q)$ as a function of Q is in well agreement within uncertainties with the world average value of $\alpha_s(M_Z) = 0.1181 \pm 0.0011$ [18] and already existing determinations performed by the CMS and other experiments.

The inclusion of the EW corrections in inclusive 2-jet event cross-sections become relevant at $H_{T,2}/2$ beyond 1 TeV. Their availability for 3-jet one and hence cross-section ratio R_{32} can improve the precision of the measurement of $\alpha_s(M_Z)$. Also as the theoretical calculations will be available for inclusive 4-jet event cross-sections, the various cross-section ratios such as $R_{43} \propto \alpha_S^1$ and $R_{42} \propto \alpha_S^2$ can be measured to extract the value of the strong coupling constant more precisely. Currently LHC is running at high center-of-mass energy of 13 TeV delivering a higher

instantaneous luminosity and this makes possible to access the extended phase space and perform the measurements with more accuracy.

Appendix A

A.1 Cross-section Ratio, R_{32}

Table A.1: Differential cross-sections ($\times 10^{-3}(\text{pb}/\text{GeV})$) and cross-section ratio R_{32} at detector level in each bin of $H_{T,2}/2$, along with statistical uncertainty (in %).

Bin	2-jet	Stat.	3-jet	Stat.	Ratio	Stat.
	cross-section	unc.	cross-section	unc.	R_{32}	unc.
300 - 330	29772.726	0.211	2640.629	0.707	0.089	+0.665 -0.661
330 - 360	16792.917	0.231	1773.485	0.704	0.106	+0.523 -0.521
360 - 390	9889.326	0.182	1176.544	0.526	0.119	+0.485 -0.483
390 - 420	5976.777	0.179	778.034	0.492	0.130	+0.206 -0.206
420 - 450	3731.760	0.067	522.624	0.180	0.140	+0.167 -0.167
450 - 480	2398.741	0.084	357.622	0.217	0.149	+0.201 -0.200
480 - 510	1570.192	0.104	246.051	0.262	0.157	+0.241 -0.241
510 - 540	1048.665	0.127	171.080	0.314	0.163	+0.288 -0.287
540 - 570	713.042	0.154	119.566	0.376	0.168	+0.344 -0.343
570 - 600	490.776	0.186	84.798	0.447	0.173	+0.407 -0.406
600 - 640	325.046	0.198	57.463	0.470	0.177	+0.427 -0.426
640 - 680	205.727	0.248	37.282	0.583	0.181	+0.529 -0.527
680 - 720	133.674	0.308	24.859	0.714	0.186	+0.646 -0.643
720 - 760	87.911	0.380	16.560	0.875	0.188	+0.791 -0.786
760 - 800	58.657	0.465	11.056	1.071	0.188	+0.968 -0.961
800 - 850	38.106	0.516	7.318	1.178	0.192	+1.063 -1.054
850 - 900	23.587	0.656	4.600	1.485	0.195	+1.339 -1.326
900 - 950	15.130	0.819	2.896	1.872	0.191	+1.694 -1.672
950 - 1000	9.696	1.023	1.812	2.366	0.187	+2.151 -2.116
1000 - 1060	6.026	1.185	1.186	2.670	0.197	+2.414 -2.371
1060 - 1120	3.668	1.518	0.716	3.436	0.195	+3.118 -3.046
1120 - 1180	2.327	1.906	0.437	4.398	0.188	+4.024 -3.903
1180 - 1250	1.419	2.260	0.265	5.227	0.187	+4.798 -4.627
1250 - 1320	0.853	2.915	0.165	6.623	0.194	+6.080 -5.811
1320 - 1390	0.477	3.898	0.080	9.492	0.169	+8.951 -8.355
1390 - 1460	0.263	5.249	0.042	13.131	0.160	+12.619 -11.449
1460 - 1530	0.192	6.143	0.029	15.811	0.151	+15.437 -13.698
1530 - 1600	0.104	8.362	0.021	18.570	0.203	+17.571 -15.536
1600 - 1680	0.060	10.314	0.009	26.726	0.149	+27.132 -22.170

A.2 Individual Sources of Jet Energy Correction Uncertainties




Figure A.1: The fractional jet energy correction (JEC) uncertainties from individual sources are shown for inclusive 2-jet (top) and 3-jet events cross-sections (middle); and cross-section ratio R_{32} (bottom). On left, JEC uncertainties are evaluated from `AbsoluteStat` (blue), `AbsoluteScale` (red), `AbsoluteMPFBias` (green) and `Fragmentation` (pink) sources whereas on right, these are evaluated from `SinglePionECAL` (blue), `SinglePionHCAL` (red) and `FlavorQCD` (green) sources.




Figure A.2: The fractional jet energy correction (JEC) uncertainties from individual sources are shown for inclusive 2-jet (top) and 3-jet events cross-sections (middle); and cross-section ratio R_{32} (bottom). On left, JEC uncertainties are evaluated from RelativeJEREC1 (blue), RelativeJEREC2 (red), RelativePtBB (green) and RelativeJERHF (pink) sources whereas on right, these are evaluated from RelativePtEC1 (blue), RelativePtEC2 (red), RelativeFSR (green) and RelativePtHF (pink) sources.




Figure A.3: The fractional jet energy correction (JEC) uncertainties from individual sources are shown for inclusive 2-jet (top) and 3-jet events cross-sections (middle); and cross-section ratio R_{32} (bottom). On left, JEC uncertainties are evaluated from RelativeStatFSR (blue), RelativeStatEC2 (red) and RelativeStatHF (green) sources whereas on right, these are evaluated from PileUpDataMC (blue), PileUpPtRef (red), PileUpPtBB (green), PileUpPtEC1 (pink), PileUpPtEC2 (orange) and PileUpPtHF (black) sources.

A.3 Experimental Uncertainties

Table A.2: Experimental uncertainties (in %), from all sources as well as the total uncertainty, affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 2-jet events.

Bin	Statistical	JEC	Unfolding	Lumi	Residual	Total
300 - 330	0.242	+2.612 -2.565	+0.948 -0.928	2.6	1.0	+3.942 -3.906
330 - 360	0.258	+2.507 -2.473	+0.976 -0.969	2.6	1.0	+3.882 -3.858
360 - 390	0.202	+2.504 -2.465	+0.779 -0.783	2.6	1.0	+3.831 -3.807
390 - 420	0.193	+2.363 -2.381	+0.905 -0.904	2.6	1.0	+3.768 -3.780
420 - 450	0.084	+2.448 -2.422	+0.904 -0.895	2.6	1.0	+3.818 -3.799
450 - 480	0.096	+2.440 -2.352	+0.797 -0.795	2.6	1.0	+3.789 -3.733
480 - 510	0.107	+2.427 -2.406	+0.728 -0.715	2.6	1.0	+3.767 -3.751
510 - 540	0.128	+2.425 -2.395	+0.835 -0.862	2.6	1.0	+3.789 -3.775
540 - 570	0.154	+2.425 -2.376	+0.687 -0.674	2.6	1.0	+3.760 -3.726
570 - 600	0.180	+2.497 -2.474	+0.839 -0.827	2.6	1.0	+3.838 -3.820
600 - 640	0.209	+2.495 -2.491	+0.744 -0.743	2.6	1.0	+3.819 -3.816
640 - 680	0.264	+2.582 -2.545	+0.912 -0.912	2.6	1.0	+3.915 -3.891
680 - 720	0.320	+2.691 -2.574	+0.763 -0.756	2.6	1.0	+3.961 -3.880
720 - 760	0.387	+2.690 -2.755	+0.705 -0.712	2.6	1.0	+3.955 -4.001
760 - 800	0.465	+2.858 -2.846	+0.859 -0.846	2.6	1.0	+4.109 -4.098
800 - 850	0.548	+2.889 -2.913	+0.783 -0.787	2.6	1.0	+4.126 -4.143
850 - 900	0.698	+3.145 -3.102	+0.961 -0.958	2.6	1.0	+4.366 -4.334
900 - 950	0.847	+3.298 -3.233	+0.828 -0.829	2.6	1.0	+4.476 -4.429
950 - 1000	1.041	+3.291 -3.330	+0.895 -0.872	2.6	1.0	+4.525 -4.549
1000 - 1060	1.268	+3.598 -3.569	+0.945 -0.956	2.6	1.0	+4.817 -4.798
1060 - 1120	1.611	+3.759 -3.756	+0.970 -0.967	2.6	1.0	+5.043 -5.040
1120 - 1180	1.985	+4.154 -4.053	+1.089 -1.080	2.6	1.0	+5.490 -5.413
1180 - 1250	2.406	+4.251 -4.313	+1.062 -1.070	2.6	1.0	+5.722 -5.770
1250 - 1320	3.101	+4.696 -4.624	+1.151 -1.144	2.6	1.0	+6.384 -6.330
1320 - 1390	4.157	+4.934 -4.979	+1.343 -1.341	2.6	1.0	+7.155 -7.186
1390 - 1460	5.270	+5.148 -5.104	+1.185 -1.177	2.6	1.0	+7.965 -7.936
1460 - 1530	6.360	+5.890 -5.652	+1.405 -1.406	2.6	1.0	+9.213 -9.063
1530 - 1600	8.183	+5.924 -6.311	+1.598 -1.590	2.6	1.0	+10.601 -10.821
1600 - 1680	10.630	+5.969 -5.655	+1.607 -1.592	2.6	1.0	+12.608 -12.461
1680 - 1760	13.864	+7.245 -7.603	+1.821 -1.839	2.6	1.0	+15.993 -16.161
1760 - 1840	18.192	+7.781 -7.820	+1.902 -1.906	2.6	1.0	+20.071 -20.087
1840 - 1920	22.612	+7.647 -7.537	+1.588 -1.590	2.6	1.0	+24.085 -24.050
1920 - 2000	29.530	+9.199 -9.469	+1.511 -1.505	2.6	1.0	+31.092 -31.172

Table A.3: Experimental uncertainties (in %), from all sources as well as the total uncertainty, affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 3-jet events.

Bin	Statistical	JEC	Unfolding	Lumi	Residual	Total
300 - 330	0.796	+3.503 -3.475	+0.564 -0.552	2.6	1.0	+4.581 -4.558
330 - 360	0.781	+3.303 -3.186	+0.640 -0.633	2.6	1.0	+4.437 -4.350
360 - 390	0.583	+3.221 -3.094	+0.490 -0.496	2.6	1.0	+4.326 -4.233
390 - 420	0.531	+3.092 -3.149	+0.584 -0.584	2.6	1.0	+4.236 -4.278
420 - 450	0.224	+3.125 -2.996	+0.604 -0.592	2.6	1.0	+4.236 -4.140
450 - 480	0.248	+2.984 -2.890	+0.531 -0.528	2.6	1.0	+4.124 -4.056
480 - 510	0.269	+2.937 -2.963	+0.511 -0.512	2.6	1.0	+4.089 -4.108
510 - 540	0.318	+3.021 -2.797	+0.592 -0.612	2.6	1.0	+4.164 -4.007
540 - 570	0.375	+2.999 -2.935	+0.506 -0.500	2.6	1.0	+4.141 -4.094
570 - 600	0.434	+2.824 -2.906	+0.646 -0.620	2.6	1.0	+4.042 -4.096
600 - 640	0.497	+2.952 -2.956	+0.598 -0.604	2.6	1.0	+4.133 -4.136
640 - 680	0.617	+3.111 -3.001	+0.777 -0.786	2.6	1.0	+4.292 -4.215
680 - 720	0.739	+3.067 -2.984	+0.642 -0.611	2.6	1.0	+4.257 -4.194
720 - 760	0.895	+3.185 -3.111	+0.595 -0.607	2.6	1.0	+4.366 -4.313
760 - 800	1.068	+3.231 -3.166	+0.763 -0.774	2.6	1.0	+4.464 -4.419
800 - 850	1.250	+3.427 -3.295	+0.674 -0.687	2.6	1.0	+4.639 -4.544
850 - 900	1.578	+3.364 -3.540	+0.903 -0.898	2.6	1.0	+4.731 -4.857
900 - 950	1.961	+3.594 -3.524	+0.792 -0.793	2.6	1.0	+5.015 -4.965
950 - 1000	2.420	+3.603 -3.783	+0.846 -0.843	2.6	1.0	+5.226 -5.351
1000 - 1060	2.844	+4.164 -4.116	+0.916 -0.940	2.6	1.0	+5.834 -5.803
1060 - 1120	3.647	+4.038 -3.815	+0.963 -0.957	2.6	1.0	+6.188 -6.044
1120 - 1180	4.607	+4.278 -4.183	+1.084 -1.087	2.6	1.0	+6.961 -6.904
1180 - 1250	5.532	+4.894 -4.771	+1.074 -1.069	2.6	1.0	+7.967 -7.891
1250 - 1320	7.141	+5.144 -5.273	+1.222 -1.217	2.6	1.0	+9.312 -9.383
1320 - 1390	10.207	+5.542 -5.642	+1.414 -1.428	2.6	1.0	+12.027 -12.076
1390 - 1460	13.831	+5.630 -5.265	+1.257 -1.256	2.6	1.0	+15.242 -15.111
1460 - 1530	15.578	+5.576 -5.491	+1.546 -1.551	2.6	1.0	+16.850 -16.822
1530 - 1600	18.729	+6.409 -7.019	+1.718 -1.716	2.6	1.0	+20.063 -20.266
1600 - 1680	26.465	+7.017 -6.255	+1.775 -1.765	2.6	1.0	+27.578 -27.393

Table A.4: Experimental uncertainties (in %), from all sources as well as the total uncertainty, affecting the measurement of cross-section ratio R_{32} , in each bin of $H_{T,2}/2$.

Bin	Statistical	JEC	Unfolding	Total
300 - 330	0.741	+1.059 -1.097	+0.754 -0.751	+1.496 -1.522
330 - 360	0.587	+0.954 -0.923	+0.685 -0.689	+1.313 -1.292
360 - 390	0.519	+0.902 -0.855	+0.594 -0.593	+1.199 -1.163
390 - 420	0.236	+0.907 -0.952	+0.439 -0.438	+1.035 -1.074
420 - 450	0.192	+0.900 -0.835	+0.360 -0.361	+0.988 -0.930
450 - 480	0.209	+0.788 -0.802	+0.307 -0.308	+0.872 -0.884
480 - 510	0.245	+0.795 -0.867	+0.254 -0.235	+0.870 -0.931
510 - 540	0.287	+0.852 -0.682	+0.264 -0.268	+0.937 -0.787
540 - 570	0.326	+0.807 -0.803	+0.193 -0.189	+0.891 -0.887
570 - 600	0.397	+0.656 -0.774	+0.199 -0.219	+0.792 -0.898
600 - 640	0.447	+0.763 -0.797	+0.150 -0.154	+0.897 -0.926
640 - 680	0.573	+0.861 -0.781	+0.153 -0.140	+1.045 -0.979
680 - 720	0.663	+0.766 -0.787	+0.147 -0.164	+1.024 -1.042
720 - 760	0.774	+0.842 -0.769	+0.118 -0.118	+1.149 -1.097
760 - 800	0.970	+0.800 -0.729	+0.115 -0.096	+1.263 -1.218
800 - 850	1.116	+0.873 -0.775	+0.115 -0.104	+1.422 -1.363
850 - 900	1.436	+0.770 -0.896	+0.069 -0.069	+1.631 -1.694
900 - 950	1.716	+0.704 -0.752	+0.050 -0.051	+1.855 -1.874
950 - 1000	2.156	+0.824 -0.897	+0.089 -0.045	+2.310 -2.336
1000 - 1060	2.554	+0.812 -0.870	+0.045 -0.040	+2.680 -2.698
1060 - 1120	3.244	+0.792 -0.658	+0.018 -0.027	+3.339 -3.310
1120 - 1180	4.121	+0.985 -0.757	+0.025 -0.043	+4.237 -4.191
1180 - 1250	4.990	+1.031 -0.848	+0.023 -0.041	+5.095 -5.062
1250 - 1320	6.456	+0.750 -1.087	+0.079 -0.079	+6.500 -6.548
1320 - 1390	8.990	+1.112 -1.144	+0.080 -0.099	+9.059 -9.063
1390 - 1460	12.699	+1.157 -0.815	+0.076 -0.078	+12.751 -12.725
1460 - 1530	13.926	+0.768 -1.235	+0.143 -0.145	+13.948 -13.981
1530 - 1600	16.903	+1.050 -1.258	+0.120 -0.127	+16.936 -16.950
1600 - 1680	28.070	+1.471 -0.859	+0.178 -0.177	+28.109 -28.084

A.4 Theoretical Uncertainties

Table A.5: Theoretical uncertainties (in %), calculated using CT10-NLO PDF set from all sources as well as the total uncertainty, affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 2-jet events.

Bin	Scale	PDF	NP	Total
300 - 330	+0.942 -6.149	+3.566 -3.090	0.825	+3.780 -6.931
330 - 360	+1.035 -6.289	+3.906 -3.342	0.736	+4.107 -7.159
360 - 390	+1.159 -6.438	+4.232 -3.573	0.696	+4.442 -7.396
390 - 420	+1.220 -6.536	+4.551 -3.794	0.723	+4.767 -7.592
420 - 450	+1.326 -6.660	+4.857 -3.997	0.745	+5.089 -7.802
450 - 480	+1.421 -6.776	+5.153 -4.186	0.765	+5.399 -8.001
480 - 510	+1.512 -6.888	+5.444 -4.365	0.782	+5.704 -8.192
510 - 540	+1.566 -6.967	+5.721 -4.527	0.797	+5.984 -8.347
540 - 570	+1.666 -7.082	+6.000 -4.682	0.810	+6.279 -8.528
570 - 600	+1.731 -7.172	+6.269 -4.825	0.822	+6.555 -8.683
600 - 640	+1.805 -7.271	+6.597 -4.979	0.833	+6.890 -8.852
640 - 680	+1.930 -7.416	+6.978 -5.143	0.845	+7.289 -9.064
680 - 720	+2.007 -7.527	+7.364 -5.295	0.856	+7.680 -9.243
720 - 760	+2.113 -7.663	+7.749 -5.437	0.865	+8.078 -9.436
760 - 800	+2.196 -7.781	+8.140 -5.569	0.873	+8.476 -9.609
800 - 850	+2.323 -7.945	+8.573 -5.706	0.881	+8.926 -9.822
850 - 900	+2.389 -8.062	+9.082 -5.863	0.889	+9.433 -10.008
900 - 950	+2.499 -8.227	+9.600 -6.018	0.896	+9.961 -10.232
950 - 1000	+2.631 -8.402	+10.134 -6.166	0.902	+10.509 -10.460
1000 - 1060	+2.738 -8.569	+10.747 -6.343	0.908	+11.127 -10.700
1060 - 1120	+2.853 -8.751	+11.431 -6.526	0.914	+11.817 -10.955
1120 - 1180	+2.992 -8.970	+12.183 -6.727	0.919	+12.579 -11.250
1180 - 1250	+3.135 -9.194	+13.019 -6.944	0.924	+13.423 -11.558
1250 - 1320	+3.324 -9.469	+14.004 -7.189	0.929	+14.423 -11.925
1320 - 1390	+3.434 -9.677	+15.080 -7.444	0.933	+15.494 -12.244
1390 - 1460	+3.629 -9.976	+16.223 -7.700	0.937	+16.650 -12.637
1460 - 1530	+3.760 -10.224	+17.505 -7.980	0.940	+17.929 -13.004
1530 - 1600	+3.894 -10.471	+18.891 -8.258	0.943	+19.311 -13.368
1600 - 1680	+4.107 -10.813	+20.496 -8.560	0.946	+20.925 -13.824
1680 - 1760	+4.421 -11.101	+22.481 -8.905	0.949	+22.931 -14.263
1760 - 1840	+4.921 -11.461	+24.654 -9.251	0.951	+25.158 -14.760
1840 - 1920	+5.404 -11.813	+27.143 -9.607	0.953	+27.692 -15.256
1920 - 2000	+5.867 -12.154	+29.986 -9.973	0.955	+30.570 -15.751

Table A.6: Theoretical uncertainties (in %), calculated using CT10-NLO PDF set from all sources as well as the total uncertainty, affecting the cross-section measurement in each bin of $H_{T,2}/2$ for inclusive 3-jet events.

Bin	Scale	PDF	NP	Total
300 - 330	+0.539 -8.294	+5.716 -4.657	1.692	+5.986 -9.662
330 - 360	+0.550 -8.577	+5.977 -4.779	1.516	+6.191 -9.935
360 - 390	+0.599 -8.709	+6.187 -4.987	1.363	+6.363 -10.128
390 - 420	+0.719 -8.948	+6.751 -5.223	1.228	+6.900 -10.433
420 - 450	+0.799 -9.145	+7.031 -5.395	1.110	+7.162 -10.676
450 - 480	+0.847 -9.247	+7.404 -5.578	1.005	+7.520 -10.845
480 - 510	+0.847 -9.294	+7.837 -5.717	0.937	+7.938 -10.951
510 - 540	+0.922 -9.436	+8.198 -5.884	0.921	+8.301 -11.158
540 - 570	+0.974 -9.566	+8.529 -6.000	0.904	+8.632 -11.328
570 - 600	+1.086 -9.786	+8.970 -6.156	0.886	+9.079 -11.595
600 - 640	+1.107 -9.852	+9.402 -6.297	0.866	+9.506 -11.724
640 - 680	+1.278 -10.101	+10.310 -6.526	0.842	+10.423 -12.055
680 - 720	+1.384 -10.342	+9.682 -6.618	0.820	+9.815 -12.305
720 - 760	+1.415 -10.404	+11.051 -6.826	0.798	+11.170 -12.469
760 - 800	+1.547 -10.615	+11.565 -7.009	0.777	+11.694 -12.744
800 - 850	+1.679 -10.804	+12.242 -7.185	0.755	+12.379 -12.997
850 - 900	+2.085 -11.134	+13.097 -7.461	0.731	+13.282 -13.422
900 - 950	+2.475 -11.432	+13.889 -7.703	0.709	+14.125 -13.804
950 - 1000	+2.655 -11.608	+14.614 -7.915	0.688	+14.869 -14.066
1000 - 1060	+3.025 -11.926	+15.576 -8.173	0.667	+15.881 -14.473
1060 - 1120	+3.299 -12.189	+14.250 -8.441	0.645	+14.641 -14.840
1120 - 1180	+3.741 -12.584	+17.984 -8.787	0.625	+18.380 -15.361
1180 - 1250	+3.969 -12.843	+19.324 -9.127	0.625	+19.737 -15.768
1250 - 1320	+4.663 -13.452	+21.246 -9.517	0.642	+21.761 -16.490
1320 - 1390	+4.878 -13.702	+22.884 -9.899	0.657	+23.407 -16.916
1390 - 1460	+5.242 -14.095	+24.854 -10.332	0.670	+25.410 -17.489
1460 - 1530	+5.582 -14.464	+27.170 -10.733	0.682	+27.746 -18.024
1530 - 1600	+6.003 -14.907	+29.741 -11.165	0.692	+30.349 -18.637
1600 - 1680	+6.503 -15.418	+32.855 -11.617	0.702	+33.500 -19.317

Table A.7: Theoretical uncertainties (in %) calculated using CT10-NLO PDF set from all sources as well as the total uncertainty, affecting the measurement of cross-section ratio R_{32} , in each bin of $H_{\text{T},2}/2$.

Bin	Scale	PDF	NP	Total
300 - 330	+0.038 -7.203	+2.458 -3.463	0.822	+2.592 -8.035
330 - 360	+0.027 -6.626	+2.317 -3.378	0.734	+2.431 -7.474
360 - 390	+0.024 -6.449	+2.149 -3.367	0.656	+2.247 -7.304
390 - 420	+0.084 -5.894	+2.411 -3.383	0.586	+2.482 -6.821
420 - 450	+0.113 -5.532	+2.345 -3.362	0.523	+2.405 -6.494
450 - 480	+0.109 -5.409	+2.390 -3.357	0.467	+2.438 -6.383
480 - 510	+0.073 -5.442	+2.506 -3.327	0.416	+2.541 -6.392
510 - 540	+0.107 -5.168	+2.559 -3.326	0.371	+2.588 -6.157
540 - 570	+0.112 -5.010	+2.586 -3.292	0.330	+2.609 -6.004
570 - 600	+0.163 -4.576	+2.729 -3.292	0.292	+2.750 -5.645
600 - 640	+0.146 -4.565	+2.824 -3.270	0.253	+2.839 -5.621
640 - 680	+0.198 -4.163	+3.368 -3.298	0.236	+3.382 -5.316
680 - 720	+0.155 -3.754	+2.352 -3.247	0.227	+2.368 -4.968
720 - 760	+0.196 -3.842	+3.267 -3.268	0.219	+3.280 -5.049
760 - 800	+0.126 -3.523	+3.366 -3.272	0.212	+3.375 -4.813
800 - 850	+0.110 -3.368	+3.596 -3.261	0.206	+3.604 -4.693
850 - 900	+0.048 -3.351	+3.909 -3.309	0.200	+3.915 -4.714
900 - 950	+0.116 -3.504	+4.148 -3.334	0.196	+4.154 -4.841
950 - 1000	+0.127 -3.511	+4.300 -3.335	0.192	+4.306 -4.846
1000 - 1060	+0.282 -3.683	+4.604 -3.357	0.204	+4.617 -4.988
1060 - 1120	+0.436 -3.779	+3.079 -3.375	0.224	+3.118 -5.071
1120 - 1180	+0.732 -3.982	+5.430 -3.452	0.241	+5.485 -5.276
1180 - 1250	+0.813 -4.031	+5.835 -3.511	0.258	+5.897 -5.352
1250 - 1320	+1.303 -4.414	+6.626 -3.591	0.275	+6.759 -5.697
1320 - 1390	+1.403 -4.471	+7.036 -3.659	0.290	+7.180 -5.785
1390 - 1460	+1.564 -4.590	+7.657 -3.778	0.304	+7.822 -5.953
1460 - 1530	+1.765 -4.738	+8.438 -3.853	0.316	+8.626 -6.115
1530 - 1600	+2.040 -4.972	+9.306 -3.962	0.328	+9.532 -6.366
1600 - 1680	+2.313 -5.179	+10.381 -4.075	0.339	+10.641 -6.599

Bibliography

- [1] CMS Collaboration, “Measurement of the ratio of the inclusive 3-jet cross section to the inclusive 2-jet cross section in pp collisions at $\sqrt{s} = 7$ TeV and first determination of the strong coupling constant in the TeV range”, Eur. Phys. J. C **73**, no. 10, 2604 (2013) doi:[10.1140/epjc/s10052-013-2604-6](https://doi.org/10.1140/epjc/s10052-013-2604-6) arXiv:[1304.7498](https://arxiv.org/abs/1304.7498) [hep-ex].
- [2] D. Griffiths, “Introduction to elementary particles; 2nd rev. version”, Wiley, New York (2008) <https://cds.cern.ch/record/111880>.
- [3] D. H. Perkins, “Introduction to high energy physics”, Cambridge (1982) ISBN : 9780521621960.
- [4] M. Herrero, “The Standard model”, NATO Sci. Ser. C **534**, 1 (1999) doi:[10.1007/978-94-011-4689-0_1](https://doi.org/10.1007/978-94-011-4689-0_1) arXiv:[hep-ph/9812242](https://arxiv.org/abs/hep-ph/9812242).
- [5] S. Weinberg, “A Model of Leptons”, Phys. Rev. Lett. **19**, 1264 (1967) doi:[10.1103/PhysRevLett.19.1264](https://doi.org/10.1103/PhysRevLett.19.1264).
- [6] M. E. Peskin and D. V. Schroeder, “An Introduction to quantum field theory”, Addison-Wesley, USA (1995).
- [7] S. L. Glashow, “Towards a Unified Theory: Threads in a Tapestry”, Rev. Mod. Phys. **52**, 539 (1980) doi:[10.1103/RevModPhys.52.539](https://doi.org/10.1103/RevModPhys.52.539).

- [8] A. Salam, “Gauge Unification of Fundamental Forces”, Rev. Mod. Phys. **52**, 525 (1980) [Science **210**, 723 (1980)] doi:[10.1103/RevModPhys.52.525](https://doi.org/10.1103/RevModPhys.52.525).
- [9] H. Georgi and S. L. Glashow, “Unity of All Elementary Particle Forces”, Phys. Rev. Lett. **32**, 438 (1974) doi:[10.1103/PhysRevLett.32.438](https://doi.org/10.1103/PhysRevLett.32.438).
- [10] CMS Collaboration, “Observation of a new boson at a mass of 125 GeV with the CMS experiment at the LHC”, Phys. Lett. B **716**, 30 (2012) doi:[10.1016/j.physletb.2012.08.021](https://doi.org/10.1016/j.physletb.2012.08.021) arXiv:[1207.7235](https://arxiv.org/abs/1207.7235) [hep-ex].
- [11] ATLAS Collaboration, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, Phys. Lett. B **716**, 1 (2012) doi:[10.1016/j.physletb.2012.08.020](https://doi.org/10.1016/j.physletb.2012.08.020) arXiv:[1207.7214](https://arxiv.org/abs/1207.7214) [hep-ex].
- [12] R. K. Ellis, W. J. Stirling and B. R. Webber, “QCD and collider physics”, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **8**, 1 (1996).
- [13] F. Halzen and A. D. Martin, “Quarks And Leptons: An Introductory Course In Modern Particle Physics”, New York, Usa: Wiley (1984) 396p.
- [14] M. Gell-Mann, “Symmetries of baryons and mesons”, Phys. Rev. **125**, 1067 (1962) doi:[10.1103/PhysRev.125.1067](https://doi.org/10.1103/PhysRev.125.1067).
- [15] K. Rabbertz, “Jet Physics at the LHC : The Strong Force beyond the TeV Scale”, Springer Tracts Mod. Phys. **268**, pp.1 (2017) doi:[10.1007/978-3-319-42115-5](https://doi.org/10.1007/978-3-319-42115-5).
- [16] G. ’t Hooft, “Dimensional regularization and the renormalization group”, Nucl. Phys. B **61**, 455 (1973) doi:[10.1016/0550-3213\(73\)90376-3](https://doi.org/10.1016/0550-3213(73)90376-3).
- [17] S. Weinberg, “New approach to the renormalization group”, Phys. Rev. D **8**, 3497 (1973) doi:[10.1103/PhysRevD.8.3497](https://doi.org/10.1103/PhysRevD.8.3497).

- [18] C. Patrignani *et al.* (Particle Data Group), “Review of Particle Physics”, Chin. Phys. C **40**, no. 10, 100001 (2016) doi:[10.1088/1674-1137/40/10/100001](https://doi.org/10.1088/1674-1137/40/10/100001).
- [19] J. C. Collins, D. E. Soper and G. F. Sterman, “Factorization of Hard Processes in QCD”, Adv. Ser. Direct. High Energy Phys. **5**, 1 (1989) doi:[10.1142/9789814503266_0001](https://doi.org/10.1142/9789814503266_0001) arXiv:[hep-ph/0409313](https://arxiv.org/abs/hep-ph/0409313).
- [20] V. N. Gribov and L. N. Lipatov, “Deep inelastic e p scattering in perturbation theory”, Sov. J. Nucl. Phys. **15**, 438 (1972) [Yad. Fiz. **15**, 781 (1972)].
- [21] Y. L. Dokshitzer, “Calculation of the Structure Functions for Deep Inelastic Scattering and e+ e- Annihilation by Perturbation Theory in Quantum Chromodynamics.”, Sov. Phys. JETP **46**, 641 (1977) [Zh. Eksp. Teor. Fiz. **73**, 1216 (1977)].
- [22] G. Altarelli and G. Parisi, “Asymptotic Freedom in Parton Language”, Nucl. Phys. B **126**, 298 (1977) doi:[10.1016/0550-3213\(77\)90384-4](https://doi.org/10.1016/0550-3213(77)90384-4).
- [23] CMS Collaboration, “Measurement of the triple-differential dijet cross section in proton-proton collisions at $\sqrt{s} = 8$ TeV and constraints on parton distribution functions”, Eur. Phys. J. C **77**, no. 11, 746 (2017) doi:[10.1140/epjc/s10052-017-5286-7](https://doi.org/10.1140/epjc/s10052-017-5286-7) arXiv:[1705.02628](https://arxiv.org/abs/1705.02628) [hep-ex].
- [24] S. Dulat *et al.*, “New parton distribution functions from a global analysis of quantum chromodynamics”, Phys. Rev. D **93**, no. 3, 033006 (2016) doi:[10.1103/PhysRevD.93.033006](https://doi.org/10.1103/PhysRevD.93.033006) arXiv:[1506.07443](https://arxiv.org/abs/1506.07443) [hep-ph].
- [25] L. A. Harland-Lang *et al.*, “Parton distributions in the LHC era: MMHT 2014 PDFs”, Eur. Phys. J. C **75**, no. 5, 204 (2015) doi:[10.1140/epjc/s10052-015-3397-6](https://doi.org/10.1140/epjc/s10052-015-3397-6) arXiv:[1412.3989](https://arxiv.org/abs/1412.3989) [hep-ph].
- [26] NNPDF Collaboration, “Parton distributions for the LHC Run II”, JHEP **1504**, 040 (2015) doi:[10.1007/JHEP04\(2015\)040](https://doi.org/10.1007/JHEP04(2015)040) arXiv:[1410.8849](https://arxiv.org/abs/1410.8849) [hep-ph].

- [27] S. Alekhin, J. Blumlein and S. Moch, “Parton Distribution Functions and Benchmark Cross Sections at NNLO”, Phys. Rev. D **86**, 054009 (2012) doi:[10.1103/PhysRevD.86.054009](https://doi.org/10.1103/PhysRevD.86.054009) arXiv:[1202.2281](https://arxiv.org/abs/1202.2281) [hep-ph].
- [28] H1 and ZEUS Collaborations, “Combination of measurements of inclusive deep inelastic $e^\pm p$ scattering cross sections and QCD analysis of HERA data”, Eur. Phys. J. C **75**, no. 12, 580 (2015) doi:[10.1140/epjc/s10052-015-3710-4](https://doi.org/10.1140/epjc/s10052-015-3710-4) arXiv:[1506.06042](https://arxiv.org/abs/1506.06042) [hep-ex].
- [29] B. Andersson, “The Lund model”, Camb. Monogr. Part. Phys. Nucl. Phys. Cosmol. **7**, 1 (1997) ISBN: 9780521420945.
- [30] G. Marchesini and B. R. Webber, “Monte Carlo Simulation of General Hard Processes with Coherent QCD Radiation”, Nucl. Phys. B **310**, 461 (1988) doi:[10.1016/0550-3213\(88\)90089-2](https://doi.org/10.1016/0550-3213(88)90089-2).
- [31] B. R. Webber, “A QCD Model for Jet Fragmentation Including Soft Gluon Interference”, Nucl. Phys. B **238**, 492 (1984) doi:[10.1016/0550-3213\(84\)90333-X](https://doi.org/10.1016/0550-3213(84)90333-X).
- [32] D. Amati and G. Veneziano, “Preconfinement as a Property of Perturbative QCD”, Phys. Lett. **83B**, 87 (1979) doi:[10.1016/0370-2693\(79\)90896-7](https://doi.org/10.1016/0370-2693(79)90896-7).
- [33] G. F. Sterman and S. Weinberg, “Jets from Quantum Chromodynamics”, Phys. Rev. Lett. **39**, 1436 (1977) doi:[10.1103/PhysRevLett.39.1436](https://doi.org/10.1103/PhysRevLett.39.1436).
- [34] G. Hanson *et al.*, “Evidence for Jet Structure in Hadron Production by e+ e- Annihilation”, Phys. Rev. Lett. **35**, 1609 (1975) doi:[10.1103/PhysRevLett.35.1609](https://doi.org/10.1103/PhysRevLett.35.1609).
- [35] G. P. Salam, “Towards Jetography”, Eur. Phys. J. C **67**, 637 (2010) doi:[10.1140/epjc/s10052-010-1314-6](https://doi.org/10.1140/epjc/s10052-010-1314-6) arXiv:[0906.1833](https://arxiv.org/abs/0906.1833) [hep-ph].

- [36] S. D. Ellis, Z. Kunszt and D. E. Soper, “The One Jet Inclusive Cross-Section at Order α_s^3 . 1. Gluons Only”, Phys. Rev. D **40**, 2188 (1989) doi:[10.1103/PhysRevD.40.2188](https://doi.org/10.1103/PhysRevD.40.2188).
- [37] G. C. Blazey *et al.*, “Run II jet physics”, arXiv:[hep-ex/0005012](https://arxiv.org/abs/hep-ex/0005012).
- [38] S. Weinzierl, “The SISCone jet algorithm optimised for low particle multiplicities”, Comput. Phys. Commun. **183**, 813 (2012) doi:[10.1016/j.cpc.2011.12.007](https://doi.org/10.1016/j.cpc.2011.12.007) arXiv:[1108.1934 \[hep-ph\]](https://arxiv.org/abs/1108.1934).
- [39] S. D. Ellis and D. E. Soper, “Successive combination jet algorithm for hadron collisions”, Phys. Rev. D **48**, 3160 (1993) doi:[10.1103/PhysRevD.48.3160](https://doi.org/10.1103/PhysRevD.48.3160) arXiv:[hep-ph/9305266](https://arxiv.org/abs/hep-ph/9305266).
- [40] S. Catani *et al.*, “Longitudinally invariant K_t clustering algorithms for hadron hadron collisions”, Nucl. Phys. B **406**, 187 (1993) doi:[10.1016/0550-3213\(93\)90166-M](https://doi.org/10.1016/0550-3213(93)90166-M).
- [41] . Catani, Y. L. Dokshitzer and B. R. Webber, “The K^- perpendicular clustering algorithm for jets in deep inelastic scattering and hadron collisions”, Phys. Lett. B **285**, 291 (1992) doi:[10.1016/0370-2693\(92\)91467-N](https://doi.org/10.1016/0370-2693(92)91467-N).
- [42] Y. L. Dokshitzer *et al.*, “Better jet clustering algorithms”, JHEP **9708**, 001 (1997) doi:[10.1088/1126-6708/1997/08/001](https://doi.org/10.1088/1126-6708/1997/08/001) arXiv:[hep-ph/9707323](https://arxiv.org/abs/hep-ph/9707323).
- [43] M. Cacciari, G. P. Salam and G. Soyez, “The Anti- $k(t)$ jet clustering algorithm”, JHEP **0804**, 063 (2008) doi:[10.1088/1126-6708/2008/04/063](https://doi.org/10.1088/1126-6708/2008/04/063) arXiv:[0802.1189 \[hep-ph\]](https://arxiv.org/abs/0802.1189).
- [44] M. Cacciari, G. P. Salam and G. Soyez, “FastJet User Manual”, Eur. Phys. J. C **72**, 1896 (2012) doi:[10.1140/epjc/s10052-012-1896-2](https://doi.org/10.1140/epjc/s10052-012-1896-2) arXiv:[1111.6097 \[hep-ph\]](https://arxiv.org/abs/1111.6097).

- [45] L. Evans and P. Bryant, “LHC Machine”, JINST **3**, S08001 (2008) doi:[10.1088/1748-0221/3/08/S08001](https://doi.org/10.1088/1748-0221/3/08/S08001).
- [46] S. Myers and E. Picasso, “The design, construction and commissioning of the CERN large Electron–Positron collider”, Contemporary Physics **31**, no. 6, 387 (1990) doi:[10.1080/00107519008213789](https://doi.org/10.1080/00107519008213789).
- [47] ALICE Collaboration, “The ALICE experiment at the CERN LHC”, JINST **3**, S08002 (2008) doi:[10.1088/1748-0221/3/08/S08002](https://doi.org/10.1088/1748-0221/3/08/S08002).
- [48] ATLAS Collaboration, “The ATLAS Experiment at the CERN Large Hadron Collider”, JINST **3**, S08003 (2008) doi:[10.1088/1748-0221/3/08/S08003](https://doi.org/10.1088/1748-0221/3/08/S08003).
- [49] CMS Collaboration, “The CMS Experiment at the CERN LHC”, JINST **3**, S08004 (2008) doi:[10.1088/1748-0221/3/08/S08004](https://doi.org/10.1088/1748-0221/3/08/S08004).
- [50] CMS Collaboration, “CMS Physics : Technical Design Report Volume 1: Detector Performance and Software”, CERN-LHCC-2006-001 (2006), CMS-TDR-8-1.
- [51] CMS Collaboration, “CMS technical design report, volume II: Physics performance”, J. Phys. G **34**, no. 6, 995 (2007) doi:[10.1088/0954-3899/34/6/S01](https://doi.org/10.1088/0954-3899/34/6/S01).
- [52] LHCb Collaboration, “The LHCb Detector at the LHC”, JINST **3**, S08005 (2008) doi:[10.1088/1748-0221/3/08/S08005](https://doi.org/10.1088/1748-0221/3/08/S08005).
- [53] LHCf Collaboration, “The LHCf detector at the CERN Large Hadron Collider”, JINST **3**, S08006 (2008) doi:[10.1088/1748-0221/3/08/S08006](https://doi.org/10.1088/1748-0221/3/08/S08006).
- [54] TOTEM Collaboration, “The TOTEM experiment at the CERN Large Hadron Collider”, JINST **3**, S08007 (2008) doi:[10.1088/1748-0221/3/08/S08007](https://doi.org/10.1088/1748-0221/3/08/S08007).

- [55] CMS Collaboration, “CMS Luminosity Based on Pixel Cluster Counting - Summer 2013 Update”, CMS Physics Analysis Summary **CMS-PAS-LUM-13-001** (2013) <http://cds.cern.ch/record/1598864>.
- [56] S. van der Meer, “Calibration of the Effective Beam Height in the ISR”, Technical Report CERN-ISR-PO-68-31 (1968).
- [57] P. Adzic *et al.*, “Energy resolution of the barrel of the CMS electromagnetic calorimeter”, JINST **2**, P04004 (2007) doi:[10.1088/1748-0221/2/04/P04004](https://doi.org/10.1088/1748-0221/2/04/P04004).
- [58] CMS Collaboration, “Performance of the CMS Hadron Calorimeter with Cosmic Ray Muons and LHC Beam Data”, JINST **5**, T03012 (2010) doi:[10.1088/1748-0221/5/03/T03012](https://doi.org/10.1088/1748-0221/5/03/T03012) arXiv:[0911.4991 \[physics.ins-det\]](https://arxiv.org/abs/0911.4991).
- [59] I. Bird *et al.*, “LHC computing Grid. Technical design report”, CERN-LHCC-2005-024 <http://cds.cern.ch/record/840543>.
- [60] CMS Collaboration, “CMS: The computing project. Technical design report”, CERN-LHCC-2005-023 (2005) <http://cds.cern.ch/record/838359>.
- [61] R. Brun and F. Rademakers, “ROOT: An object oriented data analysis framework”, Nucl. Instrum. Meth. A **389**, 81 (1997) doi:[10.1016/S0168-9002\(97\)00048-X](https://doi.org/10.1016/S0168-9002(97)00048-X).
- [62] Z. Nagy, “Three jet cross-sections in hadron hadron collisions at next-to-leading order”, Phys. Rev. Lett. **88**, 122003 (2002) doi:[10.1103/PhysRevLett.88.122003](https://doi.org/10.1103/PhysRevLett.88.122003) arXiv:[hep-ph/0110315](https://arxiv.org/abs/hep-ph/0110315).
- [63] Z. Nagy, “Next-to-leading order calculation of three jet observables in hadron hadron collision”, Phys. Rev. D **68**, 094002 (2003) doi:[10.1103/PhysRevD.68.094002](https://doi.org/10.1103/PhysRevD.68.094002) arXiv:[hep-ph/0307268](https://arxiv.org/abs/hep-ph/0307268).

- [64] D. Britzger *et al.*, “New features in version 2 of the fastNLO project”, 217 (2012) doi:[10.3204/DESY-PROC-2012-02/165](https://doi.org/10.3204/DESY-PROC-2012-02/165) arXiv:[1208.3641](https://arxiv.org/abs/1208.3641) [hep-ph].
- [65] T. Kluge, K. Rabbertz and M. Wobisch, “FastNLO: Fast pQCD calculations for PDF fits”, Proceedings DIS 2006, 14th International Workshop, 483 (2006) doi:[10.1142/9789812706706_0110](https://doi.org/10.1142/9789812706706_0110) arXiv:[hep-ph/0609285](https://arxiv.org/abs/hep-ph/0609285).
- [66] M. R. Whalley, D. Bourilkov and R. C. Group, “The Les Houches accord PDFs (LHAPDF) and LHAGLUE”, Proceedings : Workshop on the implications of HERA for LHC physics (2005) arXiv:[hep-ph/0508110](https://arxiv.org/abs/hep-ph/0508110).
- [67] A. Buckley, *et al.*, “LHAPDF6: parton density access in the LHC precision era”, Eur. Phys. J. C **75**, 132 (2015) doi:[10.1140/epjc/s10052-015-3318-8](https://doi.org/10.1140/epjc/s10052-015-3318-8) arXiv:[1412.7420](https://arxiv.org/abs/1412.7420) [hep-ph].
- [68] N. Metropolis and S. Ulam, “The Monte Carlo Method”, Journal of the American Statistical Association **44**, no. 247, 335 (1949) doi:[10.1080/01621459.1949.10483310](https://doi.org/10.1080/01621459.1949.10483310).
- [69] B. Andersson *et al.*, “Parton fragmentation and string dynamics”, Physics Reports, **97**, no. 2, 31 (1983) doi:[10.1016/0370-1573\(83\)90080-7](https://doi.org/10.1016/0370-1573(83)90080-7).
- [70] T. Sjostrand, S. Mrenna and P. Z. Skands, “PYTHIA 6.4 Physics and Manual”, JHEP **0605**, 026 (2006) doi:[10.1088/1126-6708/2006/05/026](https://doi.org/10.1088/1126-6708/2006/05/026) arXiv:[hep-ph/0603175](https://arxiv.org/abs/hep-ph/0603175).
- [71] T. Sjostrand, S. Mrenna and P. Z. Skands, “A Brief Introduction to PYTHIA 8.1”, Comput. Phys. Commun. **178**, 852 (2008) doi:[10.1016/j.cpc.2008.01.036](https://doi.org/10.1016/j.cpc.2008.01.036) arXiv:[0710.3820](https://arxiv.org/abs/0710.3820) [hep-ph].
- [72] R. Field, “Min-Bias and the Underlying Event at the LHC”, Acta Phys. Polon. B **42**, 2631 (2011) doi:[10.5506/APhysPolB.42.2631](https://doi.org/10.5506/APhysPolB.42.2631) arXiv:[1110.5530](https://arxiv.org/abs/1110.5530) [hep-ph].

- [73] CMS Collaboration, “Event generator tunes obtained from underlying event and multiparton scattering measurements”, Eur. Phys. J. C **76**, no. 3, 155 (2016) doi:[10.1140/epjc/s10052-016-3988-x](https://doi.org/10.1140/epjc/s10052-016-3988-x) arXiv:1512.00815 [hep-ex].
- [74] J. Alwall *et al.*, “MadGraph 5 : Going Beyond”, JHEP **1106**, 128 (2011) doi:[10.1007/JHEP06\(2011\)128](https://doi.org/10.1007/JHEP06(2011)128) arXiv:1106.0522 [hep-ph].
- [75] J. Alwall *et al.*, “A Standard format for Les Houches event files”, Comput. Phys. Commun. **176**, 300 (2007) doi:[10.1016/j.cpc.2006.11.010](https://doi.org/10.1016/j.cpc.2006.11.010) arXiv:hep-ph/0609017.
- [76] G. Corcella *et al.*, “HERWIG 6: An Event generator for hadron emission reactions with interfering gluons (including supersymmetric processes)”, JHEP **0101**, 010 (2001) doi:[10.1088/1126-6708/2001/01/010](https://doi.org/10.1088/1126-6708/2001/01/010) arXiv:hep-ph/0011363.
- [77] M. Bahr *et al.*, “Herwig++ Physics and Manual”, Eur. Phys. J. C **58**, 639 (2008) doi:[10.1140/epjc/s10052-008-0798-9](https://doi.org/10.1140/epjc/s10052-008-0798-9) arXiv:0803.0883 [hep-ph].
- [78] M. Bahr *et al.*, “Herwig++ 2.3 Release Note”, arXiv:0812.0529 [hep-ph].
- [79] S. Frixione, P. Nason and C. Oleari, “Matching NLO QCD computations with Parton Shower simulations: the POWHEG method”, JHEP **0711**, 070 (2007) doi:[10.1088/1126-6708/2007/11/070](https://doi.org/10.1088/1126-6708/2007/11/070) arXiv:0709.2092 [hep-ph].
- [80] P. Nason, “A New method for combining NLO QCD with shower Monte Carlo algorithms”, JHEP **0411**, 040 (2004) doi:[10.1088/1126-6708/2004/11/040](https://doi.org/10.1088/1126-6708/2004/11/040) arXiv:hep-ph/0409146.
- [81] S. Alioli *et al.*, “Jet pair production in POWHEG”, JHEP **1104**, 081 (2011) doi:[10.1007/JHEP04\(2011\)081](https://doi.org/10.1007/JHEP04(2011)081) arXiv:1012.3380 [hep-ph].
- [82] C. Oleari, “The POWHEG-BOX”, Nucl. Phys. Proc. Suppl. **205-206**, 36 (2010) doi:[10.1016/j.nuclphysbps.2010.08.016](https://doi.org/10.1016/j.nuclphysbps.2010.08.016) arXiv:1007.3893 [hep-ph].

- [83] GEANT4 Collaboration, “GEANT4: A Simulation toolkit”, Nucl. Instrum. Meth. A **506**, 250 (2003) doi:[10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8).
- [84] CMS Collaboration, “The fast simulation of the CMS detector at LHC”, J. Phys. Conf. Ser. **331**, 032049 (2011) doi:<http://dx.doi.org/10.1088/1742-6596/331/3/032049>.
- [85] T. Schörner-Sadenius, “The Large Hadron Collider : Harvest of Run 1”, Springer (2015) doi:[10.1007/978-3-319-15001-7](https://doi.org/10.1007/978-3-319-15001-7).
- [86] CMS Collaboration, “Particle-Flow Event Reconstruction in CMS and Performance for Jets, Taus, and MET”, CMS-PAS-PFT-09-001 (2009) <http://cds.cern.ch/record/1194487>.
- [87] CMS Collaboration, “Commissioning of the Particle-flow Event Reconstruction with the first LHC collisions recorded in the CMS detector”, CMS-PAS-PFT-10-001 (2010) <http://cds.cern.ch/record/1247373>.
- [88] W. Adam *et al.*, “Track reconstruction in the CMS tracker”, CERN-CMS-NOTE-2006-041 (2006) <http://cds.cern.ch/record/934067>.
- [89] CMS Collaboration, “Determination of Jet Energy Calibration and Transverse Momentum Resolution in CMS”, JINST **6**, P11002 (2011) doi:[10.1088/1748-0221/6/11/P11002](https://doi.org/10.1088/1748-0221/6/11/P11002) arXiv:[1107.4277](https://arxiv.org/abs/1107.4277) [physics.ins-det].
- [90] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, JINST **12**, P02014 (2017) doi:[10.1088/1748-0221/12/02/P02014](https://doi.org/10.1088/1748-0221/12/02/P02014) arXiv:[1607.03663](https://arxiv.org/abs/1607.03663) [hep-ex].
- [91] CMS Collaboration, “Measurement and QCD analysis of double-differential inclusive jet cross sections in pp collisions at $\sqrt{s} = 8$ TeV and cross section ratios to 2.76 and 7 TeV”, JHEP **1703**, 156 (2017) doi:[10.1007/JHEP03\(2017\)156](https://doi.org/10.1007/JHEP03(2017)156) arXiv:[1609.05331](https://arxiv.org/abs/1609.05331) [hep-ex].

- [92] C. J. Clopper and E. S. Pearson, “The Use of Confidence or Fiducial Limits Illustrated in the Case of the Binomial”, *Biometrika* **26**, 404 (1934) doi:[10.2307/2331986](https://doi.org/10.2307/2331986).
- [93] CMS Collaboration, “Jet Performance in pp Collisions at 7 TeV”, CMS Physics Analysis Summary **CMS-PAS-JME-10-003** (2010) <http://cds.cern.ch/record/1279362>.
- [94] CMS Collaboration, “Jet Identification at 8 TeV”, URL: <https://twiki.cern.ch/twiki/bin/viewauth/CMS/JetID> (accessed on 2017-10-31).
- [95] CMS Collaboration, “Jet Energy Resolution at 8 TeV”, URL: <https://twiki.cern.ch/twiki/bin/viewauth/CMS/JetResolution> (accessed on 2016-01-22).
- [96] CMS Collaboration, “Jet Energy Resolution in CMS at $\sqrt{s}=7$ TeV” CMS Physics Analysis Summary **CMS-PAS-JME-10-014** (2011) <http://cds.cern.ch/record/1339945>.
- [97] G. D’Agostini, “A Multidimensional unfolding method based on Bayes’ theorem”, *Nucl. Instrum. Meth. A* **362**, 487 (1995) doi:[10.1016/0168-9002\(95\)00274-X](https://doi.org/10.1016/0168-9002(95)00274-X).
- [98] G. D’Agostini, “Improved iterative Bayesian unfolding”, (2010) arXiv:[1010.0632 \[physics.data-an\]](https://arxiv.org/abs/1010.0632).
- [99] Tim Adye, “Unfolding algorithms and tests using RooUnfold”, Proceedings of the PHYSTAT 2011 Workshop, CERN, Geneva, Switzerland 313 (January 2011) doi:[10.5170/CERN-2011-006.313](https://doi.org/10.5170/CERN-2011-006.313) arXiv:[1105.1160 \[physics.data-an\]](https://arxiv.org/abs/1105.1160).
- [100] CMS Collaboration, “Measurement of the Inclusive Jet Cross Section in pp Collisions at $\sqrt{s} = 7$ TeV”, *Phys. Rev. Lett.* **107**, 132001 (2011) doi:[10.1103/PhysRevLett.107.132001](https://doi.org/10.1103/PhysRevLett.107.132001) arXiv:[1106.0208 \[hep-ex\]](https://arxiv.org/abs/1106.0208).

- [101] CMS Collaboration, “Jet energy scale uncertainty sources”, URL: <https://twiki.cern.ch/twiki/bin/viewauth/CMS/JECUncertaintySources> (accessed on 2017-10-31).
- [102] CMS Collaboration, “Measurements of differential jet cross sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the CMS detector”, Phys. Rev. D **87**, no. 11, 112002 (2013) Erratum: [Phys. Rev. D **87**, no. 11, 119902 (2013)] doi:[10.1103/PhysRevD.87.112002](https://doi.org/10.1103/PhysRevD.87.112002), [10.1103/PhysRevD.87.119902](https://doi.org/10.1103/PhysRevD.87.119902) arXiv:[1212.6660](https://arxiv.org/abs/1212.6660) [hep-ex].
- [103] ATLAS Collaboration, “Measurement of the inclusive jet cross section in pp collisions at $\text{sqrt}(s)=2.76$ TeV and comparison to the inclusive jet cross section at $\text{sqrt}(s)=7$ TeV using the ATLAS detector”, Eur. Phys. J. C **73**, no. 8, 2509 (2013) doi:[10.1140/epjc/s10052-013-2509-4](https://doi.org/10.1140/epjc/s10052-013-2509-4) arXiv:[1304.4739](https://arxiv.org/abs/1304.4739) [hep-ex].
- [104] ATLAS Collaboration, “Measurement of the inclusive jet cross-section in proton-proton collisions at $\sqrt{s} = 7$ TeV using 4.5 fb^{-1} of data with the ATLAS detector”, JHEP **1502**, 153 (2015) Erratum: [JHEP **1509**, 141 (2015)] doi:[10.1007/JHEP02\(2015\)153](https://doi.org/10.1007/JHEP02(2015)153), [10.1007/JHEP09\(2015\)141](https://doi.org/10.1007/JHEP09(2015)141) arXiv:[1410.8857](https://arxiv.org/abs/1410.8857) [hep-ex].
- [105] CMS Collaboration, “Measurement of the inclusive 3-jet production differential cross section in proton–proton collisions at 7 TeV and determination of the strong coupling constant in the TeV range”, Eur. Phys. J. C **75**, no. 5, 186 (2015) doi:[10.1140/epjc/s10052-015-3376-y](https://doi.org/10.1140/epjc/s10052-015-3376-y) arXiv:[1412.1633](https://arxiv.org/abs/1412.1633) [hep-ex].
- [106] CMS Collaboration, “Measurement of the inclusive jet cross section in pp collisions at $\sqrt{s} = 2.76$ TeV”, Eur. Phys. J. C **76**, no. 5, 265 (2016) doi:[10.1140/epjc/s10052-016-4083-z](https://doi.org/10.1140/epjc/s10052-016-4083-z) arXiv:[1512.06212](https://arxiv.org/abs/1512.06212) [hep-ex]].

- [107] H. L. Lai *et al.*, “New parton distributions for collider physics”, Phys. Rev. D **82**, 074024 (2010) doi:[10.1103/PhysRevD.82.074024](https://doi.org/10.1103/PhysRevD.82.074024) arXiv:[1007.2241](https://arxiv.org/abs/1007.2241) [hep-ph].
- [108] A. D. Martin *et al.*, “Parton distributions for the LHC”, Eur. Phys. J. C **63**, 189 (2009) doi:[10.1140/epjc/s10052-009-1072-5](https://doi.org/10.1140/epjc/s10052-009-1072-5) arXiv:[0901.0002](https://arxiv.org/abs/0901.0002) [hep-ph].
- [109] A. D. Martin *et al.*, “Uncertainties on alpha(S) in global PDF analyses and implications for predicted hadronic cross sections”, Eur. Phys. J. C **64**, 653 (2009) doi:[10.1140/epjc/s10052-009-1164-2](https://doi.org/10.1140/epjc/s10052-009-1164-2) arXiv:[0905.3531](https://arxiv.org/abs/0905.3531) [hep-ph].
- [110] R. D. Ball *et al.*, “Parton distributions with LHC data”, Nucl. Phys. B **867**, 244 (2013) doi:[10.1016/j.nuclphysb.2012.10.003](https://doi.org/10.1016/j.nuclphysb.2012.10.003) arXiv:[1207.1303](https://arxiv.org/abs/1207.1303) [hep-ph].
- [111] W. Hollik *et al.*, “Electroweak physics”, Acta Phys. Polon. B **35**, 2533 (2004) arXiv:[hep-ph/0501246](https://arxiv.org/abs/hep-ph/0501246).
- [112] S. Dittmaier, A. Huss and C. Speckner, “Weak radiative corrections to dijet production at hadron colliders”, JHEP **1211**, 095 (2012) doi:[10.1007/JHEP11\(2012\)095](https://doi.org/10.1007/JHEP11(2012)095) arXiv:[1210.0438](https://arxiv.org/abs/1210.0438) [hep-ph].
- [113] J. Pumplin *et al.*, “New generation of parton distributions with uncertainties from global QCD analysis”, JHEP **0207**, 012 (2002) doi:[10.1088/1126-6708/2002/07/012](https://doi.org/10.1088/1126-6708/2002/07/012) arXiv:[hep-ph/0201195](https://arxiv.org/abs/hep-ph/0201195).
- [114] CMS Collaboration, “Determination of the top-quark pole mass and strong coupling constant from the t t-bar production cross section in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Lett. B **728**, 496 (2014) Erratum: [Phys. Lett. B **738**, 526 (2014)] doi:[10.1016/j.physletb.2014.08.040](https://doi.org/10.1016/j.physletb.2014.08.040), [10.1016/j.physletb.2013.12.009](https://doi.org/10.1016/j.physletb.2013.12.009) arXiv:[1307.1907](https://arxiv.org/abs/1307.1907) [hep-ex].
- [115] CMS Collaboration, “Constraints on parton distribution functions and extraction of the strong coupling constant from the inclusive jet cross section

- in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. C **75**, no. 6, 288 (2015) doi:[10.1140/epjc/s10052-015-3499-1](https://doi.org/10.1140/epjc/s10052-015-3499-1) arXiv:[1410.6765](https://arxiv.org/abs/1410.6765) [hep-ex].
- [116] ATLAS Collaboration, “Measurement of transverse energy-energy correlations in multi-jet events in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector and determination of the strong coupling constant $\alpha_s(m_Z)$ ”, Phys. Lett. B **750**, 427 (2015) doi:[10.1016/j.physletb.2015.09.050](https://doi.org/10.1016/j.physletb.2015.09.050) arXiv:[1508.01579](https://arxiv.org/abs/1508.01579) [hep-ex].
- [117] D0 Collaboration, “Determination of the strong coupling constant from the inclusive jet cross section in $p\bar{p}$ collisions at $\text{sqrt}(s)=1.96$ TeV”, Phys. Rev. D **80**, 111107 (2009) doi:[10.1103/PhysRevD.80.111107](https://doi.org/10.1103/PhysRevD.80.111107) arXiv:[0911.2710](https://arxiv.org/abs/0911.2710) [hep-ex].
- [118] D0 Collaboration, “Measurement of angular correlations of jets at $\sqrt{s} = 1.96$ TeV and determination of the strong coupling at high momentum transfers”, Phys. Lett. B **718**, 56 (2012) doi:[10.1016/j.physletb.2012.10.003](https://doi.org/10.1016/j.physletb.2012.10.003) arXiv:[1207.4957](https://arxiv.org/abs/1207.4957) [hep-ex].
- [119] H1 Collaboration, “Measurement of multijet production in ep collisions at high Q^2 and determination of the strong coupling α_s ”, Eur. Phys. J. C **75**, no. 2, 65 (2015) doi:[10.1140/epjc/s10052-014-3223-6](https://doi.org/10.1140/epjc/s10052-014-3223-6) arXiv:[1406.4709](https://arxiv.org/abs/1406.4709) [hep-ex].
- [120] H1 Collaboration, “Measurement of Jet Production Cross Sections in Deep-inelastic ep Scattering at HERA”, Eur. Phys. J. C **77**, no. 4, 215 (2017) doi:[10.1140/epjc/s10052-017-4717-9](https://doi.org/10.1140/epjc/s10052-017-4717-9) arXiv:[1611.03421](https://arxiv.org/abs/1611.03421) [hep-ex].
- [121] ZEUS Collaboration, “Inclusive-jet photoproduction at HERA and determination of alphas”, Nucl. Phys. B **864**, 1 (2012) doi:[10.1016/j.nuclphysb.2012.06.006](https://doi.org/10.1016/j.nuclphysb.2012.06.006) arXiv:[1205.6153](https://arxiv.org/abs/1205.6153) [hep-ex].
- [122] CDF Collaboration, “Measurement of the strong coupling constant from inclusive jet production at the Tevatron $\bar{p}p$ collider”, Phys. Rev. Lett. **88**, 042001 (2002) doi:[10.1103/PhysRevLett.88.042001](https://doi.org/10.1103/PhysRevLett.88.042001) arXiv:[hep-ex/0108034](https://arxiv.org/abs/hep-ex/0108034).

- [123] R. D. Ball *et al.*, “A first unbiased global NLO determination of parton distributions and their uncertainties”, Nucl. Phys. B **838**, 136 (2010) doi:[10.1016/j.nuclphysb.2010.05.008](https://doi.org/10.1016/j.nuclphysb.2010.05.008) arXiv:[1002.4407](https://arxiv.org/abs/1002.4407) [hep-ph].
- [124] B. Schmidt and M. Steinhauser, “CRunDec: a C++ package for running and decoupling of the strong coupling and quark masses”, Comput. Phys. Commun. **183**, 1845 (2012) doi:[10.1016/j.cpc.2012.03.023](https://doi.org/10.1016/j.cpc.2012.03.023) arXiv:[1201.6149](https://arxiv.org/abs/1201.6149) [hep-ph].
- [125] K. G. Chetyrkin, J. H. Kuhn and M. Steinhauser, “RunDec: A Mathematica package for running and decoupling of the strong coupling and quark masses”, Comput. Phys. Commun. **133**, 43 (2000) doi:[10.1016/S0010-4655\(00\)00155-7](https://doi.org/10.1016/S0010-4655(00)00155-7) arXiv:[hep-ph/0004189](https://arxiv.org/abs/hep-ph/0004189).

List of Publications/Conferences

CMS Public Notes with Direct Contribution

1. CMS Collaboration, “Determination of the strong coupling constant from the measurement of inclusive multijet event cross sections in pp collisions at $\sqrt{s} = 8$ TeV”, **CMS-PAS-SMP-16-008** (2017).
-

CMS Analysis Notes with Direct Contribution

1. A. Kaur *et al.*, “Search for light scalar resonances decaying to b quarks at $\sqrt{s} = 13$ TeV”, **CMS AN-2016/384** (2016).
 2. A. Kaur *et al.*, “Measurement of inclusive multijet cross-sections in pp collisions at $\sqrt{s} = 8$ TeV”, **CMS AN-2015/102** (2015).
-

Papers in Refreed Journals

1. A. Kaur *et al.*, “Extraction of the strong coupling constant from the measurement of inclusive multijet event cross-sections in pp collisions at center of mass energy of 8 TeV”, Springer Proc. Phys. **203** (2018). doi:10.1007/978-3-319-73171-1.

2. A. Kaur *et al.*, “Measurements of event properties and multi-differential jet cross sections and impact of CMS measurements on Proton Structure and QCD parameters”, (submitted to EPJ Web of Conferences)
 3. M. Kaur and Anter P. Kaur, “Subjet Multiplicities at LHC Energies and the QCD Color Factor Ratio C_A/C_F ”, Advances in High Energy Physics, **vol. 2013**, Article ID 585809 (2013) doi:10.1155/2013/585809.
-

Papers presented in Conferences, Workshops and Symposia

1. A. Kaur *et al.*, “Measurements of event properties and multi-differential jet cross sections and impact of CMS measurements on Proton Structure and QCD parameters”, **XLVII International Symposium on Multiparticle Dynamics (ISMD2017)**, 11-15 Sepetember, 2017, Tlaxcala City, Mexico.
2. A. Kaur *et al.*, “Measurement of inclusive multijet cross sections in pp collisions using the CMS detector”, **11th Chandigarh Science Congress, CHASCON 2017**, 9-11 March, 2017, Panjab University, Chandigarh, India.
3. A. Kaur *et al.*, “Extraction of the strong coupling constant from the measurement of inclusive multijet event cross-sections in pp collisions at center of mass energy of 8 TeV”, **XXII DAE-BRNS High Energy Physics Symposium 2016**, 12-16 December, 2016, Delhi, India.
4. A. Kaur *et al.*, “Inclusive jets results from CMS”, **International Workshop on Frontiers in Electroweak Interactions of Leptons and Hadrons**, 2-6 November, 2016, Aligarh, India.
5. A. Kaur *et al.*, “HCAL Back-end Upgradation at CMS Detector”, **9th Chandigarh**

Science Congress, CHASCON 2017, 25-27 February, 2015, Panjab University, Chandigarh, India.

Schools and Workshops attended

1. **Workshop on Upgrade of the CMS Calorimeter**, 23-24 November, 2015, Tata Institute of Fundamental Research, Mumbai, India.
 2. **IX SERC SCHOOL ON EXPERIMENTAL HIGH ENERGY PHYSICS**, 2-21 December, 2013, Indian Institute of Technology Madras, Chennai, India.
 3. **CMS Data Analysis School (CMSDAS)**, 7-11 November, 2013, Saha Institute of Nuclear Physics, Kolkata, India.
-

CMS Publications

1. CMS Collaboration, “Search for heavy resonances decaying to a top quark and a bottom quark in the lepton+jets final state in proton-proton collisions at 13 TeV”, Phys. Lett. B **777**, 39 (2018).
2. CMS Collaboration, “Search for Evidence of the Type-III Seesaw Mechanism in Multilepton Final States in Proton-Proton Collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. Lett. **119**, no. 22, 221802 (2017).
3. CMS Collaboration, “Principal-component analysis of two-particle azimuthal correlations in PbPb and $p\text{Pb}$ collisions at CMS”, Phys. Rev. C **96**, no. 6, 064902 (2017).

4. CMS Collaboration, “Measurement of vector boson scattering and constraints on anomalous quartic couplings from events with four leptons and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **774**, 682 (2017).
5. CMS Collaboration, “Search for a light pseudoscalar Higgs boson produced in association with bottom quarks in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1711**, 010 (2017).
6. CMS Collaboration, “Search for supersymmetry in events with at least one photon, missing transverse momentum, and large transverse event activity in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1712**, 142 (2017).
7. CMS Collaboration, “Measurement of the differential cross sections for the associated production of a W boson and jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **96**, no. 7, 072005 (2017).
8. CMS Collaboration, “Search for direct production of supersymmetric partners of the top quark in the all-jets final state in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1710**, 005 (2017).
9. CMS Collaboration, “Search for heavy resonances that decay into a vector boson and a Higgs boson in hadronic final states at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **77**, no. 9, 636 (2017).
10. CMS Collaboration, “Constraints on anomalous Higgs boson couplings using production and decay information in the four-lepton final state”, Phys. Lett. B **775**, 1 (2017).
11. CMS Collaboration, “Search for Higgs boson pair production in the $bb\tau\tau$ final state in proton-proton collisions at $\sqrt(s) = 8$ TeV”, Phys. Rev. D **96**, no. 7, 072004 (2017).
12. CMS Collaboration, “Measurement of charged pion, kaon, and proton production in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **96**, no. 11, 112003 (2017).

13. CMS Collaboration, “Measurements of properties of the Higgs boson decaying into the four-lepton final state in pp collisions at $\sqrt{s} = 13$ TeV”, JHEP **1711**, 047 (2017).
14. CMS Collaboration, “Search for electroweak production of charginos and neutralinos in WH events in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1711**, 029 (2017).
15. CMS Collaboration, “Search for a heavy composite Majorana neutrino in the final state with two leptons and two quarks at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **775**, 315 (2017).
16. CMS Collaboration, “Measurement of the semileptonic $t\bar{t} + \gamma$ production cross section in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1710**, 006 (2017).
17. CMS Collaboration, “Measurements of jet charge with dijet events in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1710**, 131 (2017).
18. CMS Collaboration, “Particle-flow reconstruction and global event description with the CMS detector”, JINST **12**, no. 10, P10003 (2017).
19. CMS Collaboration, “Search for top squark pair production in pp collisions at $\sqrt{s} = 13$ TeV using single lepton events”, JHEP **1710**, 019 (2017).
20. CMS Collaboration, “Searches for W’ bosons decaying to a top quark and a bottom quark in proton-proton collisions at 13 TeV”, JHEP **1708**, 029 (2017).
21. CMS Collaboration, “Search for new physics in the monophoton final state in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1710**, 073 (2017).
22. CMS Collaboration, “Search for pair production of vector-like T and B quarks in single-lepton final states using boosted jet substructure in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1711**, 085 (2017).
23. CMS Collaboration, “Search for dark matter produced in association with heavy-flavor quark pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **77**, no. 12, 845 (2017).

24. CMS Collaboration, “Search for top quark partners with charge 5/3 in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1708**, 073 (2017).
25. CMS Collaboration, “Search for Low Mass Vector Resonances Decaying to Quark-Antiquark Pairs in Proton-Proton Collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. Lett. **119**, no. 11, 111802 (2017).
26. CMS Collaboration, “Measurements of $t\bar{t}$ cross sections in association with b jets and inclusive jets and their ratio using dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **776**, 355 (2018).
27. CMS Collaboration, “Combination of searches for heavy resonances decaying to WW, WZ, ZZ, WH, and ZH boson pairs in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV”, Phys. Lett. B **774**, 533 (2017).
28. CMS Collaboration, “Measurement of the B^\pm Meson Nuclear Modification Factor in Pb-Pb Collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Rev. Lett. **119**, no. 15, 152301 (2017).
29. CMS Collaboration, “Search for Supersymmetry in pp Collisions at $\sqrt{s} = 13$ TeV in the Single-Lepton Final State Using the Sum of Masses of Large-Radius Jets”, Phys. Rev. Lett. **119**, no. 15, 151802 (2017).
30. CMS Collaboration, “Search for new phenomena with the M_{T2} variable in the all-hadronic final state produced in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **77**, no. 10, 710 (2017).
31. CMS Collaboration, “Search for Charged Higgs Bosons Produced via Vector Boson Fusion and Decaying into a Pair of W and Z Bosons Using pp Collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. Lett. **119**, no. 14, 141802 (2017).
32. CMS Collaboration, “Measurement of the triple-differential dijet cross section in proton-proton collisions at $\sqrt{s} = 8$ TeV and constraints on parton distribution functions”, Eur. Phys. J. C **77**, no. 11, 746 (2017).

33. CMS Collaboration, “Search for black holes in high-multiplicity final states in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **774**, 279 (2017).
34. CMS Collaboration, “Search for supersymmetry in multijet events with missing transverse momentum in proton-proton collisions at 13 TeV”, Phys. Rev. D **96**, no. 3, 032003 (2017).
35. CMS Collaboration, “Search for physics beyond the standard model in events with two leptons of same sign, missing transverse momentum, and jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **77**, no. 9, 578 (2017).
36. CMS Collaboration, “Measurement of the top quark mass in the dileptonic $t\bar{t}$ decay channel using the mass observables $M_{b\ell}$, M_{T2} , and $M_{b\ell\nu}$ in pp collisions at $\sqrt{s}=8$ TeV”, Phys. Rev. D **96**, no. 3, 032002 (2017).
37. CMS Collaboration, “Search for $t\bar{t}$ resonances in highly boosted lepton+jets and fully hadronic final states in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1707**, 001 (2017).
38. CMS Collaboration, “Measurements of the $pp \rightarrow W\gamma\gamma$ and $pp \rightarrow Z\gamma\gamma$ cross sections and limits on anomalous quartic gauge couplings at $\sqrt{s} = 8$ TeV”, JHEP **1710**, 072 (2017).
39. CMS Collaboration, “Search for new physics with dijet angular distributions in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1707**, 013 (2017).
40. CMS Collaboration, “Search for a heavy resonance decaying to a top quark and a vector-like top quark at $\sqrt{s} = 13$ TeV”, JHEP **1709**, 053 (2017).
41. CMS Collaboration, “Measurement of the jet mass in highly boosted $t\bar{t}$ events from pp collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **77**, no. 7, 467 (2017).
42. CMS Collaboration, “Search for anomalous couplings in boosted WW/WZ $\rightarrow \ell\nu q\bar{q}$ production in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **772**, 21 (2017).

43. CMS Collaboration, “Search for associated production of dark matter with a Higgs boson decaying to $b\bar{b}$ or $\gamma\gamma$ at $\sqrt{s} = 13$ TeV”, JHEP **1710**, 180 (2017).
44. CMS Collaboration, “Search for third-generation scalar leptoquarks and heavy right-handed neutrinos in final states with two tau leptons and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1707**, 121 (2017).
45. CMS Collaboration, “Measurement of the top quark mass using single top quark events in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **77**, no. 5, 354 (2017).
46. CMS Collaboration, “Search for dark matter produced with an energetic jet or a hadronically decaying W or Z boson at $\sqrt{s} = 13$ TeV”, JHEP **1707**, 014 (2017).
47. CMS Collaboration, “Measurement of double-differential cross sections for top quark pair production in pp collisions at $\sqrt{s} = 8$ TeV and impact on parton distribution functions”, Eur. Phys. J. C **77**, no. 7, 459 (2017).
48. CMS Collaboration, “Search for standard model production of four top quarks in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **772**, 336 (2017).
49. CMS Collaboration, “Measurement of the cross section for electroweak production of $Z\gamma$ in association with two jets and constraints on anomalous quartic gauge couplings in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **770**, 380 (2017).
50. CMS Collaboration, “Measurement of prompt and nonprompt J/ψ production in pp and pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Eur. Phys. J. C **77**, no. 4, 269 (2017).
51. CMS Collaboration, “Search for associated production of a Z boson with a single top quark and for tZ flavour-changing interactions in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1707**, 003 (2017).
52. CMS Collaboration, “Study of Jet Quenching with $Z +$ jet Correlations in Pb-Pb and pp Collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Rev. Lett. **119**, no. 8, 082301 (2017).

-
- 53. CMS Collaboration, “Azimuthal anisotropy of charged particles with transverse momentum up to 100 GeV/ c in PbPb collisions at $\sqrt{s}_{NN}=5.02$ TeV”, Phys. Lett. B **776**, 195 (2018).
 - 54. CMS Collaboration, “Measurement of the inclusive energy spectrum in the very forward direction in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1708**, 046 (2017).
 - 55. CMS Collaboration, “Search for single production of vector-like quarks decaying into a b quark and a W boson in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **772**, 634 (2017).
 - 56. CMS Collaboration, “Search for single production of vector-like quarks decaying to a Z boson and a top or a bottom quark in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1705**, 029 (2017).
 - 57. CMS Collaboration, “Search for new phenomena with multiple charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **77**, no. 9, 635 (2017).
 - 58. CMS Collaboration, “Measurement of the $t\bar{t}$ production cross section using events with one lepton and at least one jet in pp collisions at $\sqrt{s} = 13$ TeV”, JHEP **1709**, 051 (2017).
 - 59. CMS Collaboration, “Search for dark matter and unparticles in events with a Z boson and missing transverse momentum in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1703**, 061 (2017). Erratum: [JHEP **1709**, 106 (2017).]
 - 60. CMS Collaboration, “Search for light bosons in decays of the 125 GeV Higgs boson in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1710**, 076 (2017).
 - 61. CMS Collaboration, “Mechanical stability of the CMS strip tracker measured with a laser alignment system”, JINST **12**, no. 04, P04023 (2017).

62. CMS Collaboration, “Search for supersymmetry in the all-hadronic final state using top quark tagging in pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **96**, no. 1, 012004 (2017).
63. CMS Collaboration, “Search for leptophobic Z’ bosons decaying into four-lepton final states in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **773**, 563 (2017).
64. CMS Collaboration, “Search for high-mass $Z\gamma$ resonances in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV using jet substructure techniques”, Phys. Lett. B **772**, 363 (2017).
65. CMS Collaboration, “Search for heavy gauge W’ boson in events with an energetic lepton and large missing transverse momentum at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **770**, 278 (2017).
66. CMS Collaboration, “Measurement of electroweak-induced production of $W\gamma$ with two jets in pp collisions at $\sqrt{s} = 8$ TeV and constraints on anomalous quartic gauge couplings”, JHEP **1706**, 106 (2017).
67. CMS Collaboration, “Search for massive resonances decaying into WW, WZ or ZZ bosons in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1703**, 162 (2017).
68. CMS Collaboration, “Measurements of the charm jet cross section and nuclear modification factor in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Lett. B **772**, 306 (2017).
69. CMS Collaboration, “Search for electroweak production of a vector-like quark decaying to a top quark and a Higgs boson using boosted topologies in fully hadronic final states”, JHEP **1704**, 136 (2017).
70. CMS Collaboration, “Searches for pair production of third-generation squarks in $\sqrt{s} = 13$ TeV pp collisions”, Eur. Phys. J. C **77**, no. 5, 327 (2017).

71. CMS Collaboration, “Search for heavy neutrinos or third-generation leptoquarks in final states with two hadronically decaying τ leptons and two jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1703**, 077 (2017).
72. CMS Collaboration, “Search for single production of a heavy vector-like T quark decaying to a Higgs boson and a top quark with a lepton and jets in the final state”, Phys. Lett. B **771**, 80 (2017).
73. CMS Collaboration, “Search for CP violation in $t\bar{t}$ production and decay in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1703**, 101 (2017).
74. CMS Collaboration, “Search for supersymmetry in events with photons and missing transverse energy in pp collisions at 13 TeV”, Phys. Lett. B **769**, 391 (2017).
75. CMS Collaboration, “Search for heavy resonances decaying to tau lepton pairs in proton-proton collisions at $\sqrt{s} = 13$ TeV”, JHEP **1702**, 048 (2017).
76. CMS Collaboration, “Measurements of the associated production of a Z boson and b jets in pp collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **77**, no. 11, 751 (2017).
77. CMS Collaboration, “Measurement of the $t\bar{t}$ production cross section using events in the $e\mu$ final state in pp collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **77**, 172 (2017).
78. CMS Collaboration, “Measurements of differential production cross sections for a Z boson in association with jets in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1704**, 022 (2017).
79. CMS Collaboration, “Search for dijet resonances in proton-proton collisions at $\sqrt{s} = 13$ TeV and constraints on dark matter and other models”, Phys. Lett. B **769**, 520 (2017). Erratum: [Phys. Lett. B **772**, 882 (2017).]
80. CMS Collaboration, “Charged-particle nuclear modification factors in PbPb and pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, JHEP **1704**, 039 (2017).

81. CMS Collaboration, “Suppression of $\Upsilon(1S)$, $\Upsilon(2S)$ and $\Upsilon(3S)$ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, Phys. Lett. B **770**, 357 (2017).
82. CMS Collaboration, “Relative Modification of Prompt ψ (2S) and J/ψ Yields from pp to PbPb Collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Rev. Lett. **118**, no. 16, 162301 (2017).
83. CMS Collaboration, “A search for new phenomena in pp collisions at $\sqrt{s} = 13$ TeV in final states with missing transverse momentum and at least one jet using the α_T variable”, Eur. Phys. J. C **77**, no. 5, 294 (2017).
84. CMS Collaboration, “Searches for invisible decays of the Higgs boson in pp collisions at $\sqrt{s} = 7$, 8, and 13 TeV”, JHEP **1702**, 135 (2017).
85. CMS Collaboration, “Search for heavy resonances decaying into a vector boson and a Higgs boson in final states with charged leptons, neutrinos, and b quarks”, Phys. Lett. B **768**, 137 (2017).
86. CMS Collaboration, “Observation of $\Upsilon(1S)$ pair production in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1705**, 013 (2017).
87. CMS Collaboration, “Search for R-parity violating supersymmetry with displaced vertices in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. D **95**, no. 1, 012009 (2017).
88. CMS Collaboration, “Search for electroweak production of charginos in final states with two τ leptons in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1704**, 018 (2017).
89. CMS Collaboration, “Search for top quark decays via Higgs-boson-mediated flavor-changing neutral currents in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1702**, 079 (2017).
90. CMS Collaboration, “Measurements of differential cross sections for associated production of a W boson and jets in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. D **95**, 052002 (2017).

91. CMS Collaboration, “Measurement of differential cross sections for top quark pair production using the lepton+jets final state in proton-proton collisions at 13 TeV”, Phys. Rev. D **95**, no. 9, 092001 (2017).
92. CMS Collaboration, “Search for anomalous Wtb couplings and flavour-changing neutral currents in t-channel single top quark production in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, JHEP **1702**, 028 (2017).
93. CMS Collaboration, “Search for high-mass $Z\gamma$ resonances in $e^+e^-\gamma$ and $\mu^+\mu^-\gamma$ final states in proton-proton collisions at $\sqrt{s} = 8$ and 13 TeV”, JHEP **1701**, 076 (2017).
94. CMS Collaboration, “Cross section measurement of *t*-channel single top quark production in pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **772**, 752 (2017).
95. CMS Collaboration, “Suppression and azimuthal anisotropy of prompt and non-prompt J/ψ production in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, Eur. Phys. J. C **77**, no. 4, 252 (2017).
96. CMS Collaboration, “Observation of charge-dependent azimuthal correlations in *p*-Pb collisions and its implication for the search for the chiral magnetic effect”, Phys. Rev. Lett. **118**, no. 12, 122301 (2017).
97. CMS Collaboration, “Search for supersymmetry in events with one lepton and multiple jets in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **95**, no. 1, 012011 (2017).
98. CMS Collaboration, “Search for long-lived charged particles in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **94**, no. 11, 112004 (2016).
99. CMS Collaboration, “Inclusive search for supersymmetry using razor variables in pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. D **95**, no. 1, 012003 (2017).
100. CMS Collaboration, “Measurement of the WZ production cross section in pp collisions at $\sqrt{s} = 7$ and 8 TeV and search for anomalous triple gauge couplings at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **77**, no. 4, 236 (2017).

101. CMS Collaboration, “Search for narrow resonances in dilepton mass spectra in proton-proton collisions at $\sqrt{s} = 13$ TeV and combination with 8 TeV data”, Phys. Lett. B **768**, 57 (2017).
102. CMS Collaboration, “Measurement of inclusive jet cross sections in pp and PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, Phys. Rev. C **96**, no. 1, 015202 (2017).
103. CMS Collaboration, “Measurement and QCD analysis of double-differential inclusive jet cross sections in pp collisions at $\sqrt{s} = 8$ TeV and cross section ratios to 2.76 and 7 TeV”, JHEP **1703**, 156 (2017).
104. CMS Collaboration, “Studies of inclusive four-jet production with two b -tagged jets in proton-proton collisions at 7 TeV”, Phys. Rev. D **94**, no. 11, 112005 (2016).
105. CMS Collaboration, “Search for high-mass diphoton resonances in proton-proton collisions at 13 TeV and combination with 8 TeV search”, Phys. Lett. B **767**, 147 (2017).
106. CMS Collaboration, “Decomposing transverse momentum balance contributions for quenched jets in PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, JHEP **1611**, 055 (2016).
107. CMS Collaboration, “The CMS trigger system”, JINST **12**, no. 01, P01020 (2017).
108. CMS Collaboration, “Measurement of the total and differential inclusive B^+ hadron cross sections in pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **771**, 435 (2017).
109. CMS Collaboration, “Measurement of the production cross section of a W boson in association with two b jets in pp collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **77**, no. 2, 92 (2017).
110. CMS Collaboration, “Measurement of the mass of the top quark in decays with a J/ψ meson in pp collisions at 8 TeV”, JHEP **1612**, 123 (2016).

111. CMS Collaboration, “Search for new phenomena in events with high jet multiplicity and low missing transverse momentum in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **770**, 257 (2017).
112. CMS Collaboration, “Measurement of the ZZ production cross section and $Z \rightarrow \ell^+ \ell^- \ell'^+ \ell'^-$ branching fraction in pp collisions at $\sqrt{s}=13$ TeV”, Phys. Lett. B **763**, 280 (2016), Erratum: [Phys. Lett. B **772**, 884 (2017)].
113. CMS Collaboration, “Measurement of electroweak production of a W boson and two forward jets in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1611**, 147 (2016).
114. CMS Collaboration, “Measurement of the WZ production cross section in pp collisions at $\sqrt(s) = 13$ TeV”, Phys. Lett. B **766**, 268 (2017)
115. CMS Collaboration, “Search for dark matter in proton-proton collisions at 8 TeV with missing transverse momentum and vector boson tagged jets”, JHEP **1612**, 083 (2016), Erratum: [JHEP **1708**, 035 (2017)].
116. CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, JINST **12**, no. 02, P02014 (2017)
117. CMS Collaboration, “Search for lepton flavour violating decays of the Higgs boson to $e\tau$ and $e\mu$ in proton–proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **763**, 472 (2016).
118. CMS Collaboration, “Observation of the decay $B^+ \rightarrow \psi(2S)\phi(1020)K^+$ in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **764**, 66 (2017)
119. CMS Collaboration, “Search for new physics in final states with two opposite-sign, same-flavor leptons, jets, and missing transverse momentum in pp collisions at $\text{sqrt}(s) = 13$ TeV”, JHEP **1612**, 013 (2016).
120. CMS Collaboration, “Measurement of the differential cross sections for top quark pair production as a function of kinematic event variables in pp collisions at $\sqrt{s}=7$ and 8 TeV”, Phys. Rev. D **94**, no. 5, 052006 (2016).

121. CMS Collaboration, “Searches for R -parity-violating supersymmetry in pp collisions at $\sqrt{s} = 8$ TeV in final states with 0-4 leptons”, Phys. Rev. D **94**, no. 11, 112009 (2016).
122. CMS Collaboration, “Evidence for collectivity in pp collisions at the LHC”, Phys. Lett. B **765**, 193 (2017)
123. CMS Collaboration, “Measurement of the transverse momentum spectra of weak vector bosons produced in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1702**, 096 (2017)
124. CMS Collaboration, “Search for Resonant Production of High-Mass Photon Pairs in Proton-Proton Collisions at $\sqrt{s} = 8$ and 13 TeV”, Phys. Rev. Lett. **117**, no. 5, 051802 (2016).
125. CMS Collaboration, “Phenomenological MSSM interpretation of CMS searches in pp collisions at $\text{sqrt}(s) = 7$ and 8 TeV”, JHEP **1610**, 129 (2016).
126. CMS Collaboration, “Measurement of the transverse momentum spectrum of the Higgs boson produced in pp collisions at $\sqrt{s} = 8$ TeV using $H \rightarrow WW$ decays”, JHEP **1703**, 032 (2017)
127. CMS Collaboration, “Search for Dark Matter and Supersymmetry with a Compressed Mass Spectrum in the Vector Boson Fusion Topology in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. Lett. **118**, no. 2, 021802 (2017)
128. CMS Collaboration, “Measurement of the W boson helicity fractions in the decays of top quark pairs to lepton + jets final states produced in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **762**, 512 (2016).
129. CMS Collaboration, “Search for top squark pair production in compressed-mass-spectrum scenarios in proton-proton collisions at $\sqrt{s} = 8$ TeV using the α_T variable”, Phys. Lett. B **767**, 403 (2017)

-
- 130. CMS Collaboration, “Coherent J/ψ photoproduction in ultra-peripheral PbPb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the CMS experiment”, Phys. Lett. B **772**, 489 (2017)
 - 131. CMS Collaboration, “Multiplicity and rapidity dependence of strange hadron production in pp, pPb, and PbPb collisions at the LHC”, Phys. Lett. B **768**, 103 (2017)
 - 132. CMS Collaboration, “Search for supersymmetry in pp collisions at $\sqrt{s} = 13$ TeV in the single-lepton final state using the sum of masses of large-radius jets”, JHEP **1608**, 122 (2016).
 - 133. CMS Collaboration, “Measurement of the double-differential inclusive jet cross section in proton–proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **76**, no. 8, 451 (2016).
 - 134. CMS Collaboration, “Search for new physics in same-sign dilepton events in proton–proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. C **76**, no. 8, 439 (2016).
 - 135. CMS Collaboration, “Search for Higgs boson off-shell production in proton-proton collisions at 7 and 8 TeV and derivation of constraints on its total decay width”, JHEP **1609**, 051 (2016).
 - 136. CMS Collaboration, “Measurement of the integrated and differential $t\bar{t}$ production cross sections for high- p_t top quarks in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. D **94**, no. 7, 072002 (2016).
 - 137. CMS Collaboration, “Search for narrow resonances in dijet final states at $\sqrt{(s)} = 8$ TeV with the novel CMS technique of data scouting”, Phys. Rev. Lett. **117**, no. 3, 031802 (2016).
 - 138. CMS Collaboration, “Pseudorapidity dependence of long-range two-particle correlations in $p\text{Pb}$ collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Rev. C **96**, no. 1, 014915 (2017)

139. CMS Collaboration, “Search for lepton flavour violating decays of heavy resonances and quantum black holes to an $e\mu$ pair in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **76**, no. 6, 317 (2016).
140. CMS Collaboration, “Evidence for exclusive $\gamma\gamma \rightarrow W^+W^-$ production and constraints on anomalous quartic gauge couplings in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* **1608**, 119 (2016).
141. CMS Collaboration, “Search for dark matter particles in proton-proton collisions at $\sqrt{s} = 8$ TeV using the razor variables”, *JHEP* **1612**, 088 (2016).
142. CMS Collaboration, “Search for two Higgs bosons in final states containing two photons and two bottom quarks in proton-proton collisions at 8 TeV”, *Phys. Rev. D* **94**, no. 5, 052012 (2016).
143. CMS Collaboration, “Measurement of the top quark mass using charged particles in pp collisions at $\sqrt{s} = 8$ TeV”, *Phys. Rev. D* **93**, no. 9, 092006 (2016).
144. CMS Collaboration, “Measurements of $t\bar{t}$ charge asymmetry using dilepton final states in pp collisions at $\sqrt{s} = 8$ TeV”, *Phys. Lett. B* **760**, 365 (2016).
145. CMS Collaboration, “Search for new physics with the M_{T2} variable in all-jets final states produced in pp collisions at $\sqrt{s} = 13$ TeV”, *JHEP* **1610**, 006 (2016).
146. CMS Collaboration, “Search for neutral resonances decaying into a Z boson and a pair of b jets or τ leptons”, *Phys. Lett. B* **759**, 369 (2016).
147. CMS Collaboration, “ $\Upsilon(nS)$ polarizations versus particle multiplicity in pp collisions at $\sqrt{s} = 7$ TeV”, *Phys. Lett. B* **761**, 31 (2016).
148. CMS Collaboration, “Search for s channel single top quark production in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, *JHEP* **1609**, 027 (2016).

-
- 149. CMS Collaboration, “Measurement of the t-tbar production cross section in the e-mu channel in proton-proton collisions at sqrt(s) = 7 and 8 TeV”, JHEP **1608**, 029 (2016).
 - 150. CMS Collaboration, “Search for heavy Majorana neutrinos in ee+ jets and e mu+jets events in proton-proton collisions at sqrt(s) = 8 TeV”, JHEP **1604**, 169 (2016).
 - 151. CMS Collaboration, “Measurement of the differential cross section and charge asymmetry for inclusive pp → W $^{\pm}$ + X production at sqrt(s) = 8 TeV”, Eur. Phys. J. C **76**, no. 8, 469 (2016).
 - 152. CMS Collaboration, “Search for direct pair production of supersymmetric top quarks decaying to all-hadronic final states in pp collisions at sqrt(s) = 8 TeV”, Eur. Phys. J. C **76**, no. 8, 460 (2016).
 - 153. CMS Collaboration, “Measurements of the t-tbar production cross section in lepton+jets final states in pp collisions at 8 TeV and ratio of 8 to 7 TeV cross sections”, Eur. Phys. J. C **77**, no. 1, 15 (2017)
 - 154. CMS Collaboration, “Search for supersymmetry in electroweak production with photons and large missing transverse energy in pp collisions at sqrt(s) = 8 TeV”, Phys. Lett. B **759**, 479 (2016).
 - 155. CMS Collaboration, “Search for heavy resonances decaying to two Higgs bosons in final states containing four b quarks”, Eur. Phys. J. C **76**, no. 7, 371 (2016).
 - 156. CMS Collaboration, “Measurement of the Zgamma → nu-bar nu gamma production cross section in pp collisions at sqrt(s) = 8 TeV and limits on anomalous ZZgamma and Zgamma-gamma trilinear gauge boson couplings”, Phys. Lett. B **760**, 448 (2016).
 - 157. CMS Collaboration, “Search for supersymmetry in the multijet and missing transverse momentum final state in pp collisions at 13 TeV”, Phys. Lett. B **758**, 152 (2016).

158. CMS Collaboration, “Measurement of dijet azimuthal decorrelation in pp collisions at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **76**, no. 10, 536 (2016).
159. CMS Collaboration, “Search for R-parity violating decays of a top squark in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *Phys. Lett. B* **760**, 178 (2016).
160. CMS Collaboration, “Combined search for anomalous pseudoscalar HVV couplings in $VH(H \rightarrow b\bar{b})$ production and $H \rightarrow VV$ decay”, *Phys. Lett. B* **759**, 672 (2016).
161. CMS Collaboration, “Search for direct pair production of scalar top quarks in the single- and dilepton channels in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **1607**, 027 (2016), Erratum: [*JHEP* **1609**, 056 (2016)].
162. CMS Collaboration, “Search for supersymmetry in pp collisions at $\text{sqrt}(s) = 8$ TeV in final states with boosted W bosons and b jets using razor variables”, *Phys. Rev. D* **93**, no. 9, 092009 (2016).
163. CMS Collaboration, “Azimuthal decorrelation of jets widely separated in rapidity in pp collisions at $\sqrt{s} = 7$ TeV”, *JHEP* **1608**, 139 (2016).
164. CMS Collaboration, “Search for massive WH resonances decaying into the $\ell\nu b\bar{b}$ final state at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **76**, no. 5, 237 (2016).
165. CMS Collaboration, “Forward–backward asymmetry of Drell–Yan lepton pairs in pp collisions at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **76**, no. 6, 325 (2016).
166. CMS Collaboration, “Measurement of inclusive jet production and nuclear modifications in pPb collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV”, *Eur. Phys. J. C* **76**, no. 7, 372 (2016).
167. CMS Collaboration, “Measurements of t t-bar spin correlations and top quark polarization using dilepton final states in pp collisions at $\text{sqrt}(s) = 8$ TeV”, *Phys. Rev. D* **93**, no. 5, 052007 (2016).

-
- 168. CMS Collaboration, “Correlations between jets and charged particles in PbPb and pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, JHEP **1602**, 156 (2016).
 - 169. CMS Collaboration, “Measurement of differential and integrated fiducial cross sections for Higgs boson production in the four-lepton decay channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, JHEP **1604**, 005 (2016).
 - 170. CMS Collaboration, “Search for supersymmetry in events with soft leptons, low jet multiplicity, and missing transverse energy in proton–proton collisions at $\sqrt{s}=8$ TeV”, Phys. Lett. B **759**, 9 (2016).
 - 171. CMS Collaboration, “Study of Z boson production in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Lett. B **759**, 36 (2016).
 - 172. CMS Collaboration, “Measurement of the inclusive jet cross section in pp collisions at $\sqrt{s} = 2.76$ TeV”, Eur. Phys. J. C **76**, no. 5, 265 (2016).
 - 173. CMS Collaboration, “Search for narrow resonances decaying to dijets in proton–proton collisions at $\sqrt(s) = 13$ TeV”, Phys. Rev. Lett. **116**, no. 7, 071801 (2016).
 - 174. CMS Collaboration, “Event generator tunes obtained from underlying event and multiparton scattering measurements”, Eur. Phys. J. C **76**, no. 3, 155 (2016).
 - 175. CMS Collaboration, “Search for dark matter and unparticles produced in association with a Z boson in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. D **93**, no. 5, 052011 (2016).
 - 176. CMS Collaboration, “Measurement of spin correlations in $t\bar{t}$ production using the matrix element method in the muon+jets final state in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **758**, 321 (2016).
 - 177. CMS Collaboration, “Search for anomalous single top quark production in association with a photon in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1604**, 035 (2016).

178. CMS Collaboration, “Search for a low-mass pseudoscalar Higgs boson produced in association with a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **758**, 296 (2016).
179. CMS Collaboration, “Measurement of top quark polarisation in t-channel single top quark production”, JHEP **1604**, 073 (2016).
180. CMS Collaboration, “Search for excited leptons in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1603**, 125 (2016).
181. CMS Collaboration, “Reconstruction and identification of τ lepton decays to hadrons and ν_τ at CMS”, JINST **11**, no. 01, P01019 (2016).
182. CMS Collaboration, “Search for a very light NMSSM Higgs boson produced in decays of the 125 GeV scalar boson and decaying into τ leptons in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1601**, 079 (2016).
183. CMS Collaboration, “Measurement of the top quark pair production cross section in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. Lett. **116**, no. 5, 052002 (2016).
184. CMS Collaboration, “Search for a light charged Higgs boson decaying to $c\bar{s}$ in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1512**, 178 (2015).
185. CMS Collaboration, “Transverse momentum spectra of inclusive b jets in pPb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Lett. B **754**, 59 (2016).
186. CMS Collaboration, “Measurement of $t\bar{t}$ production with additional jet activity, including b quark jets, in the dilepton decay channel using pp collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **76**, no. 7, 379 (2016).
187. CMS Collaboration, “Measurement of long-range near-side two-particle angular correlations in pp collisions at $\sqrt{s} = 13$ TeV”, Phys. Rev. Lett. **116**, no. 17, 172302 (2016).

188. CMS Collaboration, “Searches for a heavy scalar boson H decaying to a pair of 125 GeV Higgs bosons hh or for a heavy pseudoscalar boson A decaying to Zh, in the final states with $h \rightarrow \tau\tau$ ”, *Phys. Lett. B* **755**, 217 (2016).
189. CMS Collaboration, “Observation of top quark pairs produced in association with a vector boson in pp collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **1601**, 096 (2016).
190. CMS Collaboration, “Measurement of transverse momentum relative to dijet systems in PbPb and pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV”, *JHEP* **1601**, 006 (2016).
191. CMS Collaboration, “Search for the associated production of a Higgs boson with a single top quark in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **1606**, 177 (2016).
192. CMS Collaboration, “Search for the production of an excited bottom quark decaying to tW in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **1601**, 166 (2016).
193. CMS Collaboration, “Measurement of the $t\bar{t}$ production cross section in the all-jets final state in pp collisions at $\sqrt{s} = 8$ TeV”, *Eur. Phys. J. C* **76**, no. 3, 128 (2016).
194. CMS Collaboration, “Search for $W' \rightarrow tb$ in proton-proton collisions at $\sqrt{s} = 8$ TeV”, *JHEP* **1602**, 122 (2016).
195. CMS Collaboration, “Search for vector-like charge 2/3 T quarks in proton-proton collisions at $\text{sqrt}(s) = 8$ TeV”, *Phys. Rev. D* **93**, no. 1, 012003 (2016).
196. CMS Collaboration, “Measurement of the top quark mass using proton-proton data at $\sqrt{(s)} = 7$ and 8 TeV”, *Phys. Rev. D* **93**, no. 7, 072004 (2016).
197. CMS Collaboration, “Measurement of the inelastic cross section in proton–lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, *Phys. Lett. B* **759**, 641 (2016).
198. CMS Collaboration, “Search for single production of scalar leptoquarks in proton–proton collisions at $\sqrt{s} = 8$ TeV”, *Phys. Rev. D* **93**, no. 3, 032005 (2016), Erratum: [Phys. Rev. D **95**, no. 3, 039906 (2017)].

199. CMS Collaboration, “Search for pair production of first and second generation leptoquarks in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. D **93**, no. 3, 032004 (2016).
200. CMS Collaboration, “Measurement of differential cross sections for Higgs boson production in the diphoton decay channel in pp collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. C **76**, no. 1, 13 (2016).
201. CMS Collaboration, “Search for a charged Higgs boson in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1511**, 018 (2015).
202. CMS Collaboration, “Search for supersymmetry in the vector-boson fusion topology in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1511**, 189 (2015).
203. CMS Collaboration, “Study of B Meson Production in p+Pb Collisions at $\sqrt{s_{NN}} = 5.02$ TeV Using Exclusive Hadronic Decays”, Phys. Rev. Lett. **116**, no. 3, 032301 (2016).
204. CMS Collaboration, “Search for W’ decaying to tau lepton and neutrino in proton-proton collisions at $\sqrt(s) = 8$ TeV”, Phys. Lett. B **755**, 196 (2016).
205. CMS Collaboration, “Measurement of the charge asymmetry in top quark pair production in pp collisions at $\sqrt(s) = 8$ TeV using a template method”, Phys. Rev. D **93**, no. 3, 034014 (2016).
206. CMS Collaboration, “Search for neutral MSSM Higgs bosons decaying to $\mu^+ \mu^-$ in pp collisions at $\sqrt{s} = 7$ and 8 TeV”, Phys. Lett. B **752**, 221 (2016).
207. CMS Collaboration, “Search for supersymmetry in events with a photon, a lepton, and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **757**, 6 (2016).
208. CMS Collaboration, “Angular analysis of the decay $B^0 \rightarrow K^{*0} \mu^+ \mu^-$ from pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **753**, 424 (2016).

209. CMS Collaboration, “Measurement of the CP-violating weak phase ϕ_s and the decay width difference $\Delta\Gamma_s$ using the $B_s^0 \rightarrow J/\psi\phi(1020)$ decay channel in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **757**, 97 (2016).
210. CMS Collaboration, “Measurement of the underlying event activity using charged-particle jets in proton-proton collisions at $\text{sqrt}(s) = 2.76$ TeV”, JHEP **1509**, 137 (2015).
211. CMS Collaboration, “Search for pair-produced vectorlike B quarks in proton-proton collisions at $\sqrt{s}=8$ TeV”, Phys. Rev. D **93**, no. 11, 112009 (2016).
212. CMS Collaboration, “Limits on the Higgs boson lifetime and width from its decay to four charged leptons”, Phys. Rev. D **92**, no. 7, 072010 (2015).
213. CMS Collaboration, “Pseudorapidity distribution of charged hadrons in proton-proton collisions at $\sqrt{s} = 13$ TeV”, Phys. Lett. B **751**, 143 (2015).
214. CMS Collaboration, “Measurement of the W^+W^- cross section in pp collisions at $\sqrt{s} = 8$ TeV and limits on anomalous gauge couplings”, Eur. Phys. J. C **76**, no. 7, 401 (2016).
215. CMS Collaboration, “Inclusive and differential measurements of the $t\bar{t}$ charge asymmetry in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **757**, 154 (2016).
216. CMS Collaboration, “Search for a Higgs boson decaying into $\gamma^*\gamma \rightarrow \ell\ell\gamma$ with low dilepton mass in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **753**, 341 (2016).
217. CMS Collaboration, “Search for supersymmetry with photons in pp collisions at $\sqrt{s}=8$ TeV”, Phys. Rev. D **92**, no. 7, 072006 (2015).
218. CMS Collaboration, “Search for exotic decays of a Higgs boson into undetectable particles and one or more photons”, Phys. Lett. B **753**, 363 (2016).
219. CMS Collaboration, “Search for neutral MSSM Higgs bosons decaying into a pair of bottom quarks”, JHEP **1511**, 071 (2015).

220. CMS Collaboration, “Search for resonant $t\bar{t}$ production in proton-proton collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. D **93**, no. 1, 012001 (2016).
221. CMS Collaboration, “Search for diphoton resonances in the mass range from 150 to 850 GeV in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. B **750**, 494 (2015).
222. CMS Collaboration, “Search for a massive resonance decaying into a Higgs boson and a W or Z boson in hadronic final states in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1602**, 145 (2016).
223. CMS Collaboration, “Search for the standard model Higgs boson produced through vector boson fusion and decaying to $b\bar{b}$ ”, Phys. Rev. D **92**, no. 3, 032008 (2015).
224. CMS Collaboration, “A search for pair production of new light bosons decaying into muons”, Phys. Lett. B **752**, 146 (2016).
225. CMS Collaboration, “Search for neutral color-octet weak-triplet scalar particles in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1509**, 201 (2015).
226. CMS Collaboration, “Comparison of the Z/γ^* + jets to γ + jets cross sections in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **1510**, 128 (2015), Erratum: [JHEP **1604**, 010 (2016)].
227. CMS Collaboration, “Search for a pseudoscalar boson decaying into a Z boson and the 125 GeV Higgs boson in $\ell^+\ell^-b\bar{b}$ final states”, Phys. Lett. B **748**, 221 (2015).
228. CMS Collaboration, “Search for a Higgs boson in the mass range from 145 to 1000 GeV decaying to a pair of W or Z bosons”, JHEP **1510**, 144 (2015).
229. CMS Collaboration, “Search for Third-Generation Scalar Leptoquarks in the $t\tau$ Channel in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV”, JHEP **1507**, 042 (2015), Erratum: [JHEP **1611**, 056 (2016)].

230. CMS Collaboration, “Searches for third-generation squark production in fully hadronic final states in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **1506**, 116 (2015).
231. CMS Collaboration, “Evidence for transverse momentum and pseudorapidity dependent event plane fluctuations in PbPb and pPb collisions”, Phys. Rev. C **92**, no. 3, 034911 (2015).

*Selected
Reprint*

CMS Physics Analysis Summary

Contact: cms-pag-conveners-smp@cern.ch

2017/02/21

Determination of the strong coupling constant from the measurement of inclusive multijet event cross sections in pp collisions at $\sqrt{s} = 8$ TeV

The CMS Collaboration

Abstract

A measurement of inclusive multijet event cross sections is presented from proton-proton collisions recorded at $\sqrt{s} = 8$ TeV with the CMS detector and corresponding to an integrated luminosity of 19.7 fb^{-1} . Jets are reconstructed with the anti- k_t clustering algorithm for a jet size parameter $R = 0.7$ in a phase space region ranging up to jet transverse momenta p_T of 2.0 TeV and an absolute rapidity of $|y| = 2.5$. The inclusive 2-jet and 3-jet event cross sections are measured as a function of the average p_T of the two leading jets. The data are well described by predictions at next-to-leading order in perturbative quantum chromodynamics and additionally are compared to several Monte Carlo event generators. The strong coupling constant at the scale of the Z boson mass is inferred from a fit of the ratio of the 3-jet over 2-jet event cross section giving $\alpha_s(M_Z) = 0.1150 \pm 0.0010 \text{ (exp)} \pm 0.0013 \text{ (PDF)} \pm 0.0015 \text{ (NP)} {}^{+0.0050}_{-0.0000} \text{ (scale)}$.

1 Introduction

Inelastic collisions of protons are viewed as interactions between their constituent partons, the (anti-)quarks and gluons. Within the context of perturbative quantum chromodynamics (pQCD), the cross section of a high- p_T scattering process can be expressed as a sum of terms with increasing powers of the strong coupling constant, α_S , convoluted with the parton momentum distribution functions (PDFs) of the proton. The lowest-order α_S^2 term represents the production of two-parton final states. Terms of higher-order α_S^3, \dots in the expansion signify the existence of multi-parton final states. The theoretical description of the transition from strongly interacting, colored partons to color-neutral hadrons, which are observable in detectors, relies for this nonperturbative phase on models implemented in Monte Carlo (MC) event generators. To relate the collimated sprays of colorless hadrons to the initiating partons, jet algorithms are applied. These algorithms bundle together particles that are close in phase space and primarily move into the same direction. Hence, the constructed jets preserve energy and momentum of the initial partons so that the structure of the final jet system mirrors, to a large extent, the topology of the initial partonic system.

The inclusive jet cross section, $pp \rightarrow \text{jet} + X$, as a function of jet p_T and rapidity y is a fundamental observable providing essential information about the PDFs and the strong coupling constant. Corresponding measurements conducted by the experiments at the CERN LHC are reported in Refs. [1–11]. The investigation of inclusive multijet event cross sections $\sigma_{i\text{-jet}}$, $pp \rightarrow i\text{ jets} + X$, as suggested here, permits more elaborate tests of QCD to be performed by subdividing the observed jet event sample into classes according to the presumed minimal power in α_S necessary to describe theoretically such a topology. Moreover, the ratios of such cross sections, $R_{mn} = \frac{\sigma_{m\text{-jet}}}{\sigma_{n\text{-jet}}}$, with $m > n$, are proportional to α_S^{m-n} while at the same time numerous theoretical and experimental uncertainties cancel. Thus, they provide an ideal tool to determine the strong coupling constant $\alpha_s(M_Z)$. A previous analysis of the ratio R_{32} as a function of the average transverse momentum, $\langle p_{T1,2} \rangle$, of the two leading p_T jets in the event was performed at $\sqrt{s} = 7\text{ TeV}$ by the CMS Collaboration and lead to an extraction of $\alpha_s(M_Z) = 0.1148 \pm 0.0055$, where the dominant uncertainty stems from the estimation of higher-order corrections to the next-to-leading order (NLO) prediction [12].

In this analysis, a measurement of inclusive 2- and 3-jet event cross sections is presented using an event sample collected by the CMS experiment during 2012 at the LHC and corresponding to an integrated luminosity of 19.7 fb^{-1} of pp collisions at a centre-of-mass energy of 8 TeV . Jets are reconstructed using the infrared- and collinear-safe anti- k_t clustering algorithm [13] with a jet size parameter R of 0.7. All jets are required to satisfy $p_T > 150\text{ GeV}$ and $|y| < 5.0$. The event sample is further reduced by requiring the two leading p_T jets to lie in the central detector region of $|y| < 2.5$.

The event scale is chosen as before to be the average transverse momentum of the two leading jets, but will be referred to as $H_{T,2}/2$ in this analysis. Fits of the strong coupling constant are performed for the 2-jet and 3-jet event cross sections separately and for their ratio R_{32} .

2 Event selection and reconstruction

The measurement uses data samples which were collected with six single-jet high-level triggers (HLT) [14]. They are seeded by Level 1 (L1) triggers based on calorimetric information and require at least one jet in the event with corrected jet $p_T > 80, 140, 200, 260$, and 320 GeV . All except the highest-threshold trigger were prescaled during the 2012 run. The efficiency of each trigger is estimated using lower- p_T -threshold triggers and it is found to be more than 99% in

the $H_{\mathrm{T},2}/2$ ranges, shown in Table 1 with the corresponding effective integrated luminosities.

Table 1: Trigger regions defined as ranges of the $H_{\mathrm{T},2}/2$ for every single-jet trigger used in the inclusive multijet cross section measurement along with the effective integrated luminosities.

HLT path	$H_{\mathrm{T},2}/2$ range (GeV)	Integrated Luminosity (pb^{-1})
PFJet80	120 – 188	2.12
PFJet140	188 – 263	5.57×10
PFJet200	263 – 345	2.61×10^2
PFJet260	345 – 406	1.06×10^3
PFJet320	406 – 5000	1.97×10^4

In the CMS experiment, all particles are reconstructed and identified using a particle-flow (PF) algorithm, which combines the information from the individual subdetectors [15, 16]. The four-vectors of particle candidates, reconstructed by the above technique, are used as input to the anti- k_t jet-clustering algorithm. The clustering is performed within the FASTJET package [17] using four-momentum summation.

The reconstructed jets require additional energy corrections to account for residual nonuniformities and nonlinearities in the detector response. These jet energy corrections [18] are derived using simulated events, generated by PYTHIA 6.4 [19] with tune Z2* [20] and processed through the CMS detector simulation based on GEANT4 [21], and in situ measurements with dijet, photon+jet, and Z+jet events. The jet energy corrections, which depend on the η (pseudorapidity) and p_{T} of the jet, are applied to the jet four-momentum vector as a multiplicative factor [18]. For a jet with a p_{T} of 100 GeV, the typical correction is about 10%, and decreases with increasing p_{T} . An additional offset correction is applied to take into account the extra energy clustered into jets from additional proton-proton interactions within the same or neighbouring bunch crossings (in-time and out-of-time pileup) [18]. Pileup effects are important only for jets with low p_{T} and become negligible for jets with $p_{\mathrm{T}} > 200$ GeV. The current measurement is therefore largely insensitive to pileup effects.

Each selected event is required to have at least one offline-reconstructed vertex [22] along the beam line that is within 24 cm of the nominal interaction point. To suppress nonphysical jets, i.e. jets resulting from noise in the electromagnetic calorimeter (ECAL) and/or the brass/scintillator hadron calorimeter (HCAL), tight identification criteria [23] are applied : each jet should contain at least two particles, one of which is a charged hadron, and the jet energy fraction carried by neutral hadrons and photons should be less than 90%. These criteria have an efficiency greater than 99% for physical jets. Jets not satisfying the tight identification requirements are discarded.

A sample of multijet events is selected which has two or more jets with transverse momentum greater than 150 GeV and $|y| < 5.0$ in the event. Events, in which the two leading p_{T} jets have $|y| < 2.5$ are selected. Further jets are counted only, if they lie within the same central rapidity range of $|y| < 2.5$.

In QCD, pure jet events are balanced in p_{T} and thus exhibit a low level of missing transverse energy, which predominantly is caused by jet calibration and resolution effects of the detector. Therefore, the ratio of missing transverse energy to the total transverse energy $\frac{E_{\mathrm{T}}^{\mathrm{miss}}}{\sum E_{\mathrm{T}}}$, both derived from the reconstructed particle-flow objects, is required to be less than 0.3 to select well measured jet events.

3 Measurement of the inclusive 2-jet and 3-jet event cross sections

The inclusive differential multijet cross sections are measured as a function of the average transverse momentum, $H_{T,2}/2 = \frac{1}{2}(p_{T,1} + p_{T,2})$, where $p_{T,1}$ and $p_{T,2}$ denote the transverse momenta of the two leading jets. For inclusive 2-jet events sufficient data are available up to $H_{T,2}/2 = 2$ TeV, while for inclusive 3-jet events (and the ratio R_{32}) the accessible range in $H_{T,2}/2$ is limited to $H_{T,2}/2 < 1.68$ TeV. In the following, results for the inclusive 2-jet and 3-jet event selections will be labelled as $n_j \geq 2$ and $n_j \geq 3$, respectively.

The inclusive differential jet event cross section is defined as :

$$\frac{d\sigma}{d(H_{T,2}/2)} = \frac{1}{\epsilon \mathcal{L}_{\text{int,eff}}} \frac{N_{\text{event}}}{\Delta(H_{T,2}/2)} \quad (1)$$

where ϵ is the product of the trigger and jet selection efficiencies, which are greater than 99%, $\mathcal{L}_{\text{int,eff}}$ is the effective integrated luminosity, N_{event} is the number of 2- or 3-jet events counted in an $H_{T,2}/2$ bin, and $\Delta(H_{T,2}/2)$ are the bin widths. The measurements are reported in units of (pb/GeV).

To compare the measured cross sections with theoretical predictions at particle level, an unfolding procedure based on the iterative D'Agostini method [24] as implemented in the ROOUNFOLD software package [25] is applied. This correction for detector resolution effects is regularized by an early stopping after four iterations similarly as in a previously published 3-jet measurement [26], which prevents the buildup of large-scale correlations. The response matrix describes the mapping between the particle-level $H_{T,2}/2$ spectrum and the reconstructed $H_{T,2}/2$ spectrum. To construct the response matrix, the particle-level $H_{T,2}/2$ spectrum is taken from a fit to the theoretically predicted $H_{T,2}/2$ spectrum. The reconstructed $H_{T,2}/2$ spectrum is obtained by smearing this particle-level prediction. The resolution in $H_{T,2}/2$ is evaluated from CMS detector simulation based on the MADGRAPH5 + PYTHIA6 MC event generator using a jet-based smearing according to the jet energy resolution (JER). The JER from simulation is corrected (increased) for residual differences between data and simulation following Ref. [18].

Figure 1 shows the response matrices derived using a Toy MC procedure for inclusive 2-jet (left) and 3-jet events (right). The matrices are normalized to the number of generated events in each column and are mostly diagonal with small off-diagonal elements describing migrations between close-by $H_{T,2}/2$ bins.

Through the unfolding procedure the final statistical uncertainties become correlated among bins. The size of these correlations varies typically between 10 and 20%. As a consequence, statistical fluctuations present in data before the unfolding might affect neighbouring bins after the unfolding. The fluctuations observed in the unfolded spectrum are compatible with the statistical uncertainties of the underlying detector-level distributions and the statistical uncertainty after unfolding increases with respect to the original one of the measured data.

The unfolding procedure is affected by uncertainties of the JER. Alternative response matrices, which were built by varying the JER one standard deviation up and down [18], are used to unfold the measured spectra for comparison and introduce a corresponding uncertainty on the cross sections. In addition, to account for a model dependence of the theoretical $H_{T,2}/2$ spectrum, two different functions are assumed when fitting the theoretically predicted $H_{T,2}/2$ spectra. Finally, a supplementary uncertainty is attributed by comparison to an unfolding with a 30% reduced resolution as compared to the one extracted from simulation. This accounts for shortcomings in the detector simulation of the theory spectra leading to small nonclosures




Figure 1: Response matrices derived using a Toy MC procedure for the inclusive 2-jet (left) and 3-jet event samples (right).

observed in the unfolding. All three uncertainties are added quadratically to give the unfolding uncertainty, which increases from about 1% at low $H_{T,2}/2$ up to 2% at the high $H_{T,2}/2$ end of the cross sections.

The dominant detector-related contribution to the experimental systematic uncertainty of the measured cross sections is caused by the jet energy corrections (JEC) [18]. The JEC uncertainty ranges for inclusive 2-jet events from 3% to 10% and for inclusive 3-jet events from 3% to 8%, respectively.

The uncertainty on the integrated luminosity, which propagates directly to the cross sections, is 2.6% [27] and at low $H_{T,2}/2$ is of a similar size as the one from the JEC. To account for residual effects of small inefficiencies from triggering and jet identification, an uncorrelated uncertainty of 1% is assumed across all $H_{T,2}/2$ bins, similar as in previous CMS jet cross-section measurements [8].

The total experimental systematic uncertainty on the measured cross section is obtained by summing in quadrature the single contributions. The upper panels of Fig. 2 give an overview of all experimental uncertainties affecting the cross section measurement for inclusive 2-jet (top left) and 3-jet events (top right). The error bars indicate the statistical uncertainty after unfolding. The colored lines represent the systematic uncertainties resulting from JEC, the luminosity, residual effects, and the unfolding including JER effects. The total experimental uncertainty, indicated by dashed black lines, is calculated by adding in quadrature all the sources of uncertainty. Beyond about 1.4 (1.2) TeV the statistical uncertainty is dominating for the 2-jet (3-jet) event cross sections, respectively.

The cross section ratio R_{32} as a function of $H_{T,2}/2$ is extracted from the ratio of unfolded differential cross sections for each bin in $H_{T,2}/2$. The systematic experimental uncertainties are propagated from the cross sections to the ratio taking into account correlations. The uncertainties due to luminosity and residual effects cancel completely in this ratio. The statistical uncertainty including bin-by-bin correlations and statistical correlations between the 3-jet and 2-jet event cross sections is derived by directly unfolding the measured ratio R_{32} . Figure 2 “bottom panel” presents an overview of all experimental uncertainties affecting the cross section ratio R_{32} . The JEC and unfolding uncertainties for R_{32} amount to about 1-2% and $\approx 1\%$, respectively. The total uncertainty, calculated by adding in quadrature all individual sources of uncertainty, is dominated by statistical effects beyond about 0.8 TeV in $H_{T,2}/2$.




Figure 2: Overview of all experimental uncertainties affecting the inclusive 2-jet (top left) and 3-jet event cross sections (top right) and their ratio R_{32} (bottom). The error bars indicate the statistical uncertainty after unfolding. The colored lines represent the systematic uncertainties resulting from JEC, the luminosity, residual effects, and the unfolding including JER effects. Uncertainties due to luminosity and residual effects are cancelled completely in the ratio. The total experimental uncertainty, indicated by dashed black lines, is calculated by adding in quadrature all the sources of uncertainty.

4 Theory predictions

Predictions at NLO accuracy in pQCD are computed with the NLOJET++ program version 4.1.3 [28, 29]. The results are provided within the framework of FASTNLO version 2.3 [30] for use within fits. The renormalization and factorization scales μ_r and μ_f are chosen equal to $H_{\text{T},2}/2$. PDF sets at NLO available for a series of different assumptions on $\alpha_s(M_Z)$ via the LHAPDF6 package [31] are listed in Table 2. All sets employ a variable-flavour number scheme with at most five or six flavours apart from the ABM11 PDFs, which use a fixed-flavour number scheme with $N_F = 5$.

Out of these eight PDF sets the following three will not be considered further:

- At NLO, predictions based on ABM11 do not describe LHC jet data at small jet rapidity, cf. Refs. [4, 5, 26, 32].
- The HERAPDF2.0 set exclusively fits HERA DIS data with only weak constraints on the gluon PDF.
- The range in values available for $\alpha_s(M_Z)$ is too limited for the NNPDF3.0 set.

PDF uncertainties are evaluated according to the prescriptions given for each PDF set. Uncertainties on $\alpha_s(M_Z)$ are not considered, since this value is later on determined from a fit to the data. The PDF uncertainty as derived with the CT10 PDF set ranges from 2% to 30% for inclusive 2- and 3-jet cross sections and from 2% to 7% for R_{32} .

Table 2: NLO PDF sets available via LHAPDF6 for comparisons to data with various assumptions on the value of $\alpha_s(M_Z)$. Sets existing already in LHC Run 1 (upper rows) and newer sets for Run 2 (lower rows) are listed together with the corresponding number of flavours N_F , the assumed masses M_t and M_Z of the top quark and the Z boson, respectively, the default values of $\alpha_s(M_Z)$, and the range in $\alpha_s(M_Z)$ variation available for fits. A * behind the $\alpha_s(M_Z)$ values signifies that the parameter was fixed, not fitted.

Base set	Refs.	N_F	M_t (GeV)	M_Z (GeV)	$\alpha_s(M_Z)$	$\alpha_s(M_Z)$ range
ABM11	[33]	5	180	91.174	0.1180	0.110–0.130
CT10	[34]	≤ 5	172	91.188	0.1180*	0.112–0.127
MSTW2008	[35, 36]	≤ 5	10^{10}	91.1876	0.1202	0.110–0.130
NNPDF2.3	[37]	≤ 6	175	91.1876	0.1180*	0.114–0.124
CT14	[38]	≤ 5	172	91.1876	0.1180*	0.113–0.123
HERAPDF2.0	[39]	≤ 5	173	91.1876	0.1180*	0.110–0.130
MMHT2014	[40]	≤ 5	10^{10}	91.1876	0.1180*	0.108–0.128
NNPDF3.0	[41]	≤ 5	173	91.2	0.1180*	0.115–0.121

The uncertainty related to unknown higher orders of the perturbative series is evaluated with the conventional recipe of varying the default scale $H_{\text{T},2}/2$ chosen for μ_r and μ_f independently in the following six combinations: $(\mu_r/H_{\text{T},2}/2, \mu_f/H_{\text{T},2}/2) = (1/2, 1/2), (1/2, 1), (1, 1/2), (1, 2), (2, 1)$ and $(2, 2)$. The maximal upwards and downwards deviations in cross section from the central prediction are taken as scale uncertainty. This uncertainty ranges for inclusive 2-jet events from 5% to 13%, for inclusive 3-jet events from 11% to 17% and for their ratio R_{32} from 6% to 8%.

The computation of the NLO predictions with NLOJET++ is also subject to statistical fluctuations from the numerical integrations. For the inclusive 2-jet event cross sections this uncer-

tainty is smaller than about one per mille, while for the inclusive 3-jet event cross section it amounts to 1–9 per mille.

Higher order effects of electroweak origin affect jet cross sections at large jet p_T . These electroweak (EWK) corrections have been calculated for the inclusive 1-jet and 2-jet case, cf. Ref. [42], but are not yet known for 3-jet production. Therefore, they are considered for the 2-jet events, while for the 3-jet event cross section and for the ratio they have been neglected.

The impact of nonperturbative (NP) effects, i.e. from multiple-parton interactions (MPI) and hadronization, are evaluated by using samples obtained from different MC event generators with a simulation of parton-shower and underlying-event (UE) contributions. The leading order (LO) MC event generators HERWIG++ [43] with the default tune of version 2.3 and PYTHIA6 [19] with tune Z2* are considered, and the dijet NLO prediction from POWHEG [44–46] interfaced to PYTHIA8 with tune CUETS1 [47] for full event generation. The cross section ratios between a nominal event generation and a sample without hadronization and MPI effects are taken as correction separately for inclusive 2-, and 3-jet events, and as their ratio for R_{32} . This ratio is fitted by a power-law function. The differences in the correction factors obtained from the various MC event generators are assigned as an uncertainty. The central correction factors C^{NP} are determined by the centre of the envelope which covers all predictions and half of the spread is taken as the uncertainty.




Figure 3: Fits to the nonperturbative corrections obtained for inclusive 2-jet (top left) and 3-jet (top right) event cross sections and their ratio R_{32} (bottom) as a function of $H_{T,2}/2$ within $|y| < 2.5$ for the three investigated MC event generators.

The NP corrections are shown in Fig. 3 for the inclusive 2-jet (top left) and 3-jet event cross

sections (top right) as well for R_{32} (bottom). They amount to $\approx 4\text{--}5\%$ for inclusive 2-jet and 3-jet events and $\approx 1\%$ for R_{32} at $H_{T,2}/2 \approx 300\text{ GeV}$ and decrease for increasing $H_{T,2}/2$. The uncertainty assigned to the NP corrections is of the order of 1–2%. The non-perturbative effects are reduced in the cross section ratio.




Figure 4: Overview of theoretical uncertainties affecting the cross section prediction for inclusive 2-jet (top left) and 3-jet events (top right) and their ratio R_{32} (bottom), using the CT10 PDF set. The total uncertainty is calculated by adding in quadrature the individual sources of uncertainty. The statistical uncertainties of the NLO computations are too small to be visible and are not shown.

The total theoretical uncertainties are evaluated as the quadratic sum of the scale, PDF, NP, and statistical uncertainties. Figure 4 presents an overview of the theoretical uncertainties affecting the cross section prediction for inclusive 2-jet (top left) and 3-jet events (top right) and their ratio R_{32} (bottom), using the CT10 PDF set.

5 Comparison between measured cross sections and theory

Figure 5 shows the measured inclusive 2-jet and 3-jet event cross sections as a function of $H_{T,2}/2$ after unfolding for detector effects. On the left, the measurements are compared to the NLO-JET++ predictions using the CT10 PDF set, corrected for NP effects and in addition for EWK effects in the 2-jet case. On the right, the comparison is made to the predictions from MADGRAPH5 + PYTHIA6 with tune Z2* (MG+P6 Z2*), corrected for EWK effects in the 2-jet case. On a logarithmic scale, the data are in agreement with the NLO predictions over the whole




Figure 5: Comparison of the inclusive 2-jet and 3-jet event cross sections as a function of $H_{T,2}/2$ to theoretical predictions. On the (left), the data (points) are shown together with NLOJET++ predictions (line) using the CT10 PDF set, corrected for NP and EWK (2-jet) or only NP effects (3-jet). On the (right), the data (points) are compared to predictions from MADGRAPH5 + PYTHIA6 with tune Z2* (line), corrected for EWK effects in the 2-jet case. The error bars correspond to the total uncertainty, for which the statistical and systematic uncertainties are added in quadrature.

range of $H_{T,2}/2$ from 300 GeV up to 2.0 (2-jet) and 1.68 TeV (3-jet) respectively.

For better visibility the ratios of data over the NLOJET++ predictions using the CT10 PDF set are shown in Fig. 6. The data are well described by the predictions within their uncertainty, which is dominated at large $H_{T,2}/2$ by PDF effects in the upwards and by scale variations in the downwards direction. A trend towards an increasing systematic excess of the 2-jet data with respect to theory, starting at about 1 TeV in $H_{T,2}/2$, is remedied by the inclusion of EWK corrections. In the 3-jet case the statistical precision of the data and the reach in $H_{T,2}/2$ is insufficient to observe any effect. The alternative PDF sets MSTW2008 and NNPDF2.3 exhibit a small underestimation of the cross sections at high $H_{T,2}/2$.

As for the NP corrections, the POWHEG framework providing a NLO dijet calculation matched to the parton showers of PYTHIA8 is used for a comparison. Here, POWHEG + PYTHIA8 are employed with the CUETS1 and CUETM1 tunes. The ratios of data over theory from POWHEG + PYTHIA8 with tune CUETS1 are shown in Fig. 7. For comparison, the LO prediction from PYTHIA6 with tune Z2*, the tree-level multi-leg improved prediction by MADGRAPH5 + PYTHIA6 with tune Z2*, and the matched NLO prediction from POWHEG + PYTHIA8 with tune CUETM1 are shown as well. Significant discrepancies, which are cancelled to a large extent in the ratio R_{32} , are visible in the comparison with the LO prediction from MADGRAPH5 + PYTHIA6 with tune Z2*, in particular for small $H_{T,2}/2$. In contrast, the employed dijet MC PYTHIA8 and POWHEG + PYTHIA8 better describe the 2-jet event cross section, but fail for the 3-jet case.

The cross section ratio R_{32} as a function of $H_{T,2}/2$ is extracted from the data by dividing the differential cross sections for each bin in $H_{T,2}/2$. Figure 8 presents this ratio as obtained from unfolded data in comparison to that from NLO pQCD. The error bars correspond to the total experimental uncertainty.

For a better comparison of the behaviour of the 2- and 3-jet event cross sections and their ratio Figs. 9–11 present the respective ratios with respect to theory for varying assumptions on PDFs and $\alpha_s(M_Z)$. A small slope increasing with $H_{T,2}/2$ is visible for most PDFs in both cross




Figure 6: Ratio of data over theory using the CT10 PDF set for inclusive 2-jet (top left) and inclusive 3-jet event cross sections (top right) and their ratio R_{32} (bottom). For comparison predictions employing two other PDF sets are also shown. The error bars correspond to the statistical uncertainty of the data and the shaded rectangles to the total experimental systematic uncertainty. The shaded band around unity represents the total uncertainty of the theory.




Figure 7: Ratio of data over the prediction from POWHEG + PYTHIA8 with tune CUETS1. For comparison the alternative tune CUETM1 of POWHEG + PYTHIA8, the tree-level multi-leg improved prediction by MADGRAPH5 + PYTHIA6 with tune Z2*, and the LO MC predictions from PYTHIA6 tune Z2* are shown as well. The error bars correspond to the statistical uncertainty of the data and the shaded rectangles to the total experimental systematic uncertainty. EWK corrections have been accounted for in this comparison in the 2-jet case.




Figure 8: Cross section ratio R_{32} as a function of $H_{T,2}/2$ calculated from data (solid circles) in comparison to that from NLO pQCD (lines). The error bars correspond to the total experimental uncertainty derived as quadratic sum from all uncertainty sources. The NLO predictions using the CT10 NLO PDF set corrected with NP corrections are shown for a series of values assumed for $\alpha_s(M_Z)$ (dashed lines) together with the central prediction (solid line) where $\alpha_s(M_Z) = 0.118$. The assumption on $\alpha_s(M_Z)$ is varied in steps of 0.001 in the range of 0.112–0.127. For brevity, the relative factor of NP between data and theory has been indicated as “Data/NP” in the legend.

sections. This effect is largely cancelled in the cross section ratio. R_{32} exhibits a flat behaviour with respect to the predictions for all five PDF sets in the whole range of $H_{T,2}/2$ up to 1.68 TeV.

Moreover, the different sensitivity to $\alpha_s(M_Z)$ caused by the leading power in α_S in the expansion of the 2-jet inclusive ($\propto \alpha_S^2$) and the 3-jet inclusive cross section ($\propto \alpha_S^3$), and their ratio ($\propto \alpha_S^1$) is clearly visible from the spread between the calculations for the smallest and largest value of $\alpha_s(M_Z)$ within the same PDF set when passing through Figs. 9–11. This also demonstrates the potential of ratios R_{mn} with $m - n > 1$.

6 Fits of the strong coupling constant

As discussed in the previous section, the measured inclusive 2-jet and 3-jet event cross sections and their ratio R_{32} can be used for a determination of the strong coupling constant $\alpha_s(M_Z)$. The value of $\alpha_s(M_Z)$ is determined by minimizing the χ^2 between the experimental measurement and the theoretical predictions. The fit procedure here follows closely the one previously used in Refs. [12] and [48]. The χ^2 is defined as:

$$\chi^2 = M^T C^{-1} M, \quad (2)$$

where M is the vector of the differences between the data (D^i) and the theoretical values (T^i) in each bin i ,

$$M^i = D^i - T^i \quad (3)$$

and C is the covariance matrix including all experimental uncertainties as described in Section 3 and some theoretical uncertainties. More precisely, $C = C_{\text{exp}} + C_{\text{theo}}$ is defined as the sum of




Figure 9: Ratio of measured 2-jet inclusive event cross section (data points) over NLO theory times NP corrections for various PDF sets at their respective default value for $\alpha_s(M_Z)$ (black solid line at unity). The error bars correspond to the total experimental uncertainty. The NLO predictions have been derived with the CT10 (top left), the CT14 (top right), the MSTW2008 (middle left), the MMHT2014 (middle right) and the NNPDF2.3 PDF set (bottom) for the series of assumptions on $\alpha_s(M_Z)$ available for the respective PDF set as specified in Table 2. For brevity, the relative factor of NP between data and theory has been indicated as “Data/NP” in the legend.




Figure 10: Ratio of measured 3-jet inclusive event cross section (data points) over NLO theory times NP corrections for various PDF sets at their respective default value for $\alpha_s(M_Z)$ (black solid line at unity). The error bars correspond to the total experimental uncertainty. The NLO predictions have been derived with the CT10 (top left), the CT14 (top right), the MSTW2008 (middle left), the MMHT2014 (middle right) and the NNPDF2.3 PDF set (bottom) for the series of assumptions on $\alpha_s(M_Z)$ available for the respective PDF set as specified in Table 2. For brevity, the relative factor of NP between data and theory has been indicated as “Data/NP” in the legend.




Figure 11: Ratio of measured R_{32} ratio (data points) over NLO theory times NP corrections for various PDF sets at their respective default value for $\alpha_s(M_Z)$ (black solid line at unity). The error bars correspond to the total experimental uncertainty. The NLO predictions have been derived with the CT10 (top left), the CT14 (top right), the MSTW2008 (middle left), the MMHT2014 (middle right) and the NNPDF2.3 PDF sets (bottom) for the series of assumptions on $\alpha_s(M_Z)$ available for the respective PDF set as specified in Table 2. For brevity, the relative factor of NP between data and theory has been indicated as “Data/NP” in the legend.

covariances of experimental and theoretical sources of uncertainty as follows

$$C_{\text{exp}} = \text{Cov}^{\text{ExpStat}} + \sum \text{Cov}^{\text{JEC}} + \text{Cov}^{\text{Unfolding}} + \text{Cov}^{\text{Lumi}} + \text{Cov}^{\text{Uncor}}, \quad (4)$$

$$C_{\text{theo}} = \text{Cov}^{\text{TheoStat}} + \text{Cov}^{\text{NP}} + \text{Cov}^{\text{PDF}}, \quad (5)$$

where the labelled covariance matrices account for the following effects:

- $\text{Cov}^{\text{ExpStat}}$: the statistical uncertainty of the data including correlations introduced by the unfolding,
- Cov^{JEC} : the JEC systematic uncertainty,
- $\text{Cov}^{\text{Unfolding}}$: the unfolding systematic uncertainty including the JER,
- Cov^{Lumi} : the luminosity uncertainty,
- $\text{Cov}^{\text{Uncor}}$: a residual uncorrelated systematic uncertainty summarizing individual causes such as trigger and identification inefficiencies, time dependence of the jet p_T resolution, and uncertainty on the trigger prescale factors,
- $\text{Cov}^{\text{TheoStat}}$: the statistical uncertainty caused by numerical integrations in the cross section computations,
- Cov^{NP} : the systematic uncertainty of the NP corrections, and
- Cov^{PDF} : the PDF uncertainty.

In fits of the ratio R_{32} , the luminosity and residual uncorrelated uncertainties cancel completely. Partial cancellations between the other sources of uncertainty are taken into account in the fit. The JEC, unfolding, and luminosity uncertainties are treated as multiplicative to avoid the statistical bias that arises when estimating uncertainties from data.

The derivation of PDF uncertainties depends on each PDF set. The CT10 PDF set consists of $N_{\text{ev}} = 26$ eigenvectors with two PDF members per eigenvector k , which lead to the predictions S_k^\pm that follow from PDF variations with respect to the plus and minus directions of eigenvector k . Symmetric uncertainties as required by the use of covariance matrices are then computed by [49]:

$$(\Delta X)^2 = \frac{1}{4} \sum_{k=1}^{N_{\text{ev}}} [X(S_k^+) - X(S_k^-)]^2, \quad (6)$$

where ΔX is the uncertainty of the cross section and $X(S_k^\pm)$ is the predicted cross section for each eigenvector orientation, + or -.

Scale uncertainties of the pQCD predictions are taken into account employing the offset method, i.e. by performing separate fits with varying scale factors as described in the previous section. The largest upwards and downwards deviations from the default factors are defined as the uncertainty. At NLO such scale variations predominantly lead to smaller cross sections and also a smaller ratio R_{32} as visible in Fig. 4. As a consequence the scale uncertainty in fits is equally asymmetric, where smaller cross sections or ratios are compensated by an increase in the fitted value for $\alpha_s(M_Z)$.

First, fits to the cross sections are performed, where the range in $H_{\text{T},2}/2$ is restricted to be between 300 GeV and 1 TeV to avoid the region close to the minimal p_T threshold of 150 GeV for each jet at low p_T and the onset of electroweak effects at high p_T , which are available for

the dijet case only. The results are reported in Table 3 for the 2-jet and 3-jet event cross sections. For comparison, a simultaneous fit to both cross sections ignoring any correlations, and a fit to their ratio fully accounting for correlations are given in Table 4. Also, EWK effects are assumed to cancel in the ratio as do the luminosity and the uncorrelated uncertainty.

All cross section fits give compatible values for $\alpha_s(M_Z)$ in the range of 0.115–0.118; for the ratio R_{32} somewhat smaller values are obtained. A common issue, except for the ratio fits, is the rather small χ^2/n_{dof} . A possible explanation is an overestimation of the residual uncorrelated uncertainty of 1% that is cancelled for R_{32} . If the fits are repeated with an assumed uncertainty of 0.25% instead, the χ^2/n_{dof} values lie around unity while the $\alpha_s(M_Z)$ values are still compatible with the previous results but with slightly reduced uncertainties.

Table 3: Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross sections using five PDF sets at NLO. Only total uncertainties without scale variations are quoted. The results are obtained from a simultaneous fit to all 19 $H_{\text{T},2}/2$ bins in the restricted range of $0.3 < H_{\text{T},2}/2 < 1.0$ TeV.

PDF set	2-jets			3-jets		
	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}
CT10	0.1174	0.0032	3.0/18	0.1169	0.0027	5.4/18
CT14	0.1160	0.0035	3.5/18	0.1159	0.0031	6.1/18
MSTW2008	0.1159	0.0025	5.3/18	0.1161	0.0021	6.7/18
MMHT2014	0.1165	0.0034	5.9/18	0.1166	0.0025	7.1/18
NNPDF2.3	0.1183	0.0025	9.7/18	0.1179	0.0021	9.1/18

Table 4: Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet and 3-jet event cross sections simultaneously and from their ratio R_{32} using five PDF sets at NLO. Only total uncertainties without scale variations are quoted. The results are obtained from a simultaneous fit to all 38 (19) $H_{\text{T},2}/2$ bins in the restricted range of $0.3 < H_{\text{T},2}/2 < 1.0$ TeV. For comparison, correlations between the two cross sections are neglected in the simultaneous fit on the left, but fully taken into account in the ratio fit on the right.

PDF set	2- & 3-jets			R_{32}		
	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	$\pm \Delta \alpha_s(M_Z)$	χ^2/n_{dof}
CT10	0.1170	0.0026	8.2/37	0.1141	0.0028	19./18
CT14	0.1161	0.0029	9.1/37	0.1139	0.0032	15./18
MSTW2008	0.1161	0.0021	11./37	0.1150	0.0023	21./18
MMHT2014	0.1168	0.0025	11./37	0.1142	0.0022	19./18
NNPDF2.3	0.1188	0.0019	15./37	0.1184	0.0021	12./18

To investigate how the EWK corrections affect the fit results for $\alpha_s(M_Z)$, the range in $H_{\text{T},2}/2$ is extended to $0.3 < H_{\text{T},2}/2 < 1.68$ TeV. Table 5 reports the values obtained for $\alpha_s(M_Z)$ from fits to the 2-jet event cross section in this range with or without EWK correction factors. The largest impact is a reduction in χ^2/n_{dof} , which indicates a better agreement when EWK effects are included. In addition, a tendency to slightly smaller $\alpha_s(M_Z)$ values is observed without the EWK corrections. For the ratio R_{32} it is expected that these effects are much reduced.

From Fig. 11 follows that only the PDF sets MSTW2008 and MMHT2014 provide a large enough range in $\alpha_s(M_Z)$ values to ensure fits without extrapolation. The other three PDF sets are at the limit such that reliable fits cannot be performed for all scale settings and/or bins in scale

Table 5: Determination of $\alpha_s(M_Z)$ from the inclusive 2-jet event cross section using five PDF sets at NLO with (right) and without (left) EWK corrections. Only total uncertainties without scale variations are quoted. The results are obtained from a simultaneous fit to all 29 $H_{T,2}/2$ bins in the range of $0.3 < H_{T,2}/2 < 1.68$ TeV.

PDF set	2-jets, without EWK			2-jets, with EWK		
	$\alpha_s(M_Z)$	$\pm \Delta\alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	$\pm \Delta\alpha_s(M_Z)$	χ^2/n_{dof}
CT10	0.1163	0.0034	15./28	0.1165	0.0032	14./28
CT14	0.1137	0.0033	24./28	0.1144	0.0033	17./28
MSTW2008	0.1093	0.0028	27./28	0.1133	0.0023	19./28
MMHT2014	0.1127	0.0032	32./28	0.1141	0.0032	21./28
NNPDF2.3	0.1162	0.0024	31./28	0.1168	0.0024	23./28

$Q = H_{T,2}/2$. Tables 6–8 give the complete results for MSTW2008 and MMHT2014 for the full range in $H_{T,2}/2$ of 300 GeV up to 1.68 TeV, for scale variations in this range, and for subranges in $H_{T,2}/2$.

Using the MSTW2008 PDF set, which dates from before the LHC start, the strong coupling constant finally is determined to

$$\begin{aligned}\alpha_s(M_Z) &= 0.1150 \pm 0.0010 (\text{exp}) \pm 0.0013 (\text{PDF}) \pm 0.0015 (\text{NP})^{+0.0050}_{-0.0000} (\text{scale}) \\ &= 0.1150 \pm 0.0023 (\text{all except scale})^{+0.0050}_{-0.0000} (\text{scale}).\end{aligned}$$

The MMHT2014 PDF set, although using LHC jet data to determine the PDF parameters, leads to a very similar result of

$$\begin{aligned}\alpha_s(M_Z) &= 0.1142 \pm 0.0010 (\text{exp}) \pm 0.0013 (\text{PDF}) \pm 0.0014 (\text{NP})^{+0.0049}_{-0.0006} (\text{scale}) \\ &= 0.1142 \pm 0.0022 (\text{all except scale})^{+0.0049}_{-0.0006} (\text{scale}).\end{aligned}$$

In contrast to fits at NLO using cross sections, where the scale uncertainty recipe usually leads to a very asymmetric behaviour with the larger uncertainty towards smaller values of $\alpha_s(M_Z)$, this is inverted for the fits to the cross section ratio.

Table 9 provides in addition to the extracted $\alpha_s(M_Z)$ value for each range in $H_{T,2}/2$ the $\alpha_s(Q)$ values with total uncertainty as evolved to the respective cross-section averaged scale $\langle Q \rangle$ in that range. The evolution is performed for five flavours at 2-loop order with the RUNDEC program [50, 51]. The obtained $\alpha_s(Q)$ points are illustrated in Fig. 12 together with the world average [52] and results from other measurements of the CMS [11, 12, 26, 48, 53], ATLAS [54], D0 [55, 56], H1 [57, 58], and ZEUS [59] experiments.

7 Summary

A measurement of the inclusive 2-jet (3-jet) event cross sections has been presented in a range of $0.3 < H_{T,2}/2 < 2.0$ TeV ($0.3 < H_{T,2}/2 < 1.68$ TeV) for the average p_T of the two leading jets at central rapidity of $|y| < 2.5$. The data sample has been collected from proton-proton collisions at 8 TeV centre-of-mass energy and corresponds to an integrated luminosity of 19.7 fb^{-1} . The data are found to be well described by calculations at NLO in pQCD complemented with NP corrections that are important at low $H_{T,2}/2$. The upwards trend seen in the 2- and 3-jet data at high $H_{T,2}/2$ in comparison to the prediction at NLO QCD, is explained by the onset of EWK

Table 6: Determination of $\alpha_s(M_Z)$ from the ratio R_{32} using the two most compatible PDF sets MSTW2008 and MMHT2014 at NLO. The results are obtained from a simultaneous fit to all 29 $H_{T,2}/2$ bins in the full range of $0.3 < H_{T,2}/2 < 1.68$ TeV.

PDF set	$R_{32}: \Delta\alpha_s(M_Z) \times 1000$						
	$\alpha_s(M_Z)$	exp	PDF	NP	all exc.	scale	χ^2/n_{dof}
MSTW2008	0.1150	± 10	± 13	± 15	± 23	$^{+50}_{-0}$	26./28
MMHT2014	0.1142	± 10	± 13	± 14	± 22	$^{+49}_{-6}$	24./28

Table 7: Fitted values of $\alpha_s(M_Z)$ using R_{32} in the $H_{T,2}/2$ range from 0.3 up to 1.68 TeV at the central scale and for the six scale factor combinations for the two PDF sets MSTW2008 and MMHT2014.

$\mu_r/H_{T,2}/2$	$\mu_f/H_{T,2}/2$	MSTW2008		MMHT2014	
		$\alpha_s(M_Z)$	χ^2/n_{dof}	$\alpha_s(M_Z)$	χ^2/n_{dof}
1	1	0.1150	26./28	0.1142	24./28
1/2	1/2	0.1165	77./28	0.1160	73./28
2	2	0.1200	18./28	0.1191	18./28
1/2	1	0.1150	53./28	0.1136	48./28
1	1/2	0.1150	30./28	0.1142	28./28
1	2	0.1155	23./28	0.1147	22./28
2	1	0.1180	19./28	0.1175	19./28

Table 8: Uncertainty composition for $\alpha_s(M_Z)$ from the determination of α_S from the jet event rate R_{32} in bins of $H_{T,2}/2$. The statistical uncertainty of the NLO computation is negligible in comparison to any of the other sources of uncertainty. Electroweak corrections, significant only at high $H_{T,2}/2$, are assumed to cancel between the numerator and denominator.

$H_{T,2}/2$ (GeV)	MSTW2008: $\Delta\alpha_s(M_Z) \times 1000$					MMHT2014: $\Delta\alpha_s(M_Z) \times 1000$				
	$\alpha_s(M_Z)$	exp	PDF	NP	scale	$\alpha_s(M_Z)$	exp	PDF	NP	scale
300–420	0.1157	± 15	± 14	± 19	$^{+53}_{-0}$	0.1158	± 14	± 10	± 19	$^{+52}_{-0}$
420–600	0.1153	± 11	± 14	± 18	$^{+57}_{-0}$	0.1154	± 11	± 12	± 17	$^{+56}_{-0}$
600–1000	0.1134	± 13	± 16	± 19	$^{+52}_{-0}$	0.1140	± 12	± 12	± 18	$^{+45}_{-0}$
1000–1680	0.1147	± 29	± 17	± 18	$^{+63}_{-11}$	0.1154	± 25	± 14	± 15	$^{+56}_{-11}$
300–1680	0.1150	± 10	± 13	± 15	$^{+50}_{-0}$	0.1142	± 10	± 13	± 14	$^{+49}_{-6}$

Table 9: Evolution of the strong coupling constant between the scale of the Z boson mass and the cross-section averaged $H_{T,2}/2$ scale $\langle Q \rangle$ for the separate determinations in each respective fit range. The evolution is performed for five flavours at 2-loop order with the RUNDEC program [50, 51].

$H_{T,2}/2$ (GeV)	$\langle Q \rangle$ (GeV)	$\alpha_s(M_Z)$	$\alpha_s(Q)$	No. of data points	χ^2/n_{dof}
300–420	340	$0.1157^{+0.0060}_{-0.0030}$	$0.0969^{+0.0041}_{-0.0021}$	4	2.8/3
420–600	476	$0.1153^{+0.0062}_{-0.0025}$	$0.0928^{+0.0039}_{-0.0016}$	6	6.1/5
600–1000	685	$0.1134^{+0.0059}_{-0.0028}$	$0.0879^{+0.0035}_{-0.0017}$	9	7.1/8
1000–1680	1114	$0.1147^{+0.0074}_{-0.0040}$	$0.0841^{+0.0039}_{-0.0021}$	10	5.4/9




Figure 12: The running $\alpha_s(Q)$ as a function of the scale Q is shown as obtained by using the MSTW2008 NLO PDF set. The solid line and the uncertainty band are drawn by evolving the extracted $\alpha_s(M_Z)$ values using the 2-loop 5-flavour renormalization group equations as implemented in RUNDEC [50, 51]. The dashed line represents the evolution of the world average [52] and the black circles correspond to the $\alpha_s(Q)$ determinations presented in Table 9. Results from other measurements of CMS [11, 12, 26, 48, 53], ATLAS [54], D0 [55, 56], H1 [57, 58], and ZEUS [59] are superimposed.

corrections in the 2-jet case. For the 3-jet event cross section these correction have not yet been computed.

In the 3-jet to 2-jet cross section ratio the EWK corrections are assumed to cancel. In fact, NLO QCD provides an adequate description of R_{32} in the accessible range of $H_{T,2}/2$. In contrast, LO tree-level MC predictions exhibit significant deviations.

Based on the observed agreement, the strong coupling constant is determined in a fit to the R_{32} measurement to

$$\begin{aligned}\alpha_s(M_Z) &= 0.1150 \pm 0.0010 (\text{exp}) \pm 0.0013 (\text{PDF}) \pm 0.0015 (\text{NP}) {}^{+0.0050}_{-0.0000} (\text{scale}) \\ &= 0.1150 \pm 0.0023 (\text{all except scale}) {}^{+0.0050}_{-0.0000} (\text{scale}).\end{aligned}$$

using the MSTW2008 PDF set. Employing the MMHT2014 PDF set instead leads to very similar results. Equally compatible determinations of $\alpha_s(M_Z)$ are achieved with separate fits to the inclusive 2-jet and 3-jet event cross sections employing various PDF sets provided the range in $H_{T,2}/2$ is restricted to $0.3 < H_{T,2}/2 < 1.0$ TeV. The result for $\alpha_s(M_Z)$ is in agreement with previous determinations obtained by the ATLAS and CMS collaborations [11, 12, 26, 48, 53, 54] and with the world average value of $\alpha_s(M_Z) = 0.1181 \pm 0.0011$ derived in Ref. [52].

References

- [1] ALICE Collaboration, "Measurement of the inclusive differential jet cross section in pp collisions at $\sqrt{s} = 2.76$ TeV", *Phys. Lett. B* **722** (2013) 262, doi:10.1016/j.physletb.2013.04.026, arXiv:1301.3475.
- [2] ATLAS Collaboration, "Measurement of inclusive jet and dijet cross sections in proton-proton collisions at 7 TeV centre-of-mass energy with the ATLAS detector", *Eur. Phys. J. C* **71** (2011) 1512, doi:10.1140/epjc/s10052-010-1512-2, arXiv:1009.5908.
- [3] ATLAS Collaboration, "Measurement of inclusive jet and dijet production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector", *Phys. Rev. D* **86** (2012) 014022, doi:10.1103/PhysRevD.86.014022, arXiv:1112.6297.
- [4] ATLAS Collaboration, "Measurement of the inclusive jet cross section in pp collisions at $\sqrt{s} = 2.76$ TeV and comparison to the inclusive jet cross section at $\sqrt{s} = 7$ TeV using the ATLAS detector", *Eur. Phys. J. C* **73** (2013) 2509, doi:10.1140/epjc/s10052-013-2509-4, arXiv:1304.4739.
- [5] ATLAS Collaboration, "Measurement of the inclusive jet cross-section in proton-proton collisions at $\sqrt{s} = 7$ TeV using 4.5 fb^{-1} of data with the ATLAS detector", *JHEP* **02** (2015) 153, doi:10.1007/JHEP02(2015)153, arXiv:1410.8857.
- [6] CMS Collaboration, "Measurement of the Inclusive Jet Cross Section in pp Collisions at $\sqrt{s} = 7$ TeV", *Phys. Rev. Lett.* **107** (2011) 132001, doi:10.1103/PhysRevLett.107.132001, arXiv:1106.0208.
- [7] CMS Collaboration, "Measurement of the inclusive production cross sections for forward jets and for dijet events with one forward and one central jet in pp collisions at $\sqrt{s} = 7$ TeV", *JHEP* **06** (2012) 036, doi:10.1007/JHEP06(2012)036, arXiv:1202.0704.
- [8] CMS Collaboration, "Measurements of differential jet cross sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the CMS detector", *Phys. Rev. D* **87** (2013) 112002, doi:10.1103/PhysRevD.87.112002, arXiv:1212.6660.
- [9] CMS Collaboration, "Measurement of the ratio of inclusive jet cross sections using the anti- k_T algorithm with radius parameters $R = 0.5$ and 0.7 in pp collisions at $\sqrt{s} = 7$ TeV", *Phys. Rev. D* **90** (2014) 072006, doi:10.1103/PhysRevD.90.072006, arXiv:1406.0324.
- [10] CMS Collaboration, "Measurement of the double-differential inclusive jet cross section in proton-proton collisions at $\sqrt{s} = 13$ TeV", *Eur. Phys. J. C* **76** (2016), no. 8, 451, doi:10.1140/epjc/s10052-016-4286-3, arXiv:1605.04436.
- [11] CMS Collaboration, "Measurement and QCD analysis of double-differential inclusive jet cross-sections in pp collisions at $\sqrt{s} = 8$ TeV and ratios to 2.76 and 7 TeV", (2016). arXiv:1609.05331. Submitted to *JHEP*.
- [12] CMS Collaboration, "Measurement of the ratio of the inclusive 3-jet cross section to the inclusive 2-jet cross section in pp collisions at $\sqrt{s} = 7$ TeV and first determination of the strong coupling constant in the TeV range", *Eur. Phys. J. C* **73** (2013) 2604, doi:10.1140/epjc/s10052-013-2604-6, arXiv:1304.7498.

- [13] M. Cacciari, G. P. Salam, and G. Soyez, “The anti- k_t jet clustering algorithm”, *JHEP* **04** (2008) 063, doi:[10.1088/1126-6708/2008/04/063](https://doi.org/10.1088/1126-6708/2008/04/063), arXiv:[0802.1189](https://arxiv.org/abs/0802.1189).
- [14] CMS Trigger and Data Acquisition Group Collaboration, “The CMS high level trigger”, *Eur. Phys. J. C* **46** (2006) 605, doi:[10.1140/epjc/s2006-02495-8](https://doi.org/10.1140/epjc/s2006-02495-8), arXiv:[hep-ex/0512077](https://arxiv.org/abs/hep-ex/0512077).
- [15] CMS Collaboration, “Particle-Flow Event Reconstruction in CMS and Performance for Jets, Taus, and MET”, technical report, CERN, 2009.
- [16] CMS Collaboration, “Particle-flow commissioning with muons and electrons from J/Psi and W events at 7 TeV”, technical report, CERN, 2010.
- [17] M. Cacciari, G. P. Salam, and G. Soyez, “FastJet User Manual”, *Eur. Phys. J. C* **72** (2012) 1896, doi:[10.1140/epjc/s10052-012-1896-2](https://doi.org/10.1140/epjc/s10052-012-1896-2), arXiv:[1111.6097](https://arxiv.org/abs/1111.6097).
- [18] CMS Collaboration, “Jet energy scale and resolution in the CMS experiment in pp collisions at 8 TeV”, (2016). arXiv:[1607.03663](https://arxiv.org/abs/1607.03663). Submitted to *JINST*.
- [19] T. Sjöstrand, S. Mrenna, and P. Z. Skands, “PYTHIA 6.4 Physics and Manual”, *JHEP* **05** (2006) 026, doi:[10.1088/1126-6708/2006/05/026](https://doi.org/10.1088/1126-6708/2006/05/026), arXiv:[hep-ph/0603175](https://arxiv.org/abs/hep-ph/0603175).
- [20] CMS Collaboration, “Charged particle multiplicities in pp interactions at $\sqrt{s} = 0.9, 2.36,$ and 7 TeV”, *JHEP* **01** (2011) 079, doi:[10.1007/JHEP01\(2011\)079](https://doi.org/10.1007/JHEP01(2011)079), arXiv:[1011.5531](https://arxiv.org/abs/1011.5531).
- [21] S. Agostinelli et al., “GEANT4: A Simulation toolkit”, *Nuclear Instruments & Methods in Physics Research A* **506** (2003) 250, doi:[10.1016/S0168-9002\(03\)01368-8](https://doi.org/10.1016/S0168-9002(03)01368-8).
- [22] CMS Collaboration, “Description and performance of track and primary-vertex reconstruction with the CMS tracker”, *JINST* **9** (2014), no. 10, P10009, doi:[10.1088/1748-0221/9/10/P10009](https://doi.org/10.1088/1748-0221/9/10/P10009), arXiv:[1405.6569](https://arxiv.org/abs/1405.6569).
- [23] CMS Collaboration, “Jet Performance in pp Collisions at $\sqrt{s} = 7 \text{ TeV}$ ”, technical report, CERN, 2010.
- [24] G. D’Agostini, “A Multidimensional unfolding method based on Bayes’ theorem”, *Nucl. Instrum. Meth. A* **362** (1995) 487, doi:[10.1016/0168-9002\(95\)00274-X](https://doi.org/10.1016/0168-9002(95)00274-X).
- [25] T. Adye, “Unfolding algorithms and tests using RooUnfold”, in *Proceedings, PHYSTAT 2011 Workshop on Statistical Issues Related to Discovery Claims in Search Experiments and Unfolding*, p. 313. Geneva, Switzerland, January 17-20, 2011. arXiv:[1105.1160](https://arxiv.org/abs/1105.1160). doi:[10.5170/CERN-2011-006](https://doi.org/10.5170/CERN-2011-006).
- [26] CMS Collaboration, “Measurement of the inclusive 3-jet production differential cross section in proton-proton collisions at 7 TeV and determination of the strong coupling constant in the TeV range”, *Eur. Phys. J. C* **75** (2015) 186, doi:[10.1140/epjc/s10052-015-3376-y](https://doi.org/10.1140/epjc/s10052-015-3376-y), arXiv:[1412.1633](https://arxiv.org/abs/1412.1633).
- [27] CMS Collaboration, “CMS Luminosity Based on Pixel Cluster Counting - Summer 2013 Update”, technical report, CERN, 2013.
- [28] Z. Nagy, “Three jet cross-sections in hadron hadron collisions at next-to-leading order”, *Phys. Rev. Lett.* **88** (2002) 122003, doi:[10.1103/PhysRevLett.88.122003](https://doi.org/10.1103/PhysRevLett.88.122003), arXiv:[hep-ph/0110315](https://arxiv.org/abs/hep-ph/0110315).

- [29] Z. Nagy, “Next-to-leading order calculation of three-jet observables in hadron hadron collisions”, *Phys. Rev. D* **68** (2003) 094002, doi:10.1103/PhysRevD.68.094002, arXiv:hep-ph/0307268.
- [30] D. Britzger, K. Rabbertz, F. Stober, and M. Wobisch, “New features in version 2 of the fastNLO project”, in *Proceedings, XX. International Workshop on Deep-Inelastic Scattering and Related Subjects (DIS 2012)*, p. 217. Bonn, Germany, March 26-30, 2012. arXiv:1208.3641. doi:10.3204/DESY-PROC-2012-02/165.
- [31] A. Buckley et al., “LHAPDF6: parton density access in the LHC precision era”, *Eur. Phys. J.* **C75** (2015) 132, doi:10.1140/epjc/s10052-015-3318-8, arXiv:1412.7420.
- [32] CMS Collaboration, “Measurement of the inclusive jet cross section in pp collisions at $\sqrt{s} = 2.76 \text{ TeV}$ ”, *Eur. Phys. J. C* **76** (2016) 265, doi:10.1140/epjc/s10052-016-4083-z, arXiv:1512.06212.
- [33] S. Alekhin, J. Blümlein, and S. Moch, “Parton Distribution Functions and Benchmark Cross Sections at NNLO”, *Phys. Rev. D* **86** (2012) 054009, doi:10.1103/PhysRevD.86.054009, arXiv:1202.2281.
- [34] H.-L. Lai et al., “New parton distributions for collider physics”, *Phys. Rev. D* **82** (2010) 074024, doi:10.1103/PhysRevD.82.074024, arXiv:1007.2241.
- [35] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, “Parton distributions for the LHC”, *Eur. Phys. J. C* **63** (2009) 189, doi:10.1140/epjc/s10052-009-1072-5, arXiv:0901.0002.
- [36] A. D. Martin, W. J. Stirling, R. S. Thorne, and G. Watt, “Uncertainties on α_S in global PDF analyses and implications for predicted hadronic cross sections”, *Eur. Phys. J. C* **64** (2009) 653, doi:10.1140/epjc/s10052-009-1164-2, arXiv:0905.3531.
- [37] R. D. Ball et al., “Parton distributions with LHC data”, *Nucl. Phys. B* **867** (2013) 244, doi:10.1016/j.nuclphysb.2012.10.003, arXiv:1207.1303.
- [38] S. Dulat et al., “New parton distribution functions from a global analysis of quantum chromodynamics”, *Phys. Rev. D* **93** (2016) 033006, doi:10.1103/PhysRevD.93.033006, arXiv:1506.07443.
- [39] H1, ZEUS Collaboration, “Combination of measurements of inclusive deep inelastic $e^\pm p$ scattering cross sections and QCD analysis of HERA data”, *Eur. Phys. J. C* **75** (2015) 580, doi:10.1140/epjc/s10052-015-3710-4, arXiv:1506.06042.
- [40] L. Harland-Lang, A. Martin, P. Motylinski, and R. Thorne, “Parton distributions in the LHC era: MMHT 2014 PDFs”, *Eur. Phys. J. C* **75** (2015), no. 5, 204, doi:10.1140/epjc/s10052-015-3397-6, arXiv:1412.3989.
- [41] NNPDF Collaboration, “Parton distributions for the LHC Run II”, *JHEP* **04** (2015) 040, doi:10.1007/JHEP04(2015)040, arXiv:1410.8849.
- [42] S. Dittmaier, A. Huss, and C. Speckner, “Weak radiative corrections to dijet production at hadron colliders”, *JHEP* **11** (2012) 095, doi:10.1007/JHEP11(2012)095, arXiv:1210.0438.
- [43] M. Bähr et al., “Herwig++ Physics and Manual”, *Eur. Phys. J. C* **58** (2008) 639, doi:10.1140/epjc/s10052-008-0798-9, arXiv:0803.0883.

- [44] P. Nason, “A New method for combining NLO QCD with shower Monte Carlo algorithms”, *JHEP* **11** (2004) 040, doi:10.1088/1126-6708/2004/11/040, arXiv:hep-ph/0409146.
- [45] S. Frixione, P. Nason, and C. Oleari, “Matching NLO QCD computations with Parton Shower simulations: the POWHEG method”, *JHEP* **11** (2007) 070, doi:10.1088/1126-6708/2007/11/070, arXiv:0709.2092.
- [46] S. Alioli et al., “Jet pair production in POWHEG”, *JHEP* **04** (2011) 081, doi:10.1007/JHEP04(2011)081, arXiv:1012.3380.
- [47] CMS Collaboration, “Event generator tunes obtained from underlying event and multiparton scattering measurements”, *Eur. Phys. J. C* **76** (2015) 155, doi:10.1140/epjc/s10052-016-3988-x, arXiv:1512.00815.
- [48] CMS Collaboration, “Constraints on parton distribution functions and extraction of the strong coupling constant from the inclusive jet cross section in pp collisions at $\sqrt{s} = 7$ TeV”, *Eur. Phys. J. C* **75** (2015) 288, doi:10.1140/epjc/s10052-015-3499-1, arXiv:1410.6765.
- [49] J. Pumplin et al., “New generation of parton distributions with uncertainties from global QCD analysis”, *JHEP* **07** (2002) 012, doi:10.1088/1126-6708/2002/07/012, arXiv:hep-ph/0201195.
- [50] K. G. Chetyrkin, J. H. Kuhn, and M. Steinhauser, “RunDec: A Mathematica package for running and decoupling of the strong coupling and quark masses”, *Comput. Phys. Commun.* **133** (2000) 43, doi:10.1016/S0010-4655(00)00155-7, arXiv:hep-ph/0004189.
- [51] B. Schmidt and M. Steinhauser, “CRunDec: a C++ package for running and decoupling of the strong coupling and quark masses”, *Comput. Phys. Commun.* **183** (2012) 1845, doi:10.1016/j.cpc.2012.03.023, arXiv:1201.6149.
- [52] C. Patrignani and others (Particle Data Group), “Review of Particle Physics”, *Chin. Phys. C* **40** (2016) 100001, doi:10.1088/1674-1137/40/10/100001.
- [53] CMS Collaboration, “Determination of the top-quark pole mass and strong coupling constant from the $t\bar{t}$ production cross section in pp collisions at $\sqrt{s} = 7$ TeV”, *Phys. Lett. B* **728** (2014) 496, doi:10.1016/j.physletb.2013.12.009, arXiv:1307.1907.
- [54] ATLAS Collaboration, “Measurement of transverse energy-energy correlations in multi-jet events in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector and determination of the strong coupling constant $\alpha_s(m_Z)$ ”, *Phys. Lett. B* **750** (2015) 427, doi:10.1016/j.physletb.2015.09.050, arXiv:1508.01579.
- [55] D0 Collaboration, “Determination of the strong coupling constant from the inclusive jet cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev. D* **80** (2009) 111107, doi:10.1103/PhysRevD.80.111107, arXiv:0911.2710.
- [56] D0 Collaboration, “Measurement of angular correlations of jets at $\sqrt{s} = 1.96$ TeV and determination of the strong coupling at high momentum transfers”, *Phys. Lett. B* **718** (2012) 56, doi:10.1016/j.physletb.2012.10.003, arXiv:1207.4957.

- [57] H1 Collaboration, "Measurement of multijet production in ep collisions at high Q^2 and determination of the strong coupling α_s ", *Eur. Phys. J. C* **75** (2015) 65, doi:10.1140/epjc/s10052-014-3223-6, arXiv:1406.4709.
- [58] H1 Collaboration, "Measurement of Jet Production Cross Sections in Deep-inelastic ep Scattering at HERA", (2016). arXiv:1611.03421. Submitted to *Eur. Phys. J. C*.
- [59] ZEUS Collaboration, "Inclusive-jet photoproduction at HERA and determination of α_s ", *Nucl. Phys. B* **864** (2012) 1, doi:10.1016/j.nuclphysb.2012.06.006, arXiv:1205.6153.