

# Sohail

*by* Chairperson PublicAdmn

---

**Submission date:** 10-Dec-2017 10:57 AM (UTC+0530)

**Submission ID:** 846277614

**File name:** Chapter\_Abstract\_5\_ver1.pdf (89.68K)

**Word count:** 343

**Character count:** 1797

# Abstract

The collisions of hadrons at very high centre-of-mass energies provide a direct probe to the nature of the underlying parton-parton scattering physics. The scattering of the elementary quark and gluon constituents of the incoming hadron beams produces a spray of particles which are clustered in the form of jets. Jet production at hadron colliders provides an excellent opportunity for testing the theory of strong interactions called Quantum Chromodynamics. The inclusive multijet production cross-section is an important observable which provides the details of parton distribution functions (PDF) of the colliding hadrons and the precise measurement of the strong coupling constant  $\alpha_s$ . Instead of individual cross-sections, the cross-section ratio is a better tool to determine the value of  $\alpha_s$  as many theoretical and experimental uncertainties cancel in the ratio.

A measurement of inclusive multijet event cross-sections and the cross-section ratio is presented using data from proton-proton collisions collected at a centre-of-mass energy of 8 TeV with the CMS detector corresponding to an integrated luminosity of  $19.7 \text{ fb}^{-1}$ . Jets are reconstructed with the anti- $k_t$  clustering algorithm for a jet size parameter  $R = 0.7$ . The inclusive 2-jet and 3-jet event cross-sections as well as the ratio of the 3-jet over 2-jet event cross-section  $R_{32}$  are measured as a function of the average transverse momenta  $p_T$  of the two leading jets in a phase space region ranging up to jet  $p_T$  of 2.0 TeV and an absolute rapidity of  $|y| = 2.5$ . The measurements after correcting for detector effects are well described by predictions at next-to-leading order in perturbative quantum chromodynamics and additionally are compared to several Monte Carlo event generators. The strong coupling constant at the scale of the Z boson mass is determined from a fit of the measured  $R_{32}$  which gives  $\alpha_s(M_Z) = 0.1150 \pm 0.0010 (\text{exp}) \pm 0.0013 (\text{PDF}) \pm 0.0015 (\text{NP})^{+0.0050}_{-0.0000} (\text{scale})$  using MSTW2008 PDF set. The current measurement is in well agreement with the world average value of  $\alpha_s(M_Z) = 0.1181 \pm 0.0011$  as well as previous measurements.

## ORIGINALITY REPORT

40%

SIMILARITY INDEX

32%

INTERNET SOURCES

39%

PUBLICATIONS

7%

STUDENT PAPERS

## PRIMARY SOURCES

1

[www.osti.gov](http://www.osti.gov)

Internet Source

8%

2

F. D. Aaron. "Jet production in ep collisions at low  $Q^2$  and determination of  $\alpha_s$ ", The European Physical Journal C, 03/19/2010

Publication

7%

3

[authors.library.caltech.edu](http://authors.library.caltech.edu)

Internet Source

7%

4

[discovery.ucl.ac.uk](http://discovery.ucl.ac.uk)

Internet Source

5%

5

D. Imperio, Giulia. "Measurements of the jet and vector boson in association with jets production, extraction of  $\alpha_s$  and PDF constraints at CMS", EPJ Web of Conferences, 2015.

Publication

5%

6

[www.ec.ms.imperial.ac.uk](http://www.ec.ms.imperial.ac.uk)

Internet Source

3%

7

V. Khachatryan. "Search for Dijet Resonances

in 7 TeV pp Collisions at CMS", Physical  
Review Letters, 11/2010

Publication

3%

8

T. Affolder. "Measurement of the Strong  
Coupling Constant from Inclusive Jet  
Production at the Tevatron  $p\bar{p}$  Collider",  
Physical Review Letters, 01/2002

Publication

2%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off