

Covariance Structure and Dimension Reduction PCA

Multivariate Statistic

Made by:

**Lasse Gøransson, Marc Evald,
Anne-Charlotte Poulsen & Aske
Møller**

SDU Robotics
The Maersk Mc-Kinney Møller Institute
University of Southern Denmark

Agenda



- ▶ Motivation
- ▶ Principal Components
- ▶ Variance
- ▶ Sample Based
- ▶ Pitfalls

Motivation



- ▶ Dimension reduction
- ▶ Decorrelate data

Principal Components



$$PC_i \stackrel{def}{=} e_{i1}X_1 + \dots + e_{ip}X_p = e_i^T X$$

Principal Components



$$PC_i \stackrel{def}{=} e_{i1}X_1 + \dots + e_{ip}X_p = e_i^T X$$

- Maximize variance

Variance Explained




$$\Sigma = E\Lambda E^T$$

$$PC_i : \frac{\lambda_i}{\sum_{j=1}^p \lambda_j}$$

Visualization



Visualization



Sample Based




$$X = \begin{bmatrix} X_{11} & \dots & X_{1p} \\ \vdots & \ddots & \vdots \\ X_{n1} & \dots & X_{np} \end{bmatrix}$$

$$\hat{\Sigma} = S = \frac{1}{n-1} \sum_{j=1}^n (X_j - \bar{X})(X_j - \bar{X})^T$$

$$\frac{\sum_{i=1}^r \hat{\lambda}_i}{\sum_{j=1}^p \hat{\lambda}_j}$$

Visualization





Pitfalls



$$X = \begin{bmatrix} X_{11} & \dots & X_{1p} \\ \vdots & \ddots & \vdots \\ X_{n1} & \dots & X_{np} \end{bmatrix}$$

Pitfalls





$$\hat{\lambda}_i \sim N\left(\lambda_i, 2\frac{\lambda_i^2}{n}\right)$$

$$\frac{\hat{\lambda}_i}{1 + \sqrt{\frac{2}{n}} z_{\frac{\alpha}{2}}} < \lambda_i < \frac{\hat{\lambda}_i}{1 - \sqrt{\frac{2}{n}} z_{\frac{\alpha}{2}}}$$