

MATH 335 S2019

Electrostatics and the Fundamental Theorem of Vector Calculus

1 Electrostatics and a Green's function for the Laplacian

Let μ be a charge density contained in a set V . Recall that the electric field induced by this charge density is given by

$$\vec{E}(\vec{r}) = \frac{1}{4\pi\epsilon_0} \iiint_V \frac{\vec{r} - \vec{p}}{|\vec{r} - \vec{p}|^3} \mu(\vec{p}) dV(\vec{p}).$$

We saw previously that this field is the gradient of a potential ϕ given by

$$\phi(\vec{r}) = -\frac{1}{4\pi\epsilon_0} \iiint_V \frac{1}{|\vec{r} - \vec{p}|} \mu(\vec{p}) dV(\vec{p}).$$

Note that this means that \vec{E} is conservative.

What is the divergence of \vec{E} ? When $\vec{r} \neq \vec{p}$, we have that

$$\nabla_r \cdot \frac{\vec{r} - \vec{p}}{|\vec{r} - \vec{p}|^3} = 3 \frac{1}{|\vec{r} - \vec{p}|^3} - \frac{3}{2} \frac{2(\vec{r} - \vec{p})}{|\vec{r} - \vec{p}|^5} \cdot (\vec{r} - \vec{p}) = 0$$

where we have used the vector identity $\nabla \cdot (\vec{f}g) = (\nabla \cdot \vec{f})g + \vec{f} \cdot \nabla g$. In some sense, all of the action happens when $\vec{p} = \vec{r}$ (where the potential is not defined). We can use the divergence theorem to figure out the value of $\nabla_r \cdot \vec{E}$. First, we bring the divergence inside the integral, i.e.

$$\nabla_r \cdot \vec{E}(\vec{r}) = \frac{1}{4\pi\epsilon_0} \iiint_V \nabla_r \cdot \frac{\vec{r} - \vec{p}}{|\vec{r} - \vec{p}|^3} \mu(\vec{p}) dV(\vec{p}) = -\frac{1}{4\pi\epsilon_0} \iiint_V \nabla_p \cdot \frac{\vec{r} - \vec{p}}{|\vec{r} - \vec{p}|^3} \mu(\vec{p}) dV(\vec{p})$$

where we have used the fact that $\frac{d}{dx} f(x - y) = -\frac{d}{dy} f(x - y)$. From the result above, we see that if \vec{r} is outside of V , then the integrand is always zero. Indeed, the divergence is equal to the same integral taken over any ball which contains \vec{r} . Let $B_\delta(\vec{r})$ be the ball of radius δ around the point \vec{r} and let $S_\delta(\vec{r})$ be its boundary (a sphere). If we take δ to be small, then we can approximate μ by a constant, namely $\mu(\vec{r})$ inside. Then,

$$\nabla_r \cdot \vec{E}(\vec{r}) \approx -\frac{\mu(\vec{r})}{4\pi\epsilon_0} \iiint_{B_\delta(\vec{r})} \nabla_p \cdot \frac{\vec{r} - \vec{p}}{|\vec{r} - \vec{p}|^3} dV(\vec{p}).$$

We can now apply the divergence theorem to this expression. We get

$$\nabla_r \cdot \vec{E}(\vec{r}) \approx \frac{\mu(\vec{r})}{4\pi\epsilon_0} \iint_{S_\delta(\vec{r})} \frac{\vec{p} - \vec{r}}{|\vec{p} - \vec{r}|^3} \cdot \hat{n} dS(\vec{p}) .$$

Setting $\vec{u} = \vec{p} - \vec{r}$, we get

$$\nabla_r \cdot \vec{E}(\vec{r}) \approx \frac{\mu(\vec{r})}{4\pi\epsilon_0} \iiint_{S_\delta(0)} \frac{\vec{u}}{|\vec{u}|^3} \cdot \frac{\vec{u}}{|\vec{u}|} dV(\vec{p}) = \frac{\mu(\vec{r})}{4\pi\epsilon_0\delta^2} \iint_{S_\delta(0)} dS(\vec{p}) = \frac{\mu(\vec{r})}{\epsilon_0} .$$

In the limit as $\delta \rightarrow 0$ this expression is exact for a continuous μ .

We now have that $\nabla \cdot \vec{E} = \nabla^2 \phi = \mu/\epsilon_0$, i.e. ϕ solves Poisson's equation. If we set $G(\vec{r}, \vec{p}) = -\frac{1}{4\pi} \frac{\vec{r} - \vec{p}}{|\vec{r} - \vec{p}|^3}$ we see that

$$\nabla^2 \iiint_V G(\vec{r}, \vec{p}) \mu(\vec{p}) dV(\vec{p}) = \mu(\vec{r}) ,$$

where the derivatives are taken with respect to \vec{r} , which we will denote by ∇_r . As in the 1D case from HW3, we call such a function a Green's function for the Laplacian. Because of this, we sometimes write $\nabla_r^2 G(\vec{r}, \vec{p}) = \delta_{\vec{r}}(\vec{p})$. The δ function simply means that $\iiint_V \delta_{\vec{r}}(\vec{p}) f(\vec{p}) dV(\vec{p}) = f(\vec{r})$ for sufficiently smooth f .

The Green's function idea can be used to establish some powerful results.

2 Warm up — a Representation Theorem

Recall Green's 2nd I.D., i.e.

$$\iiint_V g \nabla^2 f - f \nabla^2 g dV = \iint_S g \partial_n f - f \partial_n g dS .$$

If we set $g(\vec{p}) = G(\vec{r}, \vec{p})$ (where G is the Green's function above) and let f be any twice continuously differentiable function, then

$$\iiint_V G \nabla^2 f - f \nabla^2 G dV = \iint_S G \partial_n f - f \partial_n G dS .$$

Rearranging, we get

$$f(\vec{r}) = \iiint_V G \nabla^2 f dV + \iint_S f \partial_n G - G \partial_n f dS .$$

A result known as Green's third identity. Note that this is rather remarkable: any twice continuously differentiable function can be determined by its Laplacian in the domain and its value and normal derivative on the boundary.

If $\nabla^2 f = 0$, then

$$f(\vec{r}) = \iint_S f \partial_n G - G \partial_n f dS ,$$

Which shows that a harmonic function f can be represented by an integral over the boundary. A similar result holds for vector fields, which we will show next.

3 Fundamental Theorem of Vector Calculus

Before we prove the main theorem, we require the following lemma, which is like the divergence theorem for the curl operator.

Lemma 1. *Let \vec{v} be a vector field defined on a domain D with boundary S . Then*

$$\iiint_V \nabla \times \vec{v} dV = \iint_S \hat{n} \times \vec{v} dS$$

Proof. We can establish the following identity for a constant vector \vec{a} and vector field \vec{v} using suffix notation. We have

$$\begin{aligned} \nabla \cdot (\vec{a} \times \vec{v}) &= \partial_i \epsilon_{ijk} a_j v_k \\ &= \epsilon_{ijk} \partial_i (a_j v_k) \\ &= a_j \epsilon_{ijk} \partial_i v_k \\ &= -a_j \epsilon_{jik} \partial_i v_k \\ &= -\vec{a} \cdot \nabla \times \vec{v} \end{aligned}$$

Applying the divergence theorem to $\vec{u} = \vec{a} \times \vec{v}$, we obtain

$$\begin{aligned} \iiint_V \nabla \cdot (\vec{a} \times \vec{v}) dV &= \iint_S \vec{a} \times \vec{v} \cdot \hat{n} dS \\ \iiint_V -\vec{a} \cdot \nabla \times \vec{v} dV &= \iint_S \vec{a} \cdot \vec{v} \times \hat{n} dS \\ \iiint_V \vec{a} \cdot \nabla \times \vec{v} dV &= \iint_S \vec{a} \cdot \hat{n} \times \vec{v} dS \end{aligned}$$

Because this holds for any \vec{a} , we can read off the desired result one entry at a time by setting $\vec{a} = e_1, e_2, e_3$. \square

In the following theorem, we will make use of some of the vector identities from Chapter 4:

- $\nabla^2 \vec{u} = \nabla(\nabla \cdot \vec{u}) - \nabla \times \nabla \times \vec{u}$
- $\nabla \cdot (f\vec{u}) = f\nabla \cdot \vec{u} + \vec{u} \cdot \nabla f$
- $\nabla \times (f\vec{u}) = f\nabla \times \vec{u} + \nabla f \times \vec{u}$.

We will also make use of the fact that $\nabla_p G(\vec{p}, \vec{r}) = -\nabla_r G(\vec{p}, \vec{r})$. Finally, we will use the divergence theorem and the fact that G is a Green's function for the Laplacian.

Theorem 1 (Fundamental Theorem of Vector Calculus). *Let \vec{F} be a twice continuously differentiable vector field defined on a domain V with boundary S . Then \vec{F} can be decomposed into a curl-free component and a divergence-free component: $\vec{F} = \nabla\Phi + \nabla \times \vec{\Psi}$, where*

$$\Phi = \iiint_V G(\vec{p}, \vec{r}) \nabla \cdot \vec{F}(\vec{r}) dV(\vec{r}) - \iint_S G(\vec{p}, \vec{r}) \hat{n} \cdot \vec{F}(\vec{r}) dS(\vec{r}) \quad (1)$$

$$\vec{\Psi} = - \iiint_V G(\vec{p}, \vec{r}) \nabla \times \vec{F}(\vec{r}) dV(\vec{r}) + \iint_S G(\vec{p}, \vec{r}) \hat{n} \times \vec{F}(\vec{r}) dS(\vec{r}) . \quad (2)$$

Proof. To start, we note that

$$\vec{F} = \nabla_p^2 \iiint_V G(\vec{p}, \vec{r}) \vec{F}(\vec{r}) dV(\vec{r}) . \quad (3)$$

Using the identity $\nabla^2 \vec{u} = \nabla(\nabla \cdot \vec{u}) - \nabla \times \nabla \times \vec{u}$, we have that

$$\vec{F} = \nabla_p \nabla_p \cdot \iiint_V G(\vec{p}, \vec{r}) \vec{F}(\vec{r}) dV(\vec{r}) - \nabla_p \times \nabla_p \times \iiint_V G(\vec{p}, \vec{r}) \vec{F}(\vec{r}) dV(\vec{r}) . \quad (4)$$

Note that

$$\begin{aligned} \nabla_p \cdot (G(\vec{p}, \vec{r}) \vec{F}(\vec{r})) &= \nabla_p G(\vec{p}, \vec{r}) \cdot \vec{F}(\vec{r}) + 0 \\ &= -\nabla_r G(\vec{p}, \vec{r}) \cdot \vec{F}(\vec{r}) \\ &= G(\vec{p}, \vec{r}) \nabla_r \cdot \vec{F}(\vec{r}) - \nabla_r \cdot (G(\vec{p}, \vec{r}) \vec{F}(\vec{r})) \end{aligned}$$

and similarly

$$\begin{aligned} \nabla_p \times (G(\vec{p}, \vec{r}) \vec{F}(\vec{r})) &= \nabla_p G(\vec{p}, \vec{r}) \times \vec{F}(\vec{r}) + 0 \\ &= -\nabla_r G(\vec{p}, \vec{r}) \times \vec{F}(\vec{r}) \\ &= G(\vec{p}, \vec{r}) \nabla_r \times \vec{F}(\vec{r}) - \nabla_r \times (G(\vec{p}, \vec{r}) \vec{F}(\vec{r})) . \end{aligned}$$

Therefore,

$$\begin{aligned} \vec{F} &= \nabla_p \iiint_V G(\vec{p}, \vec{r}) \nabla_r \cdot \vec{F}(\vec{r}) - \nabla_r \cdot (G(\vec{p}, \vec{r}) \vec{F}(\vec{r})) dV(\vec{r}) \\ &\quad - \nabla_p \times \iiint_V G(\vec{p}, \vec{r}) \nabla_r \times \vec{F}(\vec{r}) - \nabla_r \times (G(\vec{p}, \vec{r}) \vec{F}(\vec{r})) dV(\vec{r}) . \end{aligned}$$

Then, we can apply the divergence theorem and the lemma above to obtain

$$\begin{aligned}\vec{F} = & \nabla_p \left(\iiint_V G(\vec{p}, \vec{r}) \nabla_r \cdot \vec{F}(\vec{r}) dV(\vec{r}) - \iint_S G(\vec{p}, \vec{r}) \vec{F}(\vec{r}) \cdot \hat{n} dS(\vec{r}) \right) \\ & - \nabla_p \times \left(\iiint_V G(\vec{p}, \vec{r}) \nabla_r \times \vec{F}(\vec{r}) dV(\vec{r}) - \iint_S G(\vec{p}, \vec{r}) \hat{n} \times \vec{F}(\vec{r}) dS(\vec{r}) \right) .\end{aligned}$$

The above means we have explicit formulas for Φ and $\vec{\Psi}$ given \vec{F} , i.e.

$$\begin{aligned}\Phi(\vec{p}) &= \iiint_V G(\vec{p}, \vec{r}) \nabla_r \cdot \vec{F}(\vec{r}) dV(\vec{r}) - \iint_S G(\vec{p}, \vec{r}) \vec{F}(\vec{r}) \cdot \hat{n} dS(\vec{r}) \\ \vec{\Psi}(\vec{p}) &= - \iiint_V G(\vec{p}, \vec{r}) \nabla_r \times \vec{F}(\vec{r}) dV(\vec{r}) + \iint_S G(\vec{p}, \vec{r}) \hat{n} \times \vec{F}(\vec{r}) dS(\vec{r}) .\end{aligned}$$

□