Reading Group(oid) Lecture Notes



Organisers/editors:
Oscar Carballal (UCM, Madrid)
Rubén Izquierdo-López (ICMAT, Madrid)
Asier López-Gordón (IMPAN, Warsaw)
Juan Manuel López Medel (ICMAT, Madrid)
Bartosz Maciej Zawora (KMMF-UW, Warsaw)



Last update: March 23, 2026

Contents

Lectures 3

Lecture 1 – O. Carballal (UCM) 5

Lecture 2 – Bartosz M. Zawora (KMMF - UW) 11
Lie groupoids: definition and examples 11
Morphisms and subgroupoids 14

Lecture 3– Rubén Izquierdo-López (ICMAT) 17
Locally trivial Lie groupoids 17

Lecture 4 – Asier López-Gordón (IMPAN) 21
Bisections 21

Lecture 5 – Juan Manuel López Medel (ICMAT) 29
Components and transitivity 31

Lecture 6 – Paula Alba San Miguel (UNED) 35
Actions 35

Lecture 7 – [Tomasz Sobczak (KMMF-UW)] 41

Lecture 8 – [Juan Manuel López Medel (ICMAT)] 43
Linear actions and frame groupoids 43

Lecture 9 – O. Carballal (UCM) 49

Lecture 10 – Asier López-Gordón (IMPAN) 51

Lecture 11 and 12 – BARTOSZ 57

Lecture 14 – Oscar Carballal (UCM) 59
Lie Algebroid connections 60
Representations of Lie algebroids and Lie algebroids cohomology 62
The Atiyah algebroid 63
Parallel transport along A-paths 64

Lecture 15– Arnau Mas 65

Lecture 16 – Juan Manuel López Medel 67
Symplectic realization 67
Symplectic groupoid 69
Examples 71
Integrability problem 73

Bibliography 75

Lectures

Lecture 1 – O. Carballal (UCM)

Date: 7 October, 2025

Abstract: We illustrate the need for groupoids through a series of examples, primarily following [**2**, Lecture 1] and [**3**], and discuss some of their recent applications in Mathematical Physics. We also outline potential directions for future topics of our Reading Groupoid.

Since F. Klein's *Erlangen program* and the pioneer work of S. Lie regarding the definition of geometric structures through their group of automorphisms, the notion of symmetry has been studied through the theory of groups and their actions.

Symmetry $\equiv$ *Groups & actions*

Symmetry of homogeneous structures, such us homogeneous spaces, can be fully described using groups. This is no longer valid when analysing the symmetry of non-homogeneous structures.

The very first example of symmetry group is the *Euclidean group* $\mathrm{E}(n)$, consisting of rigid motions of $\mathbb{R}^n$. More concretely,

$$\mathrm{E}(n) := \{\phi : \mathbb{R}^n \to \mathbb{R}^n : \phi \text{ preserves distances}\}$$

Every $\phi \in \mathrm{E}(n)$ is univocally determined by a rotation $A \in \mathrm{O}(n)$ and a translation vector $b \in \mathbb{R}^n$ as $\phi(x) = Ax + b$ and, hence, $\mathrm{E}(n)$ is the semidirect product $\mathrm{E}(n) = \mathrm{O}(n) \ltimes \mathbb{R}^n$. Let us now consider the action $\mathrm{E}(n) \curvearrowright \mathbb{R}^n$ given by $\phi \cdot x := \phi(x)$ for every $\phi \in \mathrm{E}(n)$ and $x \in \mathbb{R}^n$. Then, we define the symmetry group of a subset Ω as the subgroup G_Ω of $\mathrm{E}(n)$ formed by those rigid motions leaving Ω invariant; that is,

$$G_\Omega := \{\phi \in \mathrm{E}(n) : \phi(\Omega) = \Omega\}.$$

Let us now adopt the following *credo*:

G_Ω is large $\equiv$ Ω is very symmetric

Example 1.1 (Weinstein's tiling [**3**])**.** Let us study the symmetry group of the following tiling of $\mathbb{R}^2$ by 2×1-rectangles:

$$\Omega := (2\mathbb{Z} \times \mathbb{R}) \cup (\mathbb{R} \times \mathbb{Z}) \subset \mathbb{R}^2.$$

Within this context, every connected component of $\mathbb{R}^2 - \Omega$ is called a *tile*. Let Λ be the lattice determined by Ω, corresponding at the borders of the tiles,

$$\Lambda = (2\mathbb{Z} \times \mathbb{R}) \cap (\mathbb{R} \times \mathbb{Z}) = 2\mathbb{Z} \times \mathbb{Z}.$$

Then, one easily sees that the symmetry group G_Ω of Ω consists of the following elements:

- translations by vectors $u \in \Lambda$, corresponding to the corner points of Λ;
- central reflections through the points of $\frac{1}{2}\Lambda = \mathbb{Z} \times \frac{1}{2}\mathbb{Z}$, the latter corresponding to the centers of the tiles, and
- axial reflections along the vertical and horizontal lines trough the points of $\frac{1}{2}\Lambda$.

Since G_Ω is a big group, following the faith in our *credo*, Ω is very symmetric. Nevertheless, there are some problems:

- Ω and the lattice Λ have the same symmetry group $G_\Omega = G_\Lambda$, although Ω and Λ look quite different;
- G_Ω contains no local information about the action $G_\Omega \curvearrowright \mathbb{R}^2$, and
- the symmetry group of the finite tiling $\widetilde{\Omega} := \Omega \cap B$, where $B := [0, 2m] \times [0, n]$ is a 'bathroom floor', is isomorphic to the Klein four group $\mathbb{Z}_2 \times \mathbb{Z}_2$ (note that it is spanned by the horizontal and the vertical reflections through the midline of B, whose composition is the 180° rotation around the centre of B). In particular, we see that Ω is very symmetric but $\widetilde{\Omega}$ is not, even though they share the same patterns and independently of the number of tiles.

To address these issues, we introduce the *action groupoid* associated with $G_\Omega \curvearrowright \mathbb{R}^n$, defined as

$$\mathcal{G}_\Omega := \{(y, \phi, x) : \phi \in G_\Omega, x, y \in \mathbb{R}^n, y = \phi(x)\}, \tag{1}$$

and endowed with the partially defined multiplication

$$(z, \psi, y)(y, \phi, x) := (z, \psi \circ \phi, x). \tag{2}$$

In order to analyse the properties of this partially defined multiplication, we introduce the *source* $\mathrm{s} : \mathcal{G}_\Omega \to \mathbb{R}^n$ and *target* $\mathrm{t} : \mathcal{G}_\Omega \to \mathbb{R}^n$ maps as

$$\mathrm{s}(y, \phi, x) := x, \qquad \mathrm{t}(y, \phi, x) := y, \qquad (y, \phi, x) \in \mathcal{G}_\Omega.$$

One easily sees that (2) satisfies the following four properties:

(1) Composition. Given $g, h \in \mathcal{G}_\Omega$, gh is only defined when $\mathrm{s}(g) = \mathrm{t}(h)$. In that case, $\mathrm{s}(gh) = \mathrm{s}(h)$ and $\mathrm{t}(gh) = \mathrm{t}(g)$.
(2) Associativity. If $g, h, k \in \mathcal{G}_\Omega$ are such that $\mathrm{s}(g) = \mathrm{t}(h)$ and $\mathrm{s}(h) = \mathrm{t}(k)$, then $(gh)k = g(hk)$.
(3) Units. For all $x \in \mathbb{R}^n$, let us denote $1_x := (x, \mathrm{id}, x) \in \mathcal{G}_\Omega$. Then, $1_{\mathrm{t}(g)}g = g = g1_{\mathrm{s}(g)}$ for all $g \in \mathcal{G}_\Omega$.
(4) Inverse. For all $g = (y, \phi, x) \in \mathcal{G}_\Omega$ there exists $g^{-1} := (x, \phi^{-1}, y)$ such that $gg^{-1} = 1_{\mathrm{t}(g)}$ and $g^{-1}g = 1_{\mathrm{s}(g)}$.

These are, exactly, the properties characterising a groupoid.

Definition 1.2 (Groupoid)**.** A *groupoid* consists of two sets, $\mathcal{G}$ and M, together with maps

- $\mathrm{s}, \mathrm{t} : \mathcal{G} \to M$ (*source* and *target* projections);
- $m : \mathcal{G}^{(2)} := \{(g, h) : \mathrm{s}(g) = \mathrm{t}(h)\} \to \mathcal{G}, (g, h) \mapsto gh$ (*multiplication*);
- $u : M \to \mathcal{G}, x \mapsto 1_X$ (*unit*), and
- $i : \mathcal{G} \to \mathcal{G}, g \mapsto g^{-1}$ (*inverse*),

satisfying the following four properties:

(1) If $z \xleftarrow{g} y \xleftarrow{h} x$, then $z \xleftarrow{gh} x$;

(2) If $z \xleftarrow{g} y \xleftarrow{h} x \xleftarrow{k} v$, then $(gh)k = g(hk)$;
(3) There exists $x \xleftarrow{1_x} x$ such that for all $y \xleftarrow{g} x$, we have $1_y g = g = 1_x$, and
(4) If $y \xleftarrow{g} x$, there exists $x \xleftarrow{g^{-1}} y$ such that $gg^{-1} = 1_y$ and $g^{-1}g = 1_x$.

From now on, we will denote the groupoid as $\mathcal{G} \rightrightarrows M$, where the parallel arrows refer to the source and target maps.

Remark 1.3. (1) A groupoid is a small category where every arrow is invertible. Within this language, given a groupoid $\mathcal{G} \rightrightarrows M$, $\mathcal{G}$ is usually referred to as the set of *arrows* of the groupoid, M is the set of *objects* or *units* (also called the *base* of the groupoid), and $\mathcal{G}^{(2)}$ is the set of *composable arrows*.
(2) We write arrows from left to right to refer to the order of composition. Everything can be done analogously writing arrows from left to right by swapping the source and target maps s and t.
(3) Groupoids can be restricted to subsets of the set of units. That is, given a groupoid $\mathcal{G} \rightrightarrows M$, *its restriction* to $N \subset M$ is the groupoid

$$\mathcal{G}|_N := \mathrm{s}^{-1}(N) \cap \mathrm{t}^{-1}(N) \rightrightarrows N.$$

Definition 1.4 (Orbit and isotropy group)**.** Let $\mathcal{G} \rightrightarrows M$ be a groupoid and let $x \in M$ be a unit.

- The *orbit of* x is $O_x := \{y \in M : \text{there exists } g \in \mathcal{G}, y \xleftarrow{g} x\}$.
- The *isotropy group of* x is $\mathcal{G}_x := \mathrm{s}^{-1}(x) \cap \mathrm{t}^{-1}(x) = \{g \in \mathcal{G} : x \xleftarrow{g} x\}$.

Example 1.5 (Weinstein's tiling continued [**3**])**.** Let us now go back to the tiling Ω of $\mathbb{R}^2$ introduced in Example 1.1. First of all, we consider the action groupoid $\mathcal{G}_\Omega \rightrightarrows \mathbb{R}^2$ associated with $G_\Omega \curvearrowright \mathbb{R}^2$, defined in (1); that is,

$$\mathcal{G}_\Omega := \{(y, \phi, x) : \phi \in G_\Omega, y \xleftarrow{\phi} x\} \rightrightarrows \mathbb{R}^2,$$

and let us now consider its restriction to the 'bathroom floor' $B = [0, 2m] \times [0, n]$, namely $\mathcal{G}_\Omega|_B \rightrightarrows B$. Then:

- $x, y \in B$ belong to the same orbit if they are similarly placed in their tiles, and
- the isotropy group of a point $x \in B$ is trivial, unless $x \in \frac{1}{2}\Lambda \cap B$, for which its isotropy is the Klein four group.

This means that $\mathcal{G}_\Omega|_B \rightrightarrows B$ detects local information.

Some applications to Mathematical Physics.

- Groupoid picture of Schwinger's quantum mechanics by F. M. Ciaglia, A. Ibort, G. Marmo and collaborators initiated in [**4**]. Recent work on the formulation of fields as functors between groupoids by A. Ibort, A. Mas and L. Schiavone [**5**].
- The material groupoid, a groupoidal approach to continuum mechanics, by M. de León, M. Epstein and V. Jiménez [**6**].
- Discrete Lagrangian and Hamiltonian mechanics on Lie groupoids by J. C. Marrero, D. Martín de Diego, E. Martínez and L. Colombo [**7, 8**].

Local Lie group actions. Let us consider the Lie group $\mathrm{PGL}(2,\mathbb{R})$ consisting of homographies of the projective line $\mathbb{RP}^1$. Explicitly, one has

$$\mathrm{PGL}(2,\mathbb{R}) = \mathrm{GL}(2,\mathbb{R})/\mathbb{R},$$

where $\mathbb{R}$ refers to the subgroup of $\mathrm{GL}(2,\mathbb{R})$ formed by non-zero diagonal scalar matrices. Clearly, $\mathrm{PGL}(2,\mathbb{R})$ acts on $\mathbb{RP}^1$ as

$$\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right] \cdot [x_0 : x_1] = [ax_0 + bx_1 : cx_0 + dx_1]. \tag{3}$$

Let us now analyse how $\mathrm{PLG}(2,\mathbb{R})$ acts on the affine line $\mathbb{R} \equiv \mathbb{RP}^1 - \{[1:0]\}$. Given an affine point $[x_0 : x_1] = [x : 1] \in \mathbb{R}$, where $x := x_0/x_1$, and $\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right] \in \mathrm{PGL}(2,\mathbb{R})$, we have that

$$\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right] \cdot [x : 1] = \left[\frac{ax+b}{cx+d} : 1\right] \in \mathbb{R}, \tag{4}$$

provided that $cx + d \neq 0$. It is natural to consider the following question:

Does formula (4) define a Lie group action on $\mathbb{R}$?

Of course, the answer to this question is negative, since (4) is only defined on an open subset of $\mathrm{PGL}(2,\mathbb{R}) \times \mathbb{R}$. Indeed, it defines what is called a *local Lie group action* (see [**9**] for more details). This means, roughly speaking, that (4) only satisfies the defining properties of a Lie group action on a neighbourhood of the unit element of $\mathrm{PGL}(2,\mathbb{R})$.

Hopefully, this problem can be tackled successfully when one considers groupoids rather than groups. First of all, let us consider the so-called action groupoid associated with the Lie group action $\mathrm{PGL}(2,\mathbb{R}) \curvearrowright \mathbb{RP}^1$ defined in (3) as the groupoid $\mathcal{G} := \mathrm{PGL}(2,\mathbb{R}) \times \mathbb{RP}^1 \rightrightarrows \mathbb{RP}^1$ with source and target maps given by

$$\mathrm{s}\left([A],[x_0 : x_1]\right) = [x_0 : x_1], \qquad \mathrm{t}\left([A],[x_0 : x_1]\right) = [A] \cdot [x_0 : x_1],$$

and multiplication defined as

$$([B],[y_0 : y_1])([A],[x_0 : x_1]) := ([BA],[x_0, x_1]),$$

provided that $[A] \cdot [x_0 : x_1] = [y_0 : y_1]$. Then, the restriction of $\mathcal{G} \rightrightarrows \mathbb{RP}^1$ to the affine line $\mathbb{R}$ is the groupoid

$$\mathcal{G}|_{\mathbb{R}} = \mathrm{s}^{-1}(\mathbb{R}) \cap \mathrm{t}^{-1}(\mathbb{R}) = \left\{ \left(\left[\begin{pmatrix} a & b \\ c & d \end{pmatrix}\right], [x : 1] \right) \in \mathrm{PGL}(2,\mathbb{R}) \times \mathbb{R} : cx + d \neq 0 \right\} \rightrightarrows \mathbb{R}.$$

Within this context, the target map of $\mathcal{G}|_{\mathbb{R}} \rightrightarrows \mathbb{R}$ is just formula (4).

Singular spaces. Let $G \curvearrowright M$ be a smooth action of a Lie group G on a manifold M (see [**10**, Chapter 2] for more details on Lie group actions). Recall that the action is said to be

- *free* if $g \cdot x = x$ for some $x \in M$, then $g = e$, and

- *proper* if the map $\Phi : G \times M \ni (g, x) \mapsto (g \cdot x, x) \in M \times M$ is a proper map. That is, for every compact subset $K \subset M \times M$ we have that $\Phi^{-1}(K) \subset G \times M$ is compact. This is equivalent to the following: if (g_n) and (x_n) are sequences in G and M, respectively, such that (x_n) and $(g_n \cdot x_n)$ are convergent, then (g_n) contains a converging subsequence.

Provided that the action $G \curvearrowright M$ is free an proper, the quotient manifold M/G has a unique smooth structure such that the projection map $\pi : M \to M/G$ is a submersion. For instance, let us consider the action $G = \mathbb{Z}^k \curvearrowright M = \mathbb{R}^k$ given by translations:

$$(n_1, \ldots, n_k) \cdot (x^1, \ldots, x^k) := (x^1 + n_1, \ldots, x^k + n_k).$$

It can be easily seen that this action is free and proper. Hence, the quotient space $\mathbb{R}^k/\mathbb{Z}^k$, which is the k-dimensional torus $\mathbb{T}^k := \mathbb{R}^k/\mathbb{Z}^k$, possesses a unique smooth structure such that the projection $\pi : M \to M/G$ is a submersion.

What happens when the action $G \curvearrowright M$ is not free and proper?

If action $G \curvearrowright M$ is not free and proper, G/M is a "singular space"

Example 1.6. Let us consider the actions $\mathrm{SO}(2) \curvearrowright \mathbb{R}^2$ and $\mathrm{SO}(3) \curvearrowright \mathbb{R}^3$ given by rotations. Both of these actions are proper, since the associated Lie groups are compact. Nevertheless, they are not free: the origin is a fixed point of every rotation. The "singular spaces" $\mathbb{R}^2/\mathrm{SO}(2)$ and $\mathbb{R}^3/\mathrm{SO}(3)$ are both homeomorphic to $[0, +\infty)$. Notwithstanding, they are not equivalent as singular spaces because their associated action groupoids $\mathrm{SO}(2) \times \mathbb{R}^2 \rightrightarrows \mathbb{R}^2$ and $\mathrm{SO}(3) \times \mathbb{R}^3 \rightrightarrows \mathbb{R}^3$ are not *Morita equivalent*. Roughly speaking, this can be justified as follows. The isotropy group of $x \in \mathbb{R}^2 - \{0\}$ is trivial, while the isotropy group of $x \in \mathbb{R}^3 - \{0\}$ is isomorphic to $\mathrm{SO}(2)$.

Lecture 2 – Bartosz M. Zawora (KMMF - UW)

Date: 14 October, 2025

Abstract: During the lecture, I will recall the definition of a groupoid from the previous session and present the definition of a Lie groupoid. I will provide many useful and important examples and prove some fundamental properties. The lecture will follow Mackenzie's book "General Theory of Lie Groupoids and Lie Algebroids".

Lie groupoids: definition and examples

Definition 2.1. A groupoid consists of two sets $\mathcal{G}$ and M, with maps,
$s : \mathcal{G} \to M$ (source),
$t : \mathcal{G} \to M$ (target),
$u : M \to \mathcal{G} \mid x \mapsto 1_x$ (unit),
$i : \mathcal{G} \to \mathcal{G} \mid g \longmapsto g^{-1}$ (inverse),
$\kappa : \mathcal{G} * \mathcal{G} \to \mathcal{G} \mid (h, g) \longmapsto hg$ where $\mathcal{G} * \mathcal{G} := \{(h, g) \in \mathcal{G} \times \mathcal{G} \mid s(h) = t(g)\}$,
satisfying

(1) $s(hg) = s(g)$ and $t(hg) = t(h)$ for all $(h, g) \in \mathcal{G} * \mathcal{G}$,
(2) $j(hg) = (jh)g$ for every $j, h, y \in \mathcal{G}$ such that $s(j) = t(h)$ and $s(h) = t(g)$,
(3) $s(1_x) = t(1_x) = x$ for any $x \in M$,
(4) $g1_{s(g)} = g$ and $1_{t(y)}g = g$ for every $g \in \mathcal{G}$,
(5) For each $g \in \mathcal{G}$, there exists two-sided inverse g^{-1} such that $s(g^{-1}) = t(g)$, $t(g^{-1}) = s(g)$, $g^{-1}g = 1_{s(g)}$, and $gg^{-1} = 1_{t(g)}$.

To simplify the notation, $\mathcal{G} \rightrightarrows M$ denotes a groupoid.

Proposition 2.2. Let $\mathcal{G} \rightrightarrows M$ and $g \in \mathcal{G}$ with $s(g) = x$ and $t(g) = y$.

(1) If $h \in \mathcal{G}$ with $s(h) = g$ and $hg = g$, then $h = 1_y$.
If $j \in \mathcal{G}$ with $t(j) = x$ and $gj = g$, then $j = 1_x$.
(2) If $h \in \mathcal{G}$ with $s(h) = y$ and $hg = 1_x$, then $h = g^{-1}$.
If $j \in \mathcal{G}$ with $t(j) = x$ and $gj = 1_y$, then $j = g^{-1}$.

Definition 2.3. Let $G \rightrightarrows M$. Then, $\mathcal{G}_x := s^{-1}(x)$ is called a s-fibre over $x \in M$, similarly $\mathcal{G}^y := t^{-1}(x)$ is a t-fibre over $y \in M$, and $\mathcal{G}_x^y := \mathcal{G}_x \cap \mathcal{G}^y = s^{-1}(x) \cap t^{-1}(y)$. Moreover, the set $\mathcal{G}_x \cap \mathcal{G}^y$ is called the vertex group. The set of identity elements 1_M is defined as $\{1_m \mid m \in M\}$.

Definition 2.4. A Lie groupoid $\mathcal{G} \rightrightarrows M$ is a groupoid $\mathcal{G} \rightrightarrows M$ with smooth manifold structures on $\mathcal{G}$ and M such that $s, t \colon \mathcal{G} \to M$ are surjective submersions, $u \colon M \to \mathcal{G}$ is smooth, and $\kappa \colon \mathcal{G} * \mathcal{G} \to \mathcal{G}$ is also smooth.

Remark 2.5. Since $s, t\colon \mathcal{G} \to M$ are submersions it follows that $\mathcal{G}_x$, $\mathcal{G}^y$, and $(s \times t)^{-1}(x, x')$ are closed embedded submanifolds for some $x, x' \in M$.

Definition 2.6. Let $\mathcal{G} \rightrightarrows M$ and $g \in \mathcal{G}$ with $s(g) = x$ and $t(g) = y$. Then, the left-translation corresponding to g is $L_g\colon h \in \mathcal{G}^x \mapsto gh \in \mathcal{G}^y$. Analogously, the right-translation corresponding to g is $R_g\colon h \in \mathcal{G}_y \mapsto hg \in \mathcal{G}_x$

Definition 2.7. A Lie groupoid is s-connected if each of its fibres is connected. Likewise, for any other property α, a Lie groupoid $\mathcal{G} \rightrightarrows M$ is s-α if each of its fibres has property α.

Proposition 2.8. Let $\mathcal{G} \rightrightarrows M$. The inverse map $i\colon \mathcal{G} \to \mathcal{G}$ is a diffeomorphism.

Proof. Note that the tangent bundle to $\mathcal{G} * \mathcal{G}$ is given by

$$\mathrm{T}(\mathcal{G} * \mathcal{G}) = \{(Y, X) \in \mathrm{T}\mathcal{G} \times \mathrm{T}\mathcal{G} \mid \mathrm{T}s(Y) = \mathrm{T}t(X)\}\,.$$

Suppose that $h, g \in \mathcal{G}$ are such that $s(h) = t(g)$, $Y \in \mathrm{T}_h(\mathcal{G}_{s(h)})$, and $X \in \mathrm{T}_g(\mathcal{G}^{t(g)})$. Then, by Leibniz rule, it follows

$$\mathrm{T}_{(h,g)}\kappa(Y, X) = \mathrm{T}_h R_g(Y) + \mathrm{T}_g L_h(X)\,. \tag{5}$$

Define $\theta\colon \mathcal{G} * \mathcal{G} \to \mathcal{G} \times_t \mathcal{G}$ as

$$\theta(h, g) = (h, hg)\,,$$

where $\mathcal{G} \times_t \mathcal{G} := \{(h, g) \in \mathcal{G} \times \mathcal{G} \mid t(h) = t(g)\}$. Then, one can simply verify that θ is bijective with $\theta^{-1}(i, j) = (i, i^{-1}j)$. To prove that θ is an immersion, suppose that $\mathrm{T}_{(h,g)}\theta(Y, X) = (0, 0)$. Moreover, define $\pi_1\colon (h, g) \in \mathcal{G} \times \mathcal{G} \mapsto g \in \mathcal{G}$ and $\pi_2\colon (h, g) \in \mathcal{G} \times \mathcal{G} \mapsto h \in \mathcal{G}$. Therefore, since $\pi_2 \circ \theta = \pi_2$, it follows that $Y = 0$. Consequently, by (5), one has $X = 0$ and θ is an immersion.

Furthermore, by the fact that s and t are submersions and $\dim \mathcal{G} * \mathcal{G} = \dim \mathcal{G} \times_t \mathcal{G}$, one obtains that θ is a diffeomorphism. For $\mu\colon h \in \mathcal{G} \mapsto (h, 1_{t(h)}\mathcal{G} \times_t \mathcal{G}$, one has

$$(\pi_1 \circ \theta^{-1} \circ \mu)(g) = (\pi_1 \circ \theta^{-1})(g, 1_{t(g)}) = \pi_1(g, g^{-1}) = g^{-1}\,.$$

Hence, $(\pi_1 \circ \theta^{-1} \circ \mu) = i$. Since $(\pi_1 \circ \theta^{-1} \circ \mu)$ is smooth, it follows that i is also smooth. In addition, since inversion is its own inverse, it is therefore a diffeomorphism. □

Remark 2.9. The unit map $u\colon M \to \mathcal{G}$ is smooth and is an immersion, by the fact that

$$t \circ u = \mathrm{id}_M\,, \qquad s \circ u = \mathrm{id}_M\,.$$

Consequently, 1_M is a closed embedded submanifold of $\mathcal{G}$.

Example 2.10 (Lie group)**.** A Lie group is a Lie groupoid with a unique unit, namely $G \rightrightarrows \{*\}$.

Example 2.11 (Manifold)**.** Any manifold M can be regarded as a Lie groupoid on itself, where $s = t = \mathrm{id}_M$.

Example 2.12 (A pair groupoid)**.** For any manifold M, one has a pair groupoid $M \times M \rightrightarrows M$ with arrows $(y, x) \in M \times M$, $s(y, x) = x$, $t(y, x) = y$, $u(x) = (x, x)$, $i(y, x) = (x, y)$, and

$$\kappa((z, y), (y, x)) = (z, x).$$

Note that $\mathcal{G}^x_x$ is trivial.

Example 2.13 (Trivial groupoid)**.** For a Lie group G and a manifold M one can construct a Lie groupoid $M \times G \times M \rightrightarrows M$ with $s(y,g,x) = x$, $t(y,g,x) = y$, $u(x) = (x,e,x)$, $i(y,g,x) = (x,g^{-1},y)$, and

$$\kappa((z,h,y),(y,g,x)) = (y,hg,x)\,.$$

Example 2.14 (Submersion groupoid)**.** For a surjective submersion $\pi\colon M \to Q$ one has a submersion groupoid

$$M \times_Q M \rightrightarrows M\,,$$

where $M \times_Q M = \{(y,x) \in M \times M \mid \pi(y) = \pi(x)\}$ and is given by the restriction of a pair groupoid $M \times M \rightrightarrows M$ to $M \times_Q M$.

Example 2.15 (Action groupoid)**.** Let $G \times M \to M$ be a Lie group action on a manifold M. Then, $G \times M \rightrightarrows M$ is an action Lie groupoid with $s(g,x) = x$, $t(g,x) = gx$, $u(x) = (e,x)$, $i(g,x) = (g^{-1},gx)$, and

$$\kappa((g_2,y),(g_1,x)) = (g_2g_1,x)\,,$$

of $y = g_1x$. Note that $\mathcal{G} \times \mathcal{M}_x^x$ is isomorphic to G_x. An action Lie groupoid is denoted as $G \triangleleft M$.

Example 2.16. ADD DRAWINGS

Example 2.17 (Fundamental groupoid)**.** Let M be a smooth manifold and let $\Pi(M)$ be the set of homotopy classes $[\gamma]$ of continuous path $\gamma\colon [0,1] \to M$, relative to fixed and points $\gamma(0)$ and $\gamma(1)$. Then, $\Pi(M) \rightrightarrows M$ is a groupoid with $s([\gamma]) = \gamma(0)$, $t([\gamma])\gamma(1)$, $u(x) = [\gamma_x]$, where γ_x is a constant path at $x \in M$, and

$$\kappa([\delta],[\gamma]) = [\delta\gamma]\,,$$

where $\delta\gamma$ is the standard concatenation of γ and δ, namely

Add

Moreover, $i([\gamma]) = [\overleftarrow{\gamma}]$, where $\overleftarrow{\gamma}$ is the reverse path, namely $\overleftarrow{\gamma}(t) = \gamma(1-t)$.

Example 2.18. Let (E,q,M) be a vector bundle. Let $\Phi(E)$ denote the set of all vector space isomorphism $\xi\colon E_x \to E_y$ for some $x,y \in M$. Then, $\Phi(E) \rightrightarrows M$ is a frame groupoid with $s(\xi) = x$, $t(\xi) = y$, $u(x) = \mathrm{id}_{E_x}$, the inverse of ξ is just an inverse map of ξ, and multiplication κ is the composition of maps.

Example 2.19 (Jet groupoid)**.** For a local diffeomorphism $\varphi\colon U \to V$ between open sets of a manifold M, and given $x \in U$, let $j^1_x\varphi$ be the one-jet of φ at $x \in U$. Then, $J^1(M,M)$, the set of all such one-jets, has a natural groupoid structure $J^1(M,M) \rightrightarrows M$ with $s(j^1_x\varphi) = x$, $t(j^1_x\varphi) = \varphi(x)$, $\kappa(j^1_{\varphi(x)}\psi, j^1_x\varphi) = j^1_x(\psi \circ \varphi)$, and the inverse is given by $i(j^1_x\varphi) = j^1_{\varphi(x)}\varphi^{-1}$.

Example 2.20 (Gauge groupoid)**.** Let (P,M,G,π) be a principal bundle and let $(u_2,u_1,g) \in P \times P \times G \mapsto (u_2g,u_1g) \in P \times P$. Let $[(u_2,u_1)]$ denote the equivalence class in $(P \times P)/G$. Then,

$$\frac{P \times P}{G} \rightrightarrows M$$

is a Lie groupoid with $s([(u_2,u_1)]) = \pi(u_1)$, $t([(u_2,u_1)]) = \pi(u_2)$, $u(x) = [(v,v)]$ for $v \in \pi^{-1}(x)$, $i([(u_2,u_1)]) = [(u_1,u_2)]$, and

$$\kappa(([(u_3,u_2)]),([(u_2,u_1)])) = [(u_3,u_1)]\,.$$

Example 2.21. Let G be a Lie group with a Lie algebra $\mathfrak{g}$. Then, $\mathrm{T}^*G \rightrightarrows \mathfrak{g}^*$ is a Lie groupoid with $s(\theta) = \theta \circ \mathrm{T}_e L_g$, $t(\theta) = \theta \circ \mathrm{T}_e R_g$ for given $\theta \in \mathrm{T}^*_g G$. Additionally, for $\varphi \in \mathrm{T}^*_h G$, one has

$$\kappa(\varphi, \theta) = \varphi \circ \mathrm{T}_{hg} R_{g^{-1}} = \theta \circ \mathrm{T}_{hg} L_{h^{-1}}$$

and $i(\mu) = \mu \in \mathrm{T}^*_e G$ with $i(\theta) = \theta \circ \mathrm{T}_e L_g \circ \mathrm{T}_{g^{-1}} R_g \in \mathrm{T}^*_{g^{-1}} G$.

Example 2.22. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. Applying tangent functor to each maps yields a Lie groupoid $\mathrm{T}\mathcal{G} \rightrightarrows \mathrm{T}M$.

Morphisms and subgroupoids

Definition 2.23. Let $\mathcal{G} \rightrightarrows M$ and $\mathcal{G}' \rightrightarrows M'$ be groupoids. A morphism of groupoids $(\mathcal{G} \rightrightarrows M) \to \mathcal{G}' \rightrightarrows M'$ is a pair of maps (F, f) such that the following diagrams commute

$$\begin{tikzcd} \mathcal{G} \arrow[r, "F"] \arrow[d, "s"] & \mathcal{G}' \arrow[d, "s'"] \\ M \arrow[r, "f"] & M' \end{tikzcd} \qquad \begin{tikzcd} \mathcal{G} \arrow[r, "f"] \arrow[d, "t"] & \mathcal{G}' \arrow[d, "t'"] \\ M \arrow[r, "f"] & M' \end{tikzcd} \qquad \begin{tikzcd} \mathcal{G} \times \mathcal{G} \arrow[r, "F \times F"] \arrow[d, "\kappa"] & \mathcal{G}' \times \mathcal{G} \arrow[d, "\kappa'"] \\ \mathcal{G} \arrow[r, "F"] & \mathcal{G}' \end{tikzcd} .$$

If $M = M'$ and $f = \mathrm{id}_M$, then F is called a base-preserving morphism, or morphism over M. If $\mathcal{G} \rightrightarrows M$ and $\mathcal{G}' \rightrightarrows M'$ are Lie groupoids, then (F, f) is a Lie groupoid morphism if F and f are smooth.

Remark 2.24. The smoothness of F already implies the smoothness of f. Moreover, The above definition implies that $F(h)F(g)$ is well defined whenever hg is.

Proposition 2.25. Let (F, f) be a groupoid morphism. Then,

(1) $F(1_x) = 1_{f(x)}$ for every $x \in M$,
(2) $F(g^{-1}) = F(g)^{-1}$ for every $g \in \mathcal{G}$.

Definition 2.26. A morphism (F, f) is an isomorphism of Lie groupoids if F and hence f are diffeomorphisms.

Example 2.27. Let $\mathcal{G} \rightrightarrows M$. Then, $\chi := (t, s) \colon \mathcal{G} \times \mathcal{G} \to M \times M \mid g \mapsto (t(g), s(g))$ is a morphism over M from $\mathcal{G} \rightrightarrows M$ and $M \times M \rightrightarrows M$. The map χ is called the anchor of $\mathcal{G}$.

Example 2.28. Let $G \triangleleft \mathfrak{g}^*$, where the action is given by $\mathrm{Ad}^*_{g^{-1}}$. Then, the left-trivialisation of

$$\lambda \colon (g, \nu) \in G \times \mathfrak{g}^* \longmapsto \nu \circ \mathrm{T}_g L_{g^{-1}} ,$$

is an isomorphism over $\mathfrak{g}^*$ of $G \triangleleft \mathfrak{g}^*$ and $\mathrm{T}^*G \rightrightarrows \mathfrak{g}^*$, see Example 2.21.

Example 2.29. For any $\mathcal{G} \rightrightarrows M$, the tangent bundle projections $p_{\mathcal{G}} \colon \mathrm{T}\mathcal{G} \to \mathcal{G}$ and $p_M \colon \mathrm{T}M \to M$ give rise to a Lie groupoid morphism from $\mathrm{T}\mathcal{G} \rightrightarrows \mathrm{T}M$ to $\mathcal{G} \rightrightarrows M$.

Definition 2.30. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. A Lie subgroupoid of $\mathcal{G} \rightrightarrows M$ is a Lie groupoid $\mathcal{G}' \rightrightarrows M'$ with injective immersions $\iota \colon \mathcal{G}' \to \mathcal{G}$ and $\iota_\circ \colon M' \to M$ such that $(\iota, \iota_\circ)$ is a Lie groupoid morphism. A Lie subgroupoid $\mathcal{G}' \rightrightarrows M'$ of $\mathcal{G} \rightrightarrows M$ is embedded if $\mathcal{G}'$ and M' are embedded sumbanifolds of $\mathcal{G}$ and M, respectively. A Lie subgroupoid $\mathcal{G}' \rightrightarrows M'$ of $\mathcal{G} \rightrightarrows M$ is wide if $M = M'$ and $\iota_\circ = \mathrm{id}_M$.

Example 2.31. If $\mathcal{G} \rightrightarrows M$ is a Lie groupoid and $N \subset M$ is an open submanifold, then $\mathcal{G}^N_N := s^{-1}(N) \cap t^{-1}(N)$ is clearly a Lie subgroupoid.

Example 2.32. If $\mathcal{G} \rightrightarrows M$ is a groupoid, the inner (set) subgroupoid is defined as $\mathcal{IG} := \{g \in \mathcal{G} \colon s(g) = t(g)\}$. In general, if $\mathcal{G} \rightrightarrows M$ is a Lie groupoid, this does not define a Lie subgroupoid.

Lecture 3– Rubén Izquierdo-López (ICMAT)

Date: 21 October, 2025

Abstract: In this lecture, I will present locally trivial Lie groupoids, and prove their "correspondence" with gauge groupoids. The lecture essentially follows section 1.3 from Mackenzie's book "General Theory of Lie groupoids and Lie algebroids"

Locally trivial Lie groupoids

Before introducing the definition of locally trivial Lie groupoid, let us recall the notion of transitive groupoid.

Definition 3.1. Let $\mathcal{G} \rightrightarrows M$ be a groupoid (not necessarily Lie groupoid). Then, it is called **transitive** if the map $(t, s)\colon \mathcal{G} \to M \times M$ is surjective.

Hence, a groupoid $\mathcal{G} \rightrightarrows M$ is transitive if for every pair of points $x, y \in M$ there is an element $g \in \mathcal{G}$ such that $s(g) = x$ and $t(g) = y$:

$$\bullet_{y} \xleftarrow{\;g\;} \bullet_{x}$$

Notice that if a **set** groupoid $\mathcal{G} \rightrightarrows M$ is transitive, then it is trivial. Indeed, fix $x \in M$ and choose a right inverse of the restriction of the target map to the s-fiber of x:

$$\alpha\colon M \to \mathcal{G}_x\,, \quad t \circ \alpha = \mathrm{id}_M\ .$$

Then, we can build the following isomorphism of *set* groupoids:

$$\Phi\colon M \times \mathcal{G}_x^x \times M \to \mathcal{G}\,, \quad \Phi(z, g, y) := s(z) \cdot g \cdot s(y)^{-1}\,. \tag{6}$$

$$\begin{tikzcd}
z & & y \arrow[ll, dashed, bend right, "s(z)\cdot g\cdot s(y)^{-1}"'] \\
& x \arrow[ul, bend left, "s(z)"] \arrow[ur, bend right, "s(y)"'] \arrow[loop above, "g"] &
\end{tikzcd}$$

What we are going to study now is how this picture translates into the theory of Lie groupoids. The adapted notion of transitivity to the smooth category is the following:

Definition 3.2. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. It is called **locally trivial** if the map $(t, s)\colon \mathcal{G} \to M \times M$ defines a surjective submersion.

Our first objective is to show that the isomorphism defined in Eq.(6) is well defined *locally*. In order to achieve this, we need to prove the following characterizations of Definition 3.2:

Proposition 3.3. Let $\mathcal{G} \rightrightarrows M$ denote a Lie groupoid and $x \in M$ be a fixed point. Then, the following statements are equivalent:

(1) The Lie groupoid $\mathcal{G} \rightrightarrows M$ is locally trivial.
(2) The restriction of the target map $t\colon \mathcal{G}_x \to M$ defines a surjective submersion.
(3) The map
$$\delta\colon \mathcal{G}_x \times \mathcal{G}_x \to \mathcal{G}\,, \quad \delta(g,h) := h \cdot g^{-1}\,,$$
is a surjective submersion.

Proof. Let us prove the equivalence of the first and last pair of conditions:

- $\boxed{(1) \iff (2)}$ (2) clearly follows from (1). Now, to prove the converse, notice that we may write $(t,s) \circ \delta = (t,t)$. Since, by hypothesis, the map on the left is a surjective submersion, it follows from the chain rule that (t,s) must be a surjective submersion as well, so that the Lie groupoid is locally trivial.
- $\boxed{(2) \iff (3)}$ Assume (2) and define $\mathcal{G}_x \times_M \mathcal{G} = \{(\eta,\xi) \in \mathcal{G}_x \times \mathcal{G} : t(\eta) = t(\xi)\}$, together with the following map:
$$\Psi\colon \mathcal{G}_x \times \mathcal{G}_x \to \mathcal{G}_x \times_M \mathcal{G}\,, \quad \Psi(h,g) := (h, gh^{-1})\,.$$
Using the fact that $t\colon \mathcal{G}_x \to M$ defines a surjective submersion, we have that $\mathcal{G}_x \times_M \mathcal{G}$ is a smooth manifold, and that Ψ defines a diffeomorphism. Hence, to show that δ is a surjective submersion, we only need to show that $\psi^{-1} \circ \delta$ is. A quick computation shows that $\psi^{-1} \circ \delta$ is the projection onto the second factor, which is clearly a surjective submersion. For the converse, it is enough to write $t \circ \delta = t \circ \pi$, where $\pi\colon \mathcal{G}_x \times_M \mathcal{G} \to \mathcal{G}_x$ denotes the projection onto the second factor. Now, the map on the left is clearly a surjective submersion, so $t\colon \mathcal{G}_x \to M$ is a surjective submersion as well.

□

Now we can show that a locally trivial Lie groupoid $\mathcal{G} \rightrightarrows M$ is, in fact, locally trivial. Indeed, choose $x \in M$ and let $\{U_i\}$ be an open cover of M, together with sections of $t\colon \mathcal{G}_x \to M$
$$s_i\colon U_i \to \mathcal{G}_x\,.$$
Now, if we define maps $\Phi\colon U_i \times \mathcal{G}_x^x \times U_i \to \mathcal{G}_{U_i}^{U_i}$ as in Eq.(6) (where the space in the right denotes the restriction groupoid), we have an isomorphism of Lie groupoids, so it is, in fact, locally trivial. The maps s_i are called **decomposing sections** and the collection $\{s_i\}$ is called **decomposing atlas**.

Let us give two examples of locally trivial Lie groupoids:

Example 3.4. Let $E \to M$ be a vector bundle. Then, the frame groupoid $\Phi(E)$, defined as the space of linear isomorphisms $E_x \to E_y$ is locally trivial.

Example 3.5. The gauge groupoid of a principal bundle is locally trivial.

In fact, we claim that every locally trivial groupoid is the gauge groupoid of a principal bundle. Indeed, by our previous discussion, we know that for fixed $x \in M$, the map
$$t\colon \mathcal{G}_x \to M$$

is a fiber bundle with standard fiber $\mathcal{G}_x^x$ (this can be seen taking a decomposing atlas). Furthermore, $\mathcal{G}_x^x$ acts on the right by multiplication:

$$\Phi\colon \mathcal{G}_x \times \mathcal{G}_x^x \to \mathcal{G}_x\,, \quad \Phi(h,g) := h \cdot g\,.$$

This action is free and defines a principal $\mathcal{G}_x^x$-bundle structure on $t\colon \mathcal{G}_x \to M$. We may wonder how does this construction depend on the choice of point $x \in M$. The relation is given as follows:

Proposition 3.6. Let $\mathcal{G} \rightrightarrows M$ be a locally trivial Lie groupoid and let $g \in \mathcal{G}$. Denote $x = s(g)$ and $y = t(g)$. Then, we have the following diffeomorphism

$$\Psi\colon \mathcal{G}_y \to \mathcal{G}_x\,, \quad \Phi(h) := h \cdot g$$

which, together with the "inner" Lie group isomorphism

$$\varphi\colon \mathcal{G}_y^y \to \mathcal{G}_x^x\,, \quad \varphi(k) = g^{-1} \cdot k \cdot g$$

defines an isomorphism of principal bundles over the identity on M.

Proof. Clearly, Ψ defines a diffeomorphism, and φ defines an isomorphism of Lie groups. To check that these two maps define an isomorphism of principal bundles, we need to show that Ψ is equivariant with respect to the action. Indeed, let $h \in \mathcal{G}_y$ and $k \in \mathcal{G}_y^y$. Then,

$$\Psi(h \cdot k) = h \cdot k \cdot g = (h \cdot g) \cdot (g^{-1} \cdot k \cdot g) = \Psi(h) \cdot \varphi(k)\,,$$

which completes the proof. □

Summarizing, to each principal bundle, we can associated its gauge groupoid, which is locally trivial; and to each locally trivial Lie groupoid we may associate a principal bundle (after making a choice $x \in M$). These operations are inverses of each other. However, the isomorphisms depend upon choices, and they cannot be chosen naturally. Nevertheless, the isomorphisms depend functorially:

Proposition 3.7. Let M be a manifold, $\pi\colon P \to M$ be a principal bundle over M, and $\mathcal{G} \rightrightarrows M$ be a locally trivial Lie groupoid.

(1) Given $u_0 \in P$ and denoting $x_0 := \pi(u_0)$, the map

$$P \to ((P \times P)/G)_{x_0}\,, \quad u \mapsto [(u, u_0)]\,,$$

is a diffeomorphism, the map

$$G \to ((P \times P)/G)_{x_0}^{x_0}\,, \quad g \mapsto [(u_0 \cdot g, u_0)]\,,$$

is an isomorphism of Lie groups, and together they define an isomorphism of principal bundles over M. Furthermore, given a morphism of principal bundles $f\colon P \to P'$, where P' is a principal bundle over M', denoting $u_0' := f(u_0)$ and $x_0' := \pi'(u_0')$, we have a commutative diagram:

$$\begin{tikzcd} ((P \times P)/G)_{x_0} \arrow[r] & ((P' \times P')/G')_{x_0'} \\ P \arrow[u] \arrow[r] & P' \arrow[u] \end{tikzcd}\,,$$

where the arrows are defined in the obvious manner.

(2) Given $x \in M$, the map

$$(\mathcal{G}_x \times \mathcal{G}_x)/\mathcal{G}_x^x \to \mathcal{G}\,, \quad [(g,h)] \mapsto gh^{-1}\,,$$

defines an isomorphism of Lie groupoids. Furthermore, given a morphism of Lie groupoids $\varphi\colon \mathcal{G} \to \mathcal{G}'$, where $\mathcal{G}'$ is a locally trivial Lie groupoid over M', if we define $x' := \varphi(x)$, we get a commutative diagram

$$\begin{tikzcd} (\mathcal{G}_x \times \mathcal{G}_x)/\mathcal{G}_x^x \arrow[r] \arrow[d] & ({\mathcal{G}'}_{x'} \times {\mathcal{G}'}_{x'})/\mathcal{G}_{x'}^{x'} \arrow[d] \\ \mathcal{G} \arrow[r] & \mathcal{G}' \end{tikzcd},$$

where the arrows are defined in the obvious manner.

Proof. The proof follows from a straight-forward computation. □

An important consequence of Proposition 3.7 is the following:

Corollary 3.8. Let $\mathcal{G} \rightrightarrows M$ denote a locally trivial Lie groupoid and $\mathcal{G}' \rightrightarrows M'$ denote a Lie groupoid (not necessarily locally trivial). Let $\varphi\colon \mathcal{G} \to \mathcal{G}'$ be a set morphism of groupoids. Then,

(1) Fix $x \in M$. If $\varphi\colon \mathcal{G}_x \to \mathcal{G}_{\varphi(x)}$ is smooth, so is $\varphi\colon \mathcal{G} \to \mathcal{G}'$.
(2) If φ is smooth in a neighborhood of the identity, φ is smooth.

Proof. (1) follows straight-forwardly using Proposition 3.7 (2). (2) follows by checking (1) for a particular $x \in M$, using right translations. □

As we mentioned already, the correspondence should not be thought of as an equivalence, given that it is not natural. However, we may use constructions from the world of principal bundles and carry them over to the world of locally trivial Lie groupoids. The most important of these constructions being the associated bundles. Given a principal G-bundle $\pi\colon P \to M$, and a left action of G on a manifold F, we can build a fiber bundle $P[F] \to M$ with standard fiber F by taking the quotient $P[F] := (P \times F)/G$, where $(p,f) \sim (p \cdot g, g^{-1} \cdot f)$. When G preserves a particular structure on F (group, vector space, hermitian...) the induced bundle $P[F]$ inherits this structure fiber wise. For instance if G acts on itself by conjugation, $P[G]$ is a Lie group bundle. When P is the principal bundle obtained from a locally trivial Lie groupoid $\mathcal{G}_x \to M$, the canonical Lie group bundle may be thought of as the inner groupoid:

Proposition 3.9. Let $\mathcal{G} \rightrightarrows M$ be a locally trivial Lie groupoid, $x \in M$, and $\mathcal{G}_x[\mathcal{G}_x^x]$ denote the canonical Lie group bundle. Then, the map

$$\mathcal{G}_x[\mathcal{G}_x^x] \to \mathcal{G}\,, \quad [(g,h)] \to g \cdot h \cdot g^{-1}$$

takes values in the inner subgroupoid $\mathcal{IG} \subset \mathcal{G}$.

Proof. The proof follows from a straight-forward computation. □

Lecture 4 – Asier López-Gordón (IMPAN)

Date: 4 November, 2025

Bisections

The main reference for this lecture are [**1**, Section 1.4] and [**11, 12**].

Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid with source $s\colon \mathcal{G} \to M$, target $t\colon \mathcal{G} \to M$, unit $u : M \to \mathcal{G} \mid x \mapsto 1_x$ and inverse $i : \mathcal{G} \to \mathcal{G} \mid g \longmapsto g^{-1}$.

On a group $G \ni g$, left-translations $L_g\colon h \mapsto gh$ and inner automorphisms $I_g\colon h \mapsto ghg^{-1}$ each form a group isomorphic to G itself, while right-tranlations $R_g\colon h \mapsto hg$ constitute a group anti-isomorphic to G.

On a groupoid, on the other hand, left-tranlations and right-tranlations only are defined on fibers of the target and source projections, respectively:

$$\begin{aligned} L_g\colon \mathcal{G}^x \ni h \mapsto gh \in \mathcal{G}^y\,, \\ R_g\colon \mathcal{G}_y \ni h \mapsto hg \in \mathcal{G}_x\,, \end{aligned}$$

where $g \in \mathcal{G}$ with $s(g) = x$ and $t(g) = y$. In order to extend these concepts to maps of the whole groupoid, we introduce the notion of bisections.

Definition 4.1. A *bisection* is a smooth section $\sigma\colon M \to \mathcal{G}$ of s (i.e., $s \circ \sigma = \mathrm{Id}_M$) such that $t \circ \sigma\colon M \to M$ is a diffeomorphism. The set of bisections of $\mathcal{G}$ is denoted by $\mathsf{Bis}(\mathcal{G})$.

In other words, a bisection is a smooth map $\sigma\colon M \to \mathcal{G}$ making the following diagram commutative:

$$\begin{tikzcd} & \mathcal{G} \arrow[dl, "t"'] \arrow[dr, "s"] & \\ M \arrow[rr, "t\circ\sigma", "\sim"'] & & M \arrow[ul, bend right, "\sigma"'] \end{tikzcd}$$

where all the maps are smooth.

Proposition 4.2. The set of bisections $\mathsf{Bis}(\mathcal{G})$ is canonically endowed with a group structure, with multiplication

$$\begin{aligned} \mathsf{Bis}(\mathcal{G}) \times \mathsf{Bis}(\mathcal{G}) \ni (\sigma, \tau) \mapsto \sigma \star \tau \in \mathsf{Bis}(\mathcal{G}) \\ \sigma \star \tau(x) := \sigma\Big((t \circ \tau)(x)\Big)\tau(x)\,, \quad \forall\, x \in M\,. \end{aligned} \tag{7}$$

The neutral element is just the unit map of the groupoid, while the inverse element σ^{-1} of σ is determined by the inverse map of the groupoid, namely,

$$\sigma^{-1}(x) \coloneqq i\Big(\sigma\Big((t\circ\sigma)^{-1}(x)\Big)\Big) = \Big(\sigma\Big((t\circ\sigma)^{-1}(x)\Big)\Big)^{-1}.$$

$$\begin{tikzcd}
z = t\circ\sigma(y) \arrow[rr, bend right, "\sigma^{-1}(z)"'] & & y = t\circ\tau(x) \arrow[ll, "\sigma(y)"'] \arrow[rr, bend right, "\tau^{-1}(y)"'] & & x = s\circ\tau(x) \arrow[ll, "\tau(x)"'] \arrow[llll, bend right, "\sigma\star\tau(x)"']
\end{tikzcd}$$

Proof. For any $\sigma,\tau\in\mathsf{Bis}(\mathcal{G})$, we have that $s(\sigma\star\tau)=s(\tau)=\mathrm{Id}_M$ and $t(\sigma\star\tau)=t(\sigma)$, which is a diffeomorphism. Thus, $\sigma\star\tau$ is also a bisection. The fact that $u\colon M\ni x\mapsto 1_x\in\mathcal{G}$ is the identity element is obvious. Finally, for any $x\in M$, we can write

$$\sigma^{-1}\Big((t\circ\sigma)(x)\Big) = \Big(\sigma(x)\Big)^{-1},$$

and thereupon

$$\sigma^{-1}\star\sigma(x) = \Big(\sigma(x)\Big)^{-1}\sigma(x) = 1_x\,.$$

Moreover,

$$\sigma\star\sigma^{-1}(x) = \sigma\Big((t\circ\sigma^{-1})(x)\Big)\sigma^{-1}(x) = \sigma\Big((t\circ\sigma^{-1})(x)\Big)\Big(\sigma\Big((t\circ\sigma)^{-1}(x)\Big)\Big)^{-1}$$

but

$$t\circ\sigma^{-1}(x) = s\circ\Big(\sigma\Big((t\circ\sigma)^{-1}(x)\Big)\Big) = (t\circ\sigma)^{-1}(x)\,,$$

and thus

$$\sigma\star\sigma^{-1}(x) = 1_x\,.$$

□

Remark 4.3. The map

$$\mathsf{Bis}(\mathcal{G})\ni\sigma\mapsto t\circ\sigma\in\mathrm{Diff}(M)$$

is a group homomorphism. Furthermore, any morphism of Lie groupoids over M

$$f\colon\ (\mathcal{G}\rightrightarrows M)\to(\mathcal{H}\rightrightarrows M)$$

induces a morphism

$$\begin{aligned}\mathsf{Bis}(f)\colon \mathsf{Bis}(\mathcal{G})&\to\mathsf{Bis}(\mathcal{H})\\ \sigma&\mapsto f\circ\sigma\,.\end{aligned}$$

Hence, we can regard Bis as a functor from the category of Lie groupoids over M to the category of groups:

$$\mathsf{Bis}\colon\mathsf{LieGroupoids}_M\to\mathsf{Groups}\,.$$

In particular, isomorphic Lie groupoids have isomorphic groups of bisections.

Remark 4.4. In principle, $\mathsf{Bis}(\mathcal{G})$ is only a group. Under certain assumptions on $\mathcal{G}$ and M, it is possible to endow it with a structure of Lie group (see [**11**] and references therein).

Example 4.5 (Lie group)**.** Given a Lie group G, we have the Lie groupoid $G \rightrightarrows \{*\}$. Then, $\mathsf{Bis}(G) \cong G$.

Example 4.6 (Manifold)**.** Given a manifold M, we can define the Lie groupoid $M \rightrightarrows M$ with $s = t = \mathrm{Id}_M$, whose group of bisections is $\mathsf{Bis}(M) = \{\mathrm{Id}_M\}$.

Example 4.7 (Pair groupoid)**.** For $M \times M \to M$, any bisection is of the form

$$\sigma\colon M \times M \to M\,, \quad \sigma(x) = \Big(\varphi(x), x\Big)\,,$$

with $\varphi \in \mathrm{Diff}(M)$. Hence,

$$\mathsf{Bis}(M \times M) \cong \mathrm{Diff}(M)\,.$$

Example 4.8 (Trivial groupoid)**.** Consider the Lie groupoid $M \times G \times M \rightrightarrows M$, for a Lie group G and a manifold M, with $s(y, g, x) = x$, $t(y, g, x) = y$, $u(x) = (x, e, x)$, $i(y, g, x) = (x, g^{-1}, y)$, and

$$\kappa((z, h, y), (y, g, x)) = (y, hg, x)\,.$$

Any bisection $\sigma\colon M \to M \times G \times M$ can be written as

$$\sigma(x) = \Big(\varphi(x), \theta(x), x\Big)\,,$$

where $\varphi \in \mathrm{Diff}(M)$ and $\theta\colon M \to G$ is a smooth map. The multiplication is given by

$$\sigma_1 \star \sigma_2(x) = \Big(\varphi_1 \circ \varphi_2(x), \Big(\theta_1 \circ \varphi_2(x)\Big)\theta_2(x), x\Big)\,, \quad \forall\, x \in M\,,$$

for any pair of bisections $\sigma_i = (\varphi_i, \theta_i, \mathrm{Id}_M) \in \mathsf{Bis}(M \times G \times M)$, $i = 1, 2$. The inverse is given by

$$\sigma^{-1}(x) = \Big(\varphi^{-1}(x), \theta^{-1}\Big(\varphi^{-1}(x)\Big), x\Big)\,, \quad \forall\, x \in M\,.$$

Example 4.9 (Action groupoid)**.** Consider an action Lie groupoid $G \triangleleft M$:

$$\begin{tikzcd}
z = hgx \arrow[rr, bend right, "(h^{-1},z)"'] & & y = gx \arrow[ll, "(h,gx)"'] \arrow[rr, bend right, "(g^{-1},y)"'] & & x \arrow[ll, "(g,x)"'] \arrow[llll, bend right, "(hg,x)"']
\end{tikzcd}$$

A bisection of G is a smooth map

$$\sigma\colon M \to G \times M\,, \quad \sigma(x) = \Big(\rho(x), x\Big)$$

such that $M \ni x \mapsto t \circ \sigma(x) = \rho(x) \cdot x \in M$ is a diffeomorphism.

The Lie group action of G on M is transitive (i.e., for any $x, y \in M$, there exists a $g \in G$ such that $y = gx$) if and only if for each $\varphi \in \mathrm{Diff}(M)$ there exists a bisection σ such that

$$\varphi(x) = t \circ \sigma(x) = \rho(x) \cdot x\,, \quad \forall x \in M\,.$$

Example 4.10 (Frame groupoid)**.** Let (E, q, M) be a vector bundle. Let us recall that the set $\Phi(E)$ of all vector space isomorphisms

$$\xi_x^y\colon E_x \to E_y\,, \quad x, y \in M\,,$$

can be equipped with the structure of a Lie groupoid, the so-called frame groupoid $\Phi(E) \rightrightarrows M$.

$$\begin{tikzcd}
z = t(\chi_y^z) \arrow[bb, bend right, "(\chi_y^z)^{-1}"'] & y = t(\xi_x^y) \arrow[l, "\chi_y^z"'] \arrow[bb, bend right, "(\xi_x^y)^{-1}"'] & x = s(\xi_x^y) \arrow[l, "\xi_x^y"'] \arrow[ll, bend right, "\chi_y^z \cdot \xi_x^y"'] \\
\end{tikzcd}$$

There is a one-to-one correspondence between bisections of $\Phi(E)$ and VB automorphisms of the VB (E, q, M). Indeed, if $\sigma \in \mathsf{Bis}(\Phi(E))$, we can define $\overline{\sigma}\colon E \to E$ by

$$\overline{\sigma}\colon E_x \ni v_x \mapsto \sigma(x)(v_x) = \xi_x^y(v_x) \in E_y\,.$$

Then,

$$\overline{\sigma \star \tau}(v_x) = \sigma \star \tau(x)(v_x) = \xi_y^z \cdot \chi_x^y(v_x) = \overline{\sigma} \circ \overline{\tau}(v_x)\,, \quad \forall v_x \in E_x\,, \quad \forall x \in M\,, \tag{8}$$

that is, the multiplication of bisections corresponds to the composition of automorphisms. The identity bisection (i.e., the unit map of the groupoid) corresponds to the identity automorphism:

$$\overline{u} = \mathrm{Id}_E\ .$$

Therefore,

$$\overline{\sigma^{-1}} = (\overline{\sigma})^{-1}\ .$$

Conversely, given a VB automorphism

$$\begin{tikzcd}
E \arrow[r, "\varphi", "\sim"'] \arrow[d, "q"] & E \arrow[d, "q"] \\
M \arrow[r, "\varphi_0", "\sim"'] & M
\end{tikzcd}$$

we can define a bisection σ by

$$\sigma(x) = \varphi_x\colon E_x \to E_{\varphi(x)}\,, \quad \forall\ \in M\,.$$

The local triviality of the VB guarantees the smoothness of σ.

Each VB automorphism $\overline{\sigma}\colon E \to E$, and thus each bisection $\sigma \in \mathsf{Bis}(\Phi(E))$, induces a map of sections of E:

$$\tilde{\sigma}\colon \Gamma(E) \to \Gamma(E)\,,$$

given by

$$\tilde{\sigma}(\mu)\Big(t \circ \sigma(x)\Big) = \overline{\sigma}_x\Big(\mu(x)\Big) = \sigma(x)\mu(x)\,, \quad \forall \mu \in \Gamma(E)\,, \quad \forall x \in M\,,$$

that is,

$$\tilde{\sigma}(\mu)(x) = \sigma\Big((t \circ \sigma)^{-1}(x)\Big)\mu\Big((t \circ \sigma)^{-1}(x)\Big)\,.$$

It also induces a map of smooth functions

$$\begin{aligned}
\hat{\sigma}\colon \mathscr{C}^\infty(M) &\to \mathscr{C}^\infty(M) \\
f &\mapsto f \circ (t \circ \sigma)^{-1}\,.
\end{aligned}$$

The map of sections inherits the $\mathbb{R}$-linerity from the associated VB automorphism. In particular,

$$\tilde{\sigma}(\mu_1 + \mu_2) = \tilde{\sigma}(\mu_1) + \tilde{\sigma}(\mu_2)\,, \quad \forall\, \mu_1, \mu_2 \in \Gamma(E)\,.$$

By equation (8), the multiplication of bisections corresponds to the composition of the associated maps of sections:

$$\widetilde{\sigma \star \mu} = \tilde{\sigma} \circ \tilde{\mu}\,.$$

Moreover,

$$\begin{aligned}\tilde{\sigma}(f\mu)(x) &= \sigma\Big((t\circ\sigma)^{-1}(x)\Big) f\Big((t\circ\sigma)^{-1}(x)\Big)\mu\Big((t\circ\sigma)^{-1}(x)\Big)\\ &= \hat{\sigma}(f)(x)\,\tilde{\sigma}(\mu)(x)\,, \quad \forall\,\mu\in\Gamma(E)\,, \quad \forall f\in\mathscr{C}^\infty(M) \quad \forall x\in M\,,\end{aligned}$$

i.e.

$$\tilde{\sigma}(f\mu) = \hat{\sigma}(f)\,\tilde{\sigma}(\mu)\,.$$

In particular, the map $\tilde{\sigma}\colon \Gamma(E)\to\Gamma(E)$ is not an automorphism of the $\mathscr{C}^\infty(M)$-module $\Gamma(E)$.

Notation 4.11. The maps $\overline{\sigma}, \hat{\sigma}$ and $\tilde{\sigma}$ are all denoted by $\overline{\sigma}$ in Mackenzie's book.

Example 4.12 (Gauge groupoid)**.** In the previous lecture, we saw an isomorphism of principal bundles over M relating (P, M, G, π) and the gauge groupoid (also known as Atiyah groupoid) $\frac{P\times P}{G} \rightrightarrows M$ (see Proposition 3.7). In particular, given $u_0\in P$ with $x_0 := \pi(u_0)$, the map

$$P \ni u \mapsto \Big[(u, u_0)\Big] \in \frac{P\times P}{G}\Big|_{x_0}$$

is a diffeomorphism. Let $\varphi\colon P\to P$ be a principal bundle automorphism, such that the diagram

$$\begin{tikzcd} P \arrow[r, "\varphi"] \arrow[d, "\pi"] & P \arrow[d, "\pi"] \\ M \arrow[r, "\varphi_0"] & M \end{tikzcd}$$

is commutative, and

$$\varphi(ug) = \varphi(u)g\,.$$

These type of principal bundle automorphisms are called *gauge transformations*. Composing the gauge transformation φ with the principal bundle isomorphism above, we can define an inner automorphism

$$\begin{aligned}\frac{P\times P}{G} &\to \frac{P\times P}{G}\\ [v,u] &\mapsto \Big[\varphi(v), \varphi(u)\Big]\,.\end{aligned}$$

Its associated bisection $\sigma\in\mathsf{Bis}((P\times P)/G)$ is given by

$$\begin{aligned}\sigma\colon M &\to \frac{P\times P}{G}\\ x=\pi(u) &\mapsto \Big[\varphi(u), u\Big]\,.\end{aligned}$$

Moreover, a locally trivial Lie groupoid $\mathcal{G}\rightrightarrows M$ and the gauge groupoid of its vertex bundle at any $x_0\in M$ are isomorphic Lie groupoids over M

$$\frac{\mathcal{G}_{x_0}\times\mathcal{G}_{x_0}}{\mathcal{G}_{x_0}^{x_0}} \underset{\sim}{\longrightarrow} \mathcal{G}\,, \quad [(g,h)]\mapsto gh^{-1}\,,$$

Therefore, their corresponding groups of bisections are isomorphic:

$$\mathsf{Bis}\,(\mathcal{G}) \cong \mathsf{Bis}\left(\frac{\mathcal{G}_{x_0} \times \mathcal{G}_{x_0}}{\mathcal{G}_{x_0}^{x_0}}\right) .$$

Hence, each bisection $\sigma \in \mathsf{Bis}(\mathcal{G})$ can be written as

$$\sigma\colon x = t(g) \mapsto \varphi(g)g^{-1} , \quad \forall g \in \mathcal{G}_{x_0} ,$$

where $\varphi\colon \mathcal{G}_{x_0} \to \mathcal{G}_{x_0}$ is an automorphism of the principal bundle $(\mathcal{G}_{x_0}, M, \mathcal{G}_{x_0}^{x_0}, t)$.

Definition 4.13. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. A bisection $\sigma \in \mathsf{Bis}(\mathcal{G})$ defines three maps $L_\sigma, R_\sigma, I_\sigma\colon \mathcal{G} \to \mathcal{G}$ as follows:

- The *left-translation*
$$L_\sigma(g) = \sigma\Big(t(g)\Big) \cdot g . \tag{9}$$
- The *right-translation*
$$R_\sigma(g) = g \cdot \sigma\Big((t \circ \sigma)^{-1}\Big(s(g)\Big)\Big) . \tag{10}$$
- The *inner automorphism*
$$I_\sigma(g) = \sigma\Big(t(g)\Big) \cdot g \cdot \Big(\sigma\Big(s(g)\Big)\Big)^{-1} . \tag{11}$$

The following properties can be proven straightforwardly:

$$\begin{aligned} L_{\sigma\star\tau} &= L_\sigma \circ L_\tau , \\ R_{\sigma\star\tau} &= R_\tau \circ R_\sigma , \\ I_{\sigma\star\tau} &= I_\sigma \circ I_\tau , \\ I_\sigma &= L_\sigma \circ R_{\sigma^{-1}} = R_{\sigma^{-1}} \circ L_\sigma , \end{aligned}$$

for any $\sigma, \tau \in \mathsf{Bis}(\mathcal{G})$. Therefore, the sets of left-translations, right-translations, and inner automorphisms each form a group under composition; the maps $\sigma \mapsto L_\sigma$ and $\sigma \mapsto I_\sigma$ are group isomorphisms, while $\sigma \mapsto R_\sigma$ is an anti-isomorphism.

Definition 4.14. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid, and let $U \subseteq M$ be an open subset. A *local bisection* is a smooth map $\sigma\colon U \to \mathcal{G}$ such that $s \circ \sigma = \mathrm{Id}_U$ and $t \circ \sigma U \to t \circ \sigma(U)$ is a diffeomorphism. The set of local bisections of $\mathcal{G}$ on U is denoted by $\mathsf{Bis}_U(\mathcal{G})$.

Each local bisection $\sigma \in \mathsf{Bis}_U(\mathcal{G})$ defines:

- The *local left-translation* $L_\sigma\colon \mathcal{G}^U \to \mathcal{G}^V$, given by (9).
- The *local right-translation* $R_\sigma\colon \mathcal{G}_V \to \mathcal{G}_U$, given by (10).
- The *local inner automorphism* $I_\sigma\colon \mathcal{G}_U^U \to \mathcal{G}_V^V$, given by (11).

Proposition 4.15. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. For any $g \in \mathcal{G}$, there exists a sufficiently small open subset $U \subseteq M$ and a local bisection $\sigma \in \mathsf{Bis}_U(\mathcal{G})$ such that $g = \sigma(U)$.

Proof. Let C be a complementary vector subspace to both $\ker \mathsf{T}_g s$ and $\ker \mathsf{T}_g t$ in $\mathsf{T}_g\mathcal{G}$, namely,

$$\mathsf{T}_g\mathcal{G} = \ker \mathsf{T}_g s \oplus C = \ker \mathsf{T}_g t \oplus C\,.$$

Since the source map is a surjective submersion, it has local sections. On a sufficiently small neighbourhood $\tilde{U} \subseteq M$ of $s(g)$, we can choose (e.g. working in local coordinates) a smooth map $\sigma\colon \tilde{U} \to \mathcal{G}$ such that $s \circ \sigma = \mathrm{Id}_{\tilde{U}}$ and

$$\mathsf{T}_{s(g)}\sigma\left(\mathsf{T}_{s(g)}M\right) = C\,.$$

Thus,

$$\mathsf{T}_{s(g)}(t \circ \sigma)\colon V \to \mathsf{T}_{s(g)}(t \circ \sigma)(V)$$

is an isomorphism. Consequently, there exists a sufficiently small neighbourhood $U \subseteq \tilde{U}$ such that $t \circ \sigma|_U$ is a diffeomorphism onto its image. $\square$

Lecture 5 – Juan Manuel López Medel (ICMAT)

Date: 18 November

Abstract: In this session we first record basic facts about local bisections and give an explicit formula for multiplication in the tangent groupoid. We continue with Section 1.5. Components and Transitivity. We show that the identity–component subgroupoid C is generated by symmetric neighbourhoods of the unit section, is open, and yields connected reductions of principal bundles by choosing components of the total space. Finally, orbits are shown to be submanifolds and transitive Lie groupoids are locally trivial.

We first start with two easy corollaries from the last proposition of the previous lecture:

Corollary 5.1. Let $G \rightrightarrows M$ be a Lie groupoid. For each $x \in M$, $t_x : G_x \to M$ is of constant rank.

Proof. Take $g, h \in G_x$. Then $j = gh^{-1}$ is defined and so there is a local bisection $\sigma \in \mathrm{Bis}_U G$ with $t(h) \in U$ and $\sigma(t(h)) = j$.
Now $L_\sigma : G_x^U \to G_x^V$, where $V = (t \circ \sigma)(U)$, maps h to g and $t_x \circ L_\sigma = (t \circ \sigma) \circ t_x$. Hence the ranks of t_x at g and h are equal. □

Corollary 5.2. Let G be a Lie groupoid on M. Then for all $x, y \in M$, G_x^y is a closed embedded submanifold of G_x, of G^y and of G. In particular, each vertex group G_x^x is a Lie group.

Proof. Since $G_x^y = t_x^{-1}(y)$ is the preimage of a point under a map of constant rank, it is a closed embedded submanifold of G_x and hence of G. A similar argument applies for $G_x^y \subseteq G^y$. It follows that $G_x^x \times G_x^x$ is a closed embedded submanifold of $G * G$ and so the restriction of the multiplication in G is smooth. □

The set of all L_σ for $\sigma \in \mathrm{Bis}_U G$ and $U \subseteq M$ open is not a pseudogroup on G since it is not closed under restriction. The following result, which will be needed in Chapter 3, shows that this is unimportant.

Proposition 5.3. Let G be a Lie groupoid on M. Let $\varphi : \mathcal{U} \to \mathcal{V}$ be a diffeomorphism from $\mathcal{U} \subseteq G$ open to $\mathcal{V} \subseteq G$ open, and let $f : U \to V$ be a diffeomorphism from $U = t(\mathcal{U}) \subseteq M$ to $V = t(\mathcal{V}) \subseteq M$, such that $s \circ \varphi = f \circ t$ and $t \circ \varphi = f \circ t$ and $\varphi(gh) = \varphi(g)h$ whenever $(g, h) \in G * G$, $g \in \mathcal{U}$ and $gh \in \mathcal{U}$. Then φ is the restriction to $\mathcal{U}$ of a unique local left–translation $L_\sigma : G^U \to G^V$ where $\sigma \in \mathrm{Bis}_U G$.

Proof. For $x \in U$ choose $g \in \mathcal{U}^x$ and define $\sigma(x) = \varphi(g)g^{-1}$; clearly $\sigma(x)$ is well defined. Since the restriction $t : \mathcal{U} \to U$ is a submersion, σ is smooth and is therefore a local bisection on U with $t \circ \sigma = f$.
That $L_\sigma(g) = \varphi(g)$ for $g \in \mathcal{U}$ is clear, as is the uniqueness. □

Thus any local diffeomorphism $\mathcal{U} \to \mathcal{V}$ which commutes with the $R_h : G_{t(h)} \to G_{s(h)}$ in the sense of last proposition is the restriction of a local left-translation $L_\sigma : G^{t(\mathcal{U})} \to G^{t(\mathcal{V})}$.

In any case, for a general Lie groupoid G we will often treat the set of local bisections as if it were a pseudogroup on M with law of composition $\star$: if $\sigma \in \mathrm{Bis}_U G$ with $(t \circ \sigma)(U) = V$ and $\tau \in \mathrm{Bis}_{V'} G$ with $(t \circ \tau)(V') = W$, then $\tau \star \sigma$ is the local bisection in $(t \circ \sigma)^{-1}(V') \cap V$ defined by

$$(\tau \star \sigma)(x) = \tau((t \circ \sigma)(x))\sigma(x),$$

providing $(t \circ \sigma)^{-1}(V') \cap V$ is not void. We will abusively refer to the set of all local bisections, together with this composition, as the *pseudogroup of local bisections of* G, and will denote it by $\mathrm{Bis}^{\mathrm{loc}}(G)$.
We will often need to consider the effect of base-preserving morphisms on bisections.

Definition 5.4. Let $\varphi : G \to G'$ be a morphism of Lie groupoids over M. Then the maps $\mathrm{Bis}G \to \mathrm{Bis}G'$, $\sigma \mapsto \varphi \circ \sigma$, and $\mathrm{Bis}^{\mathrm{loc}}G \to \mathrm{Bis}^{\mathrm{loc}}G'$, $\sigma \mapsto \varphi \circ \sigma$, are both denoted by $\tilde{\varphi}$ and called *induced morphisms* (of groups, and abusively pseudogroups, respectively).

The following important formula for the multiplication in a tangent groupoid makes essential use of the concept of bisection.

Theorem 5.5. Let $G \rightrightarrows M$ be a Lie groupoid, and denote the multiplication by κ. Let $X \in T_g(G)$ and $Y \in T_h(G)$ have $T(s)(X) = T(t)(Y) = w$. Then

$$X \bullet Y = T(\kappa)(X, Y) = T(L_\sigma)(Y) + T(R_\tau)(X) - T(L_\sigma)T(R_\tau)(T(1)(w)),$$

where σ, τ are any (local) bisections of G for which $\sigma(s(g)) = g$ and $\tau(s(h)) = h$.

Proof. First suppose that $\sigma = 1_m$. Then $X - T(1)T(s)(X)$ is defined and is annulled by $T(s)$. So $(X - T(1)T(s)(X)) \bullet 0_h$ is defined and we can write

$$X \bullet Y = (X - T(1)T(s)(X)) \bullet 0_h + T(1)T(t)(Y) \bullet Y.$$

Using (5), and remembering that $T(1)T(t)(Y)$ is an identity in TG, we have

$$X \bullet Y = T(R_h)(X - T(1)T(s)(X)) + Y. \tag{12}$$

Now consider the general case. We have $L_\sigma(1_{s(g)}) = g$; write $Z = T(L_\sigma^{-1})(X)$. Since L_σ commutes with s, we can apply (12) to get

$$Z \bullet Y = T(R_g)(Z - T(1)T(s)(Z)) + Y.$$

R_g is the restriction to $G_{th} \to G_{s(h)}$ of R_τ. Also, $X \bullet Y = T(L_\sigma)(Z) \bullet Y = T(L_\sigma)(Z \bullet Y)$, since $\kappa \circ (L_\sigma \times \mathrm{id}) = L_\sigma \circ \kappa$.
We obtain the equation of the proposition. □

This gives a complete description of the multiplicative structure of TG. In particular, if $X \in T_g G$ and σ is a (local) bisection with $\sigma(s(g)) = g$, then

$$T(\ell)(X) = T(R_\sigma^{-1})T(1)T(s)(X) + T(L_\sigma^{-1})T(1)T(t)(X) - T(R_\sigma^{-1})T(L_\sigma^{-1})(X).$$

Components and transitivity

We first extend to groupoids two elementary facts about topological groups:

- the component of the identity is a subgroup and
- that subgroup is generated by any neighbourhood of the identity.

Proposition 5.6. Let $G \rightrightarrows M$ be a Lie groupoid. For each $x \in M$ let C_x denote the connectedness component of 1_x in G_x. Then $C = C(G) = \cup_{x\in M} C_x$ is a wide Lie subgroupoid of G, called the identity-component subgroupoid of G.

Proof. By definition C contains each $1_x,\ x \in M$, so it is certainly wide. Take $g \in C^y_x$ and $h \in C^z_y$ and consider $hg = R_g(h) \in C^z_x$. Because $R_g : G_y \to G_z$ is a diffeomorphism, it maps components to components; since $g = R_g(1_y) \in R_g(C_y)$ we have $C_x \cap R_g(C_y) \neq \emptyset$ and therefore $C_x = R_g(C_y)$. Hence $hg \in C_x$. So C is closed under multiplication. Taking $g \in C^y_x$ again, we have $1_y \in R_{g^{-1}}(C_x) \cap C_y$ so $R_{g^{-1}}(C_x) = C_y$ and hence $g^{-1} = 1_x g^{-1} \in C_y$, which proves that C is closed under inversion.
Let N be a tubular neighbourhood for 1_M, and regard N as a vector bundle over M with its zero section identified with $1\colon M \to G$. Then $N \subseteq C$. Now the C_x, $x \in M$, are leaves of the foliation defined by $T^sG = \ker T(s)$ and so C is the union of the integral manifolds of T^sG which meet the open set N. Hence C is open and is therefore a Lie subgroupoid of G. □

It is implicit in this proof that the t-fibres C^y are the identity components of the t-fibres G^y of G, and that, for $g \in G^y_x$, the component of G_x containing g is $C_y g$ and the component of G^y containing g is gC_x. Clearly the various components of any one s–fibre need not be diffeomorphic.
If M is connected, then $C = (\cap_x C_x) \cap 1_M$ is connected, since each $C_x \cap 1_M$ is nonvoid. Conversely, if C is connected then $M = t(C)$ is connected.

Proposition 5.7. Let G be a Lie groupoid on M. Then the identity–component subgroupoid $C = C(G)$ is open in G.

Proof. Let $\varphi\colon \mathbb{R}^p \times \mathbb{R}^q \to \mathcal{U} \subseteq G$ be a distinguished chart for the foliation induced by T^sG, where $\mathcal{U} \cap 1_M \neq \varnothing$ and $\varphi(\{0\} \times \mathbb{R}^q) = \mathcal{U} \cap 1_M$. Then clearly $\mathcal{U} \subseteq C$. Taking the union of such $\mathcal{U}$ we obtain an open neighbourhood of 1_M in G which is contained in C. Now C is the union of those leaves of the foliation which intersect the open neighbourhood and so is itself open. □

Proposition 5.8. Let Ω be a locally trivial Lie groupoid on a connected base M. Then $C = C(\Omega)$ is also locally trivial.

Proof. Let $\{\sigma_i : U_i \to \Omega_{m_i}\}$ be a section–atlas for Ω. We can assume the U_i are connected, so each $\sigma_i(U_i)$ lies in a single component C_i of Ω_{m_i}. Choose any $m_i \in U_i$ and define $\tau_i : U_i \to C_{t(\sigma_i(m_i))}$ by $\tau_i(x) = \sigma_i(x)\sigma_i(m_i)^{-1}$.
Next consider any $t_x : \Omega_x \to M$. This is a surjective submersion and therefore open; hence $t_x(C_x)$ is open in M. Now the $t_x(C_x)$, as x ranges through M, are either disjoint or equal, and therefore are also closed. Since M is connected, we must have $t_x(C_x) = M$ for all $x \in M$. For each i we can therefore choose a $\xi_i \in C_{m_i}$ with $t\xi_i = t\sigma_i(m_i)$. Then $\{\nu_i : U_i \to C_m\}$ defined by $\nu_i(x) = \tau_i(x)\xi_i$ gives a section-atlas for C. □

Of course it is true in general that if a Lie groupoid has a subgroupoid which is locally trivial, then the ambient groupoid itself must be locally trivial. The construction of the identity component subgroupoid thus has a principal bundle analogue.
Consider a principal bundle $P(M, G, \pi)$ on a connected base. Choose any component P_o of P. Define

$$H = \{g \in G \mid P_o g = P_o\}.$$

It is clear that H is an open subgroup of G. By an argument similar to last proposition, $\pi_o = \pi|_{P_o} : P_o \to M$ is surjective; it remains a surjective submersion since P_o is open in P. Further, $P_o \times H \to P_o$ remains a free action, and its orbits are equal to the fibres of π_o, by an argument similar to that in Proposition 5.6.

Definition 5.9. The principal bundle $P_o(M, H, \pi_o)$ is a *connected reduction* of $P(M, G)$.

Example 5.10. Let P be the space $\mathbb{R} \times \mathbb{Z}$ and let G be the discrete space $\mathbb{Z} \times \mathbb{Z}$ with the semidirect product group structure

$$(m_1, n_1)(m_2, n_2) = (m_1 + m_2,\ (-1)^{m_2} n_1 + n_2).$$

Let G act on P to the right by

$$(x, p)(m, n) = (x + m, (-1)^m p + n)$$

and let $\pi : P \to M = \mathbb{S}^1$ be $\pi(x, p) = e^{2\pi i x}$. It is easy to verify that $P(M, G, \pi)$ is a principle bundle.
The component of P through $(0, 0)$ is $P_o = \mathbb{R} \times \{0\}$. Evidently, $H = \mathbb{Z} \times \{0\}$. The bundle $P_o(M, H)$ can be identified with the covering $\mathbb{R}(\mathbb{S}^1, \mathbb{Z})$.
Note that $P(M, G)$ is the pullback of the universal cover $\mathbb{R}^2(K, G)$ of the Klein bottle K along the map $\mathbb{S}^1 = \mathbb{R}/\mathbb{Z} \to K = \mathbb{R}^2/G$ induced by $\mathbb{R} \to \mathbb{R}^2,\ x \mapsto (x, 0)$.

This example illustrates that H need not be normal in G.

Proposition 5.11. Let G be a Lie groupoid on M. Let $\mathcal{U}$ be a symmetric set (that is, $\mathcal{U} \supseteq 1_M$ and $\mathcal{U}^{-1} = \mathcal{U}$) such that each $\mathcal{U}_x$ is open in G_x. Then the subgroupoid H generated by $\mathcal{U}$ has H_x open in G_x for all $x \in M$.

Proof. We prove this by induction.
Since $\mathcal{U}$ is symmetric, H is merely the set of all possible products of elements from $\mathcal{U}$. Choosing $x \in M$, the set of all n-fold products $g_n \cdots g_1$ from $\mathcal{U}$ with $sg_1 = x$ is the union of all $\mathcal{U}_{t(j)} \subseteq G_x$ where j is an $(n-1)$-fold product from $\mathcal{U}$. Since $R_j : G_{t(j)} \to G_x$ is a diffeomorphism, the set of all n–fold products from $\mathcal{U}$ which lie in G_x is open in G_x. Hence H_x is open in G_x. □

Definition 5.12. A set $\mathcal{U}$ satisfying the conditions of the previous proposition, will be called a symmetric s-neighbourhood of 1_M (or, of the base) in G.

Proposition 5.13. Let G be a Lie groupoid on M, and let H be a wide Lie subgroupoid of G. If each H_x is open in G_x, $x \in M$, then each H_x is also closed in G_x, $x \in M$.

Proof. The complement $G_x \setminus H_x$ is the union of all $H_{t(g)}g$ as g ranges over $G_x \setminus H_x$. Since $H_{t(g)}$ is open in $G_{t(g)}$ it follows that $H_{t(g)}g$ is open in G_x. □

The following result is now immediate.

Proposition 5.14. Let G be a Lie groupoid on M, and let $\mathcal{U}$ be a symmetric s-neighbourhood of 1_M in G. Then $\mathcal{U}$ generates the identity-component subgroupoid C of G.

* * * * *

As we know, a groupoid $G \rightrightarrows M$ is transitive if and only if the anchor is surjective. At the other extreme, the following case is often encountered.

Definition 5.15. A groupoid $G \rightrightarrows M$ is *totally intransitive* if the image of $(t, s) : G \to M \times M$ is the base subgroupoid Δ_M of $M \times M$.

For any $G \rightrightarrows M$, the image of the anchor $(t, s) : G \to M \times M$ is an equivalence relation on M. The equivalence class of $x \in M$, denoted $\mathcal{O}_x = \theta_x(G)$, is the *transitivity component* of G through x, or the *transitivity orbit* of G through x. Note that $\mathcal{O}_x = t_x(G_x) = s^x(G^x)$. Except where confusion is possible, we will generally just write *orbit*.

Example 5.16. The transitivity components of a fundamental groupoid $\Pi(M)$ are the connectedness components of M, and the transitivity components of an action groupoid $G < M$ are the orbits of the action. For a surjective submersion $q : M \to Q$, the transitivity components of $R(q) \subseteq M \times M$ are the fibres of q.
A Lie Groupoid Bibundle considered as a Lie groupoid is totally intransitive. Groups are both transitive and totally intransitive.

For a set groupoid $G \rightrightarrows M$, each restriction $G^{\mathcal{O}}$ to an orbit $\mathcal{O}$ is a transitive subgroupoid of G, and G can be regarded as the disjoint union of the transitive subgroupoids $G^{\mathcal{O}}$, as $\mathcal{O}$ runs through the orbits. In the case of Lie groupoids, it is clear that the smooth structure cannot be obtained so simply. A good example to keep in mind throughout this section is the cotangent groupoid example 2.21.

Theorem 5.17. Let G be a Lie groupoid on M. Then for each $x \in M$, the orbit $\mathcal{O}_x = t_x(G_x)$ is a submanifold of M.

Proof. Denote G_x by P and G^x_x by W. Then, for the same reason as in Corollary 5.2, the restriction of the groupoid multiplication to $P \times W \to P$ is a smooth action of a Lie group on a manifold. It is easily seen to be proper: if $K, L \subset P$ are compact then $\{\, w \in W \mid Kw \cap L \neq \varnothing\}$ is the image under $P \times_{t_x} P \to W$, $(h, g) \mapsto h^{-1}g$, of the closed subset $K \times_{t_x} L = (K \times L) \cap (P \times_{t_x} P)$ of the compact set $K \times L$ and is therefore compact. Since the action is also free, it follows that $\{(gw, g) \mid g \in P,\ w \in W\}$ is a closed embedded submanifold of $P \times P$ and so there is a quotient manifold structure on P/W.
Define $i : P/W \to M$ by $i(gW) = t_x(g)$. Then i is smooth and injective. Since $P \to P/W$ is a submersion, $\mathrm{rk}_g\, W(i) = \mathrm{rk}_g(t_x)$ for all $g \in P$, and so i is of constant rank with image $\mathcal{O}_x$. Now an injection of constant rank is an immersion. □

In particular, $G_x(\mathcal{O}_x, G^x_x, t_x)$ is a principal bundle. It is clear that, as sets,

$$\frac{G_x \times G_x}{G^x_x} = G^{\mathcal{O}_x}_{\mathcal{O}_x}.$$

Since t_x is of constant rank it follows, as in Proposition 3.3, that the division map $\delta_x : G_x \times G_x \to G$ is also of constant rank. Hence $\frac{G_x \times G_x}{G^x_x} \to G$ is an injection of constant rank, and therefore an immersion. Thus $G^{\mathcal{O}_x}_{\mathcal{O}_x}$ is a submanifold of G, and we have the following result:

Theorem 5.18. Let G be a Lie groupoid on M and let $\mathcal{O}$ be an orbit of M. Then there is a manifold structure on $G^{\mathcal{O}}$ with respect to which it is a submanifold of G and a Lie groupoid on $\mathcal{O}$. Further, $G^{\mathcal{O}}$ is locally trivial.

Corollary 5.19. A transitive Lie groupoid is locally trivial.

For ease of reference we state explicitly the following results, proved above.

Corollary 5.20. Let G be a Lie groupoid on M, and let $x \in M$. Then $\delta_x : G_x \times G_x \to G$ is of constant rank. Further, G is locally trivial if and only if δ_x is a surjective submersion.

Example 5.21. Let $G \times M \to M$ be a smooth action of a Lie group on a manifold and let $G < M$ be the action groupoid. Applying Corollary 5.1 shows that each evaluation map is of constant rank, applying Theorem 5.17 that the orbits are submanifolds of M, and transitive Lie groupoids are locally trivial, that if G acts transitively, then M is equivariantly diffeomorphic to a homogeneous space. The fact that transitive Lie groupoids are locally trivial, also includes the existence of local cross-sections for closed subgroups of Lie groups.

To avoid misunderstanding, we will refer to 'the orbit foliation of the base' only when the foliation is known to be regular; in general we refer to 'the orbit partition of the base'. We know that any Lie groupoid may be restricted to any open subset of the base, and that a Lie groupoid may be restricted to any of its orbits. The following general result concerning restrictions is easily proved.

Proposition 5.22. Let $G \rightrightarrows M$ be a Lie groupoid and let $\iota : S \to M$ be a submanifold such that $\iota \times \iota : S \times S \to M \times M$ and $(t, s) : G \to M \times M$ are transversal. Then G_S^S with its submanifold structure is a Lie subgroupoid of G.

A submanifold satisfying the condition is said to be transversal to G.

Lecture 6 – Paula Alba San Miguel (UNED)

Date: 25th November 2025

Abstract: In this section we discuss Lie groupoid actions on smooth manifolds, the morphisms between such actions, and their intrinsic definition. We also apply results from previous sections to the study of actions.

Actions

We are interested in defining the action of a Lie groupoid $\mathcal{G} \rightrightarrows M$ on a manifold M', so it generalizes the action of a Lie group on a manifold. The lecture follows section 1.6 from Mackenzie's book "General Theory of Lie groupoids and Lie algebroids".

In order to make use of the groupoid structure of $\mathcal{G} \rightrightarrows M$ is reasonable to expect that not every element of $\mathcal{G}$ will be able to act on any point of M'. Also, if $(h, g) \in \mathcal{G} * \mathcal{G}$, we will ask h to be able to act on gm' for every m' g can act on, and that $(hg)m' = h(gm')$.

Definition 6.1 (Action of a Lie groupoid)**.** Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid and let $f : M' \to M$ be a smooth map. Let $G * M'$ be the pullback manifold:

$$G * M' = \{(g, m') \in G \times M' | s(g) = f(m')\}$$

An *action of* $\mathcal{G}$ on $f : M' \to M$ is a smooth map $\Phi : G * M' \to M'$ that we will note $\Phi(g, m') = gm'$ such that:

(1) $f(gm') = t(g)$, $\forall (g, m') \in G * M'$.
(2) $h(gm') = (hg)m'$, $\forall (h, g) \in \mathcal{G} * \mathcal{G}$ and $m' \in M'$ such that $(g, m'), (h, m') \in G * M'$,
(3) $1_{f(m')}m' = m'$ $\forall m' \in M'$.

For $m' \in M'$, the subset $\mathcal{G}[m'] = \{gm' | g \in \mathcal{G}_{f(m')}\}$ is the orbit of m'.

Remark 6.2. In case f is a surjective submersion, its fibres, that we will denote M'_m, are closed embedded submanifolds of M' and, for each $g \in \mathcal{G}^n_m$, the map $\Phi|_{\{g\} \times M'_{s(g)}}$ is a diffeomorphism from $\{g\} \times M'_{s(g)}$ to $\{g\} \times M'_{t(g)}$.

If that is not the case, the fibres aren't necessarily manifold, so the action need not be smooth when restricted to the fibers.

Remark 6.3. Each isotropy group $\mathcal{G}^x_x$ acts on M'_x, and the action is a Lie group action when it is smooth when restricted to the fibres.

Example 6.4 (Trivial action)**.** For every groupoid $\mathcal{G} \rightrightarrows M$, we can define its action on the trivial bundle $M \times F$ as follows:

$$\begin{aligned} \mathcal{G} * (M \times F) = \bigsqcup_{m \in s(\mathcal{G})} \mathcal{G}_m \times (\{m\} \times F) &\to M \times F \\ (g, s(g), a) &\mapsto (t(g), a) \end{aligned}$$

Example 6.5 (Gauge groupoid acting on its principal bundle)**.** Given a principal bundle $P(M, G, \pi)$, the groupoid $\frac{P \times P}{G} \rightrightarrows M$ acts on $\pi : P \to M$ as follows:

$$\begin{aligned} \frac{P \times P}{G} * P &\to P \\ ([p_1, p_2], p) &\mapsto p_1 \end{aligned}$$

For $s([p_1, p_2]) = \pi(p_2) = p$, and then, there exists $g \in G$ such that $[p_1, p_2] = [p_1 g, p_2 g]$ and $p_2 g = p$.

Example 6.6 (Gauge groupoid acting on an associated fibre bundle)**.** Given a principal bundle $P(M, G, \pi)$, let $\frac{P \times P}{G} \rightrightarrows M$ be its Lie groupoid. Given $\phi : G \times F \to F$ a left action of G on a manifold F, we consider the asociated fiber bundle $\mathtt{p} : \frac{(P \times F)}{G} \to M$. Then, $\frac{P \times P}{G} \rightrightarrows M$ can act on $\frac{(P \times F)}{G}$ through the smooth map $\mathtt{p}$:

$$\begin{aligned} \frac{P \times P}{G} * \frac{P \times F}{G} &\to \frac{P \times F}{G} \\ ([p_1, p_2], \langle p_2, a \rangle) &\mapsto \langle p_1, a \rangle \end{aligned}$$

Notice that $s([p_1, p_2]) = \pi(p_2) = \mathtt{p}(\langle p_2, a \rangle)$ and $t([p_1, p_2]) = \pi(p_1) = \mathtt{p}(\langle p_1, a \rangle)$

In order to gain a better understanding of grouopid actions, we will study those maps which, while transforming the manifold on which the action takes place, remain invariant under the action.

Definition 6.7 (Equivariant map)**.** Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid, and let $\mathcal{G} * M^1 \to M^1$ and $\mathcal{G} * M^2 \to M^2$ be actions of $\mathcal{G}$ on $f_i : M^i \to M$, for $i = 1, 2$. Then, a smooth map $\psi : M^1 \to M^2$ such that $f_2 \circ \psi = f_1$ is said to be $\mathcal{G}$ *equivariant* if $\psi(gm') = g\psi(m')$ for all $(g, m') \in \mathcal{G} * M^1$.

$$\begin{tikzcd} M^1 \arrow[rr, "\psi"] \arrow[dr, "f_1"'] & & M^2 \arrow[dl, "f_2"] \\ & M & \end{tikzcd} \qquad \begin{tikzcd} \mathcal{G} * M^1 \arrow[r, "id \times \psi"] \arrow[d, "\Phi^1"'] & \mathcal{G} * M^2 \arrow[d, "\Phi^2"] \\ M^1 \arrow[r, "\psi"'] & M^2 \end{tikzcd}$$

It is also interesting to study groupoid morphisms that transform the action of one groupoid to the action of the other.

Definition 6.8 (Equivariant groupoid morphism)**.** If $\mathcal{G} \rightrightarrows M$ and $\mathcal{H} \rightrightarrows N$ are Lie groupoids and $\mathtt{p} : M' \to M$ and $\mathtt{q} : N' \to N$ are smooth maps, let $\mathcal{G} * M' \to M'$ and $\mathcal{H} * N' \to N'$ be actions on $\mathtt{p}$ and $\mathtt{q}$. Let $\varphi : \mathcal{G} \to \mathcal{H}$ be a morphism of Lie groupoids over $f : M \to N$ and let $\psi : M' \to N'$ be a smooth map such that $\mathtt{q} \circ \psi = f \circ \mathtt{p}$. Then, ψ is said to be φ *equivariant*

if $\psi(gm') = \varphi(g)\psi(m')$ for all $(g, m') \in \mathcal{G} * M'$.

$$\begin{tikzcd} M' \arrow[r, "\psi"] \arrow[d, "\mathtt{p}"'] & N' \arrow[d, "\mathtt{q}"] \\ M \arrow[r, "f"] & N \end{tikzcd} \qquad \begin{tikzcd} \mathcal{G} * M' \arrow[r, "\varphi\times\psi"] \arrow[d, "\Phi_{\mathcal{G}}"'] & \mathcal{H} * N' \arrow[d, "\Phi_{\mathcal{H}}"] \\ M' \arrow[r, "\psi"'] & N' \end{tikzcd}$$

On lecture 3, we proved that, if $\mathcal{G} \rightrightarrows M$ is locally trivial, it is isomorphic to $\frac{P\times P}{G} \rightrightarrows M$ with $P = \mathcal{G}_x$ and $G = \mathcal{G}_x^x$ with $x \in M$. Let us call that isomorphism φ.

Theorem 6.9. Let $\mathcal{G} \rightrightarrows M$ be a locally trivial groupoid, and let $\mathcal{G} * M' \to M'$ be an action of $\mathcal{G}$ on a surjective submersion $\mathtt{q} : M' \to M$. Then, for any $m \in M$ if we call $P = \mathcal{G}_m$, $G = \mathcal{G}_m^m$ and $F = M'_m$, the fibre bundle $(M', \mathtt{q}, M)$ is diffeomorphic to $(\frac{P\times F}{G}, \mathtt{p}, M)$ through a diffeomorphism of fibrations that is equivariant with respect to the isomorphism of groupoids $\varphi : \frac{P\times P}{G} = \frac{\mathcal{G}_m\times\mathcal{G}_m}{\mathcal{G}_m^m} \to \mathcal{G}$ seen in Lecture 3.

That is, all actions of locally trivial groups are of the type of example 6.6.

Proof. The isomorphism φ is defined as $\varphi([g, h]) = gh^{-1}$, so the map:

$$\begin{aligned} \psi : \frac{P \times F}{G} &\to M' \\ \langle g, a\rangle &\mapsto ga \end{aligned}$$

is equivariant with respect to φ, since $\mathtt{q} \circ \psi(\langle gh, h^{-1}a\rangle) = \mathtt{q}(ga) = t_x(g) = \mathtt{p}(\langle gh, h^{-1}a\rangle)$ and $\psi([g, h] \cdot \langle h, a\rangle) = \psi(\langle g, a\rangle) = ga = (gh^{-1})ha = \varphi([g, h])\psi(\langle h, a\rangle)$.

It only remains to proof that ψ is a diffeomorphism. □

Proposition 6.10. Let $P(M, \pi, G)$ be a principal bundle and let B and B' be two associated fibre bundles corresponding to actions $G \times F \to F$ and $G \times F' \to F'$ of G on manifolds F and F'.

(1) If $f : F \to F'$ is a G-equivariant map, then $\tilde{f} : B \to B' : \langle u, a\rangle \mapsto \langle u, f(a)\rangle$ is a well defined morphism of fibre bundles over M, and is $\frac{P\times P}{M}$-equivariant.
(2) If $\varphi : B \to B'$ is a $\frac{P\times P}{M}$-equivariant morphism of fibre bundles over M, then $\varphi = \tilde{f}$ for some G-equivariant map f.

Proof. (1) is easy to verify

For (2), if φ is a $\frac{P\times P}{M}$-equivariant morphism of fibre bundles, $\varphi([p_1, p_2]\langle p_2, a\rangle) = [p_1, p_2]\varphi(\langle p_2, a\rangle)$. This implies that $\varphi(\langle p, a\rangle) = \langle p, f(a)\rangle$. Using (1), we have proven the result. □

Example 6.11 (Inner automorphism action)**.** Given a locally trivial Lie groupoid $\mathcal{G} \rightrightarrows M$, the inner automorphism action is the map:

$$\begin{aligned} \mathcal{G} * \mathcal{IG} &\to \mathcal{IG} \\ (g, h) &\mapsto ghg^{-1} \end{aligned}$$

If $\mathcal{G}$ corresponds to a principal bundle $P(m, G, \pi)$, Theorem 6 shows that $\mathcal{IG}$ is equivariantly isomorphic as a Lie group bundle to the inner group bundle $\frac{P\times G}{G}$

Given a Lie group action $G \times M \to M$, we constructed the action groupoid $G \times M \rightrightarrows M$. Now we are going to construct the action groupoid of an action $\mathcal{G} * M' \to M'$ of the Lie groupoid $\mathcal{G} \rightrightarrows M$ on $f : M' \to M$.

It will be $\mathcal{G} * M' \rightrightarrows M'$ with $s'(g, m') = m'$, $t'(g, m') = gm'$, inclusion $i(m') = (1_{f(m')}, m')$, multiplication $\kappa((h, n'), (g, m')) = (hg, m')$ and inversion $(g, m')^{-1} = (g^{-1}, gm')$.

Definition 6.12 (Action groupoid)**.** The groupoid we constructed is noted by $\mathcal{G} \measuredangle f$ or $\mathcal{G} \measuredangle M'$ and is the action groupoid associated to the action of $\mathcal{G}$ on $f : M' \to M$

Note that we can defined a canonical morphism beetwen $\mathcal{G} \rightrightarrows M$ and $\mathcal{G} * M' \rightrightarrows M'$:

$$\begin{tikzcd} \mathcal{G} * M' \arrow[r, "\pi_1"] \arrow[d, shift left] \arrow[d, shift right] & \mathcal{G} \arrow[d, shift left] \arrow[d, shift right] \\ M' \arrow[r, "f"] & M' \end{tikzcd}$$

Example 6.13 (Action groupoid of a Lie groupoid acting on itself)**.** Any groupoid $\mathcal{G} \rightrightarrows M$ can act on itself through the target map as follows:

$$\begin{aligned} \mathcal{G} * \mathcal{G} &\to \mathcal{G} \\ (g, h) &\mapsto gh \end{aligned}$$

And, considering its action groupoid, its anchor defines an isomorphism of Lie groupoids with $\mathcal{G} \times_s \mathcal{G} = \{(g, h) | s(g) = s(h)\}$ the subgroupoid of the pair groupoid.

$$\begin{tikzcd} \mathcal{G} * \mathcal{G} \arrow[rr, "(s',t')"] \arrow[dr, "t',s'"'] & & \mathcal{G} \times_s \mathcal{G} \arrow[dl, "\bar{t},\bar{s}"] \\ & \mathcal{G} & \end{tikzcd}$$

Note that $(\bar{t}, \bar{s})(t', s')(g, h) = (\bar{t}, \bar{s})(gh, h) = (t'(g, h), s'(g, h))$.

In particular, for any Lie group $G \rightrightarrows \{x\}$, the action groupoid $G \measuredangle G$ is isomorphic to the pair groupoid $G \times G \rightrightarrows G$.

Actions groupoids can be characterized intrinsically as follows.

Definition 6.14 (Action morphism)**.** Let $\varphi : \mathcal{G}' \to \mathcal{G}$ be a morphism of Lie groupoids over $f : M' \to M$, and let φ' be:

$$\begin{aligned} \varphi' : \mathcal{G}' &\to f'\mathcal{G} = \mathcal{G} * M' \\ g' &\mapsto (\varphi(g'), s'(g')) \end{aligned}$$

Note that, since (φ, f) is a groupoid morphism, $s(\varphi(g') = f(s'(g'))$.

φ is said to be an action morphism if φ' is a diffeomorphism.

Remark 6.15. Given an action $\mathcal{G} * M' \to M'$, the canonical Lie groupoid morphism beetwen $\mathcal{G} * M' \rightrightarrows M'$ and $\mathcal{G} \rightrightarrows M$ is an action morphism.

Since we are calling φ an action groupoid, are there an action associated to it? And, is φ the canonical morphism of it?

Theorem 6.16. Let $\varphi : \mathcal{G}' \to \mathcal{G}$ be an action groupoid over a map $f : M' \to M$. Then $t' \circ (\varphi')^{-1} : f'\mathcal{G} \to M'$ is an action of $\mathcal{G}$ on $f : M' \to M$.

Proof. Since φ' is a diffeomorphism, it maps $\mathcal{G}'_{m'}$ into $\mathcal{G}_{f(m')} \times \{m'\}$, and $\varphi|_{\mathcal{G}'_{m'}} : \mathcal{G}'_{m'} \to \mathcal{G}_{f(m')}$ is a bijection, and $\varphi^{-1}(g) = (\varphi')^{-1}(g, m')$, so:

$$f(gm') = (f \circ t' \circ (\varphi')^{-1})(g, m') = (t \circ \varphi \circ (\varphi')^{-1})(g, m') = t(g)$$

Let us see that $h(gm') = (hg)m'$. If $n' = t' \circ (\varphi')^{-1}(g, m')$, $f(n') = t(g) = s(h)$ and $\varphi^{-1}(h) = (\varphi')^{-1}(h, n')$. Since $s'((\varphi')^{-1}(h, n')) = n' = t'((\varphi')^{-1}(g, m'))$, so:

$$\Big((\varphi')^{-1}(h, n'), (\varphi')^{-1}(g, m')\Big) \in \mathcal{G} * \mathcal{G}$$

And its multiplication is in $\mathcal{G}'_{m'}$, so it is mapped by φ to hg, because it is bijective. $\square$

It is easy to prove that this action and the canonical morphism associated to it are mutual inverses.

Proposition 6.17. (1) Let $\varphi : \mathcal{G}' \to \mathcal{G}'$ be an action morphism, and let $\mathcal{G} * M' \to M'$ be the associated action. Then, $\varphi' : \mathcal{G}' \to \mathcal{G} \measuredangle M'$ is an isomorphism of Lie groupoids over M', and $\pi_1 \circ \varphi' = \varphi$.

(2) Let $\mathcal{G} * M' \to M'$ be an action of a Lie groupoid on a smooth map $f : M' \to M$. Then, the action induced by the action morphism $\varphi' : f'\mathcal{G} \to \mathcal{G}$ is the original action.

Lecture 7 – [Tomasz Sobczak (KMMF-UW)]

Date: 2 December

Abstract: TBA.

Lecture 8 – [Juan Manuel López Medel (ICMAT)]

Date: 9 December

Abstract: In the eight lecture we will mainly talk about linear actions, and the relation with representation. We study a proposition that gives us a lot of examples as Riemannian frame groupoid, complex frame groupoid... Lastly, we show how are the linear actions on the fundamental groupoid

Linear actions and frame groupoids

Definition 8.1. Let (E, q, M) be a v.b. and let $\Omega * E \to E$ be an action of a locally trivial Lie groupoid Ω on q. The action is *linear* if for each $\xi \in \Omega$, the diffeomorphism $E_{s(\xi)} \to E_{t(\xi)}$ is a linear isomorphism.

Definition 8.2. Let Ω be a locally trivial Lie groupoid on M and let (E, q, M) be a v.b. Then a *representation* of Ω on (E, q, M) is a morphism $\rho : \Omega \to \Phi(E)$ of Lie groupoids over M.

Proposition 8.3. Let $\Omega \rightrightarrows M$ be a l.t.L.g. and let (E, q, M) be a v.b. If $\Omega * E \to E$ is a linear action of Ω on E then the associated map $\Omega \to \Phi(E)$, $\xi \mapsto (u \mapsto \xi u)$ is a representation; if $\rho : \Omega \to \Phi(E)$ is a representation then $(\xi, u) \mapsto \rho(\xi)(u)$ is a linear action.

Proof. One direction follows from nothing that $\Phi(E) * E \to E$ is a linear action. For the converse, suppose we have a linear action of $\Omega * E \to E$. It has to be proved that the set map $\Omega \to \Phi(E)$ is smooth. Take a chart $\psi : U \times V \to E_U$ for E and use it to write the map locally as $\Omega_U * (U \times V) \to U \times V$. It then follows easily that the associated map $\Omega_U \to U \times GL(V) \cong \Phi(E)_U$ is smooth. $\square$

Now we study invariant sections.

Definition 8.4. Let $\Omega * M' \to M'$ be an action of a l.t.L.g. Ω on a smooth surjection (M', q, M). Then a section $\mu \in \Gamma M'$ is Ω-invariant if $\xi\mu(s(\xi)) = \mu(t(\xi))$, for all $\xi \in \Omega$. The set of Ω-invariant sections of M' is denoted $(\Gamma M')^\Omega$.

If $\Omega * E \to E$ is a linear action on a vector bundle, then $(\Gamma E)^\Omega$ is an $\mathbb{R}$-vector space with respect to pointwise operations, but not usually a module over the ring of smooth functions on M. A general fibre bundle doesn't have to admit any (global) sections. In particular, a vector bundle equipped with a linear action, $(\Gamma E)^\Omega$ may consist of the zero section alone.

Proposition 8.5. Let $\Omega \rightrightarrows M$ be a locally trivial Lie groupoid and let $\Omega * E \to E$ be a linear action of Ω on a vector bundle (E, q, M). Choose $m \in M$, and write V for E_m and G for Ω^m_m. Then the evaluation map

$$(\Gamma E)^\Omega \to V^G,\ \mu \mapsto \mu(m)$$

is an isomorphism of $\mathbb{R}$-vector spaces.

Proof. The map is clearly injective. Given $v \in V^G$, define μ by $\mu(x) = \xi v$ where ξ is any element of Ω_m^x. Clearly $\mu(x)$ is well defined; μ is smooth because $T_m : \Omega_m \to M$ is a surjective submersion. $\square$

Example 8.6. Consider the principle bundle $SO(2)(S^1, \mathbb{Z}_2, p)$ where $\mathbb{Z}_2$ is embedded in $SO(2)$ as $\{1, -1\}$ and p is $z \mapsto z^2$. Let Ω be the gauge groupoid and let E be the vector bundle $SO(2) \times \mathbb{R}/\mathbb{Z}_2$, where $\mathbb{Z}_2$ acts on $\mathbb{R}$ by multiplication; this is the Möbius band. Then $(\Gamma E)^\Omega \cong \mathbb{R}^{\mathbb{Z}_2}$ is the zero space.

Proposition 8.7. Let $\Omega * E \to E$ be a linear action of a locally trivial Lie groupoid Ω on a v.b. E. For each $x \in M$, define $E^{\mathcal{J}\Omega}|_x$ to be

$$E_x^{\Omega_x^x} = \{u \in E_x;\ \lambda u = u,\ \forall \lambda \in \Omega_x^x\}.$$

Then $E^{\mathcal{J}\Omega}$ is a subvector bundle of E, and there is a natural trivialization

$$M \times V^G \to E^{\mathcal{J}\Omega}.$$

Proof. Let $\{\omega_i : U_i \to \Omega_m\}$ be a section-atlas for Ω, and write $V = E_m,\ G = \Omega_m^m$. Define $\psi_i : U_i \times V^G \to E^{\mathcal{J}\Omega}|_{U_i}$ by $(x, v) \mapsto \omega_i(x)(v)$. Then $\psi_{i,x}$ maps V^G isomorphically onto $E^{\mathcal{J}\Omega}|_x$.
To define the trivialization, note that for $x \in M$ and any two $\xi, \xi' \in \Omega_m^x$, the maps $V^G \to E^{\mathcal{J}\Omega}|_x,\ v \mapsto \xi v$ and $v \mapsto \xi' v$ are identical. $\square$

Proposition 8.8. Let (E, q, M) be a vector bundle and $n \geq 1$.

(i) The action $\Phi(E) * \mathrm{Hom}^n(E; M \times \mathbb{R}) \to \mathrm{Hom}^n(E; M \times \mathbb{R})$ defined by $\xi\varphi = \varphi \circ (\xi^{-1})^n$ is smooth.
(ii) The action $\Phi(E) * \mathrm{Hom}^n(E; E) \to \mathrm{Hom}^n(E; E)$ defined by $\xi\varphi = \xi \circ \varphi \circ (\xi^{-1})^n$ is smooth.
(iii) Let (E', p', M) be another v.b. on the same base. Then the action

$$(\Phi(E) \times_{M \times M} \Phi(E')) * \mathrm{Hom}(E; E') \to \mathrm{Hom}(E; E')$$

defined by $(\xi, \xi')\varphi = \xi' \circ \varphi \circ \xi^{-1}$ is smooth.

In (iii) the product is the product of locally trivial Lie groupoids over a fixed base in the sense of the following result, which is easily proved.

Proposition 8.9. Let Ω and Ω' be l.t.L.g. over the same base M. Let $\Omega \times_{M \times M} \Omega'$ be the pullback manifold of the two anchors $(t, s) : \Omega \to M \times M$ and $(t', s') : \Omega' \to M \times M$. Then, with the componentwise algebraic structure, $\Omega \times_{M \times M} \Omega'$ is a locally trivial Lie groupoid on M and the two projections $\Omega \times_{M \times M} \Omega' \to \Omega$ and $\Omega \times_{M \times M} \Omega' \to \Omega'$ are surjective submersions.

The Lie groupoid structure on $\Omega \times_{M \times M} \Omega'$ can be defined for weaker local triviality conditions, but the full properties of a product in the categorical sense over the fixed base M will be lost.
Remark 8.10. $\mathrm{Hom}^n(E; E')$ denotes the vector bundle on M whose fibre over $x \in M$ is the space of n-multilinear maps $E_x \times \cdots \times E_x \to E'_x$ and whose bundle structure is induced from the bundle structure of E and E' in the usual way. The actions (i) and (ii) restrict to the subbundles $\mathrm{Alt}^n(E; E')$ and $\mathrm{Sym}^n(E; E')$ of alternating and symmetric multilinear maps; further, $\mathrm{Hom}(E; E')$ in (iii) could be replaced by $\mathrm{Hom}^n(E; E')$ and the obvious action.

Lastly, there are analogous actions of $\Phi(E)$ on the tensor bundles $\otimes^r_s E$.
Before we start with the examples, we recall a past Theorem:

Theorem 8.11. Let Ω be a l.t.L.g. on M and let $\Omega * M' \to M'$ be a smooth action of Ω on the fibre bundle (M', q, M). Let $\mu \in \Gamma M'$ be an Ω-deformable section. Then the stabilizer groupoid $\Omega\{\mu\}$ of Ω at μ is a closed embedded reduction of Ω.

Example 8.12. Let (E, q, M) be a vector bundle, and let $\langle , \rangle$ be a Riemannian structure in E, regarded as a section of $\mathrm{Hom}^2(E; M \times \mathbb{R})$. Then $\langle , \rangle$ is $\Phi(E)$-deformable with respect to the action of (8.8)(i), since any two vector spaces of the same dimension with any positive-definite inner products, are isometric. Denote the stabilizer groupoid of $\langle , \rangle$ by $\Phi_{\mathcal{O}}(E)$. By Theorem (8.11), it is a locally trivial Lie groupoid on M, the *Riemannian frame groupoid* or *orthonormal frame groupoid* of $(E, \langle , \rangle)$. A section $\sigma : U \to \Phi_{\mathcal{O}}(E)_m$ of $\Phi_{\mathcal{O}}(E)$ is a moving frame for E, and the local triviality of $\Phi_{\mathcal{O}}(E)$ is equivalent to the existence of moving frames in E.

Example 8.13. A *complex structure* in a vector bundle (E, q, M) is an endomorphism $J : E \to E$ such that $J^2 = -id$. Such a J has constant rank and so, regarded as a $(1, 1)$ tensor field, J is $\Phi(E)$-deformable with respect to the action of (8.8)(ii). Denote the stabilizer groupoid of J by $\Phi_{\mathbb{C}}(E)$. By Theorem (8.11) it is a locally trivial Lie groupoid on M, the *complex frame groupoid* of (E, J).
If E further has a Hermitian metric $\langle , \rangle$ for the given J, then the argument of the previous example applies, and there is a l.t.L.g. $\Phi_U(E) \leq \Phi_{\mathbb{C}}(E)$ of Hermitian frames. $\Phi_U(E)$ is the *Hermitian frame groupoid* or *unitary frame groupoid* of $(E, J, \langle , \rangle)$.

Example 8.14. Similarly, if Δ is a determinant function in a vector bundle (E, q, M), regarded as a never-zero section of $\mathrm{Alt}^r(E; M \times \mathbb{R})$ where $r = \mathrm{rank} E$, then Δ is $\Phi(E)$-deformable and the groupoid of orientation-preserving isomorphisms between the fibres of E is a closed embedded reduction of $\Phi(E)$, denoted $\Phi^+(E)$.

Example 8.15. Let (L, q, M) be a v.b. and let $[,]$ be a section of $\mathrm{Alt}^2(L; L)$ such that each

$$[,]_x : L_x \times L_x \to L_x$$

is a Lie algebra bracket. We call such a section a *field of Lie algebra brackets* in L.

A field of Lie algebra brackets need not be $\Phi(L)$-deformable. For example, let $\mathfrak{g}$ be a non-abelian Lie algebra with bracket $[,]$ and in $L = \mathbb{R} \times \mathfrak{g}$ define $[\ ,\ ]_t = t[\ ,\ ]$. However Theorem (8.11) implies that if $[\ ,\ ]$ is a field of Lie algebra brackets in a vector bundle L and if the fibres of L are pairwise isomorphic as Lie algebras, then L admits an atlas of charts which fibrewise are Lie algebra isomorphisms. In this case, $(L, [\ ,\ ])$ is a Lie algebra bundle and we denote the stabilizer groupoid by $\Phi_{\mathrm{Aut}}(L)$.

Example 8.16. Let μ be a section of a vector bundle E on a connected base M. Then, by a similar argument, E has an atlas of charts $U \times V \to E_U$ such that the local representatives $U \to V$ of μ are constant, if and only if μ is either never zero or always zero. This result is well known in the case of tangent vector fields.

A similar treatment may be applied to any tensor structure on a vector bundle.
It is worth noting that in the previous two examples, the condition of pairwise isomorphism, or of being never zero or always zero, need hold only on each component of the base separately. A similar comment applies to the next example.

Example 8.17. Let (E^ν, q_ν, M), $\nu = 1, 2$, be vector bundles on base M, and let $\varphi : E^1 \to E^2$ be a morphism, considered as a section of $\mathrm{Hom}(E^1; E^2)$. Then φ is $\Phi(E^1) \times_{M\times M} \Phi(E^2)$-deformable if and only if it is of constant rank. Now Theorem (8.11) shows that if this is the case, there are atlases $\{\psi_i^\nu : U_i \times V^\nu \to E^\nu_{U_i}\}$, $\nu = 1, 2$, and a linear map $F : V^1 \to V^2$ such that each $\varphi : E^1_{U_i} \to E^2_{U_i}$ is represented by $(x, v) \to (x, f(v))$.

This is a vector bundle version of the standard characterization of a subimmersion (though in order to apply to general subimmersions $M \to N$ it must be extended to varying base manifolds). This result will be useful in the abstract theory of transitive Lie algebroids.

Actions of Lie groupoids on bundles where the fibres have algebraic structures of other types can handled in a similar way. We briefly consider the case of actions on Lie group bundles.

Consider an LGB (K, q, M). An action of a locally trivial Lie groupoid $\Omega \rightrightarrows M$ on K is said to be an *action by Lie group isomorhisms* if each isomorphism $\xi : K_x \to K_y$, $\xi \in \Omega$, is a Lie group isomorphism.
Let $\Phi(K)$ denote the groupoid of all Lie group isomorphisms between the fibres of K. Using the standard Lie group structure on the automorphism group of the fibre type of K, $\Phi(K)$ has a l.t.L.g. structure, and actions of a Lie groupoid Ω on K by Lie group isomorphisms can be identified with Lie groupoid morphisms $\Omega \to \Phi(K)$.

Definition 8.18. Let Ω be a locally trivial Lie groupoid on M and let (K, q, M) be an LGB. An *extension* of Ω by K is a sequence

$$K \overset{\iota}{\rightarrowtail} \Psi \overset{\pi}{\twoheadrightarrow} \Omega$$

in which Ψ is a locally trivial Lie groupoid on M, ι and π are groupoid morphisms over M, ι is an embedding, π is a surjective submersion, and $\mathrm{im}(\iota) = \ker(\pi)$.

It is easy to see that the condition that Ψ be locally trivial is superfluous.

Example 8.19. Let $K \overset{\iota}{\rightarrowtail} \Psi \overset{\pi}{\twoheadrightarrow} \Omega$ be an extension as in the definition with K an abelian LGB. For $\xi \in \Omega^y_x$, $x, y \in M$, choose $\xi' \in \Psi^y_x$ with $\pi(\xi') = \xi$ and define $\rho(\xi) : K_x \to K_y$ as $\lambda \mapsto \xi'\lambda\xi'^{-1}$, the restriction of $I_{\xi'}$. It is clear that $\rho(\xi)$ is well defined. Now $I : \Psi \to \Phi(K)$ is smooth and π is a surjective submersion so $\rho : \Omega \to \Phi(K)$ is smooth. This ρ is the representation associated to the extension $K \rightarrowtail \Psi \twoheadrightarrow \Omega$.

Example 8.20. We consider the fundamental Lie Groupoid:

$$\Pi_1(M) := \{[\gamma]; \gamma : [0, 1] \to M\}.$$

The linear action $\rho([\gamma]) : E_x \to E_y$ must be a linear isomorphism, depends only on the path, and verify:

- $\rho([\gamma_2 * \gamma_1]) = \rho([\gamma_2]) \circ \rho([\gamma_1])$,
- $\rho([c_x]) = id_{E_x}$ for a constant path c_x.

We need E to have a flat connection, an horizontal parralel transport,

$$\tau_\gamma : E_{\gamma(0)} \to E_{\gamma(1)},$$

which depends only on the class of γ.
So $\rho([\gamma]) := \tau_\gamma$ defines a linear action on $\Pi_1(M)$. The linear action of $\Pi_1(M)$ over the fibre bundle $E \to M$ is given by

$$[\gamma] : x \to y \mapsto \rho([\gamma]) : E_x \to E_y.$$

Lecture 9 – O. Carballal (UCM)

Date: 16 December, 2025

Abstract: In this talk, I will present several constructions in the category of Lie groupoids, including pullbacks, fibered products, and homotopy pullbacks. Then, some notions of equivalence between Lie groupoids will be discussed, focusing in particular on Morita equivalence, which plays a central rôle in the modern approach to differentiable stacks, foliation theory, and geometric quantisation. Several of the constructions and equivalence notions will be illustrated through examples coming from the theory of foliations.

Let us start discussing the problematic of arbitrary pull-backs in the category of smooth manifolds.

Example 9.1. Let $f, g : \mathbb{R} \to \mathbb{R}^2$ be the smooth maps defined by $f(t) := (t, 0)$ and $g(t) := (t, t^3)$. We clearly have that

Lecture 10 – Asier López-Gordón (IMPAN)

Date: 13 January, 2026

Abstract: In this talk, I will present the notion of Lie algebroid and illustrate it with several examples.

The main reference for this lecture is [**2**].

Let me start by recalling how the Lie algebra of a Lie group is defined. Let G be a Lie group, and let $\mathfrak{X}(G)$ denote the set of vector fields on G. The subset of left-invariant vector fields on G is given by

$$\mathfrak{X}_L(G) \coloneqq \{X \in \mathfrak{X}(G) \mid (L_g)_* X = X\,, \quad \forall\, g \in G\}\ .$$

Note that

$$(L_g)_*[X,Y] = [(L_g)_*X, (L_g)_*Y]\ , \quad \forall\, g \in G\,, \quad \forall X, Y \in \mathfrak{X}(G)\,.$$

Thus, $X_L(G)$ is a Lie subalgebra of $X(G)$ with respect to the Lie bracket $[\cdot,\cdot]$ of vector fields on G, namely,

$$X, Y \in \mathfrak{X}_L(G) \Longrightarrow [X,Y] \in \mathfrak{X}_L(G)\,.$$

Moreover, as a real vector space, $\mathfrak{X}_L(G)$ is isomorphic to the tangent space $\mathsf{T}_e G$ at the identity. More precisely, we can define the isomorphism

$$\begin{aligned} \Psi\colon \mathsf{T}_e G \ni v &\mapsto \overleftarrow{v} \in \mathfrak{X}_L(G) \\ \overleftarrow{v}_g &\coloneqq \mathsf{T}_e L_g(v)\,, \end{aligned}$$

with inverse

$$\Psi^{-1}(\overleftarrow{v}) = \overleftarrow{v}_e\,.$$

Indeed, for every $g, h \in G$, we have that

$$\Big((L_g)_* \overleftarrow{v}\Big)_h = \mathsf{T}_{g^{-1}\cdot h} L_g(\overleftarrow{v}_{g^{-1}\cdot h}) = \mathsf{T}_{g^{-1}\cdot h} L_g \circ \mathsf{T}_e L_{g^{-1}\cdot h}(v) = \mathsf{T}_e L_h(v) = \overleftarrow{v}_h\,.$$

Definition 10.1. The Lie algebra of a Lie group G is the vector space $\mathfrak{g} \coloneqq \mathrm{Lie}(G)$ endowed with the Lie bracket

$$\begin{aligned} [\cdot,\cdot]\colon \mathfrak{g} \times \mathfrak{g} &\to \mathfrak{g} \\ [v,w] &\coloneqq [\overleftarrow{v}\,, \overleftarrow{w}]|_e = [\Psi(v), \Psi(w)]|_e\,. \end{aligned}$$

We would now like to extend this notion to Lie groupoids.

Definition 10.2. Let $\mathcal{G} \rightrightarrows M$ be a Lie groupoid. A *left-invariant vector field* on $\mathcal{G}$ is a vector field $X \in \mathfrak{X}(\mathcal{G})$ such that:

(1) X is tangent to the t-fibers, i.e. $\mathsf{T}_g t(X_g) = 0$ for all $g \in \mathcal{G}$,

(2) $\mathsf{T}_h L_g(X_h) = X_{gh}$ for every $(g,h) \in \mathcal{G}^{(2)}$.

We shall denote the set of left-invariant vector fields by $\mathfrak{X}_L(\mathcal{G})$.

Lemma 10.3. The set of left-invariant vector fields is closed under the Lie bracket of vector fields, i.e., $\mathfrak{X}_L(\mathcal{G})$ is a Lie subalgebra of $(\mathfrak{X}(\mathcal{G}), [\cdot,\cdot])$.

Proof. A vector field $X \in \mathfrak{X}(\mathcal{G})$ is tangent to the t-fibers if and only if

$$X(f \circ t) = 0\,, \quad \forall f \in \mathscr{C}^\infty(\mathcal{G})\,.$$

Hence, for each pair of vector fields $X, Y \in \mathfrak{X}_L(\mathcal{G})$ we have that

$$[X,Y](f \circ t) = X \circ Y \circ f \circ t - Y \circ X \circ f \circ t = 0\,, \quad \forall f \in \mathscr{C}^\infty(\mathcal{G})\,,$$

which implies that $[X,Y] \in \mathfrak{X}_L(\mathcal{G})$.

For each fixed $g \in \mathcal{G}$, with $s(g) = x$ and $t(g) = y$, the left translation

$$L_g \colon t^{-1}(x) \to t^{-1}(y)$$

is a diffeomorphism. For each $X, Y \in \mathfrak{X}_L(\mathcal{G})$, we have that

$$(L_g)_* X|_{t^{-1}(x)} = X|_{t^{-1}(y)}\,, \quad (L_g)_* Y|_{t^{-1}(x)} = Y|_{t^{-1}(y)}\,,$$

and thereupon

$$\begin{aligned}(L_g)_*[X,Y]|_{t^{-1}(x)} &= (L_g)_* \left[X|_{t^{-1}(x)}, Y|_{t^{-1}(x)}\right] = \left[(L_g)_* X|_{t^{-1}(x)}, (L_g)_* Y|_{t^{-1}(x)}\right] \\ &= \left[X|_{t^{-1}(y)}, Y|_{t^{-1}(y)}\right] = [X,Y]|_{t^{-1}(y)}\,.\end{aligned}$$

Hence,

$$\mathsf{T}_h L_g [X,Y]_h = [X,Y] gh\,, \quad \forall g,h \in \mathcal{G}^{(2)}\,.$$

□

We define the following vector bundle over M:

$$\mathrm{A}(\mathcal{G}) \coloneqq \ker \mathsf{T}t|_{u(M)} = u^*(\ker \mathsf{T}t)\,,$$

and denote its space of sections by $\Gamma(\mathrm{A}(\mathcal{G}))$.

Lemma 10.4. As vector spaces, $\Gamma(\mathrm{A}(\mathcal{G}))$ and $\mathfrak{X}_L(\mathcal{G})$ are isomorphic.

Proof. The isomorphism $\varphi \colon \Gamma(\mathrm{A}(\mathcal{G})) \to \mathfrak{X}_L(\mathcal{G})$ is given by

$$\varphi(\alpha) = \overleftarrow{\alpha}\,, \quad \overleftarrow{\alpha}_g \coloneqq \mathsf{T}_e L_g(\alpha_{s(g)})\,, \quad \forall\, g \in \mathcal{G}\,,$$

and its inverse reads

$$\varphi^{-1}(X) = X|_{u(M)}\,.$$

Clearly, $\alpha \in \ker \mathsf{T}t|_{u(M)}$ is equivalent to $\overleftarrow{\alpha}$ being tangent to the t-fibers. Moreover,

$$\mathsf{T}_h L_g(\overleftarrow{\alpha}_h) = T_h L_g \circ \mathsf{T}_e L_h(\alpha_{s(h)}) = \mathsf{T}_e L_{gh}(\alpha_{s(h)}) = \mathsf{T}_e L_{gh}(\alpha_{s(gh)}) = \overleftarrow{\alpha}_{gh}\,.$$

□

The *anchor* is the vector bundle map $\rho \colon \mathrm{A}(\mathcal{G}) \to \mathsf{T}M$ given by

$$\rho(\alpha) = \mathsf{T}_{1_x} s(\alpha)\,, \quad \forall\, \alpha \in \mathrm{A}(\mathcal{G})_x = \ker(\mathsf{T}_{1_x} t)\,, \quad \forall\, x \in M\,,$$

Lemma 10.5. The bracket of sections $\Gamma(\mathrm{A}(\mathcal{G}))$ satisfies the Leibniz identity

$$[\alpha, f\beta] = f[\alpha,\beta] + \rho(\alpha)(f)\beta\,, \quad \forall\, \alpha,\beta \in \Gamma(\mathrm{A}(\mathcal{G}))\,, \quad \forall\, f \in \mathscr{C}^\infty(M)\,.$$

Proof. For each $\alpha \in \mathrm{A}(\mathcal{G})_x$, we have that

$$s_*(\overleftarrow{\alpha}) = s_*\Big(\varphi^{-1}(\alpha)\Big) = \mathsf{T}_{1_x} s(\alpha) = \rho(\alpha)\,,$$

i.e. $\overleftarrow{\alpha}$ and $\rho(\alpha)$ are s-related. In addition, for each $f \in \mathscr{C}^\infty(M)$ and for all $g \in \mathcal{G}$

$$\overleftarrow{f\alpha}_g = \varphi(f\alpha)_g = \mathsf{T}_e L_g\Big(f(x)\,\alpha(x)\Big) = f(x)\,\mathsf{T}_e L_g\Big(\alpha(x)\Big) = f(x)\overleftarrow{\alpha}_g = \Big((f\circ s)\overleftarrow{\alpha}\Big)(g)\,,$$

where $x = s(g)$, i.e.,

$$\overleftarrow{f\alpha} = s^*(f)\,\overleftarrow{\alpha}\,.$$

Hence,

$$\begin{aligned}\overleftarrow{[\alpha, f\beta]} &:= \Big[\overleftarrow{\alpha}\,, \overleftarrow{f\beta}\Big] = \Big[\overleftarrow{\alpha}\,, s^*f\,\overleftarrow{\beta}\Big] = s^*(f)\,\Big[\overleftarrow{\alpha}\,, \overleftarrow{\beta}\Big] + \overleftarrow{\alpha}\Big(s^*(f)\Big)\overleftarrow{\beta} \\ &= \overleftarrow{f[\alpha,\beta]} + \overleftarrow{\rho(\alpha)(f)\beta} = \overleftarrow{f[\alpha,\beta] + \rho(\alpha)(f)\beta}\,,\end{aligned}$$

where we have used the standard Leibniz rule for the Lie bracket of vector fields on the third step. Consequently,

$$[\alpha, f\beta] = \varphi\Big(\overleftarrow{[\alpha, f\beta]}\Big) = \varphi\Big(\overleftarrow{f[\alpha,\beta] + \rho(\alpha)(f)\beta}\Big) = f[\alpha,\beta] + \rho(\alpha)(f)\beta\,.$$

□

Definition 10.6. A *Lie algebroid* is a vector bundle $A \to M$ endowed with:

- a Lie bracket $[\cdot,\cdot]_A$ on $\Gamma(A)$,
- a vector bundle morphism

$$\begin{tikzcd} A \arrow[rr, "\rho"] \arrow[dr] & & \mathsf{T}M \arrow[dl] \\ & M & \end{tikzcd}$$

satisfying the Leibniz identity:

$$[\alpha, f\beta]_A = f[\alpha,\beta]_A + \rho(\alpha)(f)\beta\,, \quad \forall\,\alpha,\beta \in \Gamma(A)\,, \quad \forall\, f \in \mathscr{C}^\infty(M)\,.$$

The map ρ is called the *anchor* of A.

Given a Lie groupoid $\mathcal{G} \to M$, we call $\mathrm{A}(\mathcal{G})$ the *Lie algebroid of* $\mathcal{G}$. A Lie algebroid A is called integrable if $A \simeq \mathrm{A}(\mathcal{G})$ for some Lie group.

Proposition 10.7. The map $\rho\colon \Gamma(A) \to \mathfrak{X}(M)$ induced by the anchor preserves the brackets, namely,

$$\rho\Big([\alpha,\beta]_A\Big) = [\rho(\alpha), \rho(\beta)]\,, \quad \forall\,\alpha,\beta \in \Gamma(A)\,.$$

Proof. Let $\alpha, \beta, \gamma \in \Gamma(A)$ and $f \in C^\infty(M)$. Applying the Leibniz rule twice, we have that

$$\begin{aligned}[\alpha, [\beta, f\gamma]_A]_A &= [\alpha, f[\beta,\gamma]_A + (\rho(\beta)f)\gamma]_A \\ &= [\alpha, f[\beta,\gamma]_A]_A + [\alpha, (\rho(\beta)f)\gamma]_A \\ &= f[\alpha,[\beta,\gamma]_A]_A + (\rho(\alpha)f)[\beta,\gamma]_A + (\rho(\beta)f)[\alpha,\gamma]_A + (\rho(\alpha)\rho(\beta)f)\gamma\,,\end{aligned}$$

and analogously

$$[\beta, [\alpha, f\gamma]_A]_A = f[\beta,[\alpha,\gamma]_A]_A + (\rho(\beta)f)[\alpha,\gamma]_A + (\rho(\alpha)f)[\beta,\gamma]_A + (\rho(\beta)\rho(\alpha)f)\gamma$$

By the Jacobi identity,

$$\begin{aligned}[[\alpha,\beta]_A, f\gamma]_A &= [\alpha,[\beta,f\gamma]_A]_A - [\beta,[\alpha,f\gamma]_A]_A \\ &= f\left([\alpha,[\beta,\gamma]_A]_A - [\beta,[\alpha,\gamma]_A]_A\right) + \left(\rho(\alpha)\rho(\beta)f - \rho(\beta)\rho(\alpha)f\right)\gamma \\ &= f[[\alpha,\beta]_A,\gamma]_A + ([\rho(\alpha),\rho(\beta)]_{\mathfrak{X}(M)}f)\gamma\end{aligned} \tag{13}$$

On the other hand, by the Leibniz rule we have

$$[[\alpha,\beta]_A, f\gamma]_A = f[[\alpha,\beta]_A,\gamma]_A + (\rho([\alpha,\beta]_A)f)\gamma \tag{14}$$

Combining equations (13) and (14) yields

$$([\rho(\alpha),\rho(\beta)]_{\mathfrak{X}(M)}f)\gamma = (\rho([\alpha,\beta]_A)f)\gamma\,, \quad \forall\,\alpha,\beta,\gamma\in\Gamma(A)\ , \quad \forall f\in\mathscr{C}^\infty(M)\,,$$

from where the seeked identity follows. □

Example 10.8 (Tangent bundle)**.** For any manifold M, we can regard its tangent bundle $A = \mathsf{T}M \to M$ as an algebroid, with $\rho \equiv \mathrm{Id}_{\mathsf{T}M}$ and $[\cdot,\cdot]_A$ the usual Lie bracket on $\mathfrak{X}(M)$. As a matter of fact, it is an integrable Lei algebroid, because $\mathsf{T}M = \mathrm{A}(M\times M)$ for the groupoid $M\times M \rightrightarrows M$.

Example 10.9 (Involutive distribution)**.** A vector sub-bundle $D\subseteq \mathsf{T}M$ such that $[X,Y]\in\Gamma(D)$ for all $X,Y\in\Gamma(D)$ is a Lie algebroid with $\rho=\iota\colon A\hookrightarrow \mathsf{T}M$ the natural inclusion. In fact, by the Frobenius theorem, it is an integrable Lie algebroid.

Example 10.10 (Lie algebra)**.** A Lie algebra $\mathfrak{g}$ can be regarded as a trivial Lie algebroid $A=\mathfrak{g}\to\{*\}$.

Definition 10.11. Let $A\to M$ be a Lie algebroid with anchor ρ. The *isotropy Lie algebra* of A at $x\in M$ is the vector subspace

$$\mathfrak{g}_x \coloneqq \ker\rho_x \subseteq A_x$$

equipped with the Lie bracket

$$[\alpha_x,\beta_x]_{\mathfrak{g}_x} \coloneqq [\alpha,\beta]_A(x)\,.$$

Note that the bracket $[\cdot,\cdot]_{\mathfrak{g}_x}$ is well-defined, i.e., independent on the choice of extensions $\alpha,\beta\in\Gamma(A)$ for $\alpha_x,\beta_x\in\mathfrak{g}_x$. Indeed, let $\alpha,\tilde\alpha,\beta,\tilde\beta\in\Gamma(A)$ such that $\alpha(x)=\tilde\alpha(x)=\alpha_x$ and $\beta(x)=\tilde\beta(x)=\beta_x$. We can choose a basis $\{\gamma_i\}$ of local sections of A around x, and write

$$\sigma \coloneqq \alpha-\tilde\alpha = \sum_i f_i\gamma_i\,,$$

where $f_i\in\mathscr{C}^\infty(M)$ with $f_i(x)=0$. Thus,

$$[\alpha,\beta]_A - [\tilde\alpha,\beta]_A = [\alpha-\tilde\alpha,\beta]_A = [\sigma,\beta]_A = \sum_i [f_i\gamma_i,\beta]_A = \sum_i \left(f_i[\gamma_i,\beta]_A - (\rho(\beta)f_i)\gamma_i\right)\ ,$$

where we have applyed the Leibniz rule in the last step. Since $f_i(x)=0$ and $\beta_x\in\ker\rho_x$, the right-hand side vanishes at x, which implies that

$$[\alpha,\beta]_A(x) = [\tilde\alpha,\beta]_A(x)\,.$$

By the skew-symmetry of the bracket, the same argument applies to extensions of β_x, namely,

$$[\alpha,\beta]_A(x) = -[\beta,\alpha]_A(x) = -[\tilde{\beta},\alpha]_A(x) = [\alpha,\tilde{\beta}]_A(x)\,.$$

If $A = \mathrm{A}(\mathcal{G})$ for a Lie groupoid $\mathcal{G} \rightrightarrows M$, then

$$\mathfrak{g}_x = \ker \mathsf{T}_{1_x} s \cap \ker \mathsf{T}_{1_x} t\,,$$

which is the Lie algebra of the vertex group (a.k.a. the isotropy group) $\mathcal{G}_x^x = s^{-1}(x) \cap t^{-1}(x)$.

For a general Lie algebroid $A \to M$, we can define a bundle of Lie algebras

$$\mathfrak{g}(A) = \ker \rho = \bigcup_{x\in M} \ker \rho_x\,,$$

which in general is **not** smooth. We call A a *regular Lie algebroid* if ρ has constant rank, so $\mathfrak{g}(A)$ is a smooth bundle. If A is regular, we have a short exact sequence of Lie algebroids:

$$0 \longrightarrow \ker\rho \longrightarrow A \longrightarrow \operatorname{Im}\rho \longrightarrow 0\,.$$

Moreover, $\operatorname{Im}\rho \subseteq \mathsf{T}M$ is an integrable distribution, and its associated integral foliation $\mathcal{F}$ is called the *orbit foliation* of the regular algebroid. It is also possible to define a (singular) orbit foliation for non-regular Lie algebroids:

$$\mathcal{F}_A = \{\mathcal{O}_i\}\ , \quad \mathsf{T}_x\mathcal{O}_i = \operatorname{Im}\rho_x\,, \quad \forall\, x \in \mathcal{O}_i\,.$$

A Lie algebroid is called *transitive* if $\operatorname{Im}\rho = \mathsf{T}M$.

Definition 10.12. Let $A_1 \to M$ and $A_2 \to M$ be two Lie algebroids over the same base. A *morphism of Lie algebroids* is a vector bundle morphism $\Phi\colon A_1 \to A_2$ such that:

$$\Phi\Big([\alpha,\beta]_{A_1}\Big) = \Big[\Phi(\alpha),\Phi(\beta)\Big]_{A_2}\,, \quad \forall\, \alpha,\beta \in \Gamma(A_1)\,,$$

and

$$\rho_2 \circ \Phi = \rho_1\,.$$

Example 10.13 (Atiyah algebroid)**.** Let $\pi\colon P \to M$ be a G-principal bundle. Its Atiyah algebroid is the vector bundle $A := \mathsf{T}P/G \to P/G$, with the anchor $\rho\colon A \to \mathsf{T}M$ induced by $\mathsf{T}\pi\colon \mathsf{T}P \to \mathsf{T}M$:

$$\begin{tikzcd} \mathsf{T}P \arrow[loop left, "G"] \arrow[r] \arrow[d, "\mathsf{T}\pi"'] & A = \mathsf{T}P/G \arrow[dl, "\rho"] \\ \mathsf{T}M & \end{tikzcd}\,.$$

The Lie bracket $[\cdot,\cdot]_A$ is induced by the restriction of the Lie bracket of $\mathfrak{X}(P)$ to the set $\mathfrak{X}_G(P)$ of G-invariant vector fields, namely,

$$\mathfrak{X}_G(P) = \{X \in \mathfrak{X}(P) \mid g_*X = X\,, \quad \forall g \in G\}\ .$$

One can easily see that A is the Lie algebroid of the gauge groupoid (a.k.a. Atiyah groupoid) $\mathcal{G} = (P\times P)/G \rightrightarrows M$.

If $A \simeq \mathrm{A}(\mathcal{G})$ is an integrable and transitive Lie algebroid, then the anchor

$$\rho\colon\ \mathrm{A}(\mathcal{G}) = \ker \mathsf{T}t|_{u(M)} \ni \alpha \mapsto \mathsf{T}_{1_x} s(\alpha) \in \mathsf{T}M$$

is surjective, and thus the restriction of the target map $t\colon \mathcal{G}_x \to M$ is a surjective submersion. By Proposition 3.3, this in turn implies that $\mathcal{G} \rightrightarrows M$ is locally trivial. By Proposition 3.7, $\mathcal{G}$ is isomorphic to the gauge groupoid $(\mathcal{G}_x \times \mathcal{G}_x)/\mathcal{G}_x^x$. In conclusion, an integrable transitive Lie algebroid $A \to M$ is isomorphic to the Atiyah algebroid of some principal bundle $P \to M$.

Example 10.14 (Action algebroid)**.** Let $a\colon \mathfrak{g} \to \mathfrak{X}(M)$ be an infinitesimal action of a Lie algebra $\mathfrak{g}$ on a manifold G. We define the associated action algebroid as the vector bundle $A = M \times \mathfrak{g} \to M$, with the anchor

$$\rho\colon M \times \mathfrak{g} \ni (x, \xi) \mapsto a(\xi)_x \in \mathsf{T}_x M\,,$$

and the Lie bracket on $\Gamma(A) \simeq \mathscr{C}^\infty(M, \mathfrak{g})$

$$[f, g]_A(x) = [f(x), g(x)]_{\mathfrak{g}} + \left(\mathcal{L}_{a(f(x))} g\right)(x) - \left(\mathcal{L}_{a(g(x))} f\right)(x)\,.$$

As one could expect, given a Lie group action of G on M, the action groupoid has as associated Lie algebroid the action algebroid of $\mathfrak{g} = \mathrm{Lie}(G)$.

Lecture 11 and 12 – BARTOSZ

Date: 24 February, 2026

Lecture 14 – Oscar Carballal (UCM)

Date: 24 February, 2026

Abstract: One of the early glimpses of Lie algebroid connections can be traced back to Poisson geometry, when Izu Vaisman, in his study of the geometric quantisation of Poisson manifolds, introduced the notion of contravariant derivatives. This framework was subsequently developed by Rui Loja Fernandes, who employed it to study global properties of Poisson manifolds and later extended the construction to the broader setting of Lie algebroids.

In this talk, we shall introduce the notion of a Lie algebroid connection on a vector bundle. We will also discuss representations of Lie algebroids (namely, flat Lie algebroid connections) and Lie algebroid cohomology, establishing the one-to-one correspondence between Lie algebroid representations and degree +1 graded derivations of the associated exterior algebra that are coboundary operators.

Finally, we shall explain how this framework naturally gives rise to parallel transport along A-paths, thereby enabling the comparison of points along different orbits of a Lie algebroid. The theory will be illustrated through several examples.

The notion of morphism of Lie algebroids over the same base was introduced in Lecture 10 (see Definition 10.12) by Asier López-Gordón. The notion of morphism of Lie algebroids over different bases follows straightforwardly:

Definition 12.1. A morphism between two Lie algebroids $A_1 \to M_1$ and $A_2 \to M_2$ is a VB morphism $(\Phi, \phi) : (A_1 \to M_1) \to (A_2 \to M_2)$ such that the following diagrams commute

$$\begin{tikzcd} A_1 \arrow[r, "\Phi"] \arrow[d] & A_2 \arrow[d] \\ M_1 \arrow[r, "\phi"'] & M_2 \end{tikzcd} \qquad \begin{tikzcd} A_1 \arrow[r, "\Phi"] \arrow[d, "\rho_1"'] & A_2 \arrow[d, "\rho_2"] \\ \mathsf{T}M_1 \arrow[r, "\mathsf{T}\phi"'] & \mathsf{T}M_2 \end{tikzcd}$$

and satisfying $\Phi([\alpha, \beta]_{A_1}) = [\Phi(\alpha), \Phi(\beta)]_{A_2}$ for every $\alpha, \beta \in \Gamma(A_1)$.

Early work on Lie algebroid connections arose in Poisson geometry. For a Poisson manifold (M, Π) with a linear connection ∇ on $\mathsf{T}M$, the natural compatibility condition is $\nabla\Pi = 0$, which forces Π to have constant rank. Vaisman introduced the concept of contravariant connections on (M, Π) via the Lie algebroid $\mathsf{T}^*M \to M$ induced Π, and Rui Loja Fernandes employed this framework to define the holonomy of the symplectic foliation of (M, Π). Fernandes later extended these ideas to the general setting of Lie algebroids.

Let us begin by recalling the appropriate notion of paths on Lie algebroids. Throughout, I denotes the unit interval $I := [0, 1]$. For a Lie algebroid $A \to M$, we write $\tau_A : A \to M$ for the underlying VB projection.

Definition 12.2. Let $A \to M$ be a Lie algebroid. An *A-path* is a pair (a, γ_a) consisting of paths $a : I \to A$ and $\gamma_a : I \to M$ such that

$$\gamma_a(t) = \tau_A(a(t)), \qquad \rho(a(t)) = \frac{\mathrm{d}\gamma_a}{\mathrm{d}t}, \qquad t \in I.$$

In this situation, we say that a lies above γ_a, and we refer to γ_a as the base path of a.

Given an A-path (a, γ_a), define the bundle map $a\,\mathrm{d}t : \mathsf{T}I \ni \partial_t \mapsto a(t) \in A$. Then the pair $(a\,\mathrm{d}t, \gamma_a) : (\mathsf{T}I \to I) \to (A \to M)$ is a morphism of Lie algebroids. This construction yields a one-to-one correspondence

$$\left\{ \ A\text{-paths} \ \right\} \leftrightharpoons \left\{ \begin{array}{c} \text{morphisms of Lie algebroids} \\ \mathsf{T}I \to A \end{array} \right\},$$

thereby justifying that A-paths constitute the natural notion of paths in the context of Lie algebroids.

The notion on A-paths allows to define an equivalence relation $\sim_A$ on M as follows: $x \sim_A y$ if there exists an A-path (a, γ_a) such that $\gamma_a(0) = x$ and $\gamma_a(1) = y$. The corresponding equivalence classes $\mathcal{O}$ are called the orbits of A. It can be shown that each orbit $\mathcal{O}$ is an immersed submanifold of M satisfying $\mathsf{T}_x\mathcal{O} = \operatorname{Im}\rho_x$ for every $x \in \mathcal{O}$. In other words, the orbits are precisely the integral submanifolds of the generalised distribution $\operatorname{Im}(\rho) \subset \mathsf{T}M$. In particular, if $A \to M$ is transitive, then its orbits coincide with the connected components of M. Let $\mathcal{O}$ be an orbit of $A \to M$. Then, the restriction of A to $\mathcal{O}$, denoted by $A_{\mathcal{O}}$, is a transitive Lie algebroid $A_{\mathcal{O}} \to \mathcal{O}$.

Lie Algebroid connections

Definition 12.3. Let $A \to M$ be a Lie algebroid and $E \to M$ a VB. An *A-connection* on E is a $\mathbb{R}$-bilinear map

$$\nabla : \Gamma(A) \times \Gamma(E) \ni (\alpha, e) \longmapsto \nabla_\alpha e \in \Gamma(E)$$

satisfying

(i) $\nabla_{f\alpha} e = f \nabla_\alpha e$, and
(ii) $\nabla_\alpha f e = f \nabla_\alpha e + (\mathcal{L}_{\rho(\alpha)} f) e$

for all $\alpha \in \Gamma(A), e \in \Gamma(E)$ and $f \in C^\infty(M)$.

Examples 12.4. (1) Let us consider the Lie algebroid $\mathsf{T}M \to M$. A $\mathsf{T}M$-connection on a VB $E \to M$ is just an usual connection on E. Given a Lie algebroid $A \to M$ and a $\mathsf{T}M$-connection ∇ on a VB $E \to M$ we obtain an A-connection $\widetilde{\nabla}$ on E defined by

$$\widetilde{\nabla}_\alpha e := \nabla_{\rho(\alpha)} e, \qquad \alpha \in \Gamma(A), \qquad e \in \Gamma(E).$$

In particular, this shows the existence of A-connections.

(2) Let $A \to M$ be a Lie algebroid, and consider $\mathbb{R}_M := \mathbb{R} \times M \to M$ the trivial line bundle over M. Then, the anchor of A yields an A-connection ∇ over $\mathbb{R}_M$ via

$$\nabla_\alpha f = \mathcal{L}_{\rho(\alpha)} f \tag{15}$$

for all $\alpha \in \Gamma(A)$ and $f \in C^\infty(M) = \Gamma(\mathbb{R}_M)$.

(3) Let $\mathcal{F}$ be a regular foliation on a manifold M. Consider the (regular) Lie algebroid $\mathsf{T}\mathcal{F} \hookrightarrow \mathsf{T}M$ and the normal VB to the foliation, defined as the quotient VB $\nu(\mathcal{F}) := \mathsf{T}M/\mathsf{T}\mathcal{F} \to M$. Since $\Gamma(\nu(\mathcal{F})) = \mathfrak{X}(M)/\Gamma(\mathsf{T}\mathcal{F})$, every section of $\nu(\mathcal{F})$ is the equivalence class $\overline{X}$ of a vector field $X \in \mathfrak{X}(M)$. The so-called *Bott connection* $\nabla^{\mathrm{Bott}} : \Gamma(\mathsf{T}\mathcal{F}) \times \Gamma(\nu(\mathcal{F})) \to \Gamma(\nu(\mathcal{F}))$ defined by

$$\nabla^{\mathrm{Bott}}_X \overline{Y} := \overline{[X,Y]}, \qquad X \in \mathfrak{X}(M), \qquad \overline{Y} \in \Gamma(\nu(\mathcal{F})),$$

is a $\mathsf{T}\mathcal{F}$-connection on $\nu(\mathcal{F})$.

(4) Let $A \to M$ be a regular Lie algebroid. Then, $\mathfrak{g}(A) = \ker\rho \to M$ is a VB and $\operatorname{Im}\rho$ yields a regular foliation on M, the orbit foliation $\mathcal{F}_A$ associated with A, such that $\mathsf{T}\mathcal{F}_A = \operatorname{Im}\rho$. Then,

$$\nabla_\alpha \beta := [\alpha, \beta], \qquad \alpha \in \Gamma(A), \qquad \beta \in \Gamma(\mathfrak{g}(A)),$$

and

$$\nabla_\alpha \overline{X} := \overline{[\rho(\alpha), X]}, \qquad \alpha \in \Gamma(A), \qquad \overline{X} \in \nu(\mathcal{F}_A)$$

are A-connections on $\mathfrak{g}(A)$ and on $\nu(\mathcal{F}_A)$, respectively.

Definition 12.5. Let $A \to M$ be a Lie algebroid and ∇ an A-connection on a VB $E \to M$. The *curvature* of ∇ is the $\mathbb{R}$-bilinear map $R_\nabla : \Gamma(A) \times \Gamma(A) \to \Gamma(\mathrm{End}(E))$ defined by

$$R_\nabla(\alpha, \beta)(e) := \nabla_\alpha \nabla_\beta e - \nabla_\beta \nabla_\alpha e - \nabla_{[\alpha,\beta]} e$$

for all $\alpha, \beta \in \Gamma(A)$ and $e \in \Gamma(E)$. The connection ∇ is said to be flat if $R_\nabla = 0$.

Claim. The curvature $R_\nabla(\alpha, \beta)(e)$ is $C^\infty(M)$-linear in α, β and e.

Proof. Since $R_\nabla(\alpha, \beta)(e) = -R_\nabla(\beta, \alpha)(e)$, it suffices to prove the $C^\infty(M)$-linearity for β and e. Let us first prove it for β. First of all, let $f \in C^\infty(M)$, and consider

$$R_\nabla(\alpha, f\beta)(e) = \nabla_\alpha \nabla_{f\beta} e - \nabla_{f\beta} \nabla_\alpha e - \nabla_{[\alpha, f\beta]} e. \tag{16}$$

One one hand,

$$\nabla_\alpha \nabla_{f\beta} e = \nabla_\alpha (f \nabla_\beta e) = f \nabla_\alpha \nabla_\beta e + \left(\mathcal{L}_{\rho(\alpha)} f\right) \nabla_\beta e, \qquad \nabla_{f\beta} \nabla_\alpha e = f \nabla_\beta \nabla_\alpha e, \tag{17}$$

while on the other hand,

$$\nabla_{[\alpha, f\beta]} e = \nabla_{f[\alpha,\beta] + (\mathcal{L}_{\rho(\alpha)} f)\beta} e = \nabla_{f[\alpha,\beta]} e + \nabla_{(\mathcal{L}_{\rho(\alpha)} f)\beta} e = f \nabla_{[\alpha,\beta]} e + \left(\mathcal{L}_{\rho(\alpha)} f\right) \nabla_\beta e. \tag{18}$$

Inserting (17) and (18) in (16) we obtain

$$R_\nabla(\alpha, f\beta)(e) = f R_\nabla(\alpha, \beta)(e),$$

which proves that the $C^\infty(M)$-linearity in β.

Let us now prove that $C^\infty(M)$-linearity in e. Consider $f \in C^\infty(M)$, and

$$R_\nabla(\alpha, \beta)(fe) = \nabla_\alpha \nabla_\beta fe - \nabla_\beta \nabla_\alpha fe - \nabla_{[\alpha,\beta]} fe. \tag{19}$$

A routine computation shows that

$$\nabla_\alpha \nabla_\beta fe = f \nabla_\alpha \nabla_\beta e + \left(\mathcal{L}_{\rho(\alpha)} f\right) \nabla_\beta e + \left(\mathcal{L}_{\rho(\beta)} f\right) \nabla_\alpha e + \mathcal{L}_{\rho(\alpha)} \left(\mathcal{L}_{\rho(\beta)} f\right) e \tag{20}$$

and, analogously, that

$$\nabla_\beta \nabla_\alpha fe = f \nabla_\beta \nabla_\alpha e + \left(\mathcal{L}_{\rho(\beta)} f\right) \nabla_\alpha e + \left(\mathcal{L}_{\rho(\alpha)} f\right) \nabla_\beta e + \mathcal{L}_{\rho(\beta)} \left(\mathcal{L}_{\rho(\alpha)} f\right) e, \tag{21}$$

while

$$\begin{aligned}\nabla_{[\alpha,\beta]}fe &= \left(\mathcal{L}_{\rho([\alpha,\beta])}f\right)e + f\nabla_{[\alpha,\beta]}e = \left(\mathcal{L}_{[\rho(\alpha),\rho(\beta)]}f\right)e + f\nabla_{[\alpha,\beta]}e \\ &= \mathcal{L}_{\rho(\alpha)}\left(\mathcal{L}_{\rho(\beta)}f\right)e - \mathcal{L}_{\rho(\beta)}\left(\mathcal{L}_{\rho(\alpha)}f\right)e + f\nabla_{[\alpha,\beta]}e.\end{aligned} \tag{22}$$

Inserting (20)–(22) in (19), we conclude that

$$R_\nabla(\alpha,\beta)(fe) = fR_\nabla(\alpha,\beta)(e).$$

□

Representations of Lie algebroids and Lie algebroids cohomology

Let $A \to M$ be a Lie algebroid and consider a VB $E \to M$. A *p-differential A-form with values in E* is a section $\omega \in \Gamma(\wedge^p A^* \otimes E)$. Equivalently, a an alternating $C^\infty(M)$-multilinear map

$$\omega : \Gamma(A) \times \overset{p}{\cdots} \times \Gamma(A) \to \Gamma(E).$$

We denote $\Omega^0(A;E) := \Gamma(E)$ and $\Omega^p(A;E) := \Gamma(\wedge^p A^* \otimes E)$ for $p \geq 1$. Note that, if $\operatorname{rk} A = k$, then $\Omega^p(A;E) = 0$ for all $p > k$. Let now ∇ be an A-connection on $E \to M$. Then, ∇ gives rise to an operator

$$\mathrm{d}_A^\nabla : \Omega^p(A;E) \to \Omega^{p+1}(A;E), \qquad p \geq 0,$$

defined by the Koszul-type formula

$$\begin{aligned}\mathrm{d}_A^\nabla\omega(\alpha_0,\dots,\alpha_p) := \sum_{i=0}^{p}(-1)^i\nabla_{\alpha_i}(\omega(\alpha_0,\dots,\widehat{\alpha}_i,\dots,\alpha_p)) \\ + \sum_{0\leq i<j\leq p}(-1)^{i+j}\omega([\alpha_i,\alpha_j],\alpha_0,\dots,\widehat{\alpha}_i,\dots,\widehat{\alpha}_j,\dots,\alpha_p),\end{aligned}$$

for all $\omega \in \Omega^p(A;E)$ and $\alpha_0,\dots,\alpha_p \in \Gamma(A)$. The striking point is that, trough a routine yet lengthy computation, one can show that $(\mathrm{d}_A^\nabla)^2 := \mathrm{d}_A^\nabla \circ \mathrm{d}_A^\nabla$ is such that $(\mathrm{d}_A^\nabla)^2 = 0$ if and only if ∇ is flat. In such case, $(\Omega^\bullet(A;E) := \bigoplus_{p\geq 0}\Omega^p(A;E), \mathrm{d}_A^\nabla)$ is a cochain complex.

The latter shows why we are interested in flat A-connections. More specifically,

Definition 12.6. A *representation* of a Lie algebroid $A \to M$ is a pair (E,∇), where $E \to M$ is a VB and ∇ is a flat A-connection on E.

From the previous discussion, given a representation (E,∇) of a Lie algebroid $A \to M$, the cohomology $H^\bullet(A;E)$ of the cochain complex $(\Omega^\bullet(A;E), \mathrm{d}_A^\nabla)$ is referred to as the *Lie algebroid cohomology of A with coefficients in the representation* (E,∇).

Examples 12.7.
(1) Let $A \to M$ be a Lie algebroid, and consider the trivial line bundle $\mathbb{R}_M \to M$ endowed with the A-connection ∇ defined in (15). Then, ∇ is flat, hence $(\mathbb{R}_M,\nabla)$ is a representation of A, known as the *trivial representation*. In this case, one usually denotes by $(\Omega^\bullet(A), \mathrm{d}_A)$ and by $H^\bullet(A)$ the resulting cochain complex and its cohomology. Moreover, $(\Omega^\bullet(A), \mathrm{d}_A)$ and $H^\bullet(A)$ are usually referred to as the *de Rham complex* and the *de Rham cohomology* of the Lie algebroid $A \to M$.
(2) Let (E,∇) be a representation of the Lie algebroid $\mathsf{T}M \to M$. Then, $H^\bullet(\mathsf{T}M;E) = H_{\mathrm{dR}}(M;E)$ is the usual de Rham cohomology with coefficients in E. Of course, the trivial representation $E = \mathbb{R}_M$ retrives the de Rham cohomology of the manifold M.

(3) Let $\mathfrak{g}$ be a finite-dimensional real Lie algebra, considered as a Lie algebroid $\mathfrak{g} \to \{*\}$. A representation of this Lie algebroid is just a $\mathfrak{g}$-module E, and the cohomology $H^\bullet(\mathfrak{g}; E)$ is the Chevalley–Eilenberg cohomology of $\mathfrak{g}$ with coefficients in E.
(4) Let $\mathcal{F}$ be a regular foliation on M, and consider the Lie algebroid $\mathsf{T}\mathcal{F} \to M$. Then, its normal bundle $\nu(\mathcal{F})$ endowed with the Bott connection ∇^{Bott} is a representation of $\mathsf{T}\mathcal{F} \to M$. The cohomology $H^\bullet(\mathsf{T}\mathcal{F}; \nu(\mathcal{F}))$ is called the *foliated cohomology*.

Let $A \to M$ be a Lie algebroid. Then, $\Omega^\bullet(A)$ together with the wedge product of A-forms $\wedge : \Omega^p(A) \times \Omega^\ell(A) \to \Omega^{p+\ell}(A)$ is a graded-commutative algebra on which the differential $\mathrm{d}_A : \Omega^\bullet(A) \to \Omega^\bullet(A)$ is a degree $+1$ operator satisfying

$$\mathrm{d}_A(\omega \wedge \eta) = \mathrm{d}_A\omega \wedge \eta + (-1)^{|\omega|}\omega \wedge \mathrm{d}_A\eta,$$

where $|\omega|$ denotes the degree of ω. In other words, $(\Omega^\bullet(A), \wedge, \mathrm{d}_A)$ is a differential graded-commutative algebra.

More generally, let (E, ∇) be a representation of a Lie algebroid $A \to M$. Then, for every $\omega \in \Omega^\bullet(A)$ and $\eta \in \Omega^\bullet(A; E)$, we have that

$$\mathrm{d}_A^\nabla(\omega \wedge \eta) = \mathrm{d}_A\omega \wedge \eta + (-1)^{|\omega|}\omega \wedge \mathrm{d}_A^\nabla\eta, \tag{$\star$}$$

showing that $(\Omega^\bullet(A; E), \mathrm{d}_A^\nabla)$ is a differential graded module over $(\Omega^\bullet(A), \mathrm{d}_A)$.

From all these previous comments, it easily follows

Proposition 12.8. Let $A \to M$ be a Lie algebroid and $E \to M$ be a VB. There is a one-to-one correspondence

$$\left\{ \begin{array}{c} A\text{-connections on } E \to M \end{array} \right\} \leftrightharpoons \left\{ \begin{array}{c} \text{degree } +1 \text{ operators } \mathrm{d}_A^\nabla : \Omega^\bullet(A; E) \to \Omega^\bullet(A; E) \\ \text{satisfying } (\star) \end{array} \right\}.$$

Moreover (E, ∇) is a representation if and only if $(\mathrm{d}_A^\nabla)^2 = 0$.

The Atiyah algebroid

Associated with every VB $E \to M$ one has its Lie algebroid of derivations $\mathrm{Der}(E) \to M$ defined as follows: foer every $x \in M$, the fibre $\mathrm{Der}(E)_x$ of $\mathrm{Der}(E)$ over x consists of derivations of $E \to M$ ath the point $x \in M$; that is, $\mathbb{R}$-linear maps $\delta : \Gamma(E) \to E_x$ such that, for every $f \in C^\infty(M)$ and $e \in \Gamma(e)$,

$$\delta(fe) = f(x)\delta(e) + v(f)e_x$$

for a, necessarily unique, vector $v \in \mathsf{T}_xM$, called the symbol of δ. Sections of $\mathrm{Der}(E) \to M$ are derivations of E, that is, $\mathbb{R}$-linear maps $D : \Gamma(E) \to \Gamma(E)$ such that, for every $f \in C^\infty(M)$ and $e \in \Gamma(E)$,

$$D(fe) = fD(e) + X(f)e$$

for a, necessarily unique, vector field $X \in \mathfrak{X}(M)$, referred to as the *symbol of* D. The map ρ associating its symbol to every derivation is a morphism of $C^\infty(M)$-modules, and hence can be considered as a VB morphism $\rho : \mathrm{Der}(E) \to \mathsf{T}M$. Note that $\mathrm{Der}(E)$ has a natural Lie algebra structure on its space of sections given by the commutator of derivations. Moreover, if D and D' are two derivations of E with symbols $\rho(D)$ and $\rho(D')$, respectively, their commutator $[D, D'] = D \circ D' - D' \circ D$ is a derivation of E with symbol $[\rho(D), \rho(D')]$. This shows that $\mathrm{Der}(E) \to M$ is a Lie algebroid, usually referred to as the *Atiyah algebroid* of $E \to M$.

Remark 12.9. Let $E \to M$ be a VB, and let $\mathrm{Fr}(E) \to M$ be its associated frame bundle, which is an $\mathrm{GL}(\mathrm{rk}\, E, \mathbb{R})$-principal bundle. It can be proved that there exists a natural Lie algebroid isomorphism covering id_M between $\mathrm{Der}(E) \to M$ and the Atiyah algebroid of the $\mathrm{GL}(\mathrm{rk}\, E, \mathbb{R})$-principal bundle $\mathrm{Fr}(E) \to M$. This justifies why we also refer to $\mathrm{Der}(E) \to M$ as the Atiyah algebroid of $E \to M$.

Let now $A \to M$ be a Lie algebroid, and consider ∇ an A-connection on a VB $E \to M$. Then, for each $\alpha \in \Gamma(A)$, we have that $\nabla_\alpha : \Gamma(E) \to \Gamma(E)$ is a derivation of E. Specifically, ∇ gives rise to a VB morphism

$$A \ni \alpha \mapsto \nabla_\alpha \in \mathrm{Der}(E)$$

covering id_M which is compatible with the anchors of $A \to M$ and $\mathrm{Der}(E) \to M$. Note that it is a Lie algebroid morphism if and only if (E, ∇) is a representation of $A \to M$.

As a consequence of the previous discussion, we have the following

Proposition 12.10. Let $A \to M$ be a Lie algebroid, and consider $E \to M$ a VB. There is a one-to-one correspondence

$$\left\{ \text{ representations } (E, \nabla) \text{ of } A \to M \right\} \leftrightharpoons \left\{ \text{ Lie algebroid morphisms } A \to \mathrm{Der}(E) \right\}.$$

Parallel transport along A-paths

Definition 12.11. Let $(\Phi, \phi) : (A_1 \to M_1) \to (A_2 \to M_2)$ be a Lie algebroid morphism, and consider an A_2-connection ∇ on a VB $E \to M_2$. The map $\Phi^*\nabla : \Gamma(A_1) \times \Gamma(\phi^* E) \to \Gamma(\phi^* E)$ defined by

$$(\Phi^*\nabla)_\alpha(\phi^* e) := \nabla_{\Phi(\alpha)} e, \qquad \alpha \in \Gamma(A), \qquad e \in \Gamma(E)$$

is an A_1-connection on the pull-back VB $\phi^* E \to M_1$, known as the pull-back connection of ∇ along (Φ, ϕ).

Let now (a, γ_a) be an A-path on a Lie algebroid $A \to M$, and consider an A-connection ∇ on a VB $E \to M$. Since $(a\, \mathrm{d}t, \gamma_a) : (\mathsf{T}I \to I) \to (A \to M)$ is a Lie algebroid morphism, let ∇^* denote the pull-back $\mathsf{T}I$-connection on the pull-back VB $\gamma_a^* E \to M$. Note that sections of $\gamma_a^* E$ are paths $u : I \to E$ such that $\tau_E(u(t)) = \gamma_a(t)$ for all $t \in E$. Then, the *A-derivative of $u \in \Gamma(\gamma_a^* E)$ along a* is defined as

$$(D_a u)(t) := \nabla^*_{\frac{\partial}{\partial t}} u \in \Gamma(\gamma_a^* E)$$

As usual, a path $u \in \Gamma(\gamma_a^* E)$ is said to be *parallel along a* if $D_a u \equiv 0$. In such case, given $e_0 \in E_{\gamma_a(0)}$, there exists a unique path $\tilde{u} \in \Gamma(\gamma_a^* E)$ parallel along a such that $\tilde{u}(0) = e_0$. We then define the *parallel transport along a* as the map

$$P_{\gamma_a} : E_{\gamma_a(0)} \ni e_0 \mapsto \tilde{u}(1) \in E_{\gamma_a(1)}.$$

Lecture 15– Arnau Mas

Date: 3 March, 2026

Lecture 16 – Juan Manuel López Medel

Date: 10 March, 2026

Abstract: In this lecture we study the problem of integration a Poisson manifold. Given a manifold (M, π), can we find a surjective submersion $\mu : (S, \omega) \to (M, \pi)$ being (S, ω). This is called a symplectic realization of the Poisson manifold.

Moreover, we are going to see that there is a natural structure of groupoid on the symplectic manifold (S, ω) related by the multiplicative form. This is called a symplectic groupoid.

Symplectic realization

The notion of Symplectic Groupoid is connected with the Poisson manifolds.

Definition 14.1. A symplectic realization of a Poisson manifold (M, π) $\mu : (S, \omega) \to (M, \pi)$ consits of

- a symplectic manifold (S, ω),
- a surjective submersion $\mu : S \to M$ which is a Poisson map.

Lemma 14.2. If $(S.\omega)$ is a symplectic realization of (M, π) then

$$\dim(S) \geq 2\dim(M) - \operatorname{rank} \pi_x, \ \forall x \in M.$$

Example 14.3. In view of submersion theorem, the problem of finding a symplectic realization takes the following form *locally*:

Given a Poisson structure π in local coordinates $(x^1, \ldots, x^m)$ with brackets

$$\{x^i, x^j\} = \pi^{ij}(x).$$

We need to define new brackets

$$\{x^i, u^a\} = \theta^{ia}(x, u), \quad \{u^a, u^b\} = \varphi^{ab}(x, u),$$

which are smooth functions in $(x, u) = (x^1, \ldots, x^m.u^1, \ldots, u^n)$ such that

$$\Pi = \sum_{i<j} \pi^{ij}(x) \frac{\partial}{\partial x^i} \wedge \frac{\partial}{\partial x^j} + \sum_{i,a} \theta^{ia}(x) \frac{\partial}{\partial x^i} \wedge \frac{\partial}{\partial u^a} + \sum_{a<b} \varphi^{ab}(x) \frac{\partial}{\partial u^a} \wedge \frac{\partial}{\partial u^b}$$

is nondegenerate and a Poisson structure.

In the general case, we can't find a Symplectic realization, but under some properties, as we will see in this lecture, we can find it.

Example 14.4. *Symplectic realization of a linear Poisson structure.*
In the linear case, the problem have complete solution, something that doesn't happen on

the general case.

We consider the linear Poisson structure on $\mathbb{R}^2$ associated with the nonabelian, two dimensional Lie algebra $\mathfrak{g} = \mathfrak{aff}(1, \mathbb{R})$:

$$\{x, y\} = x.$$

Because Poisson structure vanishes at origin, the smallest possible dimension of a symplectic realization is four.
We consider the map $\mu : \mathbb{R}^4 \to \mathbb{R}^2,\ \mu(x, y, u, v) = (x, y)$.
We look for a Poisson bracket on $\mathbb{R}^4$ for which μ is a Poisson map.
The only nonzero structure functions are:

$$\{x, y\} = x, \quad \{y, v\} = 1, \quad \text{and } \{x, u\} = \phi,$$

where ϕ is a function to determine.
The Jacobi identity is satisfied if $\phi = e^{-v}$, so

$$\{x, y\} = x, \quad , \{x, u\} = e^{-v}, \quad , \{y, v\} = 1, \quad , \{x, v\} = \{y, u\} = \{u, v\} = 0.$$

The underlying bivector is nondegenerate. The symplectic form is

$$\omega = e^v(du \wedge dx + xdu \wedge dv) + dv \wedge dy,$$

and can be written as $\omega = -d(x\theta_1 + y\theta_2)$ where

$$\theta_1 = e^v du, \quad \theta_2 = dv.$$

Form a coframe on $\mathbb{R}^2$ and satisfy $d\theta_1 = -\theta_1 \wedge \theta_2, d\theta_2 = 0$.

An example of a symplectic structure that admits a symplectic reduction is

$$\{x, y\} = x^2 + y^2.$$

Given a Poisson manifold (M, π), we search for a symplectic manifold (S, ω) and a submersion μ.
But, we can also think of that problem in this way:

Given a surjective submersion $\mu : (S, \omega) \to M$ defined on a symplectic manifold, does there exist a Poisson structure π on M so that μ becomes a symplectic reduction?
The answer of this question depends on the distribution that is symplectic orthogonal to the fibers of μ.

Theorem 14.5 (Libermann)**.** Let $\mu : (S, \omega) \to M$ a surjective submersion with connected fibers defined on a symplectic manifold.
Then M admits a Poisson structure π such that μ is a symplectic realization if and only if the ω-orthogonal to the fibers of μ

$$(\ker d\mu)^{\perp_\omega} \subset TS,$$

is an involutive distribution. Moreover, in this case π is unique.

Symplectic groupoid

Now, we add the structure of groupoid for the manifold of symplectic manifold in the realization problem.

It is known that a Lie groupoid has an associated Lie algebroid. We'll look at groupoids whose associated Lie algebroid is isomorphic to the cotangent Lie algebroid of a Poisson structure.
These should be thought of as the group-like objects "integrating" Poisson brackets. They are characterized by the presence of a symplectic structure on the space of arrows compatible with the multiplication.

We first discuss "compatibility" between a differential form and the groupoid structure. We consider $\mathcal{G} \rightrightarrows M$ and a degree 0-form $f \in C^\infty(\mathcal{G})$. The compatibility condition is that $f : (\mathcal{G}, \cdot) \to (\mathbb{R}, +)$ must be a homomorphism of groupoids

$$f(g \cdot h) = f(g) + f(h),\ \forall (g,h) \in \mathcal{G}^{(2)}.$$

This equation can be seen as an equation of 0-forms on $\mathcal{G}^{(2)}$,

$$m^* f = pr_1^* f + pr_2^* f.$$

Definition 14.6. A multiplicative form on $\mathcal{G} \rightrightarrows M$ is $\omega \in \Omega^k(\mathcal{G})$ satisfying

$$\boxed{m^*\omega = pr_1^*\omega + pr_2^*\omega.}$$

Definition 14.7. A symplectic groupoid, denoted as $\Sigma \rightrightarrows M$, is a Lie groupoid together with a multiplicative symplectic form $\Omega \in \Omega^2(\Sigma)$.

Example 14.8 (Pair groupoid)**.** Let (M, ω) be a symplectic manifold. The pair groupoid $\Sigma := M \times M \rightrightarrows M$ with the symplectic and multiplicative form

$$\Omega := pr_1^*\omega - pr_2^*\omega \in \Omega(M \times M),$$

makes $(M \times M, \Omega) \rightrightarrows M$ a symplectic groupoid.

Example 14.9. We consider the cotangent bundle $\Sigma = T^*M$ with canonical symplectic form $\Omega = \omega_{can}$. The addition on the fibers makes T^*M a bundle of abelian Lie groups over M, which is actually a symplectic groupoid:

$$(T^*M, \omega_{can}) \rightrightarrows M.$$

The multiplicative condition of ω_{can} follows from the multiplicativity of Liouville 1-form θ_L, as follows.

The multiplicative on $\Sigma^{(2)} = T^*M \oplus T^*M$ is given by $m : \Sigma^{(2)} \to T^*M,\ m(\alpha, \beta) = \alpha + \beta$.

If $(v_1, v_2) \in T_{(\alpha,\beta)}\Sigma^{(2)}$ then

$$dpr(v_1) = dpr(v_2) = v \in TM,$$

so

$$\theta_L(dm(v_1, v_2)) = (\alpha + \beta)(v) = \alpha(v) + \beta(v) = \theta_L(dpr_1(v_1, v_2)) + \theta_L(dpr_2(v_1, v_2)).$$

This said that θ_L is multiplicative.

Example 14.10. Let G be a Lie groupo and $\Sigma = G \ltimes \mathfrak{g}^* \rightrightarrows \mathfrak{g}^*$ the action groupoid associated with the coadjoint action.

Using the isomorphism $l : T^*G \simeq G \times \mathfrak{g}^*$ induced by left translation. We consider the symplectic form

$$\Omega := -l_*\omega_{can} = d(l_*\theta_L) \in \Omega^2(G \ltimes \mathfrak{g}^*).$$

It is multiplicative and then $(G \ltimes \mathfrak{g}^*, \Omega) \rightrightarrows \mathfrak{g}^*$ is a symplectic groupoid.

We show that $\theta := l_*(\theta_L) \in \Omega^1(G \ltimes \mathfrak{g}^*)$ is multiplicative,

$$\theta : T_{(g,\xi)}(G \ltimes \mathfrak{g}^*) \to \mathbb{R},\ (v_g, \eta) \mapsto \langle \xi, dL_{g^{-1}}(v_g)\rangle.$$

We identify $\Sigma^{(2)} = G \times G \times \mathfrak{g}^*$:

$$((g,\eta),(h,\xi)) \leftrightarrow (g,h,\xi),\ \text{where } \eta = \mathrm{Ad}_h^*\xi.$$

Then,

$$m(g,h,\xi) = (gh,\xi), \quad pr_1(g,h,\xi) = (g, \mathrm{Ad}_h^*\xi), \quad pr_2(g,h,\xi) = (h,\xi).$$

It follows that

$$m^*\theta_{(g,h,\xi)}(v,w,\eta) = \langle \xi, dL_{(gh)^{-1}}(dR_h v + dL_g w)\rangle = \langle \xi, \mathrm{Ad}_{h^{-1}}(dL_{g^{-1}}v) + dL_{h^{-1}}w\rangle,$$

and

$$pr_1^*\theta_{(g,h,\xi)}(v,w,\eta) = \langle \mathrm{Ad}_h^*\xi, dL_{g^{-1}}v\rangle, \qquad pr_2^*\theta_{(g,h,\xi)}(v,w,\eta) = \langle \xi, dL_{h^{-1}}w\rangle.$$

There is a more geometric interpretation of the multiplicative condition.

Lemma 14.11. A form $\Omega \in \Omega^2(\Sigma)$ on a Lie groupoid $\Sigma \rightrightarrows M$ is multiplicative if and only if the graph of the multiplication

$$Graph(m) = \{(g,h,g\cdot h); (g,h) \in \Sigma^{(2)}\} \subseteq \Sigma \times \Sigma \times \Sigma$$

is isotropic in $(\Sigma,\Omega) \times (\Sigma,\Omega) \times (\Sigma,-\Omega)$

Proposition 14.12. A symplectic form $\Omega \in \Omega^2(\Sigma)$ on a Lie groupoid $\Sigma \rightrightarrows M$ is multiplicative if and only if $Graph(m) \subseteq (\Sigma,\Omega) \times (\Sigma,\Omega) \times (\Sigma,-\Omega)$ is a Lagrangian submanifold.

Moreover, any symplectic groupoid (Σ,Ω) has the following properties

(1) The t-fibers and s-fibers are symplectic orthogonal

$$(\ker dt)^{\perp_\Omega} = \ker ds.$$

(2) The unit section $u : M \hookrightarrow \Sigma$ is a Lagrangian embedding:

$$(TM)^{\perp_\Omega} = TM.$$

(3) The inverse map $i : \Sigma \to \Sigma$ is antisymplectic, $i^*\Omega = -\Omega$.

In particular, $\dim \Sigma = 2 \dim M$.

Proof. To do... □

Theorem 14.13. $(\Sigma,\Omega) \rightrightarrows M$. There exists a unique Poisson structure on M such that the target map is Poisson. Moreover:

$$t \text{ is a } \textit{complete} \text{ symplectic realization.}$$

PROOF.
$$\begin{tikzcd} T_g\Sigma \arrow[r, "d_g t"] \arrow[d, "\Omega_g^\flat"] & T_xM \\ T_g^*\Sigma \arrow[u, bend left, "(\Omega_g^{-1})^\#"] & T_x^*M \arrow[l, "(d_g t)^*"'] \arrow[u, "(\pi_x^g)^\#"'] \end{tikzcd}$$
□

Examples

Now we present some examples of Poisson manifold and we'll try to see them as the base if a symplectic groupoid.

Given a Poisson manifold (M, π), the search for a symplectic manifold $(\Sigma, \Omega) \rightrightarrows M$ inducing the given π on the base is part of the search for symplectic realization of (M, π). The target map of Σ will give such realization. This is the conexion of symplectic groupoids and realizations, and it is due to the next Theorem:

Theorem 14.14 (Coste, Dazord, Weinstein)**.** Let $\mu : (S, \omega) \to (M, \pi)$ be a symplectic realization with connected fibers and a Lagrangian section $u : M \to S$. Then there exists at most one symplectic groupoid structure on $(S, \omega) \rightrightarrows M$ with target map $t = \mu$ and unit section $u = u$. This exists when the symplectic realization is complete and each leaf of symplectic orthogonal foliation $(\ker d\mu)^{\perp_\omega}$ intersects the Lagrangian section in precisely one point.

The proof gives us a recipe to construct the groupoid structure (as we will see in the example).

Example 14.15. We are going to consider the previous linear example. The linear Poisson structure on $\mathbb{R}^2$ given by $\{x, y\} = x$, with symplectic realization $\mu : \mathbb{R}^4 \to \mathbb{R}^2,\ (x, y, u, v) \mapsto (x, y)$, and

$$\omega = e^v(du \wedge dx + xdu \wedge dv) + dv \wedge dy$$

the symplectic form. This is a symplectic realization with connected fibers and it admits the gloval Lagrangian section $u : \mathbb{R}^2 \to \mathbb{R}^4,\ U(x, y) = (x, y, 0, 0)$.

We calculate the orbit foliation, which is spanned by

$$\mathfrak{a}(dx) = (\omega^{-1})^\#(dx) = x\frac{\partial}{\partial y} + e^{-v}\frac{\partial}{\partial u}$$

and

$$\mathfrak{a}(dy) = (\omega^{-1})^\#(dy) = -x\frac{\partial}{\partial x} + \frac{\partial}{\partial v}.$$

These vector fields have flows given by

$$\phi^t(x, y, u, v) = (x, y + xt, u + e^{-v}t, v)$$

and

$$\psi^t(x, y, u, v) = (e^{-t}x, y, u, t + v).$$

Let $f : \mathbb{R}^4 \to \mathbb{R}$ be a smooth constant function on the orbits of $\mathfrak{a}$. Using $t = -v$ for the second flow and $t = -u$ for the first, we obtain that

$$f(x, y, u, v) = f(xe^v, y, u, 0) = f(xe^v, y - xue^v, 0, 0).$$

So the orbit foliation coincides with the fibers of the submersion

$$\mathbb{R}^4 \to \mathbb{R}^2,\ (x,y,u,v) \mapsto (xe^v, y - xue^v).$$

Now, we choose μ to be the source map (because is the simpler) and this last for the target map.

Since the source map is anti-Poisson, we are forced to change the sign of the symplectic form on $\mathbb{R}^4$:

$$\Omega := -\omega = e^v(dx \wedge du + xdv \wedge du) + dy \wedge dv.$$

To find the multiplication, we consider two composable "arrows":

$$s(x',y',u',v') = t(x,y,u,v) \iff x' = xe^v,\ y' = y - xue^v.$$

The expression for $t(x',y',u',v')$ then yields:

$$x'e^{v'} = xe^{v+v'}, \quad y' - x'u'e^{v'} = y - x(ue^{-v'} + u')e^{v+v'}.$$

It follows that the product arrow

$$(x,y,u,v) = (x',y',u',v') \cdot (x,y,u,v)$$

can be depicted as:

$$(x',y',u',v') \longleftarrow (x',y') \longleftarrow (x,y)$$

with

$$(x',y') = (xe^{yv}, ye^{-xu}),$$

and

$$(x'e^{y'v'},\ y'e^{-x'u'}) = \Big(xe^{y(v+e^{-xu}v')},\ ye^{-x(u+e^{yv}u')}\Big).$$

which suggests taking

$$U = u + e^{yv}u', \qquad V = v + e^{-xu}v'.$$

So the structure of symplectic groupoid on

$$\mathbb{R}^4 \underset{s}{\overset{t}{\rightrightarrows}} \mathbb{R}^2$$

is given by

$$\begin{cases} \Omega = -d(xu) \wedge d(yv) + dx \wedge du + dy \wedge dv, \\ s(x,y,u,v) = (x,y), \\ t(x,y,u,v) = (xe^{yv},\ ye^{-xu}), \\ (x',y',u',v') \cdot (x,y,u,v) = \Big(x, y,\ u + e^{yv}u',\ v + e^{-xu}v'\Big). \end{cases}$$

This is a symplectic groupoid for which the induced Poisson structure on the base is $\{x,y\} = x$.

Also, as a groupoid, this is an *action groupoid*. $\mathcal{G} = \mathbb{R}^2$ is the group with multiplication

$$(u',v') \cdot (u,v) = (ue^{-v'} + u', v + v'),$$

acting on $\mathbb{R}^2$ from the left by

$$(u,v) \cdot (x,y) = (xe^v, y - xue^v).$$

$\mathcal{G}$ integrates the 2-dimensional, nonabelian Lie algebra $\mathfrak{g}$ and this action is the coadjoint action of $\mathcal{G}$ on $\mathfrak{g}^*$ which matches with $\Sigma = G \ltimes \mathfrak{g}^* \rightrightarrows \mathfrak{g}^*$ example.

Example 14.16. Given a symplectic manifold (M, ω), the pair groupoid $M \times M \rightrightarrows M$ is a symplectic groupoid for the symplectic form

$$\Omega := t^*\omega - s^*\omega.$$

The target map $t = pr_1 : (M \times M, \Omega) \to (M, \omega)$ is Poisson.

The Lie algebroid of $M \times M$ is the tangent bundle TM. For a concrete example, we consider $M = \mathbb{R}^2$ with the canonical Poisson structure $\{p, q\} = 1$.

symplectic groupoid on

$$\mathbb{R}^4 \underset{s}{\overset{t}{\rightrightarrows}} \mathbb{R}^2$$

is given by

$$\begin{cases} \Omega = dq^1 \wedge dp_1 - dq^2 \wedge dp_1, \\ s(q^1, p_1, q^2, p_2) = (q^2, p_2), \\ t(q^1, p_1, q^2, p_2) = (q^1, p_1), \\ (q^1, p_1, q^2, p_2) \cdot (q^2, p_2, q^3, p_3) = (q^1, p_1, q^3, p_3). \end{cases}$$

Actually for any Lie groupoid integrating TM the 2-form is symplectic and multiplicative. In particular, the homotopy groupoid $\Pi(M) \rightrightarrows M$ is a symplectic groupoid.

Integrability problem

Given a Poisson manifold (M, π), is there a symplectic groupoid $(\Sigma, \Omega) \rightrightarrows M$ inducing π?

Whenever a solution exists, we say that (M, π) is an integrable Poisson manifold. A solution (Σ, Ω) is called a symplectic integration of (M, π). One should think of it as a "symplectic desingularization".

For some Poisson structures, we can't find a symplectic integration.

Example 14.17. Considering the 3-sphere S^3 with zero Poisson structure. It has the symplectic integration

$$T^*S^3 \simeq S^3 \times \mathbb{R}^3 \to S^3,$$

which is a bundle of Lie groups with addition on the fibers and multiplicative symplectic form $\Omega = \omega_{can}$. The bundle of Lie groups $\mathcal{G} = S^3 \times \mathbb{T}^3 \to S^3$ does not carry any symplectic form. This follows because every class $c \in H^2(S^3 \times T^3)$ squares to zero: $c^2 = 0$.

So T^*M of a Poisson manifold is always a Lie algebroid.

Bibliography

[1] K. C. H. Mackenzie, *General Theory of Lie Groupoids and Lie Algebroids*. in London Mathematical Society Lecture Note Series. Cambridge: Cambridge University Press, 2005. DOI:10.1017/CBO9781107325883. (cited in p. 21)

[2] R. L. Fernandes. Lecture notes from *"Lie Groupoids and Lie Algebroids"* 2021 course at U. Illinois. Available at https://publish.illinois.edu/ruiloja/math-595-lie-groupoids-and-lie-algebroids-fall-2021/. (cited in pp. 5 and 51)

[3] A. Weinstein. Groupoids: Unifying Internal and External Symmetry. A Tour through Some Examples. *Notices Amer. Math. Soc.* **43** (1996) 744–752. (cited in pp. 5 and 7)

[4] F. M. Ciaglia, A. Ibort and G. Marmo. Schwinger's picture of quantum mechanics I: Groupoids.*Int. J. Geom. Methods Mod. Phys.* **16** (2019) 1950119. DOI:10.1142/S0219887819501196 (cited in p. 7)

[5] A. Ibort, A. Mas and L. Schiavone. "Fields" in classical and quantum field theories. *Preprint* arXiV: 2506.15327 [math-ph]. (cited in p. 7)

[6] M. de León, M. Epstein and V. Jiménez.*Material Geometry: Groupoids in Continuum Mechanics*. (Singapore: World Scientific) 2021. DOI:10.1142/12168 (cited in p. 7)

[7] J. C. Marrero, D. M. de Diego and E. Martínez, Discrete Lagrangian and Hamiltonian mechanics on Lie groupoids. *Nonlinearity* **19** (2006) 1313. DOI:10.1088/0951-7715/19/6/006 (cited in p. 7)

[8] L. Colombo, D. M. de Diego. Second-order variational problems on Lie groupoids and optimal control applications. *Discrete Contin. Dyn. Syst.* **36** (2016) 6023–6064. DOI:10.3934/dcds.2016064 (cited in p. 7)

[9] R. Palais. A global formulation of the Lie theory of transformation groups. *Mem. Amer. Math. Soc.* **22** (1957). DOI:10.1090/memo/0022 (cited in p. 8)

[10] J.-P. Ortega and T. Ratiu. *Momentum Maps and Hamiltonian Reduction*. (Boston: Birkhäuser) 2004. DOI:10.1007/978-1-4757-3811-7 (cited in p. 8)

[11] A. Schmeding and C. Wockel, "The Lie group of bisections of a Lie groupoid," *Annals of Global Analysis and Geometry*, vol. 48, no. 1, pp. 87–123, Jun. 2015. DOI:10.1007/s10455-015-9459-z. (cited in pp. 21 and 22)

[12] I. Moåçerdijk and J. Mrcun, *Introduction to Foliations and Lie Groupoids*. in Cambridge Studies in Advanced Mathematics. Cambridge: Cambridge University Press, 2003. DOI:10.1017/CBO9780511615450. (cited in p. 21)