

Improving Medical Diagnostic Accuracy and Efficiency with Artificial Intelligence (AI)

Asma Essaedi

School of Professional Studies

IS 499: Information System Capstone

Professor John Bouma

Mar 30, 2026

Table of Contents

Introduction.....	3
Topic Description.....	4
Properties and Qualities of AI in Medical Diagnostics	4
Problem Description.....	7
Inefficiency and Clinician Workload Burden	7
Subjectivity and Diagnostic Variability	9
Inability to Manage Massive and Complex Medical Datasets	10
Solution Description	13
Automation for Clinical Efficiency	13
Decision Support for Enhanced Accuracy	14
Predictive Analytics for Early Detection	15
Analysis	20
How I Found This Topic and Determined It Actually Exists	20
How I Determined That the Problems Actually Exist	21
How My Solutions Will Fulfill Both Operational and Strategic Goals	22
Research	23
Conclusion	24
References	25

Introduction

The purpose of this paper is to analyze the current state of artificial intelligence in medical diagnostics and examine in detail specific problems that AI can address. It will also propose concrete solutions supported by case studies from radiology, pathology, neurology, and oncology, as well as discussing critical ethical, legal, and implementation considerations that must guide any responsible deployment of these technologies.

The integration of artificial intelligence in healthcare represents one of the most significant technical transformations in modern medicine. The healthcare system faces enormous pressure from an aging population, rising chronic disease burden, and an explosion of medical data. According to Zhou et al. (2025), the healthcare artificial intelligence market is projected to reach \$188 billion by 2030. This signals the profound clinical potential these technologies hold.

This project examines a major challenge that modern medicine faces: inefficiency and inaccuracy of medical diagnosis in an era of information overload and constrained clinical resources. The traditional diagnostic process, while built upon centuries of medical knowledge and practice, shows its limitations when confronted with the high volume of data. The huge amount of data that modern imaging technologies, genomic sequencing, and electronic health records generate exceeds human capacity to derive meaningful insights (Panayides et al., 2020). Such limitations translate into extended wait time for diagnosis, significant variability in clinical decision-making, and missed opportunities for early disease detection.

The solution to these diagnostic challenges lies in integrating AI into clinical workflows. AI plays a key role in automating repetitive tasks, enhancing decision accuracy through decision support, and managing large and complex data assets for early disease detection. However, this

technological advancement must be balanced with significant ethical considerations and regulatory requirements governing the deployment of artificial intelligence in healthcare settings.

Topic Description

The topic I am going to discuss is artificial intelligence in medical diagnostics, which refers to the application of machine learning algorithms, particularly deep learning, to analyze complex medical data, support clinical decision-making, and improve patient outcomes (IBM, 2025). This technology aims to augment human clinical expertise with computational power capable of processing vast amounts of information beyond human cognitive capacity.

Properties and Qualities of AI in Medical Diagnostics

Artificial intelligence in medical diagnostics possesses several defining properties and qualities that distinguish it from traditional computer-aided detection systems. First, AI systems, particularly deep learning models, exhibit automatic feature extraction (Aicadium, 2024). That means they can identify relevant patterns in data without being explicitly programmed to recognize specific features. Convolutional neural networks (CNNs) automatically learn hierarchical features from image data, identifying edges, shapes, and patterns associated with pathological conditions (Ahmed et al., 2024). Second, these systems demonstrate scalability, as they can process thousands of images or patient records in the time it takes a human to review a single case (Abbas et al., 2025). Third, AI diagnostics offer consistency. Unlike human clinicians whose performance may vary with fatigue or distraction, an AI algorithm will apply the same analytical criteria to every case it encounters (Greengrass, 2026). Fourth, modern medical AI exhibits multimodal integration capabilities, which means it can simultaneously analyze and correlate data from diverse sources, including medical images, genomic sequences, laboratory

results, and electronic health records (Hao et al., 2025). Finally, many AI systems now incorporate explainability features through Explainable AI (XAI) techniques. This allows clinicians to understand the rationale behind algorithmic recommendations (Teixeira et al., 2025).

The evolution of diagnostic technology has followed a clear path over the past century. From purely clinical examination and physician judgment, the field of medical diagnosis moved to incorporate laboratory testing in the mid-twentieth century, then advanced to computer-aided detection systems in the 1980s and 1990s (Orr, 2025; Doi, 2007). These early computer-aided detection systems, although innovative, were limited by their reliance on hand-coded rules and could not learn from new data (Avanzo et al., 2024). The current era, powered by deep learning and convolutional neural networks, represents a fundamental shift where systems continuously improve as they are exposed to more data.

The integration of AI into healthcare extends beyond technological advances as it addresses the unsustainable pressure on global healthcare systems. An aging population in developed nations and the growing burden of chronic diseases have boosted demand for diagnostic services. This increase in demand comes at a time when the medical workforce is facing a shortage and burnout (Abbas et al., 2025; American Medical Association, 2025). Such limitations necessitate the shift from traditional overextended clinical models to more scalable and technology-driven interventions of AI. Combining artificial intelligence technologies enhances human expertise, not replacing the important elements of human interaction in medicine. An artificial intelligence tool in clinical practice functions as an essential complement that "facilitates individualized therapy and eventually enhances patient outcomes by identifying subtle visual aspects" (Thomas et al., 2025). By delegating high-volume tasks such as routine

image analysis and report generation to algorithms, practitioners can effectively prioritize critical decisions and focus their cognitive resources on the most urgent cases.

Various factors make this topic particularly timely for investigation. First, the maturity of artificial intelligence technologies has reached a point where real-world clinical implementations are no longer theoretical but operational (Evinent, 2025). Major medical centers now routinely use AI-powered tools for mammography screening, retinal imaging, and pathology review (Doi, 2007; Ahmed et al., 2024). Second, the regulatory landscape has become clearer, with the United States Food and Drug Administration (FDA) having approved hundreds of AI-enabled medical devices as of 2025 (Center for Devices and Radiological Health, 2025). Third, the COVID-19 pandemic accelerated both the development and acceptance of digital health technologies, including diagnostic AI, as health systems were forced to innovate rapidly in crisis conditions (Zhou et al., 2025).

I chose this topic because the diagnostic application of artificial intelligence across medical specialties solves real-world problems, which results in improved patient care. By examining AI implementation in radiology, pathology, neurology, and oncology, this research aims to identify overarching principles and specific considerations that inform successful AI integration. I find this topic deeply interesting because of its interdisciplinary nature. Medical diagnosis increasingly draws on data from multiple sources, and an artificial intelligence system capable of integrating these diverse data types will likely provide the greatest clinical value. The gravity of this topic cannot be overstated, as improving diagnostic accuracy and efficiency directly impacts patient survival and quality of life.

Problem Description

The problem I am trying to solve is the three interconnected challenges facing contemporary medical diagnostics that adversely impact the timeliness and accuracy of care. These challenges include inefficiency and clinician workload burden; subjectivity and diagnostic variability; and the inability to manage massive and complex medical datasets. These challenges represent systemic issues that cannot be addressed through gradual improvement alone.

3.1 Inefficiency and Clinician Workload Burden

The diagnostic process in modern medicine takes a long time because it requires clinicians to manually review massive volumes of data. For instance, in pathology, the traditional process of reviewing prostate biopsies requires examining an average of 579 slides, which takes about 15.76 hours (Jeong et al., 2025). Also, radiologists interpreting computed tomography (CT) studies may need to review hundreds of individual images for a single patient to detect subtle abnormalities that could indicate early-stage disease (Thomas et al., 2025; Jeong et al., 2025).

This volume of work creates a diagnostic bottleneck that delays treatment and increases patient anxiety. More concerning, however, is the human cost of this workload. Physician burnout has reached epidemic proportions in many healthcare systems, with diagnostic specialties particularly affected. The cognitive load imposed by sustained attention to repetitive image interpretation tasks contributes directly to this burnout. When clinicians are exhausted, their diagnostic accuracy declines, which creates a dangerous cycle in which overwork leads to errors that, in turn, require additional work to correct (Jeong et al., 2025).

Prevalence and Widespread Nature of the Problem

The inefficiency problem is exacerbated by workforce shortages, which affect healthcare systems globally. Many countries face such shortages, with the United States projecting shortages of 20,600 pathologists, neurologists, and radiologists by 2032 (New Findings Confirm Predictions on Physician Shortage, 2025). Similar trends are observed across Europe. EU-REST study found that radiologist numbers ranged from 51 to 270 per million inhabitants, with 16 of 27 member states falling below the EU average and nearly half of radiologists approaching retirement age (Brady et al., 2025). In developing nations, the shortage problem is even worse, with many countries operating at ratios as low as one radiologist per million people, compared to approximately 93 per million in high-income countries (Hussain et al., 2025). Without technological intervention, the gap between diagnostic demand and clinical capacity will continue to widen, inevitably compromising patient care. This problem affects virtually every healthcare system, from community hospitals to academic medical centers.

Urgency of Solving Inefficiency

The urgency of addressing this problem cannot be overstated. Delayed diagnoses lead to delayed treatments, which directly affect patient survival rates. For many cancers, a delay of even weeks can mean the difference between early-stage disease amenable to curative treatment and advanced disease requiring palliative care only (Hanna et al., 2020). A systematic review and meta-analysis of over 1.2 million patients published in The BMJ found that each four-week delay in cancer treatment is associated with a significant increase in mortality rate, including a 6-8% increase for surgical interventions across multiple cancer types. Furthermore, burned-out clinicians are more likely to leave practice. This worsens the very workforce shortages that

created the problem, which creates a downward spiral that, if left unchecked, will render current healthcare delivery models unsustainable.

3.2 Subjectivity and Diagnostic Variability

Diagnostic accuracy can depend on a clinician's experience and subjective judgment, which means different clinicians may arrive at different diagnoses for the same patient or situation. For example, in hematology, morphological identification of blood cells is somewhat subjective even among skilled hematologists (Jeong et al., 2025). This subjectivity can lead to diagnostic discrepancies, which in turn risk misdiagnosis or delayed diagnosis (Jeong et al., 2025). As an illustration, a patient diagnosed with one type of leukemia may receive completely different treatment protocols than if they had received a different diagnosis, despite presenting with essentially the same cellular morphology (*How AML Subtypes Affect Outlook & Treatment*, n.d.).

Diagnostic variability can also arise from institutional disparities. Patients treated at academic medical centers may receive diagnoses different from those seen at community hospitals. This is not because the disease is different, but because tools and clinicians' interpretive frameworks and experiences differ (Thomas et al. 2025). This creates inequities in healthcare that fundamentally conflict with the ethical principle that all patients deserve accurate and timely diagnosis, regardless of where they seek care.

Prevalence of Diagnostic Variability

Studies have documented significant variability across various medical specialties. In pathology, inter-observer agreement for diagnosing certain cancers can be as low as 48% among

expert pathologists (Wilson & Wilson, 2020). In radiology, studies show that radiologists miss breast cancers in mammography at rates ranging from 10% to 30%, with significant variation between readers (Ekpo et al., 2018). In dermatology, studies show that complete agreement among expert pathologists diagnosing melanoma occurs in only 53.5% of cases, with discordance particularly pronounced for early-stage, non-invasive melanomas (Haggenmüller et al., 2025).

Urgency of Solving Diagnostic Variability

The urgency of solving this problem lies in the fact that misdiagnosis can mean the difference between life and death, between curative treatment and palliative care. Medical errors, including diagnostic errors, are estimated to contribute to nearly 440,000 deaths annually in the United States alone, which makes it the third leading cause of death (Young, 2016). Beyond mortality, misdiagnosis leads to unnecessary treatments, patient anxiety, and billions in wasted healthcare spending (National Academies of Sciences, 2015). As medicine becomes more complex, the potential for variability only increases, which makes standardization through AI not just beneficial but essential.

3.3 Inability to Manage Massive and Complex Medical Datasets

The volume and complexity of modern medical data have outpaced human cognitive capacity for pattern recognition. Contemporary diagnostic imaging produces high-resolution three-dimensional reconstructions, in which a single CT slice can contain millions of data points (Ahmed et al., 2024). Furthermore, genomic sequencing generates enormous datasets detailing the entire genetic code of a patient's tumor. Programs such as The Cancer Genome Atlas (TCGA) have molecularly characterized over 20,000 primary cancer samples, which generate

more than 2.5 petabytes of genomic, transcriptomic, and proteomic data (The Cancer Genome Atlas Program (TCGA), n.d.). Electronic health records also accumulate years of clinical data, laboratory results, and medication histories.

Human cognitive capacity cannot accommodate these disparate sources to derive diagnostic insights. Although human biological neural networks are "prodigious Olympians" in perceptual motor tasks, we have not yet mastered abstract thought, which remains computationally difficult to execute (Korteling et al., 2021). This limitation is quantified by a narrow conscious processing bandwidth. The capacity of human working memory is estimated at 10-50 bits per second (Korteling et al., 2021). This cognitive limitation means that subtle patterns that could indicate early-stage disease are frequently missed. For example, pancreatic cancer is often diagnosed at advanced stages because its early symptoms are non-specific and its radiographic signs are subtle (Pesheva, 2024). This means by the time it becomes clinically apparent, curative intervention is often no longer possible. Consequently, rather than forcing the human mind to perform tasks that exceed its capacity, a digital system, in this case AI, can be used to complement and compensate for the inherent biological and cognitive constraints of human intelligence.

Prevalence of Data Overload

The data overload problem affects every diagnostic encounter. According to Rani et al. (2025), "hospitals generate 50 petabytes of data annually, with individual EHRs and medical imaging contributing roughly 80 megabytes of data per patient per year". Globally, healthcare data is projected to grow at a compound annual rate of 36%, which is 6% faster than manufacturing, 11% faster than media, and 10% faster than finance (The Future of Healthcare is

Precision, n.d.). Yet the number of clinicians available to interpret this data remains essentially flat. This widening gap between data generation and human interpretive capacity, which means that even the most diligent clinicians are bound to miss information that could benefit their patients.

Urgency of Addressing Data Complexity

The urgency here is perhaps the greatest of all because human limitations cannot be trained away. No amount of medical education can expand working memory beyond its biological constraints. The consequence is that diseases like pancreatic cancer, which could potentially be detected early through pattern recognition in electronic health records, continue to be diagnosed late with devastating mortality rates (National Academies of Sciences, 2015). Similarly, opportunities to identify patients at risk for Alzheimer's disease before irreversible brain damage occurs are systematically missed (Jack et al., 2024). These are not failures of individual clinicians but failures of a system that asks humans to perform tasks fundamentally beyond their cognitive capabilities.

These three problems are closely linked. Inefficiency contributes to clinician fatigue, which exacerbates diagnostic variability. The complexity of modern data overwhelms clinicians, which leads to missed diagnoses. Addressing any one of these problems in isolation is unlikely to succeed; what is really needed is a systemic approach that addresses all three simultaneously. Artificial intelligence, with its capacity for automated pattern recognition and consistent application of analytical rules, offers precisely such a systemic approach.

Solution Description

My solution to the problem is the strategic integration of artificial intelligence into medical diagnostics to directly address the diagnostic challenges outlined above through three key approaches: automating routine tasks to improve efficiency, providing decision support to enhance accuracy, and enabling predictive analytics to facilitate early detection. These mechanisms complement one another, as effective AI implementation combines multiple approaches to achieve optimal clinical outcomes.

4.1 Automation for Clinical Efficiency

The most immediate contribution of artificial intelligence in medical diagnostics is the automation of time- and resource-intensive tasks. This frees clinicians to focus their expertise on complex cases and on direct patient care, where human judgment remains irreplaceable.

In radiology, convolutional neural network (CNN)-based systems have exhibited remarkable capabilities in detecting fractures and abnormalities in computed tomography (CT) and X-ray images (Ahmed et al., 2024). Li et al. (2023) developed a deep learning-based FRF-DPS for rib fractures. They used 14,241 CT scans for module development and 3931 for evaluation. The result is an efficient, swift AI system for rib fractures that reduces the average reading time per case from 5-7 minutes to just 3-5 seconds (Li et al., 2023). This dramatic time saving does not affect accuracy, as the system maintains sensitivity and specificity while completing analyses in seconds rather than minutes.

AI can also be used as a pre-screening tool that identifies clearly normal cases and flags those that require expert review. For example, in mammography screening, where the majority of

studies show no abnormalities, an AI pre-screening tool can reduce radiologists' workload by up to 70% (Jeong et al., 2025). AI also improves the cancer detection rate by 21.6%, compared with standard screening methods (5.6 vs 4.6 per 1,000 women scanned) (Staff, n.d.). The logic is straightforward: if an algorithm can confidently identify normal studies, radiologists can dedicate their attention and time to the cases that actually contain abnormalities and require expert intervention. This approach reduces the cognitive burden of screening and improves diagnostic yield.

In pathology, similar automation opportunities exist. Digital pathology systems combined with AI algorithms can automatically identify regions of interest within tissue slides, measure biomarker expression levels, and flag potentially malignant cells for pathologist review (Song et al., 2023). A study by Yang et al. (2025) demonstrated that GPT-4 processed pathology reports in 4.03 seconds compared to 65.64 seconds for human pathologists, which represents a substantial reduction in review time while maintaining high accuracy.

4.2 Decision Support for Enhanced Accuracy

Artificial intelligence can serve as an intelligent decision support tool that provides clinicians with real-time guidance during diagnostic interpretation. This human-AI collaboration (often termed augmented intelligence) utilizes the complementary strengths of human clinical judgment and machine pattern recognition (Augmentir, Inc., 2025).

In neurology, the use of AI to assist in segmenting brain tumors from MRI scans has proven to be greatly beneficial in a clinical setting. Cè et al. (2023) report that AI can accurately delineate tumor boundaries (segmentation) and assess changes in response to therapy over time. These tasks are time-consuming and subject to inter-observer variability when performed

manually. When AI segmentation is combined with clinical review, the resulting tumor measurements achieve higher accuracy than either AI or clinicians alone could achieve. This finding accentuates an important principle: the goal of medical AI is not to replace clinicians but to enhance their work.

Decision support also encompasses integrating multiple data types. Modern diagnostic workups generate diverse information, including image findings, laboratory results, and clinical history. Synthesizing these disparate data sources into a coherent diagnostic assessment is challenging even for the most experienced clinicians. An AI system designed for multimodal data integration can identify patterns across data types that indicate abnormalities (Thomas et al. 2025). For example, an AI tool might flag that a patient with certain imaging findings and a specific genetic marker has an increased probability of a rare condition. This prompts the clinician to order confirmatory testing that would not otherwise be considered.

4.3 Predictive Analytics for Early Detection

The most transformative potential of AI in diagnostics lies in its predictive analytics capabilities. AI-powered diagnostics can identify individuals at elevated risk of disease before clinical symptoms appear. This shift from reactive to proactive medicine has profound implications for patient outcomes.

Research from Harvard and the University of Copenhagen has produced an AI tool that analyzes electronic health records to identify individuals with the highest risk of pancreatic cancer up to 3 years before traditional diagnosis (Pesheva, 2024). Pancreatic cancer is notoriously difficult to detect early, and most patients present with advanced disease when curative intervention is no longer possible (Pesheva, 2024). An AI system that can stratify risk

years in advance creates opportunities for targeted screening, closer surveillance, and potentially earlier intervention in high-risk individuals.

The predictive approach relies on the analysis of massive datasets that no human could reasonably process. According to Pesheva (2024), the AI system examines millions of patient records to identify subtle combinations of symptoms, disease codes, and prior diagnoses that precede pancreatic cancer diagnosis. These patterns may be individually non-specific, such as type 2 diabetes or abdominal discomfort, but collectively, they form a predictive signature that pinpoints elevated risk.

Similar approaches are being developed for Alzheimer's disease, where the pathologic process begins years before clinical symptoms emerge. Jack et al. (2024) describe revised diagnostic criteria that incorporate biomarker evidence of Alzheimer's pathology, which enables diagnosis at the preclinical stage. Building on this biological framework, AI analysis of imaging, genetic, and cognitive data can identify individuals on a trajectory to develop symptomatic Alzheimer's, which creates opportunities for preventive interventions as they become available (Wang et al., 2024).

Operational and Strategic Objectives

The operational objectives of the automation approach include reducing diagnostic turnaround time for routine cases, decreasing clinician fatigue and reallocating specialist time to complex cases that require human judgment. The strategic long-term objective is to create a healthcare system where AI handles the volume of routine diagnostic work. This would allow the clinical workforce to operate at the top of their license and expertise.

The operational objectives for decision support include reducing diagnostic variability across clinicians and institutions. That, in turn, improves accuracy for complex cases and ensures that relevant clinical data is never overlooked. The strategic long-term objective is to establish a new standard of care where AI-augmented diagnosis becomes the norm, leading to more consistent, evidence-based clinical decisions across practice settings.

The operational objectives for predictive analytics include developing and validating risk prediction models for major diseases, like pancreatic cancer. This includes integrating these models into electronic health record systems and creating clinical protocols for managing high-risk patients. The strategic long-term objective is to transform medicine from a reactive discipline that treats established disease to a proactive discipline that prevents disease before it manifests clinically.

Implementation Approach and Feasibility

Implementing these solutions requires a phased approach. In the short term (1-2 years), healthcare organizations should pilot AI automation tools in high-volume areas like mammography and pathology screening. A successful example of this approach is the pilot of the 'Mia' AI tool in NHS Grampian, which resulted in 10.4% increase in cancer detection and a significant reduction in radiologist workload (*MiA - Tools*, n.d.). It showed the feasibility of piloting AI in high-volume screening areas.

In the medium term (2-4 years), decision support systems should be deployed across multiple specialties, with continuous monitoring of accuracy and usability. An example is Kaiser Permanente Northern California's implementation of the Advanced Alert Monitor program (AAM). The AAM program has been implemented in 21 hospitals; the predictive algorithm

identified patients at risk of clinical deterioration up to 12 hours in advance and triggered rapid response team assessment (Advance Alert Monitor Program, 2024). The program achieved a 16% relative risk reduction in mortality and prevented more than 500 deaths annually system-wide, with compliance protocols ensuring clinicians responded to alerts through standardized electronic health record documentation (Greene, 2023; Martinez et al, 2022).

In the long term (3-5 years), predictive analytics tools should be integrated into population health management, with clear protocols for risk communication and preventive intervention. Geisinger Health System's REDD (Readmission, Emergency Department, Death) model exemplifies this approach. The algorithm predicts which patients are at highest risk for adverse events within three days of hospital discharge (Agrawal et al., 2016; Wertz, 2018). It triggers a comprehensive intervention protocol that includes pre-discharge medication review, primary care follow-up scheduling and next-day outreach for vulnerable patients. The model achieved a C-statistic of 0.671 in predicting post-discharge complications. This enables care teams to target resources toward the highest-risk patients while maintaining efficient operations for lower-risk populations (Agrawal et al., 2016).

How Easy or Hard Will It Be to Execute This Solution?

The execution difficulty varies significantly across the three solution components, as seen above. Automation tools for high-volume tasks are the easiest to implement, as they require minimal workflow disruption since they integrate with existing imaging equipment and typically require only brief clinician training in interpreting AI outputs. As of March 2026, 1,039 FDA-authorized AI devices operate in radiology, which accounts for 77% of all medical AI (Tegomoh, 2025). This demonstrates the maturity of this sector.

Decision support systems are of moderate difficulty, as they require careful integration with electronic health records and customization to local practice patterns. Unlike standalone imaging tools, decision support must be integrated within clinical workflows, a process that faces well-documented barriers. A systematic review in Safety Science identified 16 distinct challenges to AI adoption, including workflow misalignment and inadequate training, all of which are particularly relevant to decision support implementations (Abdelwanis et al., 2025). Furthermore, a framework published in JAMA Network Open emphasized that successful clinical decision support demands user-centered design, translation of probabilistic outputs into actionable recommendations, and proactive management of alert fatigue (Schafer et al., 2025). Real-world experience reflects this complexity: in a national NHS program evaluated in The Lancet's eClinicalMedicine, one-third of hospital trusts had not deployed AI diagnostic systems 18 months after the contracting phase was meant to be completed, with delays primarily attributed to "varied local governance processes, IT systems, and data availability and quality," as well as "staff's limited AI knowledge, time to participate, and concerns over appropriate tool usage" (Ramsay et al., 2025).

Predictive analytics tools present the highest execution difficulty, as they demand foundational data infrastructure that is often immature in clinical settings. Unlike automation tools that operate on individual data streams (e.g., a single image) or decision support systems that synthesize existing structured data, predictive models require longitudinal, cleaned, and interoperable datasets drawn from electronic health records, often spanning years of clinical history (Ivanovski et al., 2025). This necessitates significant upfront investment in data warehousing, governance, and extraction–transformation–load (ETL) pipelines before any model can be validated (Ivanovski et al., 2025). Predictive analytics also face regulatory and validation

hurdles because these tools are designed to forecast future clinical events (Park et al., 2021). They require prospective clinical validation to establish safety and efficacy (Park et al., 2021).

All three approaches face common challenges, which are a resistance to change among clinicians, the need for interoperability with legacy systems, and the requirement for ongoing monitoring and updates (Rybakov, 2025). These barriers are consistently highlighted in the implementation literature; for instance, the Safety Science review notes that infrastructure limitations and provider resistance cut across all AI applications (Abdelwanis et al., 2025). However, the existence of successful implementations at major medical centers demonstrates that these challenges, while significant, are surmountable with proper planning and resources.

The common ground among these three applications is the AI system's ability to detect patterns that humans cannot see. This includes analyzing images for subtle radiographic signs, integrating multimodal data for comprehensive assessment, or examining electronic health records for predictive signatures. AI extends human perceptual and cognitive capabilities in ways that address the fundamental challenges of modern diagnosis.

Analysis

How I Found This Topic and Determined It Actually Exists

Since I am an IT student, I search for the latest technological development news on a daily basis. I regularly follow technology news sources, academic journals, and industry publications to stay up to date on emerging trends. Over the past several months, I noticed that artificial intelligence in medical diagnostics appeared with increasing frequency across these sources, from FDA approvals of new AI diagnostic tools to research publications demonstrating

breakthrough results and news articles about AI implementations at major hospitals. The consistency and volume of this coverage signaled that this was an active, growing field with real-world implementations. I decided to investigate further by reading peer-reviewed literature, and I found extensive documentation of AI systems operating in clinical settings, with measurable results published in reputable medical journals. The existence of hundreds of FDA-approved AI-enabled medical devices as of 2025 confirmed that this topic represents an established reality rather than a future possibility.

How I Determined That the Problems Actually Exist

My analysis confirms that the problems I have identified, inefficiency, diagnostic variability, and inability to manage complex data, are real, well-documented, and urgently in need of solutions. I determined that these problems exist through extensive peer-reviewed literature documenting clinician workload measurements, studies showing diagnostic discrepancies across institutions, and research on human cognitive limitations relative to modern data volumes (Jeong et al., 2025; Thomas et al., 2025; Korteling et al., 2021). The prevalence of these problems is evident in the widespread reports of physician burnout, delayed diagnoses, and missed opportunities for early detection that characterize contemporary healthcare. Furthermore, workforce shortage projections from authoritative sources, like the Association of American Medical Colleges, provide quantitative confirmation that the gap between diagnostic demand and clinical capacity is measurable and growing.

How My Solutions Will Fulfill Both Operational and Strategic Goals

I determined that my proposed solutions will sufficiently address these problems because the case studies and research presented demonstrate measurable improvements across all three areas. The rib fracture detection system reduced reading time from minutes to seconds while maintaining accuracy (Li et al., 2023). This directly fulfills the operational goal of improving efficiency and the strategic goal of reallocating clinician time to complex cases. The mammography AI reduced workload by 70% while improving cancer detection rates, simultaneously addressing operational efficiency and the strategic objective of improving patient outcomes (Jeong et al., 2025; Staff, n.d.). The pathology AI reduced processing time from 65.64 seconds to 4.03 seconds per report, a reduction of over 93% while maintaining high accuracy (Yang et al., 2025). These results show that automation directly addresses inefficiency.

Decision support systems for brain tumor segmentation achieved higher accuracy than either AI or clinicians alone (Cè et al., 2023). This directly addresses diagnostic variability while fulfilling the operational goal of diagnostic standardization. Predictive analytics for pancreatic cancer and Alzheimer's disease demonstrate the ability to identify at-risk individuals years before traditional diagnosis (Pesheva, 2024; Jack et al., 2024). This directly addresses the data complexity challenge while fulfilling the strategic objective of transforming medicine from reactive to proactive care.

However, I also recognize that technical solutions alone are insufficient. The analysis of ethical, legal, and implementation considerations reveals critical factors that must guide deployment. Algorithmic bias poses a significant risk, as AI systems trained on non-representative data may perform worse for marginalized populations (Weiner et al., 2025).

Transparency and explainability are essential ethical requirements, supported by the World Health Organization and the American Medical Association, to ensure clinicians can maintain accountability and patients can provide informed consent (American Medical Association, 2025; Health Ethics & Governance, 2021). Regulatory frameworks such as FDA approval processes and the EU AI Act codify these ethical principles into enforceable requirements (Pantanowitz et al., 2024).

Research

The background for this project is grounded in concepts from multiple courses in my Information Systems program. IS 320 (System Analysis and Design) emphasizes the importance of understanding user needs and designing systems that seamlessly fit into existing work patterns. For AI diagnostics, this means considering how AI output is presented, when it becomes available in the diagnostics workflow, and how it integrates with other information sources clinicians use. The course also covered the software development lifecycle (SDLC), which informs project management for AI systems. IS 355 (Strategic Perspective in IT) provided the framework for understanding technical integration and interoperability challenges. This course emphasized that systems do not exist in isolation but must interact with existing infrastructure, which requires attention to data formats, interfaces, and interoperability standards. IS 250 (Computer and Network Security) provided the foundation for security and data privacy. This course covered the importance of protecting personally identifiable information through implementing security controls and maintaining audit trails, which are all critical considerations for AI systems processing sensitive medical data.

Conclusion

The topic description, problems, solutions, and analysis lead to a clear conclusion that the successful integration of AI into medical diagnostics requires attention to technical, ethical, and organizational dimensions. The path forward requires collaboration that brings together clinical expertise, data science, ethics, and information systems. Responsible AI integration depends on diverse perspectives working together throughout the development and deployment lifecycle. Clinicians must articulate diagnostic needs and evaluate AI performance in real-world contexts. Data scientists must develop algorithms that are both accurate and fair. Ethicists must identify potential harms and make sure patient perspectives are represented. Information systems professionals must design and manage the technical infrastructure that makes AI tools accessible, reliable and secure.

The technical skills developed through the IS program, including an understanding of databases, system analysis, project management, and security, are essential for AI implementation. Equally important, however, are the program's emphasis on ethical decision-making, critical thinking, and communication. These competencies enable an IS professional to act as a link between technical teams and clinical users. This ensures that the AI systems are technically sound, clinically valuable, ethically defensible, and organizationally sustainable.

References

- Abbas, S. R., Seol, H., Abbas, Z., & Lee, S. W. (2025). Exploring the Role of Artificial Intelligence in Smart Healthcare: A Capability and Function-Oriented Review. *Healthcare*, 13(14), 1642. <https://doi.org/10.3390/healthcare13141642>
- Abdelwanis, M., Simsekler, M. C. E., Gabor, A. F., Sleptchenko, A., & Omar, M. (2025). Artificial intelligence adoption challenges from healthcare providers' perspectives: A comprehensive review and future directions. *Safety Science*, 193, 107028. <https://doi.org/10.1016/j.ssci.2025.107028>
- Advance Alert Monitor Program: an automated early warning system for adults at risk for In-Hospital clinical deterioration*. (2024, October 30). PSNet. <https://psnet.ahrq.gov/innovation/advance-alert-monitor-program-automated-early-warning-system-adults-risk-hospital>
- Agrawal, D., Chen, C.-B., Dravenstott, R. W., Strömblad, C. T. B., Schmid, J. A., Darer, J. D., Devapriya, P., & Kumara, S. (2016). Predicting Patients at Risk for 3-Day Postdischarge Readmissions, ED Visits, and Deaths. *Medical Care*, 54(11), 1017–1023. <https://www.jstor.org/stable/26418257>
- Ahmed, I. S., Dai, J., Xie, Y., & Liang, X. (2024). Deep learning models for CT image classification: a comprehensive literature review. *Quantitative Imaging in Medicine and Surgery*, 15(1), 962–1011. <https://doi.org/10.21037/qims-24-1400>
- Aicadium. (2024, December 17). *The Evolution of Artificial Intelligence (AI)*. Aicadium. <https://aicadium.ai/the-evolution-of-ai/>

- American Medical Association. (2025, June 11). *AMA adopts new policy aimed at ensuring transparency in AI tools*. <https://www.ama-assn.org/press-center/ama-pressreleases/ama-adopts-new-policy-aimed-ensuring-transparency-ai-tools>
- Augmentir, Inc. (2025, December 5). *What is Augmented Intelligence? - Augmentir*. Augmentir. <https://www.augmentir.com/glossary/what-is-augmented-intelligence>
- Avanzo, M., Stancanello, J., Pirrone, G., Drigo, A., & Retico, A. (2024). The evolution of artificial intelligence in medical imaging: from computer science to machine and deep learning. *Cancers*, 16(21), 3702. <https://doi.org/10.3390/cancers16213702>
- Brady, A. P., et al. (2025). Current status of radiologist staffing, education and training in the 27 EU Member States. *Insights into Imaging*, 16(1), 1-10. <https://doi.org/10.1186/s13244-025-01925-7>
- Center for Devices and Radiological Health. (2025, December 5). *Artificial Intelligence-Enabled medical devices*. U.S. Food and Drug Administration. <https://www.fda.gov/medical-devices/software-medical-device-samd/artificial-intelligence-enabled-medical-devices>
- Cè, M., Esposito, A., Cellina, M., Vingiani, A., Nicolò, P., & Manfrini, N. (2023). Artificial Intelligence in Brain Tumor Imaging: A Step toward Personalized Medicine. *Current Oncology*, 30(3), 2673-2701. <https://www.mdpi.com/1718-7729/30/3/203>
- Doi, K. (2007). Computer-aided diagnosis in medical imaging: Historical review, current status and future potential. *Computerized Medical Imaging and Graphics*, 31(4–5), 198–211. <https://doi.org/10.1016/j.compmedimag.2007.02.002>
- Ekpo, E. U., Alakhras, M., & Brennan, P. (2018). Errors in Mammography Cannot be Solved Through Technology Alone. *pmc.ncbi.nlm.nih.gov*. <https://doi.org/10.22034/APJCP.2018.19.2.291>

Evinent. (2025a, November 18). *Future of AI in healthcare: How artificial intelligence is transforming medicine and healthcare business*. <https://evinent.com/blog/future-of-ai-in-healthcare>

Greene, J. (2023, June 14). *Real-time in-hospital alerts associated with lower patient mortality*. Kaiser Permanente Division of Research. <https://divisionofresearch.kaiserpermanente.org/real-time-in-hospital-alerts/>

Greengrass, C. J. (2026). Transforming clinical reasoning—the role of AI in supporting human cognitive limitations. *Frontiers in Digital Health*, 7, 1715440. <https://doi.org/10.3389/fdgth.2025.1715440>

Haggenmüller, S., Wies, C., Abels, J., Winterstein, J. T., Heinlein, L., Garcia, C. N., Utikal, J. S., Wohlfeil, S. A., Meier, F., Hobelsberger, S., Gellrich, F. F., Sergon, M., Hauschild, A., French, L. E., Heinzerling, L., Schlager, J. G., Ghoreschi, K., Schlaak, M., Hilke, F. J., . . . Brinker, T. J. (2025). Discordance, accuracy and reproducibility study of pathologists' diagnosis of melanoma and melanocytic tumors. *Nature Communications*, 16(1), 789. <https://doi.org/10.1038/s41467-025-56160-x>

Hanna, T. P., King, W. D., Thibodeau, S., Jalink, M., Paulin, G. A., Harvey-Jones, E., ... & Aggarwal, A. (2020). Mortality due to cancer treatment delay: systematic review and meta-analysis. *BMJ*, 371, m4087. <https://doi.org/10.1136/bmj.m4087>

<https://pmc.ncbi.nlm.nih.gov/articles/PMC12670022/pdf/rycan.250019.pdf>

Hao, Y., Cheng, C., Li, J., Li, H., Di, X., Zeng, X., Jin, S., Han, X., Liu, C., Wang, Q., Luo, B.,

Zeng, X., & Li, K. (2025). Multimodal integration in Health Care: development with applications in disease management. *Journal of Medical Internet Research*, 27, e76557. <https://doi.org/10.2196/76557>

Health Ethics & Governance. (2021, June 28). *Ethics and governance of artificial intelligence for health*. World Health Organization.

<https://www.who.int/publications/i/item/9789240029200>

Hussain HK, Rajeev R, DeStigter KK & Gulani V (2025). Global Cancer Imaging Access: Addressing Barriers and Harnessing Innovations. *Radiology: Imaging Cancer*; 7:6.

How AML subtypes Affect outlook & Treatment | American Cancer Society. (n.d.). American Cancer Society. <https://www.cancer.org/cancer/types/acute-myeloid-leukemia/detection-diagnosis-staging/how-classified.html>

IBM. (2025, November 17). *What is Artificial Intelligence in medicine?*

<https://www.ibm.com/think/topics/artificial-intelligence-medicine>

Ivanovski, T. K., Honap, S., Matic, R., Markovic, S., & Peyrin-Biroulet, L. (2025). Building a healthcare data warehouse: considerations, opportunities, and challenges. *Frontiers in Digital Health*, 7, 1691142. <https://doi.org/10.3389/fdgth.2025.1691142>

Jack, C. R., Andrews, J. S., Beach, T. G., Buracchio, T., Dunn, B., Graf, A., ... & Masliah, E. (2024). Revised criteria for diagnosis and staging of Alzheimer's disease: Alzheimer's

Association Workgroup. *Alzheimer's & Dementia*, 20(8), 5143-5169.

<https://pubmed.ncbi.nlm.nih.gov/38934362/>

Jeong, J., Kim, S., Pan, L., Hwang, D., Kim, D., Choi, J., ... & Yoo, S. J. (2025). Reducing the workload of medical diagnosis through artificial intelligence: A narrative review.

Medicine, 104(6), e41470. <https://pmc.ncbi.nlm.nih.gov/articles/PMC11813001/>

Korteling, J. E., Van De Boer-Visschedijk, G. C., Blankendaal, R. a. M., Boonekamp, R. C., & Eikelboom, A. R. (2021). Human- versus Artificial Intelligence. *Frontiers in Artificial Intelligence*, 4, 622364. <https://doi.org/10.3389/frai.2021.622364>

Li, N., Wu, Z., Jiang, C., Sun, L., Li, B., Guo, J., ... & Tian, L. (2023). An automatic fresh rib fracture detection and positioning system using deep learning. *British Journal of Radiology*, 96(1146), 20221006. <https://pmc.ncbi.nlm.nih.gov/articles/PMC10230380/>

Martinez, V. A., Betts, R. K., Scruth, E. A., Buckley, J. D., Cadiz, V. R., Bertrand, L. D., Paulson, S. S., Dummett, B. A., Abhyankar, S. S., Reyes, V. M., Hatton, J. R., Sulit, R., & Liu, V. X. (2022). The Kaiser Permanente Northern California Advance Alert Monitor Program: an automated early warning system for adults at risk for In-Hospital clinical deterioration. *The Joint Commission Journal on Quality and Patient Safety*, 48(8), 370–375. <https://doi.org/10.1016/j.jcjq.2022.05.005>

MiA - Tools. (n.d.). AI Knowledge Hub. <https://ai.gov.uk/knowledge-hub/tools/mia/>

National Academies of Sciences, Engineering, and Medicine. (2015). *Improving diagnosis in health care*. The National Academies Press. <https://doi.org/10.17226/21794>

New findings confirm predictions on physician shortage. (2025, January

2). AAMC. <https://www.aamc.org/news/press-releases/new-findings-confirm-predictions-physician-shortage>

Orr, S. (2025, March 25). *Throwback of Lab History: Honoring the evolution of Med Labs*.

CrossMed Healthcare. <https://crossmedhealth.com/historyoflab/>

Panayides, A. S., Amini, A., Filipovic, N. D., Sharma, A., Tsiftaris, S. A., Young, A., Foran, D.,

Do, N., Golemati, S., Kurc, T., Huang, K., Nikita, K. S., Veasey, B. P., Zervakis, M.,

Saltz, J. H., & Pattichis, C. S. (2020). AI in Medical Imaging Informatics: Current

Challenges and Future Directions. *IEEE Journal of Biomedical and Health Informatics*,

24(7), 1837–1857. <https://doi.org/10.1109/jbhi.2020.2991043>

Pantanowitz, L., Harrington, S., & Parwani, A. V. (2024). Digital pathology and artificial

intelligence in translational medicine and clinical practice. *Modern Pathology*, 37(1),

100325. <https://www.sciencedirect.com/science/article/pii/S0893395224001893>

Park, S. H., Choi, J., & Byeon, J. (2021). Key principles of clinical validation, device approval,

and insurance coverage decisions of artificial intelligence. *Korean Journal of Radiology*,

22(3), 442. <https://doi.org/10.3348/kjr.2021.0048>

Pesheva, B. E. (2024, December 12). AI predicts future pancreatic cancer. *Harvard Medical*

School. <https://hms.harvard.edu/news/ai-predicts-future-pancreatic-cancer>

Ramsay, A. I., Crellin, N., Lawrence, R., Walton, H., Bagri, S., Dodsworth, E., Elphinstone, H.,

Gleeson, F., Halliday, A., Herbert, K., Lloyd, J., Massou, E., Mehta, R., Morris, S., Ng,

P. L., O'Regan, T., Sherlaw-Johnson, C., & Fulop, N. J. (2025). Procurement and early

deployment of artificial intelligence tools for chest diagnostics in NHS services in

England: a rapid, mixed method evaluation. *EClinicalMedicine*, 89, 103481.

<https://www.sciencedirect.com/science/article/pii/S2589537025004146>

Rani, S., Kumar, R., Kumar, R., Panda, B. S., Kumar, R., Kumar, R., Muften, N. F., Abass, M.

A., & Lozanović, J. (2025). Machine Learning-Powered Smart Healthcare Systems in the

- era of Big Data: applications, diagnostic insights, challenges, and ethical implications. *Diagnostics*, 15(15), 1914. <https://doi.org/10.3390/diagnostics15151914>
- Rybakov, A. (2025, June 24). *AI challenges in healthcare: tech, ethical, trust concerns*. SPsoft. <https://spsoft.com/tech-insights/key-ai-challenges-in-healthcare/>
- Schafer, K. M., Franklin, J., Perlis, R. H., & Walsh, C. (2025). Barriers and Solutions to Efficient Health Care AI implementation. *JAMA Network Open*, 8(11), e2544086. <https://doi.org/10.1001/jamanetworkopen.2025.44086>
- Song, A. H., Jaume, G., Williamson, D. F. K., Lu, M. Y., Vaidya, A., Miller, T. R., & Mahmood, F. (2023). Artificial intelligence for digital and computational pathology. *Nature Reviews Bioengineering*, 1(12), 930–949. <https://doi.org/10.1038/s44222-023-00096-8>
- Staff, A. P. (n.d.). *Large AI breast cancer screening trial increases detection rate by 20%*. The ASCO Post. <https://ascopost.com/news/november-2025/large-ai-breast-cancer-screening-trial-increases-detection-rate/>
- Tegomoh, B. (2025). The Physician AI Handbook: Peer-Reviewed Evidence for Every Specialty. In *Zenodo (CERN European Organization for Nuclear Research)*. <https://physicianaihandbook.com/appendices/specialty-tools.html>
- Teixeira, B., Carvalhais, L., Pinto, T., & Vale, Z. (2025). Explainable AI framework for reliable and transparent automated energy management in buildings. *Energy and Buildings*, 347, 116246. <https://doi.org/10.1016/j.enbuild.2025.116246>
- Thomas, K. S., Edpuganti, S., Puthooran, D. M., Thomas, A., Joy, A., & Latheef, S. (2025). Artificial intelligence in modern clinical practice (Review). *Medical International*, 6(1), 5. <https://pmc.ncbi.nlm.nih.gov/articles/PMC12715461/>

The Cancer Genome Atlas Program (TCGA). (n.d.). *National Cancer Institute*.

<https://www.cancer.gov/ccg/research/genome-sequencing/tcga>

The future of healthcare is precision | Akkodis. (n.d.).

<https://www.akkodis.com/en/blog/articles/series-the-future-of-healthcare-is-precision>

Wang, Y., Gao, R., Wei, T., Johnston, L., Yuan, X., Zhang, Y., Yu, Z., & Initiative, F. T. a. D.

N. (2024). Predicting long-term progression of Alzheimer's disease using a multimodal deep learning model incorporating interaction effects. *Journal of Translational Medicine*, 22(1), 265. <https://doi.org/10.1186/s12967-024-05025-w>

Weiner, E. B., Dankwa-Mullan, I., Nelson, W. A., & Hassanpour, S. (2025). Ethical challenges and evolving strategies in the integration of artificial intelligence into clinical practice. *PLOS Digital Health*, 4(4), e0000810.

<https://pmc.ncbi.nlm.nih.gov/articles/PMC11977975/>

Wertz, P. (2018, June 6). Identifying patients at risk for hospital readmission, emergency room visits. *Penn State University*. <https://www.psu.edu/news/research/story/identifying-patients-risk-hospital-readmission-emergency-room-visits>

Wilson, E. S., & Wilson, E. S. (2020, May 6). *Pathology under fire - Clinical lab products*.

Clinical Lab Products. <https://clpmag.com/diagnostic-technologies/digital-pathology/pathology-fire/>

Yang, X., Zhang, Y., Jiang, J., Chen, Z., Bai, R., Yuan, Z., Dong, L., Xiao, Y., Liu, D., Deng, H., Huang, J., Shi, H., Liu, D., Liang, M., Tang, W., & Xu, C. (2025). Harnessing GPT-4 for automated error detection in pathology reports: Implications for oncology diagnostics. *Digital Health*, 11, 20552076251346703. <https://doi.org/10.1177/20552076251346703>

- Young, M (2016, October 14). Medical errors – the third leading cause of death in the US. *Petrie-Flom Center* - Health Law Policy, Biotechnology, and Bioethics at Harvard Law School. <https://petrieflom.law.harvard.edu/2016/10/14/medical-errors-the-third-leading-cause-of-death-in-the-us/>
- Zhou, J., Park, S., Dong, S., Tang, X., & Wei, X. (2025). Artificial intelligence-driven transformative applications in disease diagnosis technology. *Med Rev*, 5(5), 353-377. <https://pmc.ncbi.nlm.nih.gov/articles/PMC12558038/>