

Optimization Issues

Starting with the following,

$$\Psi(\lambda, h, g) = \frac{1}{x_{max} - x_{min}} \left(\sum_x \left[\ln \frac{1}{1 - q_x} - \frac{h}{g} e^{gx} (e^g - 1) - \lambda \right] \right)^2$$

For simplicity, Set

$$R = x_{max} - x_{min} , \quad Q = \sum_x \ln \frac{1}{1 - q_x} , \quad H = \frac{h}{g} (e^g - 1)$$

So,

$$\Psi(\lambda, h, g) = \frac{1}{R} (Q - R\lambda - H \sum_x e^{gx})^2$$

Note that λ here has minimal interaction with any of the observations. It's actually quite trivial to show that to minimize Ψ on λ we set λ to minimum of its range (i.e. $\lambda \rightarrow 0$). This is infact what we get when we run optimization functions in R or visualize Ψ through brute force.

Additionally, if we took the exponent of equation (10) in your paper:

$$\ln\left(\frac{1}{1 - q_x}\right) = \frac{h}{g} (e^g - 1) e^{gx} + \lambda$$

We get an expression analgous to:

$$y = ae^{bx} + c$$

This represent the model that we would need to solve, and unfourtunately the gradient of this expression at $c = \lambda$ is just 1, making it impossible to optimize.