

CM0246 Discrete Structures

Relations and Their Properties

Andrés Sicard-Ramírez

Universidad EAFIT

Semestre 2014-2

Preliminaries

Textbook

Kenneth H. Rosen ([1988] 2004). Matemática Discreta y sus Aplicaciones.

Convention

The numbers and page numbers assigned to chapters, examples, exercises, figures, quotes, sections and theorems on these slides correspond to the numbers assigned in the textbook.

Relations

Recall the definition of Cartesian product

Let A and B be sets. The Cartesian product of A and B is

$$A \times B = \{ (a, b) \mid a \in A \wedge b \in B \}.$$

Example

Let $A = \{a, b\}$ and $B = \{1, 2\}$. Then

$$A \times B = \{(a, 1), (a, 2), (b, 1), (b, 2)\}.$$

Relations

Definition

Let A and B be sets. A **binary relation** from A to B is a subset of $A \times B$.

Relations

Definition

Let A and B be sets. A **binary relation** from A to B is a subset of $A \times B$.

Example (trivial relations)

$R = \emptyset$ and $R = A \times B$ are relations from A to B .

Relations

Definition

Let A and B be sets. A **binary relation** from A to B is a subset of $A \times B$.

Example (trivial relations)

$R = \emptyset$ and $R = A \times B$ are relations from A to B .

Example

See whiteboard.

Relations

Definition

Let A and B be sets. A **binary relation** from A to B is a subset of $A \times B$.

Example (trivial relations)

$R = \emptyset$ and $R = A \times B$ are relations from A to B .

Example

See whiteboard.

Notation

We shall use $(a, b) \in R$ and $a R b$.

The slides for the 6th ed. of Rosen's textbook use $\langle a, b \rangle$.

Relations

Relations and functions

The functions are relations with additional constraints.

Relations

Definition

A **relation on** a set A is a relation from A to A .

Relations

Definition

A **relation on** a set A is a relation from A to A .

Example

Some relations on $\mathbb{Z}$:

$$R_1 = \{ (a, b) \in \mathbb{Z} \times \mathbb{Z} \mid a \leq b \},$$

$$R_2 = \{ (a, b) \in \mathbb{Z} \times \mathbb{Z} \mid a > b \},$$

$$R_3 = \{ (a, b) \in \mathbb{Z} \times \mathbb{Z} \mid a = b \vee a = -b \},$$

$$R_4 = \{ (a, b) \in \mathbb{Z} \times \mathbb{Z} \mid a = b \},$$

$$R_5 = \{ (a, b) \in \mathbb{Z} \times \mathbb{Z} \mid a = b + 1 \},$$

$$R_6 = \{ (a, b) \in \mathbb{Z} \times \mathbb{Z} \mid a + b \leq 3 \}.$$

Properties of Relations

Definition

Let R be a relation on a set A . The relation R is

reflexive iff $\forall x(xRx)$,

symmetric iff $\forall x\forall y(xRy \rightarrow yRx)$,

antisymmetric iff $\forall x\forall y((xRy \wedge yRx) \rightarrow x = y)$ and

transitive iff $\forall x\forall y\forall z((xRy \wedge yRz) \rightarrow xRz)$.

Properties of Relations

Example

See slides § 8.1, p. 5 for the 6th ed. of Rosen's textbook.

Combing Relations

See slides § 8.1, pp. 6-8 for the 6th ed. of Rosen's textbook.

Combing Relations

Definition

Let R be a relation from A to B and let S be a relation from B to C .

The **composition** of S with R , denoted $S \circ R$, is the relation from A to C where if $(a, b) \in R$ and $(b, c) \in S$ then $(a, c) \in S \circ R$.

Combing Relations

Example (composition of relations)

Let $A = \{1, 2, 3\}$, $B = \{1, 2, 3, 4\}$ and $C = \{0, 1, 2\}$.

Let R and S be the relations from A to B and B to C , respectively, given by

$$R = \{(1, 1), (1, 4), (2, 3), (3, 1), (3, 4)\},$$

$$S = \{(1, 0), (2, 0), (3, 1), (3, 2), (4, 1)\},$$

then $S \circ R$ is the relation from A to C , given by

$$S \circ R = \{(1, 0), (1, 1), (2, 1), (2, 2), (3, 0), (3, 1)\}.$$

Combing Relations

Problem 31 (p. 448)

Let R be the relation on the set of people consisting of pairs (a, b) , where a is a parent of b . Let S be the relation on the set of people consisting of pairs (a, b) , where a and b are siblings (brothers or sisters).

Combing Relations

Problem 31 (p. 448)

Let R be the relation on the set of people consisting of pairs (a, b) , where a is a parent of b . Let S be the relation on the set of people consisting of pairs (a, b) , where a and b are siblings (brothers or sisters).

What are $S \circ R$ and $R \circ S$?

Combing Relations

Problem 31 (p. 448)

Let R be the relation on the set of people consisting of pairs (a, b) , where a is a parent of b . Let S be the relation on the set of people consisting of pairs (a, b) , where a and b are siblings (brothers or sisters).

What are $S \circ R$ and $R \circ S$?

- $(a, b) \in S \circ R$ if exists c such that $(a, c) \in R$ (a is parent of c) and $(c, b) \in S$ (c is sibling of b), that is

$$S \circ R = \{ (a, b) \mid a \text{ is a parent of } b \text{ and } b \text{ has a sibling} \}.$$

Combing Relations

Problem 31 (p. 448)

Let R be the relation on the set of people consisting of pairs (a, b) , where a is a parent of b . Let S be the relation on the set of people consisting of pairs (a, b) , where a and b are siblings (brothers or sisters).

What are $S \circ R$ and $R \circ S$?

- $(a, b) \in S \circ R$ if exists c such that $(a, c) \in R$ (a is parent of c) and $(c, b) \in S$ (c is sibling of b), that is

$$S \circ R = \{ (a, b) \mid a \text{ is a parent of } b \text{ and } b \text{ has a sibling} \}.$$

- $(a, b) \in R \circ S$ if exists c such $(a, c) \in S$ (a is sibling of c) and $(c, b) \in R$ (c is parent of b), that is

$$R \circ S = \{ (a, b) \mid a \text{ is an aunt or uncle of } b \}.$$

Combing Relations

Definition

Let R be a relation on the set A . The **powers** R^n , for $n \in \mathbb{Z}^+$ are defined recursively by

$$\begin{aligned}R^1 &= R, \\ R^{n+1} &= R^n \circ R.\end{aligned}$$

Combing Relations

Definition

Let R be a relation on the set A . The **powers** R^n , for $n \in \mathbb{Z}^+$ are defined recursively by

$$\begin{aligned}R^1 &= R, \\ R^{n+1} &= R^n \circ R.\end{aligned}$$

Example

See slides § 8.1, pp. 9-10 for the 6th ed. of Rosen's textbook.

Combing Relations

Theorem 1 (p. 446)

Let R be a relation on a set A . The relation R is transitive iff $R^n \subseteq R$ for $n \in \mathbb{Z}^+$.

Proved on next slides

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.
2. Basis step $P(1)$: $R^1 = R \subseteq R$

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.
2. Basis step $P(1)$: $R^1 = R \subseteq R$
3. Inductive step:
Inductive hypothesis $P(k)$: if R is transitive implies $R^k \subseteq R$

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.

2. Basis step $P(1)$: $R^1 = R \subseteq R$

3. Inductive step:

Inductive hypothesis $P(k)$: if R is transitive implies $R^k \subseteq R$

Let's prove $P(k+1)$:

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.

2. Basis step $P(1)$: $R^1 = R \subseteq R$

3. Inductive step:

Inductive hypothesis $P(k)$: if R is transitive implies $R^k \subseteq R$

Let's prove $P(k+1)$:

3.1 Let $(a, b) \in R^{k+1}$.

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.

2. Basis step $P(1)$: $R^1 = R \subseteq R$

3. Inductive step:

Inductive hypothesis $P(k)$: if R is transitive implies $R^k \subseteq R$

Let's prove $P(k+1)$:

3.1 Let $(a, b) \in R^{k+1}$.

3.2 Exists $x \in A$ such that $(a, x) \in R^k$ and $(x, b) \in R$ (definition of $\circ$).

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.

2. Basis step $P(1)$: $R^1 = R \subseteq R$

3. Inductive step:

Inductive hypothesis $P(k)$: if R is transitive implies $R^k \subseteq R$

Let's prove $P(k+1)$:

3.1 Let $(a, b) \in R^{k+1}$.

3.2 Exists $x \in A$ such that $(a, x) \in R^k$ and $(x, b) \in R$ (definition of $\circ$).

3.3 $(a, x) \in R$ (IH).

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.

2. Basis step $P(1)$: $R^1 = R \subseteq R$

3. Inductive step:

Inductive hypothesis $P(k)$: if R is transitive implies $R^k \subseteq R$

Let's prove $P(k+1)$:

3.1 Let $(a, b) \in R^{k+1}$.

3.2 Exists $x \in A$ such that $(a, x) \in R^k$ and $(x, b) \in R$ (definition of $\circ$).

3.3 $(a, x) \in R$ (IH).

3.4 If $(a, x) \in R$ and $(x, b) \in R$ then $(a, b) \in R$ (R is transitive).

Combing Relations

Proof of $\Rightarrow$ (if R is transitive implies $R^n \subseteq R$ for $n \in \mathbb{Z}^+$).

By induction on $n \in \mathbb{Z}^+$.

1. $P(n)$: if R is transitive implies $R^n \subseteq R$.

2. Basis step $P(1)$: $R^1 = R \subseteq R$

3. Inductive step:

Inductive hypothesis $P(k)$: if R is transitive implies $R^k \subseteq R$

Let's prove $P(k+1)$:

3.1 Let $(a, b) \in R^{k+1}$.

3.2 Exists $x \in A$ such that $(a, x) \in R^k$ and $(x, b) \in R$ (definition of $\circ$).

3.3 $(a, x) \in R$ (IH).

3.4 If $(a, x) \in R$ and $(x, b) \in R$ then $(a, b) \in R$ (R is transitive).

3.5 $R^{k+1} \subseteq R$. ■

Continued on next slide

Combing Relations

Proof of $\Leftarrow$ (if $R^n \subseteq R$ for $n \in \mathbb{Z}^+$ implies R is transitive).

- 1 Suppose that $(a, b) \in R$ and $(b, c) \in R$.
- 2 $(a, c) \in R^2$. (def. of R^2)
- 3 $(a, c) \in R$. ($R^2 \subseteq R$)
- 4 Therefore, R is transitive. ■

Inverse and Complementary Relations

Definition

Let R be a relation from A to B . The **inverse relation** from B to A , denoted by R^{-1} , is the set of ordered pairs

$$R^{-1} = \{ (b, a) \in B \times A \mid (a, b) \in R \}.$$

Inverse and Complementary Relations

Definition

Let R be a relation from A to B . The **complementary relation** from A to B , denoted by $\overline{R}$, is the set of ordered pairs

$$\overline{R} = \{ (a, b) \in A \times B \mid (a, b) \notin R \}.$$

References

Kenneth H. Rosen [1988] (2004). Matemática Discreta y sus Aplicaciones. Trans. by José Manuel Pérez Morales, Julio Moro Carreño, Ana Isabel Lías Quintero and Pedro Antonio Ramos Alonc. 5th ed. McGraw-Hill (cit. on p. 2).