

CM0246 Discrete Structures

Boolean Algebras

Andrés Sicard-Ramírez

Universidad EAFIT

Semestre 2014-2

(Last modification: 8th April 2026)

Preliminaries

Textbook

Kenneth H. Rosen ([1988] 2004). Matemática Discreta y sus Aplicaciones.

Convention

The numbers and page numbers assigned to chapters, examples, exercises, figures, quotes, sections and theorems on these slides correspond to the numbers assigned in the textbook.

Boolean Functions

Boolean operations

We define the following operations in the set $B = \{0, 1\}$:

- Boolean sum

$$0 + 0 = 0, 0 + 1 = 1, 1 + 0 = 1 \text{ and } 1 + 1 = 1.$$

- Boolean product

$$0 \cdot 0 = 0, 0 \cdot 1 = 0, 1 \cdot 0 = 0 \text{ and } 1 \cdot 1 = 1.$$

- Complement

$$\bar{0} = 1 \text{ and } \bar{1} = 0.$$

Boolean Functions

Boolean operations

We define the following operations in the set $B = \{0, 1\}$:

- Boolean sum

$$0 + 0 = 0, 0 + 1 = 1, 1 + 0 = 1 \text{ and } 1 + 1 = 1.$$

- Boolean product

$$0 \cdot 0 = 0, 0 \cdot 1 = 0, 1 \cdot 0 = 0 \text{ and } 1 \cdot 1 = 1.$$

- Complement

$$\overline{0} = 1 \text{ and } \overline{1} = 0.$$

Precedence (highest to lowest): Complement, Boolean product and Boolean sum.

Boolean Functions

Boolean operations

We define the following operations in the set $B = \{0, 1\}$:

- Boolean sum

$$0 + 0 = 0, 0 + 1 = 1, 1 + 0 = 1 \text{ and } 1 + 1 = 1.$$

- Boolean product

$$0 \cdot 0 = 0, 0 \cdot 1 = 0, 1 \cdot 0 = 0 \text{ and } 1 \cdot 1 = 1.$$

- Complement

$$\overline{0} = 1 \text{ and } \overline{1} = 0.$$

Precedence (highest to lowest): Complement, Boolean product and Boolean sum.

Example

Whiteboard.

Boolean Functions

From Boolean operations/logical operators to logical operators/Boolean operations

Boolean operations	logic operators
$\cdot$	$\wedge$
$+$	$\vee$
$-$	$\neg$
0	F
1	T

Boolean Functions

From Boolean operations/logical operators to logical operators/Boolean operations

Boolean operations	logic operators
$\cdot$	$\wedge$
$+$	$\vee$
$-$	$\neg$
0	F
1	T

Example (from equality/logical equivalence to logical equivalence/equality)
Whiteboard.

Boolean Functions

Definition

Let $B = \{0, 1\}$. A function from B^n to B is called a **Boolean function of degree n** .

Boolean Functions

Definition

Let $B = \{0, 1\}$. A function from B^n to B is called a **Boolean function of degree n** .

Example

Whiteboard.

Boolean Functions

Theorem (Example 5, p. 280)

If $|A| = m$ and $|B| = n$ then $|\{f : A \rightarrow B\}| = n^m$.

Boolean Functions

Theorem (Example 5, p. 280)

If $|A| = m$ and $|B| = n$ then $|\{f : A \rightarrow B\}| = n^m$.

Example

There are 16 Boolean functions of degree 2.

Boolean Functions

Definition

Let $x_1, x_2, \dots, x_n$ be Boolean variables. The **Boolean expressions** are inductively defined by

- Basis step: 0 , 1 and $x_1, x_2, \dots, x_n$ are Boolean expressions.
- Inductive step: If E_1 and E_2 are Boolean expressions then $\overline{E_1}$, $(E_1 \cdot E_2)$ and $(E_1 + E_2)$ are Boolean expressions.

Boolean Functions

Definition

Let $x_1, x_2, \dots, x_n$ be Boolean variables. The **Boolean expressions** are inductively defined by

- Basis step: 0, 1 and $x_1, x_2, \dots, x_n$ are Boolean expressions.
- Inductive step: If E_1 and E_2 are Boolean expressions then $\overline{E_1}$, $(E_1 \cdot E_2)$ and $(E_1 + E_2)$ are Boolean expressions.

Each Boolean expression represents a Boolean function.

Example

Whiteboard.

Logical Equivalences

Identity laws

$$p \wedge \text{T} \equiv p$$

$$p \vee \text{F} \equiv p$$

Domination laws

$$p \wedge \text{F} \equiv \text{F}$$

$$p \vee \text{T} \equiv \text{T}$$

Idempotent laws

$$p \wedge p \equiv p$$

$$p \vee p \equiv p$$

Double negation law

$$\neg(\neg p) \equiv p$$

Commutative laws

$$p \wedge q \equiv q \wedge p$$

$$p \vee q \equiv q \vee p$$

Associate laws

$$(p \wedge q) \wedge r \equiv p \wedge (q \wedge r)$$

$$(p \vee q) \vee r \equiv p \vee (q \vee r)$$

Distributive laws

$$p \wedge (q \vee r) \equiv (p \wedge q) \vee (p \wedge r)$$

$$p \vee (q \wedge r) \equiv (p \vee q) \wedge (p \vee r)$$

De Morgan's laws

$$\neg(p \wedge q) \equiv \neg p \vee \neg q$$

$$\neg(p \vee q) \equiv \neg p \wedge \neg q$$

Absorption laws

$$p \wedge (p \vee q) \equiv p$$

$$p \vee (p \wedge q) \equiv p$$

Negation laws

$$p \wedge \neg p \equiv \text{F}$$

$$p \vee \neg p \equiv \text{T}$$

Boolean Identities

Identity laws

$$x \cdot 1 = x$$

$$x + 0 = x$$

Domination laws

$$x \cdot 0 = 0$$

$$x + 1 = 1$$

Idempotent laws

$$x \cdot x = x$$

$$x + x = x$$

Double complement law

$$\overline{\overline{x}} = x$$

Commutative laws

$$x \cdot y = y \cdot x$$

$$x + y = y + x$$

Associate laws

$$(x \cdot y) \cdot z = x \cdot (y \cdot z)$$

$$(x + y) + z = x + (y + z)$$

Distributive laws

$$x \cdot (y + z) = (x \cdot y) + (x \cdot z)$$

$$x + (y \cdot z) = (x + y) \cdot (x + z)$$

De Morgan's laws

$$\overline{x \cdot y} = \overline{x} + \overline{y}$$

$$\overline{(x + y)} = \overline{x} \cdot \overline{y}$$

Absorption laws

$$x \cdot (x + y) = x$$

$$x + x \cdot y = x$$

Complement laws

$$x \cdot \overline{x} = 0$$

$$x + \overline{x} = 1$$

Boolean Identities

Each Boolean identity can be proved using a table.

Example

Whiteboard.

Boolean Algebras

Definition

Let $\wedge$ and $\vee$ be two binaries operations, $\neg$ a unary operation and 0 and 1 two constants. A **Boolean algebra** is an algebraic structure $(B, \wedge, \vee, \neg, 0, 1)$, which satisfy the following **axioms** for all x, y and z in B :

Identity laws

$$x \wedge 1 = x$$

$$x \vee 0 = x$$

Complement laws

$$x \wedge \bar{x} = 0$$

$$x \vee \bar{x} = 1$$

Associate laws

$$(x \wedge y) \wedge z = x \wedge (y \wedge z)$$

$$(x \vee y) \vee z = x \vee (y \vee z)$$

Commutative laws

$$x \wedge y = y \wedge x$$

$$x \vee y = y \vee x$$

Distributive laws

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

$$x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$$

Boolean Algebra

Example

$(B, \wedge, \vee, -, 0, 1)$	Set theory	Propositional logic	$(\{0, 1\}, \cdot, +, -, 0, 1)$
B	U	set of formulae	$\{0, 1\}$
$\wedge$	$\cap$	$\wedge$	$\cdot$
$\vee$	$\cup$	$\vee$	$+$
$-$	$-$	$\neg$	$-$
0	$\emptyset$	F	0
1	U	T	1

Boolean Algebra

Definition

The **dual** of any statement in a Boolean algebra $(B, \wedge, \vee, -, 0, 1)$ is the statement obtained by interchanging $\wedge$ and $\vee$, and interchanging 0 and 1.

Boolean Algebra

Definition

The **dual** of any statement in a Boolean algebra $(B, \wedge, \vee, -, 0, 1)$ is the statement obtained by interchanging $\wedge$ and $\vee$, and interchanging 0 and 1.

Example

The dual of $x \wedge (y \vee 0)$ is $x \vee (y \wedge 1)$.

Boolean Algebra

Definition

The **dual** of any statement in a Boolean algebra $(B, \wedge, \vee, -, 0, 1)$ is the statement obtained by interchanging $\wedge$ and $\vee$, and interchanging 0 and 1.

Example

The dual of $x \wedge (y \vee 0)$ is $x \vee (y \wedge 1)$.

Theorem (Principle of duality. Problem 38, p. 660)

The dual of any theorem in a Boolean algebra is also an theorem.

Boolean Algebra

Definition

The **dual** of any statement in a Boolean algebra $(B, \wedge, \vee, -, 0, 1)$ is the statement obtained by interchanging $\wedge$ and $\vee$, and interchanging 0 and 1.

Example

The dual of $x \wedge (y \vee 0)$ is $x \vee (y \wedge 1)$.

Theorem (Principle of duality. Problem 38, p. 660)

The dual of any theorem in a Boolean algebra is also an theorem.

Proof.

Similar to the proof of the principle of duality for lattices. ■

Boolean Algebra

Problem 31 (p. 660)

Let $\mathbb{B} = (B, \wedge, \vee, ^-, 0, 1)$ be a Boolean algebra. To prove that $\mathbb{B}$ satisfy the idempotent laws $x \vee x = x$ and $x \wedge x = x$, for every element x .

Boolean Algebra

Problem 31 (p. 660)

Let $\mathbb{B} = (B, \wedge, \vee, -, 0, 1)$ be a Boolean algebra. To prove that $\mathbb{B}$ satisfy the idempotent laws $x \vee x = x$ and $x \wedge x = x$, for every element x .

Proof.

$$\begin{aligned}x \vee x &= (x \vee x) \wedge 1 && \text{(identity law)} \\&= (x \vee x) \wedge (x \vee \bar{x}) && \text{(complement law)} \\&= x \vee (x \wedge \bar{x}) && \text{(distributive law)} \\&= x \vee 0 && \text{(complement law)} \\&= x && \text{(identity law)}\end{aligned}$$

Boolean Algebra

Problem 31 (p. 660)

Let $\mathbb{B} = (B, \wedge, \vee, ^-, 0, 1)$ be a Boolean algebra. To prove that $\mathbb{B}$ satisfy the idempotent laws $x \vee x = x$ and $x \wedge x = x$, for every element x .

Proof.

$x \vee x = (x \vee x) \wedge 1$	(identity law)	$x \wedge x = (x \wedge x) \vee 0$
$= (x \vee x) \wedge (x \vee \bar{x})$	(complement law)	$= (x \wedge x) \vee (x \wedge \bar{x})$
$= x \vee (x \wedge \bar{x})$	(distributive law)	$= x \wedge (x \vee \bar{x})$
$= x \vee 0$	(complement law)	$= x \wedge 1$
$= x$	(identity law)	$= x$



Boolean Algebra

Problem 31 (p. 660)

Let $\mathbb{B} = (B, \wedge, \vee, ^-, 0, 1)$ be a Boolean algebra. To prove that $\mathbb{B}$ satisfy the idempotent laws $x \vee x = x$ and $x \wedge x = x$, for every element x .

Proof.

$x \vee x = (x \vee x) \wedge 1$	(identity law)	$x \wedge x = (x \wedge x) \vee 0$
$= (x \vee x) \wedge (x \vee \bar{x})$	(complement law)	$= (x \wedge x) \vee (x \wedge \bar{x})$
$= x \vee (x \wedge \bar{x})$	(distributive law)	$= x \wedge (x \vee \bar{x})$
$= x \vee 0$	(complement law)	$= x \wedge 1$
$= x$	(identity law)	$= x$



Remark: $x \wedge x = x/x \vee x = x$ also follows from $x \vee x = x/x \wedge x = x$ by the principle of duality.

Boolean Algebra

Problem 34 (p. 660)

Let $\mathbb{B} = (B, \wedge, \vee, ^-, 0, 1)$ be a Boolean algebra. To prove that $\mathbb{B}$ satisfy the double complement law, i.e. $\forall x(x = \overline{\overline{x}})$.

Boolean Algebra

Problem 34 (p. 660)

Let $\mathbb{B} = (B, \wedge, \vee, ^-, 0, 1)$ be a Boolean algebra. To prove that $\mathbb{B}$ satisfy the double complement law, i.e. $\forall x(x = \overline{\overline{x}})$.

Hint: From the complement laws, we have

$$\overline{x} \wedge \overline{\overline{x}} = 0,$$

$$\overline{x} \vee \overline{\overline{x}} = 1.$$

Proved on next slide

Boolean Algebra

Proof (Lipschutz 1994).

$x = x \vee 0$	(identity law)
$= x \vee (\overline{x} \wedge \overline{\overline{x}})$	(complement law)
$= (x \vee \overline{x}) \wedge (x \vee \overline{\overline{x}})$	(distributive law)
$= 1 \wedge (x \vee \overline{\overline{x}})$	(complement law)
$= (\overline{x} \vee \overline{\overline{x}}) \wedge (x \vee \overline{\overline{x}})$	(complement law)
$= (\overline{\overline{x}} \vee \overline{x}) \wedge (\overline{\overline{x}} \vee x)$	(commutative law)
$= \overline{\overline{x}} \vee (\overline{x}) \wedge x$	(distributive law)
$= \overline{\overline{x}} \vee (x \wedge \overline{x})$	(commutative law)
$= \overline{\overline{x}} \vee 0$	(complement law)
$= \overline{\overline{x}}$	(identity law) 

References

- Seymour Lipschutz (1994). Teoría de Conjuntos y Temas Afines. Serie Schaum. McGraw-Hill (cit. on p. 29).
- Kenneth H. Rosen [1988] (2004). Matemática Discreta y sus Aplicaciones. Trans. by José Manuel Pérez Morales, Julio Moro Carreño, Ana Isabel Lías Quintero and Pedro Antonio Ramos Alonc. 5th ed. McGraw-Hill (cit. on p. 2).