

CM0832 Elements of Set Theory

4. Natural Numbers

Andrés Sicard-Ramírez

Universidad EAFIT

Semester 2017-2

Preliminaries

Convention

The number assigned to chapters, examples, exercises, figures, sections, and theorems on these slides correspond to the numbers assigned in the textbook [Enderton 1977].

Defining the Natural Numbers

Approaches for introducing mathematical objects

- Axiomatic
- Definitional

Defining the Natural Numbers

Approaches for introducing mathematical objects

- Axiomatic
- Definitional

Definitional approach for introducing natural numbers

- We shall define natural numbers in terms of sets.
- We shall prove the properties of natural numbers from properties of sets.

Defining the Natural Numbers

Approaches for introducing mathematical objects

- Axiomatic
- Definitional

Definitional approach for introducing natural numbers

- We shall define natural numbers in terms of sets.
- We shall prove the properties of natural numbers from properties of sets.

Question

How to define natural numbers in terms of sets?

Inductive Sets

von Neumann's construction

Informal idea: A natural number is the set of all smaller natural numbers

$$0 := \emptyset,$$

$$1 := \{0\} = \{\emptyset\},$$

$$2 := \{0, 1\} = \{\emptyset, \{\emptyset\}\},$$

$$3 := \{0, 1, 2\} = \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\},$$

$$\vdots$$

Inductive Sets

von Neumann's construction

Informal idea: A natural number is the set of all smaller natural numbers

$$\begin{aligned}0 &:= \emptyset, \\1 &:= \{0\} = \{\emptyset\}, \\2 &:= \{0, 1\} = \{\emptyset, \{\emptyset\}\}, \\3 &:= \{0, 1, 2\} = \{\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}\}, \\&\vdots\end{aligned}$$

Some 'extra' properties

$$0 \in 1 \in 2 \in 3 \in \cdots \quad \text{and} \quad 0 \subseteq 1 \subseteq 2 \subseteq 3 \subseteq \cdots.$$

Inductive Sets

A wrong impredicative definition

$$n := \{0, 1, \dots, n - 1\}.$$

‘We cannot just say that a set n is a natural number if its elements are all the smaller natural numbers, because such a “definition” would involve the very concept being defined.’ [Hrbacek and Jech (1978) 1999, p. 40]

Inductive Sets

Definition

Let a be a set. The **successor** of a is

$$a^+ := a \cup \{a\}.$$

Inductive Sets

Definition

Let a be a set. The **successor** of a is

$$a^+ := a \cup \{a\}.$$

Example

$$0 = \emptyset,$$

$$1 = \emptyset^+,$$

$$2 = \emptyset^{++},$$

$$3 = \emptyset^{+++},$$

$$\vdots$$

Inductive Sets

Definition

A set A is **inductive** iff

- $\emptyset \in A$ and
- if $a \in A$ then $a^+ \in A$.

Inductive Sets

Definition

A set A is **inductive** iff

- $\emptyset \in A$ and
- if $a \in A$ then $a^+ \in A$.

Observation

An inductive is an infinite set.

Inductive Sets

Definition

A set A is **inductive** iff

- $\emptyset \in A$ and
- if $a \in A$ then $a^+ \in A$.

Observation

An inductive is an infinite set.

Question

Are there inductive sets?

Infinite Axiom

Infinity axiom

There exists an inductive set, that is,

$$\exists A [\emptyset \in A \wedge \forall a (a \in A \rightarrow a^+ \in A)].$$

The Set of Natural Numbers

Definition

A **natural number** is a set that belongs to every inductive set.

The Set of Natural Numbers

Definition

A **natural number** is a set that belongs to every inductive set.

Theorem 4A

There is a set whose members are exactly the natural numbers.

The Set of Natural Numbers

Definition

A **natural number** is a set that belongs to every inductive set.

Theorem 4A

There is a set whose members are exactly the natural numbers.

Proof.

Let A be an inductive set. By the subset axiom scheme, there is a set

$$\{x \in A \mid x \in I \text{ for every inductive set } I\}.$$



The Set of Natural Numbers

Definition

The set of **all natural numbers**, denoted by ω , is defined by

$$\omega := \{ x \in A \mid x \in I \text{ for every inductive set } I \}.$$

That is,

$$x \in \omega \quad \text{iff} \quad x \text{ is a natural number.}$$

The Set of Natural Numbers

Theorem 4B

The set ω is inductive, and it is a subset of every other inductive set.

The Set of Natural Numbers

Theorem 4B

The set ω is inductive, and it is a subset of every other inductive set.

Observation

The set ω is the **smallest** inductive set.

The Set of Natural Numbers

Observation

Since that the collection of all inductive sets is not a set but a proper class, using class we could define the set of natural numbers by

$$\omega := \bigcap \{ A \mid A \text{ is an inductive set} \}.$$

The Set of Natural Numbers

Observation

Since that the collection of all inductive sets is not a set but a proper class, using class we could define the set of natural numbers by

$$\omega := \bigcap \{ A \mid A \text{ is an inductive set} \}.$$

Observation

Mendelson [(1973) 2008] in the proof of Theorem ZFC 8 defines the set ω as an intersection of some inductive sets.

Induction Principle for Natural Numbers

Induction principle for ω (p. 69)

Any inductive subset of ω coincides with ω .

Induction Principle for Natural Numbers

Induction principle for ω (p. 69)

Any inductive subset of ω coincides with ω .

Induction principle for ω (other version) [Hrbacek and Jech (1978) 1999]

Let $P(x)$ be a property. Assume that

- (i) $P(0)$ holds,
- (ii) for all $n \in \omega$, $P(n)$ implies $P(n^+)$.

Then P holds for all natural numbers n .

Proof.

'This is an immediate consequence of our definition of ω . The assumptions i) and ii) simply say that the set $A = \{n \in \omega \mid P(n)\}$ is inductive. $\omega \subseteq A$ follows.' [Hrbacek and Jech (1978) 1999, p. 42] 

Defining Natural Numbers as Sets

Observation

So far, we defined natural numbers on terms of sets. A different point of view is stated by some authors (see, e.g. Benacerraf [1965]).



‘Thus inductive definability is a notion intermediate in strength between predicate and fully impredicative definability. It would be interesting to formulate a coherent conceptual framework that made induction the principal notion. There are suggestions of this in the literature, but the possibility has not yet been fully explored.’ [Aczel 1977, p. 780]

Transitive Sets

Definition

Let A be a set. The set A is a **transitive set** iff every member of a member of A is itself a member of A , that is,

$$x \in a \in A \text{ implies } x \in A.$$

Transitive Sets

Definition

Let A be a set. The set A is a **transitive set** iff every member of a member of A is itself a member of A , that is,

$$x \in a \in A \text{ implies } x \in A.$$

Example

Whiteboard.

Transitive Sets

Theorem

A set A is a transitive set iff $\bigcup A \subseteq A$.

(continued on next slide)

Transitive Sets

Proof.

i) (Only if) Let A be a transitive set. Then

$$\begin{aligned}x \in \bigcup A &\Rightarrow \exists b (x \in b \wedge b \in A) \\&\Rightarrow x \in A\end{aligned}$$

(by definition of $\bigcup A$)
(because A is transitive)

ii) (If) Let $\bigcup A \subseteq A$. Then

$$\begin{aligned}x \in a \wedge a \in A &\Rightarrow x \in \bigcup A \\&\Rightarrow x \in A\end{aligned}$$

(by definition of $\bigcup A$)
(because $\bigcup A \subseteq A$)



Transitive Sets

Theorem

A set A is a transitive set iff $a \in A$ implies $a \subseteq A$.

Transitive Sets

Theorem

A set A is a transitive set iff $a \in A$ implies $a \subseteq A$.

Proof.

- i) (Only if) Let A be a transitive set and let $a \in A$. If $x \in a$ implies $x \in A$ because A is transitive.
- ii) (If) Let $a \in A$ implies $a \subseteq A$. If $x \in a \wedge a \in A$ implies $x \in A$ because $a \subseteq A$.



Transitive Sets

Theorem

A set A is a transitive set iff $A \subseteq \mathcal{P}A$.

Transitive Sets

On transitive sets

Let A be a set. Transitive sets can be defined using any of the followings equivalent affirmations:

- (i) $x \in a \in A$ implies $x \in A$,
- (ii) $\bigcup A \subseteq A$,
- (iii) $a \in A$ implies $a \subseteq A$,
- (iv) $A \subseteq \mathcal{P}A$.

Transitive Sets

Theorem 4E

If a is a transitive set, then $\bigcup(a^+) = a$.

Transitive Sets

Theorem 4E

If a is a transitive set, then $\bigcup(a^+) = a$.

Theorem 4F

Every natural number is a transitive set.

Transitive Sets

Theorem 4E

If a is a transitive set, then $\bigcup(a^+) = a$.

Theorem 4F

Every natural number is a transitive set.

Theorem 4G

The set ω is a transitive set.

Recursion on Natural Numbers

Recursion theorem on ω (p. 73)

Let A be a set, $a \in A$ and $F : A \rightarrow A$. Then there exists a unique function h such that

$$\begin{aligned}h &: \omega \rightarrow A \\h(0) &= a, \\h(n^+) &= F(h(n)), \text{ for all } n \in \omega.\end{aligned}$$

Arithmetic

Idea

We shall apply the recursion theorem to define addition and multiplication on ω .

Arithmetic

Example

We want to define the function

$$A_5 : w \rightarrow w$$
$$A_5(n) = \text{addition of } 5 \text{ to } n.$$

Arithmetic

Example

We want to define the function

$$\begin{aligned} A_5 : w &\rightarrow w \\ A_5(n) &= \text{addition of } 5 \text{ to } n. \end{aligned}$$

Let $F : \omega \rightarrow \omega := n \mapsto n^+$. By the recursion theorem there exists a unique function

$$\begin{aligned} A_5 : w &\rightarrow w \\ A_5(0) &= 5, \\ A_5(n^+) &= (A_5(n))^+. \end{aligned}$$

Arithmetic

Example

Let $m \in \omega$. By the recursion theorem there exists a unique function

$$\begin{aligned}A_m &: w \rightarrow w \\A_m(0) &= m, \\A_m(n^+) &= (A_m(n))^+.\end{aligned}$$

Arithmetic

Definition

Let m and n be natural numbers. We define the **addition** of m and n by

$$\begin{aligned} (+) : w \times w &\rightarrow w \\ m + n &= A_m(n). \end{aligned}$$

Arithmetic

Theorem 4I

Let m and n be natural numbers. Then

$$m + 0 = m,$$

$$m + n^+ = (m + n)^+.$$

Arithmetic

Example

Let $m \in \omega$. By the recursion theorem there exists a unique function

$$M_m : \omega \rightarrow \omega$$

$$M_m(0) = 0,$$

$$M_m(n^+) = M_m(n) + m.$$

Arithmetic

Definition

Let m and n be natural numbers. We define the **multiplication** of m and n by

$$\begin{aligned}(\cdot) : w \times w &\rightarrow w \\ m \cdot n &= M_m(n).\end{aligned}$$

Arithmetic

Theorem 4J

Let m and n be natural numbers. Then

$$m \cdot 0 = 0,$$

$$m \cdot n^+ = (m \cdot n) + m.$$

Ordering on Natural Numbers

Strong induction principle for ω (p. 87)

Let A be a subset of ω , and assume that for every n in ω ,

$$m < n \rightarrow m \in A \quad \text{implies} \quad n \in A.$$

Then $A = \omega$.

References



Aczel, Peter (1977). An Introduction to Inductive Definitions. In: Handbook of Mathematical Logic. Ed. by Barwise, Jon. Vol. 90. Studies in Logic and the Foundations of Mathematics. Elsevier. Chap. C.7. DOI: [10.1016/S0049-237X\(08\)71120-0](https://doi.org/10.1016/S0049-237X(08)71120-0) (cit. on p. 26).



Benacerraf, Paul (1965). What Numbers Could not Be. The Philosophical Review 74.1, pp. 47–73. DOI: [10.2307/2183530](https://doi.org/10.2307/2183530) (cit. on p. 25).



Enderton, Herbert B. (1977). Elements of Set Theory. Academic Press (cit. on p. 2).



Hrbacek, Karel and Jech, Thomas [1978] (1999). Introduction to Set Theory. Third Edition, Revised and Expanded. Marcel Dekker (cit. on pp. 8, 23, 24).



Mendelson, Elliott [1973] (2008). Number Systems and the Foundations of Analysis. Republication of the 1985 reprint. Dover Publications (cit. on pp. 21, 22).