MT8002 Elements of Set Theory
Chapter 7. Ordered Sets
Section 7.2. Linearly Ordered Sets

Andrés Sicard-Ramírez

Universidad EAFIT

Semester 2026-2

Preliminaries

Textbook

Derek Goldrei (1996). Classic Set Theory. A Guided Independent Study.

Convention

The numbers and page numbers assigned to chapters, examples, exercises, figures, quotes, sections and theorems on these slides correspond to the numbers assigned in the textbook.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Introduction

- We can use binary relations to order some (partial order) or all (linear order) the elements of a set.
- Orders are binary relations that satisfy certain properties.
- Weak orders and strict orders can be interchangeably defined.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Binary Relations

Definition (p. 164)

Let X be a set. A set R is a **binary relation on** X if $R \subseteq X \times X$.

Notation

Let R be a binary relation on a set X.

(a) We write $(x, y) \in R$ or xRy, if $x \in X$ is related to $y \in X$ by the relation R.

(b) We write $(x, y) \notin R$ or $x \not R y$, if $x \in X$ is not related to $y \in X$ by the relation R.

Weak Orders

Definitions (p. 164)

A binary relation R on a set X is a **weak partial order** on the set X (or the set X is **weakly partially ordered** by the relation R), if it has the following properties:

(a) **reflexive**: for all $x \in X$, xRx;

(b) **anti-symmetric**: for all $x, y \in X$, if xRy and yRx then $x \doteq y$;

(c) **transitive**: for all $x, y, z \in X$, if xRy and yRz then xRz.

Weak Orders

Definitions (p. 164)

A binary relation R on a set X is a **weak partial order** on the set X (or the set X is **weakly partially ordered** by the relation R), if it has the following properties:

(a) **reflexive**: for all $x \in X$, xRx;

(b) **anti-symmetric**: for all $x, y \in X$, if xRy and yRx then $x \doteq y$;

(c) **transitive**: for all $x, y, z \in X$, if xRy and yRz then xRz.

If additionally the binary relation R has the following property:

(d) **linear**: for all $x, y \in X$, xRy or yRx,

then the relation R is a **weak linear order** on the set X (or the set X is **weakly linearly ordered** by the relation R).

Weak Orders

Examples (informal, p. 164)

The usual relations $\leq$ and $\geq$ on the sets $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$ and $\mathbb{R}$ are weak linear orders on those sets.

Weak Orders

Examples (informal, p. 164)

The usual relations $\leq$ and $\geq$ on the sets $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$ and $\mathbb{R}$ are weak linear orders on those sets.

Example

Let X be a set. The relation $\subseteq$ on $\mathcal{P}(X)$ is a weak partial order.

Weak Orders

Exercise

Let X be a set. Is the relation $\subseteq$ on $\mathcal{P}(X)$ a weak linear order?

Weak Orders

Exercise

Let X be a set. Is the relation $\subseteq$ on $\mathcal{P}(X)$ a weak linear order?

Answer

No! For example the relation $\subseteq$ on the power set of $\{a, b\}$ is no linear because $\{a\}, \{b\} \in \mathcal{P}(\{a, b\})$ but neither $\{a\} \nsubseteq \{b\}$ not $\{b\} \nsubseteq \{a\}$.

Weak Orders

Description

The **restriction** of a weak partial order to a subset is the weak partial order obtained by discarding all comparisons involving elements outside of the subset.

Weak Orders

Description

The **restriction** of a weak partial order to a subset is the weak partial order obtained by discarding all comparisons involving elements outside of the subset.

Definition (p. 164)

Let R be a weak partial order on a set X and let $A \subseteq X$. The **restriction** of the relation R to the set A, denoted $R|_{A\times A}$, is the relation defined by

$$R|_{A\times A} := R \cap A \times A \doteq \{(a,b) \in R : a, b \in A\}.$$

Weak Orders

Description

The **restriction** of a weak partial order to a subset is the weak partial order obtained by discarding all comparisons involving elements outside of the subset.

Definition (p. 164)

Let R be a weak partial order on a set X and let $A \subseteq X$. The **restriction** of the relation R to the set A, denoted $R|_{A\times A}$, is the relation defined by

$$R|_{A\times A} := R \cap A \times A \doteq \{(a,b) \in R : a, b \in A\}.$$

Theorem

Let R be a weak partial (respectively linear) order on a set X and let $A \subseteq X$. Then $R|_{A\times A}$ is a partial (respectively linear) order on the set A (Exercise 7.2).

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Strict Orders

Definitions (p. 165)

A binary relation S on a set X is a **strict partial order** on X (or the set X is **strictly partially ordered** by the relation S) if it has the following properties:

(a) **irreflexive**: for all $x \in X$, $x \not S x$;

(b) **transitive**: for all $x, y, z \in X$, if xSy and ySz then xSz.

Strict Orders

Definitions (p. 165)

A binary relation S on a set X is a **strict partial order** on X (or the set X is **strictly partially ordered** by the relation S) if it has the following properties:

(a) **irreflexive**: for all $x \in X$, $x \not S x$;

(b) **transitive**: for all $x, y, z \in X$, if xSy and ySz then xSz.

If additionally the binary relation S has the following property:

(c) **linear**: for all $x, y \in X$, xSy or $x \doteq y$ or ySx,

then the relation S is a **strict linear order** on the set X (or the set X is **strictly linearly ordered** by the relation S).

Strict Orders

Examples

The usual relations $<$ and $>$ on the sets $\mathbb{N}$, $\mathbb{Z}$, $\mathbb{Q}$ and $\mathbb{R}$ are strict linear orders on those sets.

Example

Let X be a set. The relation $\subset$ on $\mathcal{P}(X)$ is a strict partial order.

Strict Orders

Theorem (relationship between weak and strict partial orders)

(a) Let R be a weak partial order on a set X. The binary relation S defined by

$$xSy := xRy \text{ and } x \neq y$$

is a strict partial order on the set X.

Strict Orders

Theorem (relationship between weak and strict partial orders)

(a) Let R be a weak partial order on a set X. The binary relation S defined by

$$xSy := xRy \text{ and } x \neq y$$

is a strict partial order on the set X.

(b) Let S be a strict partial order on a set X. The binary relation R defined by

$$xRy := xSy \text{ or } x \doteq y$$

is a weak partial order on the set X.

Strict Orders

Theorem (relationship between weak and strict partial orders)

(a) Let R be a weak partial order on a set X. The binary relation S defined by

$$xSy := xRy \text{ and } x \neq y$$

is a strict partial order on the set X.

(b) Let S be a strict partial order on a set X. The binary relation R defined by

$$xRy := xSy \text{ or } x \doteq y$$

is a weak partial order on the set X.


See Exercise 7.4.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Orders on Natural Numbers

An "extra" property of natural numbers



Orders on Natural Numbers

Theorem 3.6

The set of natural numbers $\mathbb{N}$ is strictly linearly ordered by the binary relation $\in$.

Orders on Natural Numbers

Theorem 3.6

The set of natural numbers $\mathbb{N}$ is strictly linearly ordered by the binary relation $\in$.

That is, the binary relation $\in$ on $\mathbb{N}$ has the following properties:

(a) **irreflexive**: for all $\mathbf{n} \in \mathbb{N}$, $\mathbf{n} \notin \mathbf{n}$;

(b) **transitive**: for all $\mathbf{m}, \mathbf{n}, \mathbf{p} \in \mathbb{N}$, if $\mathbf{m} \in \mathbf{n}$ and $\mathbf{n} \in \mathbf{p}$ then $\mathbf{m} \in \mathbf{p}$;

(c) **linear**: for all $\mathbf{m}, \mathbf{n} \in \mathbb{N}$, $\mathbf{m} \in \mathbf{n}$ or $\mathbf{m} \doteq \mathbf{n}$ or $\mathbf{n} \in \mathbf{m}$.

Orders on Natural Numbers

Example (p. 42)

Although the binary relation $\in$ is transitive on the set $\mathbb{N}$ it does not mean that the relation is transitive on any set. For example, the relation is not transitive on the set

$$\{\mathbf{0}, \{\mathbf{0}\}, \{\{\mathbf{0}\}\}\}$$

because $\mathbf{0} \in \{\mathbf{0}\}$ and $\{\mathbf{0}\} \in \{\{\mathbf{0}\}\}$ but $\mathbf{0} \notin \{\{\mathbf{0}\}\}$.

Orders on Natural Numbers

Definition (p. 41)

For all $\mathbf{m}, \mathbf{n} \in \mathbb{N}$,

$\mathbf{m} < \mathbf{n} := \mathbf{m} \in \mathbf{n}$,
$\mathbf{m} \leq \mathbf{n} := \mathbf{m} < \mathbf{n}$ or $\mathbf{m} \doteq \mathbf{n}$.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Conventions

- "*We shall often write the order relation on X as $<_X$ or $\leq_X$, depending on whether the order is strict or weak.*" *(p. 166)*
- Sometimes a (weakly or strictly; partially or linearly) ordered set X by a relation R will we denote by the pair (X, R).†
- Joining above conventions we shall write:
  - "Let $(X, \leq_X)$ be a partially (respectively linearly) ordered set" when $\leq_X$ is a weak partial (respectively linear) order on the set X.
  - "Let $(X, <_X)$ be a partially (respectively linearly) ordered set" when $<_X$ is a strict partial (respectively linear) order on the set X.

†Note that the textbook does not use this convention.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Order-Isomorphism

We need a precise way to express when two ordered sets are “the same”.

Order-Isomorphism

We need a precise way to express when two ordered sets are "the same".

Definitions (p. 166)

Let $(X, <_X)$ and $(Y, <_Y)$ be partially ordered sets. The ordered sets are **order-isomorphic**, denoted $X \cong Y$ or $(X, <_X) \cong (Y, <_Y)$, if there is a bijection $f : X \longrightarrow Y$ such that for all $x_1, x_2 \in X$,

$$x_1 <_X x_2 \quad \text{if and only if} \quad f(x_1) <_Y f(x_2).$$

Order-Isomorphism

We need a precise way to express when two ordered sets are "the same".

Definitions (p. 166)

Let $(X, <_X)$ and $(Y, <_Y)$ be partially ordered sets. The ordered sets are **order-isomorphic**, denoted $X \cong Y$ or $(X, <_X) \cong (Y, <_Y)$, if there is a bijection $f : X \longrightarrow Y$ such that for all $x_1, x_2 \in X$,

$$x_1 <_X x_2 \quad \text{if and only if} \quad f(x_1) <_Y f(x_2).$$

The map f is an **order-isomorphism**.

Order-Isomorphism

We need a precise way to express when two ordered sets are "the same".

Definitions (p. 166)

Let $(X, <_X)$ and $(Y, <_Y)$ be partially ordered sets. The ordered sets are **order-isomorphic**, denoted $X \cong Y$ or $(X, <_X) \cong (Y, <_Y)$, if there is a bijection $f : X \longrightarrow Y$ such that for all $x_1, x_2 \in X$,

$$x_1 <_X x_2 \quad \text{if and only if} \quad f(x_1) <_Y f(x_2).$$

The map f is an **order-isomorphism**.

Remark

The above definition on weak partial orders is similar.

Order-Isomorphism

Exercise (informal)

Let $<$ and $>$ be the usual orders on $\mathbb{Z}$. Show that $(\mathbb{Z}, <)$ and $(\mathbb{Z}, >)$ are order-isomorphic.

Order-Isomorphism

Exercise (informal)

Let $<$ and $>$ be the usual orders on $\mathbb{Z}$. Show that $(\mathbb{Z}, <)$ and $(\mathbb{Z}, >)$ are order-isomorphic.

Solution

Let $f : \mathbb{Z} \longrightarrow \mathbb{Z}$ be the function $n \longmapsto -n$. We can show that the function f is a bijection and that it preserves the order, that is, for all $m, n \in \mathbb{Z}$,

$$m < n \quad \text{if and only if} \quad -m > -n.$$

Order-Isomorphism

Theorem

The binary relation $\cong$ is an equivalence relation on strictly partially ordered sets.

Order-Isomorphism

Theorem

The binary relation $\cong$ is an equivalence relation on strictly partially ordered sets.

That is, let $(X, <_X)$, $(Y, <_Y)$ and $(Z, <_Z)$ be partially ordered sets, then the binary relation $\cong$ satisfies the following properties (Exercise 7.6):

(a) **reflexive**: for all X, $X \cong X$;

(b) **symmetric**: for all X and Y, if $X \cong Y$ then $Y \cong X$;

(c) **transitive**: for all X, Y and Z, if $X \cong Y$ and $Y \cong Z$ then $X \cong Z$.

Order-Isomorphism

Theorem

The binary relation $\cong$ is an equivalence relation on strictly partially ordered sets.

That is, let $(X, <_X)$, $(Y, <_Y)$ and $(Z, <_Z)$ be partially ordered sets, then the binary relation $\cong$ satisfies the following properties (Exercise 7.6):

(a) **reflexive**: for all X, $X \cong X$;

(b) **symmetric**: for all X and Y, if $X \cong Y$ then $Y \cong X$;

(c) **transitive**: for all X, Y and Z, if $X \cong Y$ and $Y \cong Z$ then $X \cong Z$.

Technical remark

Note that the equivalence relation $\cong$ should be defined on the universe $\mathcal{V}$, that is, it is not a relation within ZFC.

Order-Isomorphism

Definition (p. 168)

An **order-theoretic invariant** is a property of an ordered set preserved under order-isomorphism.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Order-Embedding

We also need to define when an ordered set fits inside another while preserving its order structure entirely.

Order-Embedding

We also need to define when an ordered set fits inside another while preserving its order structure entirely.

Definition (p. 167)

Let $(X, <_X)$ and $(Y, <_Y)$ be partially ordered sets. A function $f : X \longrightarrow Y$ is an **order-embedding** of X into Y (or of $(X, <_X)$ into $(Y, <_Y)$) if f is one-one and, for all $x_1, x_2 \in X$,

$$x_1 <_X x_2 \quad \text{if and only if} \quad f(x_1) <_Y f(x_2).$$

Order-Embedding

Exercise (informal)

Let $\mathbb{N}$ and $\mathbb{R}$ with the usual order $<$. Show an order-embedding of $(\mathbb{N}, <)$ into $(\mathbb{R}, <)$

Order-Embedding

Exercise (informal)

Let $\mathbb{N}$ and $\mathbb{R}$ with the usual order $<$. Show an order-embedding of $(\mathbb{N}, <)$ into $(\mathbb{R}, <)$

Solution

Let $f : \mathbb{N} \longrightarrow \mathbb{R}$ be the function $n \longmapsto n$. We can show that the function f is one-one and that it preserves the order, that is, for all $m, n \in \mathbb{N}$,


$$m < n \quad \text{if and only if} \quad f(m) < f(n).$$

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Representation of Linearly Ordered Sets

Let $(X, <_X)$ be a linearly ordered set and let $x, x' \in X$ with $x <_x x'$. We picture x to the left of x'. The red dots mean that an element does not belong to the set X.



"*Of course, we must not read too much into the picture. The picture is very suggestive of the usual real line, but for instance there might not be any elements of X between x and x' in the picture above.*" *(p. 167)*

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Maximum and Minimum Elements

Definitions (p. 168)

Let $(X, \leq_X)$ be a linearly ordered set and let $a, b \in X$.

Maximum and Minimum Elements

Definitions (p. 168)

Let $(X, \leq_X)$ be a linearly ordered set and let $a, b \in X$.

(a) The element b is the **maximum** (or **greatest**) element of X, if $x \leq_X b$ for all $x \in X$.

Maximum and Minimum Elements

Definitions (p. 168)

Let $(X, \leq_X)$ be a linearly ordered set and let $a, b \in X$.


(a) The element b is the **maximum** (or **greatest**) element of X, if $x \leq_X b$ for all $x \in X$.

(b) The element a is the **minimum** (or **least**) element of X, if $a \leq_X x$ for all $x \in X$.

Maximum and Minimum Elements

Example (informal, p. 168)

Let $A = \left\{1 - \frac{1}{n+1} : n \in \mathbb{N}\right\}$ and $B = A \cup \{1\}$, subsets of $\mathbb{Q}$, with the usual order $\leq$.



The set A has minimum element, 0, but no maximum element.
The set B has minimum element, 0, and maximum element, 1.

Maximum and Minimum Elements

In the next theorem we shall see that the maximum and the minimum are order-theoretic invariants.

Maximum and Minimum Elements

In the next theorem we shall see that the maximum and the minimum are order-theoretic invariants.

Theorem 7.1

Let $(X, \leq_X)$ and $(Y, \leq_X)$ be linearly ordered sets and let $f : X \longrightarrow Y$ be an order-isomorphism from X to Y. Then

(a) if X has a maximum element then so does Y;

(b) if X has a minimum element then so does Y.

Maximum and Minimum Elements

Example (informal)

Let $A = \left\{1 - \frac{1}{n+1} : n \in \mathbb{N}\right\}$ and $B = A \cup \{1\}$, subsets of $\mathbb{Q}$, with the usual order $\leq$.

By Theorem 7.1 the ordered sets $(A, \leq)$ and $(B, \leq)$ are not order-isomorphic because the set B has maximum element, but the set A does not.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Limit Points

Definition (p. 169)

Let $(X, <_X)$ be a linearly ordered set and let $c \in X$. The element c is a **limit point** of X if

(a) there exits $x \in X$ such that $x <_X c$ and

(b) for all $x \in X$ with $x <_X c$, there exits $y \in X$ such that $x <_X y <_X c$.

Limit Points

In the next theorem we shall see that limit points are order-theoretic invariants.

Limit Points

In the next theorem we shall see that limit points are order-theoretic invariants.

Theorem 7.2

Let $(X, <_X)$ and $(Y, <_X)$ be linearly ordered sets, let $f : X \longrightarrow Y$ be an order-isomorphism from X to Y and let $c \in X$. If c is a limit point of X then $f(c)$ is a limit point of Y (Exercise 7.11).

Limit Points

Exercise (informal, p. 170)

Let A and D be subsets of $\mathbb{Q}$, with the usual order $<$, defined by


$$A = \left\{1 - \frac{1}{n+1} : n \in \mathbb{N}\right\},$$
$$D = \left\{1 - \frac{1}{n+1} : n \in \mathbb{N}\right\} \cup \left\{2 - \frac{1}{n+1} : n \in \mathbb{N}\right\}.$$

Are $(A, <)$ and $(D, <)$ order-isomorphic?

(continued on next slide)


Limit Points

Solution



Limit Points

Solution



(a) The element $0 = 1 - \frac{1}{0+1}$ is the minimum (so 0 is no a limit point) of the sets A and D.

(b) The set A has not limit points.

(c) The element $1 = 2 - \frac{1}{0+1}$ is a limit point of set D.

Therefore, since $(D, <)$ has one limit point but $(A, <)$ does not have, the ordered sets are not order-isomorphic by Theorem 7.2.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Successors and Predecessors

Definition (p. 170)

Let $(X, <_X)$ be a linearly ordered set and let $x, d \in X$. The element d is the **successor** of x, denoted x^+, and x is the **predecessor** of d, if

(a) $x <_X d$ and

(b) for all $y \in X$, if $x <_X y$ then $d \doteq y$ or $d <_X y$.

Successors and Predecessors

Definition (p. 170)

Let $(X, <_X)$ be a linearly ordered set and let $x, d \in X$. The element d is the **successor** of x, denoted x^+, and x is the **predecessor** of d, if

(a) $x <_X d$ and

(b) for all $y \in X$, if $x <_X y$ then $d \doteq y$ or $d <_X y$.

That is, d is the successor of x (and x is the predecessor of d) if d is the first element of X greater than x.

Successors and Predecessors

Questions

In a strictly linearly ordered set

(a) can a limit point also be a successor?

Successors and Predecessors

Questions

In a strictly linearly ordered set

(a) can a limit point also be a successor?

(b) can a successor also be a limit point?

Successors and Predecessors

Exercise 7.12

Let $(X, <_X)$ be a linearly ordered set and suppose that an element $c \in X$ is neither minimum element of X nor a successor. Show that c is a limit point.

Successors and Predecessors

Examples (informal)

(a) In $\mathbb{R}$ with the usual order $<$, every real number is a limit point, so $\mathbb{R}$ has no successors.

Successors and Predecessors

Examples (informal)


(a) In $\mathbb{R}$ with the usual order $<$, every real number is a limit point, so $\mathbb{R}$ has no successors.

(b) In $\mathbb{N}$ with the usual order $<$, every natural number different from zero is a successor, so $\mathbb{N}$ has no limit points.

Successors and Predecessors

Example (informal)

Let $D = \left\{1 - \frac{1}{n+1} : n \in \mathbb{N}\right\} \cup \left\{2 - \frac{1}{n+1} : n \in \mathbb{N}\right\}$ be a subset of $\mathbb{Q}$, with the usual order $<$.



(a) The element $0 = 1 - \frac{1}{0+1}$ is the minimum (0 is neither a successor nor a limit point).

(b) Every element $1 - \frac{1}{n+1}$, with $n > 0$, is the successor of the element $1 - \frac{1}{n}$.

(c) The element $1 = 2 - \frac{1}{0+1}$ is a limit point.

(d) Note that for all $x \in D$, with $x < 1$, x has a successor x^+ and the set $\{x \in D : x < 1\}$ is infinite (see Exercise 7.14).

(e) Every element $2 - \frac{1}{n+1}$, with $n > 0$, is the successor of the element $2 - \frac{1}{n}$.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Finite Strictly Linearly Ordered Sets

Definition (p. 188)

Let $(X, <_X)$ be a linearly ordered set. The relation $<_X$ is a **well-order** (or the set X is **well-ordered** by the relation $<_x$) if every non-empty subset A of X has a least element, i.e., there is some element $a \in A$ such that $a \leq_X x$, for all $x \in A$.

Finite Strictly Linearly Ordered Sets

Definition (p. 188)

Let $(X, <_X)$ be a linearly ordered set. The relation $<_X$ is a **well-order** (or the set X is **well-ordered** by the relation $<_x$) if every non-empty subset A of X has a least element, i.e., there is some element $a \in A$ such that $a \leq_X x$, for all $x \in A$.

Theorem 3.9

The set $\mathbb{N}$ is well-ordered by the relation $\in$.

Finite Strictly Linearly Ordered Sets

The following theorem will state that every strictly linearly ordered finite set is order-isomorphic to some natural number $\mathbf{n}$ ordered by $\in$.

Finite Strictly Linearly Ordered Sets

The following theorem will state that every strictly linearly ordered finite set is order-isomorphic to some natural number $\mathbf{n}$ ordered by $\in$.

Theorem

Let $\mathbf{n} \in \mathbb{N}$ and let $(X, <_X)$ be a linearly ordered finite set with $\mathbf{n}$ elements.
Then $(X, <_X) \cong (\mathbf{n}, \in)$ (Exercise 7.15(a)).

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

Upper Bounds and Least Upper Bounds

Definition (p. 175)

Let $(X, \leq_X)$ be a linearly ordered set and let A be a non-empty subset of X. An element $x \in X$ is the **least upper bound** (or the **supremum**) of the set A, denoted $\operatorname{lub} A$, if (Exercise 7.19)

(a) the element x is an **upper bound** of the set A, i.e., $y \leq x$ for all $y \in A$;

(b) the element x is the least such upper bound of the set A, i.e., if z is an upper bound of the set A, then $x \leq z$.

Upper Bounds and Least Upper Bounds

Definition (p. 175)

Let $(X, \leq_X)$ be a linearly ordered set and let A be a non-empty subset of X. An element $x \in X$ is the **least upper bound** (or the **supremum**) of the set A, denoted $\operatorname{lub} A$, if (Exercise 7.19)

(a) the element x is an **upper bound** of the set A, i.e., $y \leq x$ for all $y \in A$;

(b) the element x is the least such upper bound of the set A, i.e., if z is an upper bound of the set A, then $x \leq z$.

Example

In the whiteboard.

Outline

Introduction
Weak Orders
Strict Orders
Orders on Natural Numbers
Conventions
Order-Isomorphism
Order-Embedding
Representation of Linearly Ordered Sets
Maximum and Minimum Elements
Limit Points
Successors and Predecessors
Finite Strictly Linearly Ordered Sets
Upper Bounds and Least Upper Bounds
References

References

Derek Goldrei (1996). Classic Set Theory. A Guided Independent Study. Chapman & Hall (cit. on p. 2).