

Notes

Mainly to Elements of Set Theory

Lin

August 24, 2026

$$B \subseteq C \implies \mathcal{P}(B) \subseteq \mathcal{P}(C)$$

Proof. By def $\mathcal{P}(B) = \{x \mid x \subseteq B\}$

$\forall x, x \in \mathcal{P}(B)$, Then, $x \subseteq B \subseteq C$

$$\implies x, x \subseteq C$$

$$\implies x \in \mathcal{P}(C) \text{ By def } \mathcal{P}C = \{x \mid x \subseteq C\}$$

Thus, $\mathcal{P}(B) \subseteq \mathcal{P}(C)$

□

$$A \text{ is a finite set, } |A| = n \implies |\mathcal{P}(A)| = 2^n$$

Proof. Consider taking i elements from A to build a subset.

Then $i \in \{0, \dots, n\}$, and the number of that kind of subsets is $\binom{n}{i}$.

(if $i = 0$, the subset is $\emptyset$; if $i = n$, the subset is A itself)

$$\implies |\mathcal{P}(A)| = \sum_{i=0}^n \binom{n}{i}$$

$$\implies |\mathcal{P}(A)| = 2^n$$

□

$$x, y \in B \implies \{\{x\}, \{x, y\}\} \in \mathcal{P}(\mathcal{P}(B))$$

Proof. $x, y \in B$

$$\implies \{x\}, \{x, y\} \in \mathcal{P}(B)$$

$$\implies \{\{x\}, \{x, y\}\} \subseteq \mathcal{P}(B)$$

Thus, by def $\{\{x\}, \{x, y\}\} \in \mathcal{P}(\mathcal{P}(B))$

□

Remark 1. A simple way to calculate the rank of set (which finite) is to count the deepest nested brackets until meet atom or $\emptyset$.

$$\forall \alpha \in \mathbb{N}, \alpha \leq 3, V_{\alpha+1} = A \cup \mathcal{P}(V_\alpha)$$

Proof. $V_0 = A$

By def, $V_{n+1} = V_n \cup \mathcal{P}(V_n)$

$$V_1 = V_0 \cup \mathcal{P}(V_0) = A \cup \mathcal{P}(V_0)$$

$$V_2 = V_1 \cup \mathcal{P}(V_1) = A \cup \mathcal{P}(V_0) \cup \mathcal{P}(V_1) = A \cup \mathcal{P}(V_1)$$

$$\mathcal{P}(V_0) \cup \mathcal{P}(V_1) = \mathcal{P}(V_1), \text{ By } \mathcal{P}(V_0) \subseteq \mathcal{P}(V_1) \text{ since } V_0 \subseteq V_1$$

$$V_3 = V_2 \cup \mathcal{P}(V_2) = A \cup \mathcal{P}(V_1) \cup \mathcal{P}(V_2) \text{ Similarly,}$$

$$V_3 = A \cup \mathcal{P}(V_2)$$

□

Remark 2. I guess we can use mathematical induction to make it to $\forall \alpha \in \mathbb{N}$