COMPENDIUM ON BASIC MODELS OF CARDIOVASCULAR DISEASE

Animal Models to Study Cardiac Arrhythmias

Daniel J. Blackwell, Jeffrey Schmeckpeper, Bjorn C. Knollmann

ABSTRACT: Cardiac arrhythmias are a significant cause of morbidity and mortality worldwide, accounting for 10% to 15% of all deaths. Although most arrhythmias are due to acquired heart disease, inherited channelopathies and cardiomyopathies disproportionately affect children and young adults. Arrhythmogenesis is complex, involving anatomic structure, ion channels and regulatory proteins, and the interplay between cells in the conduction system, cardiomyocytes, fibroblasts, and the immune system. Animal models of arrhythmia are powerful tools for studying not only molecular and cellular mechanism of arrhythmogenesis but also more complex mechanisms at the whole heart level, and for testing therapeutic interventions. This review summarizes basic and clinical arrhythmia mechanisms followed by an in-depth review of published animal models of genetic and acquired arrhythmia disorders.

Key Words: atrial fibrillation ■ cardiomyopathies ■ channelopathies ■ models, animals ■ morbidity

ardiac arrhythmias affect ≈2% of community-dwelling adults, with an incidence of ≈0.5% per year. Arrhythmias can manifest as relatively benign entities, such as atrial and ventricular premature beats, or as life-threatening arrhythmias such as ventricular tachycardia (VT) and ventricular fibrillation (VF), which can lead to sudden cardiac death (SCD), accounting for 10-15% of all deaths in the United States. Atrial fibrillation (AF) accounts for the greatest arrhythmia burden and is associated with stroke and heart failure, fueling huge health care costs. Arrhythmia treatment approaches focused on risk factor reduction, drug therapy, catheter ablation, device implantation, or a combination of these strategies has improved morbidity and mortality over the last 20 years, but treatment with antiarrhythmic drugs is often ineffective or increases mortality long term.^{2,3} A more thorough understanding of the pathophysiology of arrhythmia initiation and maintenance is important for improving clinical outcomes.

The mechanisms underlying arrhythmogenesis at the cellular level involve ion channels and electrogenic transporters that are altered via biogenic (synthesis, processing, trafficking, and degradation), biochemical (posttranslation modification, phosphorylation), and biophysical (gating, permeation) processes (reviewed here in study by Delisle et al4). The interplay between ion channels and transporters controls the action potential duration (APD), effective refractory period, and Ca2+ cycling to coordinate excitation-contraction coupling and normal myocyte function; dysregulation leads to abnormal cardiomyocyte electrical activity.5 Structural and hemodynamic parameters contribute to further cardiac remodeling, increasing the risk for arrhythmia development and maintenance. 6-9

To study underlying arrhythmia mechanisms and evaluate treatment approaches, multiple in vitro systems and in vivo models have been developed. This review focuses on animal models that have informed our understanding of arrhythmia pathophysiology and have been used to develop new therapeutic approaches. An ideal model would recapitulate human anatomic, electrophysiological, and hemodynamic parameters. Currently, no single model can accomplish this feat. However, animal models have enabled the discovery of new treatment strategies for humans with genetic arrhythmia disorders. For example, mouse models demonstrated the efficacy of flecainide in catecholaminergic polymorphic ventricular tachycardia¹⁰ and mexiletine in long QT type 3.11 When choosing an animal model of cardiac arrhythmia, researchers must consider the most appropriate model to address a specific scientific question based on cost, complexity, ease of handling, access to diagnostic and surgical expertise, and the ability for genetic modification.

Correspondence to: Björn C. Knollmann, MD, PhD, William Stokes Professor of Medicine and Pharmacology, Director, Vanderbilt Center for Arrhythmia Research and Therapeutics (VanCART), Vanderbilt University School of Medicine, Medical Research Bldg IV, Rm. 1265, 2215B Garland Ave, Nashville, TN 37232-0575. Email bjorn. knollmann@vumc.org*

D.J. Blackwell and J. Schmeckpeper contributed equally.

For Sources of Funding and Disclosures, see page 1951.

© 2022 American Heart Association, Inc.

Circulation Research is available at www.ahajournals.org/journal/res

ΔΤ

Nonstandard Abbreviations and Acronyms

AF atrial fibrillation
AP action potential

APD action potential duration

ARVC arrhythmogenic right ventricular

cardiomyopathy atrial tachycardia

BrS Brugada syndrome
CHB complete heart block

CPVT catecholaminergic polymorphic ventricu-

lar tachycardia

CRDS Ca²⁺ release deficiency syndrome

DAD delayed afterdepolarizationDCM dilated cardiomyopathy

DSC2 desmocollin-2 DSG2 desmoglein-2

EAD early afterdepolarization **ECM** extracellular matrix

HCM hypertrophic cardiomyopathy

LQTS long QT syndrome

MAT multifocal atrial tachycardia

MPO myeloperoxidase

PES programmed electrical stimulation

PKP2 plakophilin-2PLB phospholamban

PVC premature ventricular contraction

SCD sudden cardiac death SOTS short QT syndrome SR sarcoplasmic reticulum **SVT** supraventricular tachycardia TAC transverse aortic constriction **TdP** Torsades de Pointes **TGF** transforming growth factor VF ventricular fibrillation VT ventricular tachycardia

Here, we provide the reader with a brief overview of basic and clinical cardiac electrophysiology, followed by an in-depth review of existing animal models of cardiac arrhythmias. Animal models are classified as either genetic (ie, arrhythmia risk caused by gene mutation) or acquired (ie, arrhythmia risk caused by nongenetic heart diseases such as myocardial infarction, metabolic abnormalities, or cardiac hypertrophy).

PRINCIPLES OF CARDIAC ELECTROPHYSIOLOGY

The Cardiac Conduction System

Normal heart rhythm is generated and regulated in the specialized cardiac conduction system, which consists

of the sinoatrial node, the atrioventricular (AV) node, and the HIS-Purkinje system (Figure 1). Electrical impulses are initiated in the sinoatrial node and spread through the atria to the AV node. After a slight delay (0.12-0.20 seconds), excitation continues through the bundle of His, the right and left bundle branches, and finally the Purkinje fibers, which then excite the working myocardium. The delay in the AV node allows the atria to contract earlier than the ventricles and provides adequate time for optimal ventricular filling. 12 The specialized cells within the sinoatrial node, AV node, and His-Purkinje system are capable of spontaneous depolarization that is regulated by both the sympathetic and parasympathetic nervous system. Conduction through the heart depends on electrical coupling between cells, which is mediated by gap junctions.

Species Differences in the Cardiac Action Potential and Cardiac Ca²⁺ Handling

The cardiac action potential (AP) results from the opening and closing of ion channels and electrogenic transporters in the plasma membrane of individual cardiomyocytes (see study by Varró et al¹³ for details). Figure 2 illustrates AP wave forms and underlying membrane currents for ventricular cardiomyocytes of humans and mice.

When choosing an animal model for arrhythmia research, it is important to recognize species differences in cardiac AP and membrane currents, which are the result of species-specific expression of ion channels and transporters. For example, unlike humans, mice and rats have a low AP plateau at ≈−40 mV membrane potential (Figure 2). This is primarily the result of differential expression in repolarizing transient K-currents, as illustrated in Figure 2. On the other hand, rabbits and guinea pigs have a more positive AP plateau analogous to humans.¹⁴ For a more detailed comparison of ionic currents in different species, the reader is referred to here.¹⁵

As with the AP, there are important species differences in cardiac Ca²+ handling. For example, mice and rats primarily (>90%) utilize sarcoplasmic reticulum (SR)-mediated Ca²+ cycling (via the cardiac ryanodine receptor [RyR2] and the SR Ca uptake pump [SERCA2a]) for excitation-contraction coupling, whereas in humans, dogs, and rabbits the SR accounts for $\approx\!65\%$, with the remainder coming from outside the cell via the L-type Ca channel (Ca_1.2) and the NCX (Na/Ca exchanger). For a more detailed comparison of species differences in Ca²+ handling, the reader is referred to here. 17

Pathophysiology of Cardiac Arrhythmias

The main mechanisms of arrhythmogenesis can be divided into either abnormal impulse generation or abnormal impulse propagation. Disorders of impulse generation and propagation, regulation of the AP duration, and cellular




Figure 1. Schematic of the cardiac conduction system and clinical classification of cardiac arrhythmias.

AF indicates atrial fibrillation; AV, atrioventricular; AVNRT, AV nodal reentry tachycardia; AVRT, AV reciprocating tachycardia; PAC, premature atrial;

substrates can all contribute to 3 categories of arrhythmias; enhanced automaticity, triggered ectopic beats, and reentry. Each arrhythmia category is explained briefly below. A more detailed review can be found in here.⁵

and SA, sinoatrial. Illustration credit: Ben Smith.

Automaticity

Automaticity is the ability of cells to generate their own AP.¹⁸ The intrinsic depolarization rate of the sinoatrial node is faster than the rest of the cardiac conduction system and overdrives pacemaking in the AV node and His-Purkinje system. However, automaticity in the AV node and His-Purkinje system can become dominant in sinoatrial nodal dysfunction. The sinoatrial node is more sensitive to increased sympathetic and parasympathetic tone, leading to sinus tachycardia and bradycardia, respectively. Under normal conditions, atrial and ventricular cardiomyocytes display either no or very slow intrinsic depolarization that are easily suppressed by the faster, coordinated impulses from the sinoatrial node through the conduction system. Increased automaticity in the atria can lead to focal and multifocal atrial tachycardia and AF. Specifically, the pulmonary vein sleeve, where the left atria myocytes transition to the tunica media of the pulmonary veins, is known to harbor tissue with increased automaticity, and is a target for catheter based ablation by pulmonary vein isolation for AT and AF. Increased automaticity in the ventricle is less common but can lead to VT or accelerated idioventricular rhythms.

Afterdepolarizations and Triggered Arrhythmia

Triggered arrhythmias are because of spontaneous membrane depolarization of atrial or ventricular myocytes that precede the next sinus beat. Membrane depolarizations that occur within or follow the cardiac AP are referred to as afterdepolarizations. Two classes are traditionally recognized: early and delayed. An early afterdepolarization (EAD) interrupts the repolarization during phase 2 or early phase 3 of the cardiac AP, whereas a delayed afterdepolarization (DAD) occurs after full repolarization in Phase 4. When an EAD or DAD brings the membrane to its threshold potential, a spontaneous AP is referred to as a triggered response. These triggered events can give rise to premature extrasystolic complexes in the atria or the ventricle (PVCs), precipitating tachyarrhythmias. In general, any unbalanced increased inward current (ie, gain-of-function




Figure 2. The ventricular action potential and ionic currents in humans and mice.

Note the differences in action potential shape, which is caused primarily by differences in ionic currents circled in red. $I_{Ca(L)}$ indicates L-type Ca current; $I_{Ca(L)}$, T-type Ca current; I_{Nal} , Na current; I_{NaCa} , Na-Ca-Exchange current; I_{Nal} , NaK-ATPase pump current; I_{K1} , Inward rectifier K-current; I_{K2} rapidly activating delayed rectifier K-current; I_{K2} slowly activating delayed rectifier K-current; I_{K2} rapidly activating steady-state K-current; I_{K2} rapidly activating that current densities (pA/pF) measured in single cells vary drastically with experimental conditions and voltage clamp protocols. Current densities were chosen to reflect relative contributions to the AP.

mutations in Na⁺ or Ca²⁺ channels) or decreased outward currents (ie, loss-of-function mutations in K⁺ channels) will depolarize the cell membrane and can lead to EADs or DADs.²⁰ Specifically, a major cause of triggered arrhythmia is spontaneous RyR2-mediated SR Ca²⁺ release, driving inward Na⁺ current via the NCX, leading to EAD and in particular DAD formation, which are important cellular arrhythmia mechanisms in AF, VT, and SCD.²¹

Reentrant Arrhythmia

In reentry, a group of myocardial cells that are not activated during the early stage of depolarization can resume excitability before the impulse vanishes. In this situation, they may connect to re-excite zones that were previously depolarized but were recovered from the refractory period of the initial wave. Two crucial factors predisposing reentry are prolonged conduction time and shortened refractory period. Reentry is the dominant mechanism of arrhythmias in the clinical setting and occurs due to anatomic and functional factors.²²

Classification of Clinical Arrhythmias

Clinically, cardiac arrhythmias are usually classified as bradyarrhythmias and tachyarrhythmias (Figure 1). Both types can reduce cardiac output, resulting in hypotension and ultimately can cause death but have different underlying mechanisms. Figure 1 lists the major clinical types of brady and tachyarrhythmias. Briefly, bradyarrhythmias reduce the heart rate either by reducing spontaneous depolarization within the sinoatrial node, slowing conduction through the conduction system, or increasing parasympathetic tone. Sinus bradycardia, sinus node exit block, sinus arrest, and asystole are caused by dysfunction within the sinus node itself, due to destruction of the pacemaker cells, fibrosis of the sinoatrial node, or increased parasympathetic tone. AV nodal block prolongs the conduction above, within or below the AV node. Depending on the severity of AV block, it is classified as first-degree, second-degree, or complete (third-degree) heart block (Figure 1).

Tachyarrhythmias are accelerated rhythms that originate from either above (supraventricular tachycardia) or below the AV node (ventricular arrhythmia). The most common supraventricular tachycardias are sinus tachycardia and AF. Premature atrial contractions and AT are commonly caused by automatic foci within the atria. Reentrant atrial arrhythmias include atrial flutter, AV nodal reentry tachycardia, and AV reciprocating tachycardia. Atrial flutter is a macroreentrant loop, typically involving the tricuspid annulus limited by anatomic

barriers such as the superior and inferior cava veins, the coronary sinus and crista terminalis.²³ AV nodal reentry tachycardia is a microreentry related to differences in the refractory period of the slow and fast pathway within the AV node.²⁴ AV reciprocating tachycardia, also known as preexcitation syndrome, occurs due to the presence of an accessory pathway, most notably the Bundle of Kent leading to Wolf-Parkinson-White syndrome, which can prematurely conduct impulses between the atria and ventricles.

Ventricular arrhythmias include premature ventricular contractions (PVCs), VT, and VF. PVCs are single premature beats due to EADs or DADs in myocardial cells and benign, unless they trigger VT or VF. VT and VF are usually reentrant arrhythmias, and if not treated rapidly, can lead to sudden cardiac death. While a majority of cases of VT are because of reentry around the scar in structural heart disease, 10% of VT occurs in structurally normal hearts due to nonreentrant mechanisms such as catecholaminergic polymorphic VT (CPVT), fascicular VT, left or right outflow tract VT, mitral and tricuspid annular VT, long QT, and Brugada syndrome.²⁵

Animal Models

An important consideration for selecting an animal model to study cardiac arrhythmias is how closely the species resembles human cardiac physiology. Caenorhabditis elegans and Drosophila melanogaster both develop heart tubes and have been primarily used to screen gene function and examine development and cardiac structure. Zebrafish have a 2 chambered heart with some similarities in AP electrophysiology to humans and provide advantages for understanding cardiogenesis. Zebrafish embryos are transparent, enabling optical viewing, fluorescent protein expression, and optogenetic pacing; they have large clutch sizes with a rapid embryonic stage lasting only 3 to 4 days postfertilization; are amenable to drug absorption; and genes are easily manipulated. Mouse hearts are anatomically similar to human hearts with 4 chambers and comparable development,26 albeit differences in coronary anatomy.²⁷ However, there are major differences in heart rate, cardiac AP, and membrane currents (Figure 2). These differences influence ion channel function, refractoriness, and arrhythmia susceptibility. In addition, the small size of the mouse heart may contribute to the frequently observed self-termination of reentrant arrhythmias or lack of spontaneous arrhythmias in many models. Nevertheless, the mouse has been the primary animal model for cardiac arrhythmia studies of inherited cardiomyopathies and channelopathies, and many models faithfully capture cardiac disease. Rabbits more closely recapitulate the human AP compared with rats and mice. The rabbit AP has a sustained Ca2+ current-driven plateau phase, and the major repolarizing K+ currents are similar to humans. Rabbit heart size and

beating rate is between that of mice and humans. Dogs, pigs, and goats have a similar cardiac anatomy, size, and beating rate (slightly higher in dogs) as humans. Their cardiac electrophysiology, APs, and ionic currents are all fairly comparable to humans, and their primary limitations as an animal model for research come from their cost, size, and time to breed and reach sexual maturity.

ANIMAL MODELS OF GENETIC ARRHYTHMIA DISORDERS

Genetic arrhythmia disorders are either caused by or associated with identifiable gene mutations. Genetic arrhythmia syndromes can be subdivided into channelopathies without structural heart disease (eg, CPVT, long QT syndrome [LQTS], Brugada Syndrome) and genetic arrhythmia syndromes associated with structural heart disease (eg, hypertrophic cardiomyopathy [HCM], dilated cardiomyopathy [DCM], action potential duration [ARVC], AF). Tables 1 and 2 list published animal models of genetic arrhythmia syndromes. Most animal arrhythmia models are mice, given the ease of genetic manipulation. Despite differences between rodent and human cardiac electrophysiology (Figure 2), mouse models have enabled the study of human genetic diseases, identification of pathogenic mutations, characterization of disease pathophysiology, and testing/screening of therapeutic interventions. Advances in gene editing technology such as CRISPR/Cas9 have provided faster, easier, and cheaper methods to develop genetic models in animals and cells. Compared with cellular models such as human-induced pluripotent stem cells, animal models provide a distinct advantage in modeling cardiac arrhythmias where the anatomy of the heart is relevant.²¹⁷ In the following section, each of the major genetic arrhythmia syndromes are introduced, followed by examples of animal models that have informed disease pathophysiology and treatment approaches.

Channelopathies

Channelopathies are caused by mutations in ion channel genes or genes that regulate ion channels and generate arrhythmia risk in the structurally normal heart. However, overlap syndromes caused by mutations in channelopathy genes (eg, SCN5A) can also be associated with alterations in cardiac structure, which are discussed in Section genetic arrhythmia syndromes associated with structural heart disease.

Catecholaminergic Polymorphic Ventricular Tachycardia

CPVT is characterized by arrhythmogenesis evoked by elevated catecholamines during stress or exercise. CPVT is caused by gain-of-function mutations in proteins that constitute the intracellular Ca²⁺ release unit of the SR. These mutations result in spontaneous Ca²⁺

COMPENDIUM ON BASIC MODELS OF CARDIOVASCULAR DISEASE

Table 1. Animal Models of Genetic Arrhythmia Channelopathies

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref |
|-----------------------|-------------------------------|---|---|-----|
| CPVT | | | | _ |
| CPVT1 | | | | |
| Mouse | RYR2 (RyR2) | R4496C± | Catecholamine/exercise-induced arrhythmia; phenotype not as penetrant as some other models | 28 |
| Mouse | RYR2 (RyR2) | V2475F± | Homs are embryonic lethal, catecholamine-induced arrhythmia | 29 |
| Mouse | RYR2 (RyR2) | R2474S± | Seizures, exercise-induced arrhythmia, sudden death | 30 |
| Mouse | RYR2 (RyR2) | P2328S± | Iso-induced PVCs, VT, VF | 31 |
| Mouse | RYR2 (RyR2) | R2474S/V3599K | GoF + LoF; protective- no arrhythmia | 32 |
| Mouse | RYR2 (RyR2) | R4496C+/- with E4872Q+/- | GoF + LoF; protective- no arrhythmia | 33 |
| Mouse | RYR2 (RyR2) | L433P+/-, N2386I+/- | Mutations from patients with CPVT; no report of CPVT in mouse models; mice also develop AF with pacing protocol | 34 |
| Mouse | RYR2 (RyR2) | R176Q± | CPVT-like phenotype; AF | 35 |
| Mouse | RYR2 (RyR2) | exon3del± | Homs embryonic lethal, hets did not develop CPVT; patients with exon3del have exercise-induced VT | 36 |
| C. elegans | RYR2 (RyR2), CASQ2 (Casq2) | R4743C, Casq2 KO | Enables optogenetic pacing, observed defect with mutations | 37 |
| CPVT2 | 1 | | | |
| Mouse | CASQ2 (Casq2) | КО | Severe CPVT, ultrastructure changes, catecholamine/exercise-induced arrhythmia | 38 |
| Mouse | CASQ2 (Casq2) | D307H/D307H, D309deltaE9/ D309deltaE9 (Casq2-/-) | Both mutations result in loss of Casq2 protein, catecholamine/ exercise-induced arrhythmia | 39 |
| Mouse | CASQ2 (Casq2) | Ventricular/Purkinje KO | Subtissue-selective knockout | 40 |
| Mouse | CASQ2 (Casq2) | Ventricular/Purkinje KO | Subtissue-selective knockout | 41 |
| Mouse | CASQ2 (Casq2) | R33Q/R33Q | Reduction of Casq2 protein, ultrastructure changes, arrhythmia | 42 |
| Mouse | CASQ2 (Casq2) | K180R± | Autosomal dominant inheritance; iso-induced arrhythmias | 43 |
| CPVT4 | | | | |
| Zebrafish | CALM (CaM) | N53I and N97S | Increased iso-induced HR (indicated dominant negative effect) | 44 |
| Zebrafish | CALM (CaM) | E105A | Arrhythmias, tachycardia, altered RyR2 binding | 45 |
| CPVT5 | | | | |
| Mouse | TRDN (Triadin) | КО | Ultrastructure changes; iso-induced arrhythmias; overlap syndrome with LQT? | 46 |
| CPVT? | | | | |
| Mouse | KCNJ2 (Kir2.1) | R67Q+/, cardiac-specific | Structurally normal heart, iso-induced VT, no LQT | 47 |
| CRDS | | | | |
| Mouse | RYR2 (RyR2) | A4860G± | VF with sympathetic stimulation, homs embryonic lethal | 48 |
| Mouse | RYR2 (RyR2) | D4646A± | RyR2 Ca2+ release deficiency syndrome (CRDS); homs are embryonic lethal | 49 |
| LQTS | | | | |
| LQT1 | | | | |
| Mouse | KCNQ1 (Kv7.1) | КО | Deaf, shaker/waltzer phenotype; LQT | 50 |
| Mouse | KCNQ1 (Kv7.1) | Truncated isoform, cardiac-specific overexpression | LQT, sinus node dysfunction, occasional AV block | 51 |
| Mouse | KCNQ1 (Kv7.1) | A340V—/— | Homs have LOT, hets do with gene dose dependence; PVCs after feeding (linked to diabetes) | 52 |
| Rabbit | KCNQ1 (Kv7.1) | Y315S cardiac-specific overex- pression | LQT; sympathetic stimulation induces EADs and VT; rabbits die within 3 weeks of AV node ablation | 53 |
| LQT2 | | | | |
| Rabbit | KCNH2 (Kv11.1/hERG) | G628S cardiac-specific overex- pression | LQT, spontaneous PVT, sudden death; prolonged APD | 53 |
| Zebrafish | KCNH2 (Kv11.1/hERG) | I462R, M521K | 2:1 block (phenotype) | 54 |
| Zebrafish | KCNH2 (Kv11.1/hERG) | I59S-/- | 2:1 block (phenotype); prolonged APD; EADs | 55 |

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref |
|--------------------------|-------------------------------|--|---|-----|
| Zebrafish | KCNH2 (Kv11.1/hERG) | Morpholino KD of WT + expression of hERG ± mutants | Tested 40 pathogenic and 10 nonpathogenic hERG mutants in the zERG background | 56 |
| LQT3 | | | | |
| Mouse | SCN5A (Nav1.5) | ΔΚΡΟ/+ | 1505–1507 deletion; prolonged QT/QTc; prominent T wave; prolonged APD; arrhythmias; sudden death | 57 |
| Mouse | SCN5A (Nav1.5) | ΔQΚΡ/+ | 1507–1509 deletion; long QTc, wide QRS, AV block; spontaneous PVCs, VT, and VF with sudden death; no atrial arrhythmias | 58 |
| Mouse | SCN5A (Nav1.5) | N1325S cardiac-specific overex- pression | LQT; arrhythmia; sudden death; also other non-LQT3 features like shorter PR and elevated heart rate | 59 |
| Mouse | SCN5A (Nav1.5) | 1795insD | LQT and Brugada in family; homs embryonic lethal; sinus node dysfunction, conduction slowing, bradycardia, and LQT | 60 |
| Guinea pig | SCN5A (Nav1.5) | Cellular model, isolated cells tx | Isolated cardiomyocytes treated with anthopleurin; rescued by mexillitine | 61 |
| Minor types of L | QTS | | | |
| Mouse | ANK2 (Ankyrin-B) | КО | LQT with HR deceleration, sinus node dysfunction | 62 |
| Mouse | ANK2 (Ankyrin-B) | ± | Bradycardia, variable HR, slow conduction, AV dissociation, long QTc; iso/exercise-induced PVT & death | 63 |
| Mouse | KCNA1 (Kv1.1) | N-term fragment overexpression | Long QTc; spontaneous PVC, couplets, ventricular tachycardia | 64 |
| Mouse | KCNA5 (Kv1.5) | W461F cardiac-specific overex- pression | Long QTc | 65 |
| Mouse | KCNB1 (Kv2.1) | N216 cardiac-specific overex- pression | Long QTc | 66 |
| Mouse | KCND2 (Kv4.2) | W362F cardiac-specific overex- pression | Subtle bradycardia, prolonged QTc | 67 |
| Mouse | KCND2×KCNA4 | W362F×KO | QRS widening, prolonged QTc, AV block/dropped beats, spontaneous ventricular arrhythmia | 68 |
| Rabbit | KCNE1 (minK) | G52R dominant negative overex- pression | Long QTc; drug-induced arrhythmia (Torsades) | 69 |
| Mouse | KCNE1 (minK) | КО | LQT with HR deceleration | 70 |
| Mouse | KCNE2 (MiRP1) | КО | Long QTc with age, mice had hyperkalemia | 71 |
| Mouse | KCNE3 (MiRP2) | КО | Females have long QTc at 9 mo (hyperaldosteronism); increased susceptibility to IR arrhythmias | 72 |
| Mouse | KCNJ2 (Kir2.1) | КО | Bradycardia, LQT | 73 |
| Mouse | KCNJ2 (Kir2.1) | T75R cardiac-specific overex- pression | Long QTc; spontaneous VT; iso-induced PVCs, VT, atrial flutter/fibrillation | 74 |
| Mouse | KCNIP2 (KChIP2) | КО | Significant reduction in Ito; elevated ST segment, atrial flutter and VT with PES; prolonged APD in cells | 75 |
| Mouse | CACNA1C (Cav1.2) | -/-, ± | KO embryonic lethal, hets survive and are just like WT | 76 |
| Mouse | CACNA1C (Cav1.2) | G406R cardiac-specific overex- pression | Long QTc, exercise-induced PVCs and Torsades; crossing with AKAP150 KO protected against all phenotypes of the G406R mutant | 77 |
| Zebrafish | CALM (CaM) | D129G | Bradycardia; conduction abnormality; LQT | 78 |
| Mouse | SCN1B (Scn β1) | КО | Bradycardia, prolonged APD/QTc, slowed repolarization; sodium channel expression increased | 79 |
| Mouse | SLC18A2 (VMAT2) | ± | LOT | 80 |
| Mouse | ATP1A3 (NaK ATPase α3) | Human isoform overexpression | LQT, steeper QT rate dependence, T wave alternans, VT | 81 |
| Short QT syndrom | e (SQTS) | | | |
| SQTS1 | | | | |
| Rabbit | KCNH2 (Kv11.1/hERG) | N588K cardiac-specific overex- pression | Shortened APs and QTc, normal T wave height, ex vivo perfused hearts inducible VT/VF | 82 |
| Zebrafish | KCNH2 (Kv11.1/hERG) | L499P | SQT | 83 |
| SQTS8 | | | | |
| Zebrafish | SLC4A3 (AE3) | Knockdown | SQT and systolic duration; WT SLC4A3 expression rescued phenotype, but R370H did not | 84 |
| | | | | |

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref. |
|-----------------------|-------------------------------|------------------------------------|--|------|
| Mouse | SLC8A1 (NCX) | КО | SQT | 85 |
| Mouse | CAV3 (Caveolin 3) | WT cardiac-specific overexpression | Short QTc, bradycardia, prolonged PR | 86 |
| Kangaroo | Unknown | None reported | Kangaroos have LV hypertrophy, SQT, and are highly susceptible to VF and sudden death, especially under light anesthesia | 87 |
| BrS | | | | |
| Pig | SCN5A (Nav1.5) | E558X± | Conduction abnormalities, QRS widening, reduced conduction velocity, ex vivo hearts have increased susceptibility to VF | 237 |
| Mouse | SCN5A (Nav1.5) | ± | KO is embryonic lethal; hets have sick sinus, slowed conduction, pacing-induced VT; QTS widening and fibrosis with age | 88 |
| Mouse | SCN5A (Nav1.5) | 1795insD | LQT and BrS in family; hom mice embryonic lethal; sinus node dysfunction, conduction slowing, bradycardia, and LQT | 60 |
| Mouse | SCN5A (Nav1.5) | ΔSIV/+ | C-term truncation; SA, AV, and His conduction slowing; one human patient with V2016M diagnosed with Brugada | 89 |

AF indicates atrial fibrillation; APD, action potential duration; AV, atrioventricular; caff, caffeine; BrS, Brugada Syndrome; CPVT, catecholaminergic polymorphic ventricular tachycardia; CRDS, calcium release deficiency syndrome; DAD, delayed afterdepolarization; DKO, double knockout; DN, dominant negative; EAD, early afterdepolarization; GoF, gain of function; het, heterozygous; HF, heart failure; hom, homozygous; HR, heart rate; IR, ischemia-reperfusion; iso, isoproterenol; KO, knockout; LoF, loss of function; LQTS, long QT syndrome; MVT, monomorphic ventricular tachycardia; NSVT, nonsustained ventricular tachycardia; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RBBB, right bundle branch block; SVT, supraventricular tachycardia; VF, ventricular fibrillation; and WT, wild type.

release from RYR2 SR Ca²+ release channels, raising diastolic Ca²+ and generating membrane depolarizations and delayed afterdepolarizations via Ca²+ extrusion through the electrogenic Na+- Ca²+ exchanger. Pathological mutations in RYR2 make up more than half of identified CPVT cases and are inherited in an autosomal dominant fashion. Mutations in regulatory partners of RyR2 are rarer and include calsequestrin, calmodulin, and triadin. Evidence suggests that KCNJ2 mutations may also be involved in CPVT.

Mice have been the primary animal model for studying CPVT and have been valuable not only to establish arrhythmia pathophysiology but also to identify new drug therapy that proved efficacious in humans.²¹⁸ At least 1 attempt to generate a rabbit model overexpressing human RYR2-R4497C was unsuccessful, likely due to the selected promotor and size of the transgene.²¹⁹ The first animal model of CPVT generated was an RYR2-R4496C± knock-in mouse, 28 which had exercise- and catecholamine-inducible ventricular arrhythmias. Since then, several mouse models have been designed that carry RYR2 mutations identified from patients with CPVT (see Table 1). When loss-of-function mutations are combined with gain-of-function mutations, mice are protected from CPVT.33 Identifying the pathogenicity of specific mutations is important because RYR2 mutations are implicated in several other arrhythmia disorders, as detailed below. Interestingly, a RYR2-exon3 deletion was identified in patients with a severe form of CPVT, but the corresponding mouse model failed to reproduce the CPVT phenotype.36

Cardiac calsequestrin (CASQ2) mutations are inherited in an autosomal recessive manner, leading to loss of function. Casq2 serves as a high-capacity Ca²⁺ buffer

in the SR and regulates RYR2 gating. In the mouse, CASQ2 deletion causes a severe exercise- and/or catecholamine-induced arrhythmia phenotype consistent with patients lacking Casq2 expression. Observations from CASQ2 mouse models carrying mutations identified from patients show these commonly lead to loss of Casq2 expression in the heart. Recent evidence, however, suggests that Casq2 mutations could also be inherited in an autosomal dominant manner.²²⁰ A knock-in mouse model expressing Casq2-K180R± reported catecholamine-induced arrhythmias validating this inheritance pattern.²²¹

Triadin is another protein in the SR Ca2+ release unit where autosomal recessive inheritance has been reported in patients with CPVT. When triadin was knocked out in mice, they developed substantial ultrastructural changes in the junctional SR and reduced expression of proteins comprising the Ca2+ release unit, leading to catecholamine-inducible arrhythmias.46 Calmodulin genes (CALM1/2/3) encode 3 identical proteins (CaM) and some mutations are associated with CPVT.²²² Zebrafish models have been generated to examine the pathogenesis of overexpressing CALM mutations, and they have successfully demonstrated cardiac arrhythmias (Table 1). Finally, KCNJ2 loss of function mutations, which commonly have been linked to the Anderson-Tawil Syndrome and cause reduced IK1 current (see section Long QT syndrome), have also been identified in patients with CPVT. A knock-in mouse carrying KCNJ2- R67Q± had significant evoked arrhythmias without any QT prolongation (Table 1).

The CASQ2 knockout (KO) mouse is an excellent model to investigate CPVT. It has a severe and highly penetrant phenotype, something not always observed

with RYR2 mouse models of CPVT (Table 1). The age of onset for CPVT is only a few weeks old and mice have spontaneous arrhythmias under normal housing conditions.³⁸ CASQ2 KO mice were used to establish proof of principle for the efficacy of gene-therapy in CPVT.²²³ In what may be the only example of its kind for an arrhythmia syndrome, the CASQ2 KO mouse was used to establish the therapeutic efficacy of an existing FDA-approved drug, flecainide, in CPVT.²¹⁸ CASQ2 KO mice were instrumental to demonstrate that in vivo, RYR2 block is the principal mechanism of flecainide's antiarrhythmic action.²²⁴ Since its discovery in CASQ2 KO mice, flecainide has become the standard of care for preventing arrhythmias in CPVT patients when beta-blockers are insufficient.^{225,226}

CASQ2 KO mice have also recently been used to establish a novel tissue mechanism responsible for ventricular ectopy in CPVT (Figure 3). To determine the cellular origin of ventricular arrhythmias in CPVT, Blackwell et al⁴¹ used conditional murine models with Casq2 expression only in ventricular myocardium or in the specialized conduction system, utilizing the contactin-2 promoter to drive Cre expression and control tissue-specific Casq2 expression. CPVT occurred when Casq2 was deleted in the ventricular myocardium but still expressed in the conduction system. Moreover, catecholamine challenge did not elicit any arrhythmias when Casq2 was deleted in the conduction system but still expressed in the ventricular myocardium. Additional experiments determined that the subendocardial ventricular myocardium juxtaposed to Purkinje fibers is the only cellular source for focal ventricular arrhythmias in CPVT. To understand why that was the case, in silico modeling demonstrated an intriguing phenomenon whereby subthreshold DADs in ventricular myocardium elicit full-blown APs in the conduction system to generate arrhythmias, identifying the Purkinje-myocardial junction as the tissue origin of ventricular ectopy in CPVT (Figure 3). This discovery could shape treatment and may be critical to our understanding of arrhythmogenesis in other ventricular arrhythmia syndromes where DADs are the cellular arrhythmia mechanism.

Calcium Release Deficiency Syndrome

Whereas RYR2 gain-of-function mutations have been associated with CPVT, arrhythmogenic cardiomyopathies, and AF, loss-of-function mutations in RYR2 can cause an arrhythmia syndrome recently termed Ca²⁺ release deficiency syndrome (CRDS).⁴⁹ In patients with CRDS, exercise stress tests do not provoke arrhythmias. Consequently, this syndrome can escape clinical diagnosis and often presents as sudden cardiac death. In CRDS, arrhythmias are thought to develop because of electrophysiological remodeling. Given the substantial number of benign RYR2 mutations observed, predicting pathogenicity without in vitro or in vivo examination is challenging. CRDS has only been recently described, but 1 knock-in mouse model was generated carrying the

patient-specific RYR2-D4646A± mutant allele.49 Exercise and catecholamine challenge did not invoke arrhythmias, as predicted. Importantly, this animal model enabled the authors to establish a burst pacing protocol that induced arrhythmias, supporting a possible new clinical diagnostic tool, which was recently confirmed in a clinical study. A previously reported loss-of-function mutation, RYR2-A4860G±, was identified in a patient with idiopathic VF and subsequently knocked-in to a mouse, but arrhythmias were not reported in vivo. 48 Ex vivo hearts did develop VF in response to isoproterenol but had no arrhythmias in the presence of the RyR2 agonist caffeine. These data hint that this RYR2 loss-of-function mutation may be part of the calcium release deficiency syndrome. Prior work on RYR2 has focused on gain-offunction mutations. CRDS is a relatively new syndrome and animal models provide an opportunity to advance our understanding of CRDS pathophysiology and determine the impact of loss-of-function mutations.

Long QT Syndrome

Congenital LQTS often presents as a multiorgan syndrome caused by mutations in proteins responsible for the repolarization of the heart and can cause seizures, syncope, arrhythmia, and sudden death. Alterations in repolarization manifest as prolongation of the AP and, consequently, the QT interval, predisposing the heart to EAD, DAD, and reentrant circuits. Ca²⁺, Na⁺, and many K⁺ channels play a role in cardiac repolarization; accordingly, the genes involved in this syndrome vary and LQTS can be inherited in autosomal dominant or recessive forms. LQTS is traditionally described as a distinct disease manifestation; however, long QT intervals are observed in many overlap syndromes (see Table 1). Approximately 80% of LQTS cases are caused by mutations in KCNQ1 or KCNH2, with SCN5A constituting 7% to 10% of cases.²²⁷ It should be reiterated that the currents responsible for mouse and human repolarization are guite different (Figure 2), and mice are inadequate to model human disease involving mutations in delayed rectifier K+ channels (eg, LQT1, LQT2).

An aspect of understanding LQT pathogenesis is the observed sex differences in patients, such as QT interval, the onset of cardiac events, and sudden death. Animal models allow for significant insight into these investigations since most in vitro models fail to capture the hormonal factors that influence development and regulation. It is noteworthy that sexual dimorphism in QT is not captured by mouse models and may require other animal models such as rabbits. Stradiol treatment in ovariectomized rabbits led to prolongation of the QT interval and changes in proteins responsible for repolarization, whereas dihydrotestosterone did not. Another study in rabbits recapitulated findings in patients showing that QT interval was more prolonged in females following treatment with erythromycin.

LQT1 is caused by loss-of-function mutations in KCNQ1 (Kv7.1). Knockout of KCNQ1 in mice leads to characteristics of Jervell and Lange-Nielsen syndrome, causing deafness and prolonged QT interval.⁵⁰ Other models have examined distinct mutations with varying phenotypes; however, caution should be used when reviewing the mouse LQT1 literature. Rabbits are a much better model for examining human-like mechanisms of altered repolarization. However, only 1 transgenic rabbit model is reported, which carries cardiac-specific overexpression of KCNQ1-Y315S, leading to LQT and inducible EADs and VT with sympathetic stimulation.⁵³

LQT2 is caused by loss-of-function mutations in KCNH2 (K,11.1). In contrast to humans, KCNH2 contributes little to repolarization in mice (Figure 2). As such, mouse models are not realistic for modeling LQT2 and should not be used. A rabbit transgenic model was generated with cardiac-specific overexpression of KCNH2-G628S and developed LQT, spontaneous PVT, and sudden death. Arguably, the most interesting LQT2 model is the work in zebrafish.⁵⁶ In 1 study, the endogenous ortholog of hERG (zERG) was knocked down using morpholinos and then various hERG mutants were expressed in its place. The phenotype data (prolonged APD and/or 2:1 AV block) correctly identified 39/39 pathogenic mutants and 9/10 nonpathogenic polymorphisms. Further work using this model could establish the pathogenicity of other variants of uncertain significance.

LQT3 is caused by gain-of-function mutations in SCN5A (Nav1.5). Several mouse models have been generated that recapitulate the LQT phenotype. Both the SCN5A-ΔKPQ/+ and SCN5A-ΔQKP/+ mouse models report many characteristic phenotypes such as prolonged QT interval, a more pronounced T wave, prolonged APD, arrhythmias, and sudden death, which make them suitable for examining LQT3 mechanisms and pathogenesis. 57,58 A successful bench-to-bedside study, conducted in ventricular myocytes isolated from guinea pigs, demonstrated that mexiletine restored the APD in cells treated with the Na+ channel inactivation inhibitor, anthopleurin.⁶¹ Mexiletine is now used routinely in the treatment of patients with LQT3. In rare cases, patients have been identified linking LQTS with mutations in the beta accessory proteins for SCN5A. An SCN1B knockout mouse was generated and had long QT, bradycardia, and delayed repolarization⁷⁹; interestingly, it was found that these mice had increased Na+ channel expression. Nav1.5 channel gating was unaffected, but peak and persistent currents were increased in isolated cardiomyocytes.

Andersen-Tawil syndrome is a rare LQTS associated with physical abnormalities and hypokalemic periodic paralysis and is primarily caused by loss of function mutations in KCNJ2 (Kir2.1), resulting in reduced IK1 current. Neonatal (1 day old) KCNJ2 knockout mice were characterized by long QT and bradycardia, before dying from complete cleft palate and inability to feed.⁷³

A more useful arrhythmia phenotype—LQT and spontaneous VT—was observed in the KCNJ2-T75R cardiac-specific overexpression mouse model.⁷⁴

The remaining LQTS types result from mutations in K+ channels, Ca2+ channels, and key regulators of ion channel function (Table 1). The ankyrin-B syndrome is characterized as an overlap syndrome, with long QT, sinus node dysfunction, conduction abnormalities, exerciseinduced arrhythmia, VF, and VT. The ankyrin-B knockout mouse demonstrated long QT and sinus node dysfunction⁶²; however, a more severe and faithful phenotype was reported in heterozygous mice, capturing many of the same observations from humans.63 KCNE1 knockout mice developed long QT, but only when heart rate deceleration occurred.70 A transgenic rabbit model overexpressing dominant negative KCNE1-G52R developed long OT and increased susceptibility to drug-induced arrhythmia by accelerating IKs and IKr deactivation kinetics.⁶⁹ This model could be useful for examining the proarrhythmic liability of drugs. CACNA1C (Ca.1.2) gainof-function mutations cause LQTS that can be associated with extracardiac manifestations known as Timothy syndrome. In mice, CACNA1C knockout is embryonic lethal, but heterozygous mice survived without any cardiac phenotypes. Cardiac-specific overexpression of the G406R mutation in mice led to long QTc and exerciseinduced PVCs and Torsades de Pointes (TdP).77 It was hypothesized that this mutant had altered interaction with AKAP150; in fact, when G406R-overexpressing mice were crossed with AKAP150 KO mice, they were protected against all phenotypes of the G406R mutant.

Investigators have also knocked out many of the potassium channel genes in mice to determine their effects on cardiac electrophysiology. When a dominantnegative fragment of KCNA1 was expressed, mice developed LQT and spontaneous ventricular arrhythmias.64 Interestingly, it was later shown by Glasscock et al¹⁷³ that KCNA1 is preferentially expressed (≈10-fold higher) in atria over ventricles. Programmed electrical stimulation (PES) induced AF in KCNA1 knockout mice but did not lead to any ventricular arrhythmias and no differences in QT interval were observed. Dominant negative overexpression of KCNA5, KCNB1, or KCND2 all prolonged the QT interval without any other overt electrophysiological changes. Mice do not express KCNE3 in the adult heart, but deletion led to long QTc in aged female mice because of hyperaldosteronism.⁷²

Mouse models of LQTS should be viewed with caution when the repolarizing current of interest does not reflect the human AP. As transgenic rabbit models become more common, they may find strong ground in advancing our understanding LQT pathogenesis. Rabbits accurately capture sex differences and express a similar repolarization ion channel gene profile as humans. An area of value will be evaluating predisposition to drug-induced LQT and arrhythmia, ²³⁴ which rabbit models will be best suited to answer.

Short QT Syndrome

Short QT syndrome (SQTS) is an extremely rare disorder; only a few 100 cases have been identified to date. SQTS is caused by the shortening of the cardiac AP. Like long QT syndrome, alterations in cardiac repolarization alter the QT interval. Because of the abbreviated QT interval, the refractory period is also shortened, leaving the heart susceptible to reentrant arrhythmias. Symptoms associated with short QT syndrome include both atrial and ventricular fibrillation, palpitations, and sudden cardiac death. Gain-of-function mutations in KCNH2, KCNQ1, and KCNJ2 or loss-of-function mutations in CACNA1C, CACNB2, CACNA2D1, SCN5A, and SLC4A3 have all been associated with SQTS.

There are few genetic animal models available to examine SQTS in vivo. Zebrafish carrying the KCNH2-L499P mutation have a shortened QT interval.⁸³ Knockdown of SLC4A3 in zebrafish led to a short QT interval that was rescued by expressing WT SLC4A3 but not by SLC4A3-R370H, identified from a patient with SQTS.⁸⁴ To appreciably capture the human cardiac AP, a rabbit model was engineered to overexpress KCNH2 carrying the N588K mutation.⁸² Transgenic rabbits had shortened AP and QTc but a normal T wave height. Ex vivo perfused hearts had inducible VT and VF; this model is arguably the best available and, as discussed above, rabbit models more accurately typify human cardiac repolarization.

Brugada Syndrome

Brugada syndrome (BrS) is a disorder characterized by elevated ST segment, partial bundle branch block, arrhythmia, and sudden cardiac death. It is commonly caused by mutations in SCN5A; however, ≈20 genes are now associated with BrS. Brugada ECG patterns are sometimes observed in overlap syndromes, such as with long QT syndrome, as SCN5A mutations are associated with several arrhythmia disorders. In mice, SCN5A deletion is embryonic lethal, but heterozygous mice survived and developed slowed conduction, pacing-induced VT, and fibrosis with age.88 Interestingly, the authors observed variability in phenotype penetrance that correlated with NaV1.5 expression levels.²³⁵ A mouse model for an overlap syndrome of LQT and Brugada was generated to carry 1795insD in SCN5A.60 Homozygous mice were embryonic lethal, but heterozygous mice developed sinus node dysfunction, slowed conduction, bradycardia, and QT prolongation. Interestingly, CaMKII-dependent phosphorylation of wild-type Nav1.5 appears to phenocopy this mouse model, as late current predominates at slower heart rates. Phosphomimetic and phosphoablation mouse models demonstrated that phosphorylation and oxidation modulate Nav1.5 current and susceptibility to arrhythmias. 137,236 A transgenic pig model was generated to better understand BrS disease mechanisms. Pigs were designed to carry the orthologous SCN5A-E558X/+ mutation identified in a patient diagnosed with BrS and developed conduction abnormalities and QRS widening but had no elevated ST segment, arrhythmias, or sudden death through 2 years of age. However, ex vivo hearts had increased susceptibility to VF with programmed stimulation. One debate surrounding BrS is whether many of the associated genes cause BrS or simply increase susceptibility to developing BrS. SCN5A mutations have variable and incomplete penetrance and differing phenotypes. Moreover, SCN5A mutations are prevalent in the general population and discerning pathogenesis can be difficult. Given the failure of animal BrS models to reproduce the full clinical syndrome, their utility studying BrS pathogenesis and treatment options remains to be determined.

Genetic Arrhythmia Syndromes Associated With Structural Heart Disease

Animal models have been beneficial for understanding the pathogenesis of arrhythmias caused by mutations in nonion channel genes that result in structural heart disease. The section below discusses the major arrhythmia syndromes associated with a cardiomyopathy phenotype (ARVC, HCM, and DCM) and microscopic structural disease such as AF, sick sinus syndrome, heart block, and preexcitation syndromes.

Arrhythmogenic Right Ventricular Cardiomyopathy/ Dysplasia

ARVC/dysplasia is a disease manifesting as fibrofatty replacement of the right ventricular myocardium and widespread electrophysiological remodeling, predisposing individuals to ventricular arrhythmias and increased risk of sudden death. Approximately half of the ARVC cases are caused by mutations in desmosomal proteins, which make up cell-cell mechanical junctions. Arrhythmias often arise during periods of exercise or stress, suggesting that catecholamines contribute to arrhythmogenesis. Accordingly, one of the primary phenotypes in animal models is catecholamine-induced ventricular arrhythmias.

The primary genetic model used to study ARVC has been the mouse. Of note, mice do not get fatty infiltration in the heart, one of the primary phenotypes of ARVC in humans. However, numerous cardiac phenotypes have been found, from remodeling to arrhythmogenesis. The first genetic mouse models to examine the importance of desmosomal proteins were congenital knockouts (Table 2). Mouse models with global knockout of plakoglobin, desmoplakin, plakophilin-2, and desmoglein-2 are all embryonic lethal. However, heterozygotes survived and developed varying degrees of fibrosis and arrhythmia phenotypes (Table 2). Subsequent animal models were based on patient mutations and frequently resulted in haploinsufficiency. This seems to be the primary cause, along with repression of the Wnt signaling pathway. DSC2 (desmocollin-2) mutations are rare, and the link between

pathogenesis and this protein is not well understood because data are lacking and conflicting. DSC2 knockout mice are viable but did not develop any cardiac phenotypes. However, DSC2 knockdown reportedly led to altered ultrastructure and contractile dysfunction, while overexpression also led to fibrotic remodeling. Germline knockout of DSG2 (desmoglein-2) results in embryonic lethality in mice, but the cardiac-specific knockout led to dilation, fibrosis, and electrophysiological remodeling. Finally, the inhibitor of apoptosis-stimulating protein of p53 has been linked to ARVC; knockout mouse model developed dilation, arrhythmia, and sudden death. 110

The PKP2 (plakophilin-2) mouse model is well-suited for understanding ARVC mechanisms and disease progression. Heterozygote PKP2 mice survived and developed arrhythmias in the absence of overt structural remodeling,98 which was validated in mice carrying a PKP2 truncation mutant.99 These findings raised an interesting question about the cause of arrhythmias: how do arrhythmias arise in the absence of structural changes to the heart? Because PKP2 knockout is embryonic lethal, a cardiac-specific inducible PKP2 knockout mouse was generated, which had many of the characteristic ARVC phenotypes: fibrosis, remodeling, reduced ejection fraction, arrhythmia, and sudden death. 100 In this model, PKP2 signaling regulates transcription of many genes involved in Ca2+ homeostasis and proteins of the intracellular Ca2+ release unit are downregulated, causing proarrhythmic RyR2 activity. The authors discovered that flecainide, a class Ic antiarrhythmic that inhibits RyR2 channels, effectively prevented arrhythmias in these mice. Exercise exacerbated RyR2 hyperactivity, and it was shown that membrane-permeable beta-blockers had greater efficacy than nonpermeable beta-blockers.²³⁸

ARVC has been observed in dogs and cats, generally of unknown cause. In these models, fibro-fatty replacement is commonly observed alongside syncope, arrhythmias, and sudden death. Dog models could be considered when disease mechanisms, as they relate to human cardiac (electro)physiology, are essential. Nondesmosomal proteins, such as phospholamban, RyR2, lamin A/C, transmembrane protein 43, and integrin linked kinase have all been associated with ARVC in patients; however, their role is less well understood due to the spectrum of phenotypes in ARVC. For example, some mutations in phospholamban cause DCM, which is a differential diagnosis, but may also present with ARVC. Because of the difficulty of identifying pathogenic mutations, diagnostic criteria rely on interpreting cardiac imaging and ECG data. An important line of investigation is understanding the molecular mechanisms that drive the development of the disease, as early disease progression escapes detection and the heterogeneity of genes involved makes prognosticating a diagnosis difficult.

Dilated Cardiomyopathy

Congenital DCM is characterized by ventricular dilation and is commonly caused by mutations in cytoskeletal

or myofibrillar proteins leading to remodeling, reduced ejection fraction, conduction abnormalities, arrhythmia, and sudden death. Many animal models have been created to investigate the structural and functional consequences of these mutations. Here we focus on animal models where arrhythmias are a prominent feature of the reported phenotype (Table 2). Of note, there is significant overlap with the clinical diagnosis of arrhythmogenic (right ventricular) cardiomyopathy.

A widely studied DCM mutant is SCN5A-D1275N. A wide spectrum of phenotypes in different families carrying this mutation have been reported, including AF, conduction defects, and sinus dysrhythmia. ^{239,240} It is speculated that the variation in genetic background between the families contributes to these phenotypes. Homozygous SCN5A-D1275N mice developed many of the characteristic arrhythmia phenotypes observed in humans, while heterozygous mice did not. ¹³⁸ Many changes in the ECG parameters reflected a gene dose-dependent effect and these features were also observed in zebrafish. ¹³⁹

Mutations in LMNA cause severe DCM and sudden death. Several mouse models have been generated that carry mutations identified from patients (Table 2). A variety of symptoms have been reported and spontaneous arrhythmias occur frequently. Mutations in RMB20 also have a severe phenotype in mice, commonly leading to spontaneous ventricular arrhythmias and sudden death. Duchenne muscular dystrophy has been studied in pig, dog, and mouse models, each having electrophysiological changes (Table 2).

Hypertrophic Cardiomyopathy

HCM is characterized by enlargement of the left ventricle primarily caused by mutations in sarcomeric proteins, leading to increased risk of arrhythmia and sudden death.²⁴¹ HCM is commonly caused by mutations in either the beta myosin heavy chain or myosin binding protein C, together accounting for nearly half of all cases. Mice express beta myosin heavy chain during cardiogenesis, but rapidly switch from the beta to the alpha isoform, postnatal. Numerous HCM animal models have been generated; however, the predominant focus has been the study of underlying changes in sarcomere function, contractility, and hypertrophy, without detailed examination of electrophysiological phenotypes. Many studies have reported on arrhythmia susceptibility in ex vivo hearts but only reports examining in vivo arrhythmia phenotypes are discussed here.

Alpha myosin heavy chain mutations are commonly associated with HCM, but the only reports of arrhythmias in an animal model come from the R403Q mutant, which was identified in MYH7 from a human patient. The knock-in mouse model was designed to carry R403Q in MYH6.¹⁵⁵ Mice developed HCM but only had a modest arrhythmia phenotype. A rabbit model was generated to overexpress the transgene but did not exhibit an arrhythmia phenotype.²⁴² Troponin mutations are also associated

with HCM and a transgenic mouse was generated to carry TNNI3-G203S.¹⁵³ Mice had conduction defects, but no other arrhythmias. However, when these mice were crossed with MYH6-R403Q mice, they developed a more severe phenotype with conduction defects, LQT, and catecholamine-induced VT.¹⁵⁴ Knockout of MYBPC caused long QT and spontaneous VT in mice.¹⁵⁸

Several troponin T models have been designed to carry mutations identified in patients. Knollmann et al¹⁴⁹ showed that TNNT2-I79N mice had tachycardia, isoproterenol-inducible ectopy, and spontaneous nonsustained VT, despite having no hypertrophy or fibrosis. Subsequent work in mice demonstrated that Ca²⁺sensitizing TNNT2 mutations cause inducible arrhythmias, whereas nonsensitizing mutants (R278C) do not leave the heart susceptible to arrhythmias. 151 Myofilament Ca2+ sensitization increases cytosolic Ca2+-binding affinity, alters intracellular Ca2+ homeostasis, and causes pause-dependent Ca2+-triggered arrhythmia.243 In addition, myofilament Ca2+ sensitization causes focal energy deprivation, which further increases arrhythmia susceptibility in mice.²⁴⁴ Hence, myofilament sensitization per se, caused by drugs, mutations, or posttranslational modifications after myocardial infarction, 245 is a novel arrhythmia mechanism.²⁴⁶ These reports illustrate the power of murine HCM models for discovering new arrhythmia mechanisms and identifying therapeutic targets.

Atrial Fibrillation

AF is the most common arrhythmia and is characterized by rapid, abnormal atrial rhythms, with symptoms manifesting as palpitations, syncope, stroke, and heart failure, among others. The etiology of AF is multifactorial, stemming from environmental factors, diet, lifestyle, family history, medication, and surgery. For an in-depth review of acquired AF and various animal models available, the reader is referred to this review. Many genetic animal models develop AF alongside their primary disease phenotype (see Tables 1 and 2), but the models discussed in this section are more directly related to AF as a primary pathology.

One trigger for AF is thought to be hypersensitive and leaky RyR2 channels. Thus, animal models of CPVT (both gain-of-function RyR2 and loss-of-function Casq2) have been used to investigate arrhythmogenic mechanisms and screen therapeutic modalities. ¹⁶⁵ It was demonstrated that RyR2 inhibition can attenuate AF in these models. Other CPVT models are susceptible to AF with PES. A more severe phenotype was seen with CREM mice, which developed atrial dilation and spontaneous paroxysmal and persistent AF that worsened with age. ¹⁶⁷ A second trigger for AF may be channelopathies that accentuate excitability. KCNE1, SCN5A, KCNQ1, SK2, and SK3 mutations have all been introduced into

mice, with varying phenotypes (Table 2). The KCNE1 knockout mouse develops spontaneous AF.¹⁷² SCN5A models show overt structural changes, fibrosis, conduction abnormalities, and mitochondrial injury. Proteins associated with development, signaling pathways, and transcription regulation have been found to induce AF. Atrial-specific or complete knockout of LKB1 in mice caused electrical and structural remodeling and mice developed spontaneous AF.^{179,180} A goat model overexpressing constitutively active TGF-β1 had atrial fibrosis and AF inducibility with PES.¹⁸¹

Genome-wide association studies have identified many gene loci associated with increased AF risk.²⁴⁸ AF loci include genes known to affect ion channel function, cardiogenesis, or cell-cell conduction, although in many cases, candidate genes have not been determined. There are several animal models of inherited AF, mostly in mice. A primary challenge with mouse models of AF is that they do not commonly develop spontaneous AF, instead, only uncovering the phenotype with PES. Moreover, AF typically lasts for a brief period (on the order of seconds) before resolving. However, some mouse models have more severe and protracted AF that occurs spontaneously (Table 2). AF manifests in a wide range of cardiac diseases and resolving causative versus correlative pathogenesis is ongoing for some genetic models. Despite these shortcomings, investigators have successfully captured AF phenotypes in many different mouse models for genes identified in genomewide association studies studies or laboratory testing.

Genome-wide association studies have identified loss of function (PITX2, TBX5, GJA1) and gain of function (KCNN3) variants associated with patients with AF. PITX2 heterozygous mice had normal cardiac parameters except reduced transpulmonary flow; however, ex vivo hearts were more susceptible to atrial pacing-induced arrhythmia. 176 Subsequent work showed AF in vivo with PES and several groups have used this model to study AF and explore treatment, 249 especially since PITX2 is the most common risk locus identified in patients with AF. Inducible deletion of the TBX5 transcription factor led to a much more severe phenotype of spontaneous AF and electrophysiological remodeling within 2 weeks of tamoxifen treatment.177 Gene profiling identified numerous changes in the expression of Ca2+ handling proteins and ion channels, including PITX2. These data provide a link between TBX5-PITX2 activity and electrophysiological protein regulation in the heart. AF may also be promoted by delayed conduction. Mice carrying the GJA1-G60S/+ mutation are more susceptible to pacing-induced AF.178 Mice overexpressing KCNN3 had bradyarrhythmias, heart block, abnormal AV node morphology, and sudden death.²¹²

Many other models have explored mutations or deletion of other ion channels, transcription factors,




Figure 3. Tissue-targeted CASQ2 knock-out mice help decipher the anatomic origin of ventricular ectopy in catecholaminergic polymorphic ventricular tachycardia (CPVT).

Based on a combination of tissue-targeting and in silico modeling, the PMJ was identified as the likely origin of ventricular ectopy in CPVT. DAD indicates delayed afterdepolarization; LV, left ventricular; PMJ, Purkinje myocardial junction; and RV, right ventricular. Illustration credit: Ben Smith.

developmental pathways, or hormones to examine AF (Table 2). Data from patients have frequently confirmed the downregulation of various proteins, providing yet another link to pathogenesis. AF is a complex multifactorial disease; each animal model is an opportunity to extend our understanding of this disease.

Sick Sinus Syndrome

Sick sinus syndrome occurs when the sinoatrial node is not capable of generating a normal rhythm. In heterozygous SCN5A± mice (see also Brugada Syndrome), investigators identified sick sinus syndrome phenotypes 186 and sex differences in sinoatrial node function with age.²⁵⁰ The hyperpolarization-activated cyclic nucleotide-gated channels (HCN) play an essential role in generating spontaneous pacemaker activity. Various models have examined the deletion of HCN1-4. HCN4 is the most abundant isoform, and congenital deletion in mice is embryonic lethal. However, embryonic hearts had sinus bradycardia. 194 Inducible cardiac-specific knockout led to severe bradycardia, block, and sudden death within days. 197 Mice with germline deletion of HCN1 or HCN2 survived and developed sinus dysrhythmia. 192,193 Deletion of HCN3 did not lead to bradycardia or arrhythmia, but mice had subtle perturbations in their ECG morphology at lower heart rates.²⁵¹ Perhaps the most elegant and accurate mouse model of sick sinus syndrome is an inducible HCN4-specific cellular ablation mouse.¹⁹⁸ This model accurately captured many of the phenotypes observed in patients, such as fibrosis of nodal tissue, sinoatrial dysrhythmia, supraventricular tachycardia, VT. and sudden death.

Atrioventricular Block

AV block is partial or complete disruption of impulse propagation from the atria to the ventricles. As the single site for collating atrial depolarization and passing it to the ventricle, aberrant AV node function may prevent or delay sinus rhythm from conducting into the ventricle. The AV node makes up part of the conduction system and, as such, expresses many of the same ion channels and gap junctions as the sinoatrial node, bundle of His, and Purkinje cells. AV block is observed in many of the models described above. However, some genes seem to specifically alter AV node conduction and effect block without altering sinus node rhythm. The NKX2-5 gene appears to be definitely related to AV block in patients, and mouse models carrying 2 different mutations independently validated the role of this gene in disease progression. 199,200

Preexcitation Syndrome

Preexcitation occurs when the ventricles are activated prematurely via an accessory pathway. The most common mouse models generated to study preexcitation syndrome have mutations in PRKAG2, a regulatory subunit of 5′ AMP-activated protein kinase.^{204–206} Mutations

Table 2. Animal Models of Genetic Arrhythmia Syndromes With Structural Heart Disease

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref. |
|--------------------------|-------------------------------|---|--|------|
| Arrhythmogenic rig | ht ventricular cardiomyopathy | , | | |
| Desmosomal | | | | |
| Mouse | JUP (Plakoglobin) | ± | Homs embryonic lethal; hets develop arrhythmia, RV structure change, etc | 90 |
| Mouse | JUP (Plakoglobin) | c.2037-2038delTG | Truncated form, LoF, mortality, fibrosis | 91 |
| Mouse | JUP (Plakoglobin) | 23654del2 overexpression | Repressed Wnt signaling; fibrosis, dysfunction, death | 92 |
| Zebrafish | JUP (Plakoglobin) | Knockdown | Smaller heart size, blood reflux between chambers, reduced heart rate | 93 |
| Zebrafish | JUP (Plakoglobin) | 2057del2 | Structural change, mortality | 94 |
| Mouse | DSP (Desmoplakin) | ± | Homs are embryonic lethal; hets have fibrosis, dysfunction, arrhythmias; Wnt signaling implicated | 95 |
| Mouse | DSP (Desmoplakin) | Conduction-specific KO (HCN4) | Migrating atrial pacemakers, sinus rhythm dysfunction, all without cardiac remodeling | 96 |
| Mouse | DSP (Desmoplakin) | R2834H cardiac-specific over- expression | Cardiac fibrosis, other small changes, no dysplasia | 97 |
| Mouse | PKP2 (Plakophilin 2) | ± | Homs embryonic lethal; hets histologically normal, but electrical remodeling, ultrastructure changes, arrhythmia susceptibility | 98 |
| Mouse | PKP2 (Plakophilin 2) | S329X cardiac-specific over- expression | Histologically normal, but electrical remodeling, ultrastructure changes, arrhythmia susceptibility | 99 |
| Mouse | PKP2 (Plakophilin 2) | Inducible cardiac KO | Fibrosis, arrhythmia, remodeling, death, reduced lvEF | 100 |
| Mouse | PKP2 (Plakophilin 2) | R735X AAV9-expression | RV dysfunction with exercise | 101 |
| Zebrafish | PKP2 (Plakophilin 2) | Knockdown | Structural defects, signaling reduced | 102 |
| Zebrafish | DSC2 (Desmocollin 2) | Knockdown | Altered ultrastructure, contractile dysfunction | 103 |
| Mouse | DSC2 (Desmocollin 2) | G790del/G790del, +/G790del | No phenotype | 104 |
| Mouse | DSC2 (Desmocollin 2) | WT overexpression | Necrosis, acute inflammation and patchy cardiac fibrotic remodeling | 105 |
| Mouse | DSG2 (Desmoglein 2) | КО | Embryonic lethal | 106 |
| Mouse | DSG2 (Desmoglein 2) | Cardiac-specific KO | Dilation, fibrosis, electrophysiological remodeling | 107 |
| Mouse | DSG2 (Desmoglein 2) | N271S cardiac-specific over- expression | Biventricular dilatation; spontaneous ventricular arrhythmias, cardiac dysfunction, sudden death | 108 |
| Mouse | DSG2 (Desmoglein 2) | Q558X | Fibrosis, decrease in desmosomal size and number, and reduced Wnt signaling | 109 |
| Mouse | PPP1R13L (IASPP) | КО | Inducible Ppp1r13l knockout mouse model, dilation, arrhythmia, sudden death | 110 |
| Mouse | SORBS2 (ArgBP2) | КО | QRS widening, RBBB, spontaneous PVCs, NSVT, VT | 111 |
| Nondesmosoma | I | | | |
| Mouse | RYR2 (RyR2) | R176Q± | Patients with R176Q also have T250M and present with ARVC; but mouse 176Q alone gives CPVT-like phenotype; no ARVC/D, but slight end-diastolic changes | 35 |
| Mouse | RYR2 (RyR2) | Inducible cardiac-specific KO | Sinus bradycardia, block, ventricular tachycardia, sudden death | 112 |
| Mouse | ITGB1 (Integrin β1) | Inducible cardiac specific KO | Beta1d isoform is reduced in ARVC patients; KO mice had iso/caff-inducible VT | 113 |
| Mouse | TMEM43 (Luma) | S358L± | Fibro-fatty replacement, structural abnormalities, arrhythmia, sudden death | 114 |
| Mouse | ILK (ILK) | Cardiac-specific KO | Arrhythmia, sudden death; arrhythmogenic cardiomyopathy in some patients with missense mutations in ILK | 115 |
| Mouse | RPSA (LAMR1) | 1031 bp insertion (spontaneous) | ARVC; conduction abnormalities (QRS widening); no examination of inducible arrhythmias | 116 |
| Unknown | | | | |
| Dog | Unknown | Autosomal dominant inheritance | Boxers; fatty replacement of RV myocardium, ventricular arrhythmia, syncope, sudden death | 117 |
| Dog | Unknown | Unknown | Weimaraner; syncope, ventricular arrhythmias, and sudden death, with histopathologic fatty tissue infiltration | 118 |
| Dog | Unknown | Unknown | English bulldog | 119 |

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref. |
|-----------------------|-------------------------------|--|--|------|
| Cat | Unknown | Unknown | SVT, VT, PVT, RBBB | 120 |
| Dilated cardiomyo | pathy (DCM) | | | |
| Mouse | LMNA (Lamin A/C) | ± | DCM, arrhythmia | 121 |
| Mouse | LMNA (Lamin A/C) | H222P/H222P | Chamber dilation; slowed conduction, AV block, spontaneous PVCs | 122 |
| Mouse | LMNA (Lamin A/C) | N195K/N195K | Bradycardia, exit block, AV block, arrhythmia, sudden death | 123 |
| Mouse | LMNA (Lamin A/C) | G609G/G609G | Truncating splice variant; bradycardia, QRS widening, SA block, LQT | 124 |
| Pig | LMNA (Lamin A/C) | G609G/+ | Bradycardia, SA block, short QTc; spontaneous PVT, 3rd degree block at death | 125 |
| Zebrafish | DES (Desmin) | KO or aggregating mutant | Embryonic tachycardia, arrhythmia; however, no reports of electrophysiological changes or arrhythmias in two independent KO mouse models | 126 |
| Mouse | DES (Desmin) | R349P± | Human R350P; DCM, ARVC; slowed conduction and AV block; spontaneous and induced atrial fibrillation, PVCs, and VT | 127 |
| Zebrafish | ACTN2 (F-Actin) | Knockdown | DCM, bradycardia | 128 |
| Mouse | LDB3 (ZASP) | S196L cardiac-specific over- expression | DCM, arrhythmia | 129 |
| Mouse | CDH2 (Cadherin) | cardiac-specific KO | DCM, arrhythmia, conduction defects | 130 |
| Mouse | LMOD2 (Leiomodin) | КО | DCM; LQT | 131 |
| Rat | RMB20 (RMB20) | ко | DCM; QRS widening, AV block, susceptibility to arrhythmias with PES, sudden death | 132 |
| Mouse | RMB20 (RMB20) | КО | DCM; slowed conduction, LQT; changes in ion channel and calcium- handling proteins; spontaneous calcium release in isolated cardio- myocytes | 133 |
| Mouse | RMB20 (RMB20) | S637A/S637A | DCM; spontaneous AF, spontaneous VT/VF with syncope, sudden death; much more severe cardiomyopathy than KO | 134 |
| Mouse | PLN (Phospholamban) | R14del/R14del | Mice develop severe DCM; no arrhythmias in vivo; explanted hearts have induced ventricular arrhythmias | 135 |
| Mouse | SCN5A (Nav1.5) | Cardiac-specific knockdown | Slow conduction, sudden death | 136 |
| Mouse | SCN5A (Nav1.5) | S571E/S571E | Phosphomimetic CaMKII target; LV dilation; LQT, iso-induced PVCs and VT | 137 |
| Mouse | SCN5A (Nav1.5) | D1275N/+ or D1275N/ D1275N | Homs have slow conduction, heart block, AF, VT, and DCM without significant fibrosis or myocyte disarray | 138 |
| Zebrafish | SCN5A (Nav1.5) | D1275N | Bradycardia, sinus pause, AV block, sudden death; no AF or VT observed | 139 |
| Pig | DMD (Dystrophin) | KO (exon52del) | Fibrosis, low voltage areas, and sudden death | 140 |
| Dog | DMD (Dystrophin) | X-linked DMD | Short PR interval, sinus arrest, spontaneous ventricular arrhythmias | 141 |
| Mouse | DMD (Dystrophin) | KO (mdx strain) | Tachycardia; short PR, QRS, and QTc | 142 |
| Mouse | DMD (Dystrophin) | KO (5cv strain) | Short PR interval, inducible VT | 143 |
| Mouse | VCL (Vinculin) | Cardiac-specific KO | Normal sinus rhythm, AV block, spontaneous PVT, sudden death (before onset of DCM phenotype) | 144 |
| Mouse | VASP (VASP) | Cardiac-specific overexpression | Bradycardia, AV block, sudden death | 145 |
| Mouse | REST (NRSF) | DN cardiac-specific overex- pression | Prolonged PQ, AV block, spontaneous VT, sudden death (observed as VT/VF with asystole) | 146 |
| Dog | Unknown | Autosomal dominant | Doberman Pinscher, DCM with age, PVCs on Holter monitor | 147 |
| Dog | Unknown | X-linked recessive inheritance | Great Dane; DCM; AF | 148 |
| НСМ | , | ' | | |
| Mouse | TNNT2 (TnT) | I79N cardiac-specific | Elevated diastolic Ca with elevated HR; iso-inducible ectopy; spontaneous NSVT; no hypertrophy or fibrosis | 149 |
| Mouse | TNNT2 (TnT) | +/ΔK210 and ΔK210/ΔK210 | Cardiac enlargement; HF; TdP, VF, sudden death; homs worse than hets but both had phenotype | 150 |
| Mouse | TNNT2 (TnT) | F110I, R278C, or slow skeletal isoform TG | Iso-induced PVCs and VT in mice with F110I or skeletal isoform; VT inducibility with PES in ex vivo hearts; R278C had no arrhythmias compared to control | 151 |
| Rat | TNNT2 (TnT) | Trunc transgenic overexpression | VT, VF | 152 |

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref. |
|--------------------------|-------------------------------|--|--|------|
| Mouse | TNNI3 (TnI) | G203S cardiac-specific over- expression | PR prolongation, conduction delay; no arrhythmias, but later shown to have AF | 153 |
| Mouse | TNNI3 (TnI)×MYH6 (α- MHC) | G203S×R403Q | Bradycardia, slow conduction (PR and QRS), long QTc, catecholamine-induced VT | 154 |
| Mouse | MYH6 (α-MHC) | R403Q/+ | Right axis deviation, prolonged ventricular repolarization and pro- longed sinus node recovery times; programmable VT more in males than females | 155 |
| Mouse | MYH6 (α-MHC) | R403Q overexpression | | 156 |
| Mouse | MYPBC3 (MyBP-C) | trunc/trunc | Arrhythmias with PES | 157 |
| Mouse | MYPBC3 (MyBP-C) | КО | Prolonged QTc, spontaneous PVCs and VT | 158 |
| Cat | MYPBC3 (MyBP-C) | A31P | HCM | 159 |
| Cat | MYPBC3 (MyBP-C) | R820W | HCM | 160 |
| Mouse | HRAS (H-Ras) | Cardiac-specific overexpression | Sinus arrest, idioventricular rhythm, VT, block, and AF; phenotype stronger and more penetrant in females | 161 |
| Mouse | RYR2 (RyR2) | P1124L/P1124L | Mild HCM; bradycardia, iso/caff-induced VT | 162 |
| Mouse | OBSCN (Obscurin) | R4344Q/R4344Q | Tachycardia, spontaneous PVCs and VT; all without structural remodeling | 163 |
| Kangaroo | Unknown | Unknown | Kangaroos have LV hypertrophy, short QT intervals, and are highly susceptible to VF and sudden death, especially under light anesthesia | 87 |
| AF | 1 | 1 | | |
| SR Calcium Rele | ease | | | |
| Mouse | RYR2 (RyR2) | L433P+/ , N2386I+/, R2474S± | Mutations from CPVT patients. Also develop AF with atrial PES | 34 |
| Mouse | FKBP1B (FKBP12.6) | КО | No ECG abnormalities or spontaneous arrhythmias; AF with PES | 164 |
| Mouse | CASQ2 (Casq2) | КО | AF with PES | 165 |
| Mouse | JPH2 (Junctophilin 2) | E169K/+ (non AF-associated A399S ctrl) | E169K identified in HCM family with AF; before hypertrophy- AF with PES only in E169K mice | 166 |
| Mouse | CREM (CREM) | lb∆C-X | Atrial dilation; 100% of mice developed paroxysmal and persistent AF with age | 167 |
| Mouse | NLRP3 (NLRP3) | A350V/+ | Constitutively active; normal conduction, spontaneous PACs, pacing-induced AF | 168 |
| Mouse | SLN (Sarcolipin) | КО | Cellular APD prolongation; atrial fibrosis; AF with age | 169 |
| Ion Channel | | | | |
| Mouse | SCN5A (Nav1.5) | F1759A± | Atrial and ventricular enlargement, myofibril disarray, fibrosis and mitochondrial injury, and electrophysiological dysfunction | 170 |
| Mouse | SCN5A (Nav1.5) | ΔΚΡΩ/+ | Atrial enlargement; increased susceptibility to AF with PES | 171 |
| Mouse | KCNE1 (minK) | КО | Spontaneous AF | 172 |
| Mouse | KCNA1 (Kv1.1) | КО | AF with PES | 173 |
| Mouse | KCNJ2 (Kir2.1) | T75R cardiac-specific overex- pression | Long QTc; spontaneous VT; iso-induced PVCs, VT, atrial flutter/ fibrillation | 74 |
| Mouse | KCNE5 (MiRP4) | КО | Inducible PVCs, atrial arrhythmia, PVT | 174 |
| Mouse | KCNN2 (KCa2.2) | -/- and ± | Sinus and AV node dysfunction, AF with PES | 175 |
| Structural | - | | | |
| Mouse | PITX2 (PITX2) | ± | Normal cardiac parameters except reduced transpulmonary flow (pulmonary valve narrowing), ex vivo hearts were more susceptible to atrial pacing-induced arrhythmia | 176 |
| Mouse | TBX5 (TBX5) | KO, inducible | Spontaneous AF within 2 wk post-induction, substantial arrhythmogenesis and cardiac remodeling starting after 3 wk, calcium-handling protein expression changes | 177 |
| Mouse | GJA1 (Cx43) | G60S/+ | Highly susceptible to inducible AF | 178 |
| Mouse | STK11IP (LKB1) | КО | Spontaneous AF, AV block, atrial flutter, electrical and structural remodeling | 179 |
| Mouse | STK11IP (LKB1) | Inducible atrial-specific KO | Spontaneous AF | 180 |

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref |
|-----------------------|-------------------------------|--|---|-----|
| Mouse | CALCR (calcitonin receptor) | КО | Atrial fibrosis, inducible AF | 180 |
| Goat | TGFB1 (TGF-β1) | C33S cardiac-specific overex- pression | Constitutively active TGF-beta; atrial fibrosis, prolonged P wave, AF inducible with PES, no spontaneous or persistent AF | 181 |
| Mouse | TGFB1 (TGF-β1) | C33S cardiac-specific overex- pression | Constitutively active TGF-beta; atrial fibrosis, atrial inducibility with PES | 182 |
| Mouse | MAP2K4 (MKK4) | Atrial-specific KO | Regulator of TGF-beta; reduced P wave amplitude, spontaneous atrial tachycardia, polymorphic atrial beats, AF induced ex vivo with PES | 183 |
| Mouse | ACE (ACE) | Cardiac-restricted expression | Atrial enlargement, mild fibrosis; low QRS voltage, spontaneous AF, sudden death (escape rhythm preceded death in observed cases) | 184 |
| Mouse | JDP2 (JDP2) | Cardiac-specific overexpression | QRS widening, AV block, spontaneous paroxysmal AF; atrial hypertrophy, fibrosis | 185 |
| Dog | Unknown | Unknown | Great Danes with DCM; AF | 148 |
| Sick sinus syndror | me | | , | |
| Mouse | SCN5A (Nav1.5) | ± | Bradycardia, slowed conduction, exit block | 186 |
| Mouse | SCN3B (Scn β3) | КО | Bradycardia, sinus conduction slowing, exit block, AV block | 187 |
| Mouse | NOTCH1 (notch receptor 1) | Inducible intracellular domain (Notch activation) | Bradycardia, sinus pauses, reduced conduction velocity; atrial ar- rhythmias with PES; Nkx2-5, Tbx2, Tbx5 expression altered | 188 |
| Zebrafish | SMO (smoothened) | Unreported, homozygous | Bradycardia, reduced spontaneous hyperpolarizing current | 189 |
| Mouse | SLC8A1 (NCX) | Atrial-specific KO | Bradycardia, no P waves, junctional escape rhythm (His) | 190 |
| Zebrafish | SLC8A1 (NCX) | Truncation | Embryonic lethal; embryos have atrial fibrillation/bradycardia/tachycardia; some VF but mostly silent ventricle; cardiac morphological defects | 191 |
| Mouse | HCN1 (HCN1) | КО | Bradycardia, sinus dysrhythmia, prolonged SA node recovery time, increased SA conduction time, and recurrent sinus pauses | 192 |
| Mouse | HCN2 (HCN2) | КО | Sinus dysrhythmia | 193 |
| Mouse | HCN4 (HCN4) | КО | Global or cardiac HCN4-/- embryonic lethal, but embryos have sinus bradycardia, isolated cells have no spontaneous pacemaker activity | 194 |
| Mouse | HCN4 (HCN4) | R669Q± | Homs embryonic lethal, but hets survive and develop SA exit block during exercise | 195 |
| Mouse | HCN4 (HCN4) | 573X inducible cardiac-specific overexpression | Bradycardia, but no dysrhythmia | 196 |
| Mouse | HCN4 (HCN4) | Inducible cardiac-specific KO | Bradycardia, reduced iso response, AV block, sudden death | 197 |
| Mouse | HCN4 (HCN4) | Inducible HCN4+ cell ablation | Nodal tissue fibrosis, bradycardia, exit block, SVT, VT, complete block, sudden death | 198 |
| Atrioventricular blo | ock (AV block) | | | |
| Mouse | NKX2-5 (NKX2.5) | I183P cardiac-specific overex- pression | PR prolongation, worsening AV block with age | 199 |
| Mouse | NKX2-5 (NKX2.5) | R52G± | PR prolongation, AV node smaller, AV block | 200 |
| Mouse | DMPK (DMPK) | _/_ and ± | PR prolongation as mice age, ± mice develop 1st degree block, -/- mice develop third-degree block | 201 |
| Mouse | GJA5 (Cx40) | _/_ | Conduction delay (first-degree block) | 202 |
| Mouse | TRPM4 (TRPM4) | КО | AV block (prolonged PR and QRS widening), Wenckebach | 203 |
| Preexcitation synd | rome | | | - |
| Mouse | PRKAG2 (AMPK γ2) | R302Q cardiac-specific over- expression | Hypertrophy, preexcitation, accessory pathway, QRS widening, and inducible reentrant arrhythmia | 204 |
| Mouse | PRKAG2 (AMPK γ2) | N488I overexpression | Hypertrophy, sinus bradycardia, accessory pathway, preexcitation | 205 |
| Mouse | PRKAG2 (AMPK γ2) | R531G cardiac-specific over- expression | Hypertrophy, impaired contractile function, and electrical conduction abnormalities | 206 |
| Mouse | TBX2 (TBX2) | Cardiac-specific KO | Accessory pathway, preexcitation | 207 |
| Other cardiac con | | · · · · · · · · · · · · · · · · · · · | | |
| Mouse | GJA1 (Cx43) | D378stop cardiac-specific inducible | Conducting truncation; germline deletion die right after birth; inducible model die 16 days after tamoxifen; 2-3-fold QRS widening, BBB, spontaneous MVT/PVT/VF, sudden cardiac death | 208 |

| Disease, type, animal | Human ortholog gene (protein) | Mutation | Notes | Ref. |
|-----------------------|-------------------------------|---|--|------|
| Mouse | MAP2K4 (MKK4) | Cardiac-specific KO | Reduced Cx43 expression; ~55% QRS widening, long QTc, VT with PES | 209 |
| Zebrafish | KCNJ3 (Kir3.1) | N83H cardiac-specific overex- pression | Atrial dilation; sinus arrest, sinus bradycardia, SA block, and AV block; patient mutations associated with AF | 210 |
| Mouse | CACNA1D/G (Cav1.3, Cav3.1) | CACNA1D KO or CACNA1D/ CACNA1G DKO | Sinus bradycardia, slow conduction; DKO also had 3rd degree block, escape rhythms, spontaneous VT | 211 |
| Mouse | KCNN3 (KCa2.3) | WT overexpression | Bradyarrhythmias, AV block, abnormal AV node, sudden death | 212 |
| Mouse | IRX3 (IRX-1) | КО | His-Purkinje transcription factor; normal PR, wide QRS, notched R wave, block; spontaneous PVCs and VT; iso- and exercise-induced VT | 213 |
| Developmental | | | | |
| Mouse | TBX3 (TBX3) | КО | Ectopic atrial pacemakers | 214 |
| Mouse | TBX5 (TBX5) | ± | Hypoplasia, arrhythmias, see above in atrial fibrillation | 215 |
| Mouse | MECP2 (MeCp2) | КО | X-linked; long QTc, QRS widening, pacing-inducible VT, asystole/sudden death; neuronal KO was similar | 216 |

Models are separated by disease, listing the animal species, orthologous human gene and protein, mutation, notable arrhythmia phenotypes/findings, and reference. AF indicates atrial fibrillation; APD, action potential duration; AV, atrioventricular; caff, caffeine; DAD, delayed afterdepolarization; DKO, double knockout; DN, dominant negative; EAD, early afterdepolarization; GoF, gain of function; HCM, hypertrophic cardiomyopathy; het, heterozygous; HF, heart failure; hom, homozygous; HR, heart rate; IR, ischemia-reperfusion; iso, isoproterenol; KO, knockout; LoF, loss of function; MVT, monomorphic ventricular tachycardia; NSVT, nonsustained ventricular tachycardia; PES, programmed electrical stimulation; PVC, premature ventricular complex; PVT, polymorphic ventricular tachycardia; RBBB, right bundle branch block; SVT, supraventricular tachycardia; VF, ventricular fibrillation; and WT, wild type.

in the TBX2 transcription factor also led to the development of an accessory pathway in mice.²⁰⁷

ANIMAL MODELS OF ACQUIRED ARRHYTHMIA DISORDERS

Acquired heart disease is the most common cause of increased arrhythmia risk in humans. Acquired heart disease develops over the course of a person's life becasue of structural remodeling associated with hypertension,²⁹³ coronary artery disease, 225 nonischemic cardiomyopathy,294 primary and secondary valvular disease,295 autoimmune rheumatic diseases, 294 and myocarditis. 296 Acute cardiac stress or injury leads to activation of specific cell signaling pathways, 297,298 mitochondrial dysfunction,^{299,300} altered Ca²⁺ handling,³⁰¹⁻³⁰³ and a switch in metabolism³⁰⁴ from fatty acid oxidation to glycolysis.³⁰⁵ This leads to chronic changes in gene expression of trophic and mitotic factors, inflammation, and ultimately to myocyte hypertrophy and fibrosis. Pathological fibrosis is characterized by excessive proliferation of cardiac fibroblasts and ECM (extracellular matrix) protein deposition. Several key profibrotic factors have been identified, including TGF (transforming growth factor)-β, angiotensin II and aldosterone, which contribute to the development of cardiac fibrosis regardless of the underlying pathology.306 However, the relative contribution of a distinct molecular pathway depends on the type and the degree of the initial cardiac injury. Various animal models have been developed to study aspects of these acute and chronic changes leading to arrhythmia risk and are discussed below (Table 3).

Transverse Aortic Constriction

The transverse aortic constriction (TAC) model mimics chronic hypertension or aortic stenosis by causing a stricture in the thoracic aorta.²⁵² Left ventricular and atrial pressure overload increases the wall tension leading to hypertrophy, chamber dilation, and fibrosis. 307,308 This can be done in various methods, including sutures, 252,253 inflatable cuffs³⁰⁹ or intravascular stents.²⁵⁸ Within hours of injury, myocyte hypertrophy is induced by activation of p38 MAP Kinase,310 ERK1/2,311 and PI3K/AKT signaling.³¹² Increased TGF-β production due to cytokine³¹³ and adrenergic receptor activation314,315 leads to fibroblast proliferation and collagen deposition. The reninangiotensin-aldosterone system plays an important role in developing hypertrophy and arrhythmia, as treatment with ACE inhibitors316 and spironolactone317 reduces fibrosis and improving conduction velocity. This initially leads to ventricular hypertrophy, later followed by chamber dilation.313 While recruitment of Ly6ClowCXCR1+ macrophages has been found in the LV early after TAC,318 there is histologically less inflammation in this model than in other cardiac injury models.313 Notable in this model is the upregulation of NCX³¹⁹ and downregulation of SERCA2a in myocytes over time, which is also seen in explant human hearts with reduced ejection fraction. While restoring SERCA2 has shown promise in improving systolic dysfunction320,321 and suppression of ventricular arrhythmias^{322,323} in animal models, the CUPID2 trial in patients with human hearts with reduced ejection fraction showed neutral results.324

In mice with TAC, multiple investigators have shown an increase conduction time, AP duration, and AV nodal

COMPENDIUM ON BASIC MODELS OF CARDIOVASCULAR DISEASE

Downloaded from http://ahajournals.org by on September 26, 2022

Table 3. Animal Models of Acquired Arrhythmia Disorders

| Animal | Notes | Ref. |
|------------------------|---|---------|
| Transverse aortic con | striction | |
| Mouse | Severity of injury dependent on strain (BALB/c>C57BL/6>129S1/SvlmJ) and substrain (C57BL/6Tac>C57BL/6NCrl>C57BL/6J). Model requires PES to induce in vivo arrhythmias | 252-254 |
| Rat | Develop spontaneous arrhythmias with catecholamine challenge | 255 |
| Guinea Pig | High mortality, develops catecholamine-induced arrhythmias | 256 |
| Rabbit | In vivo injury, but arrhythmia studies completed ex vivo with Langendorff system | 257 |
| Pig | Model of heart failure with preserved ejection fraction, no study of arrhythmias | 258 |
| Sheep | Model of heart failure with reduced ejection fraction, no study of arrhythmias | 259 |
| Myocardial ischemia | | |
| Mouse | Single left coronary artery leads to variation in severity of injury. Model requires PES with catecholamine challenge to induced arrhythmias | 260-262 |
| Rat | Requires PES and catecholamine challenge to induce ventricular arrhythmias reliably but rare spontaneous arrhythmias | 263 |
| Rabbit | PES done as ex vivo in Langendorff system | 264 |
| Dog | Atrial ischemia extensive studied | 265 |
| Pig | High mortality due to poor collateral circulation, early spontaneous VT/VF during injury and late model of SCD | 266 |
| Sheep | High mortality with spontaneous ventricular arrhythmia | 267 |
| AV node ablation | | |
| Rat | High mortality, particularly in male rats. Requires specialized surgical equipment and skill | 268 |
| Rabbit | Studied completed ex vivo in Langendorff system | 269 |
| Dog | Can be induced minimally invasively with low mortality | 270,271 |
| Sheep | Can be induced minimally invasively with low mortality, model of TdP | 272 |
| Chronic atrial pacing | | |
| Rat | Model for AF | 273 |
| Rabbit | In vivo pacing, but PES studied completed ex vivo with Langendorff system | 274 |
| Dog | AF and spontaneous VT model, recapitulates tachycardia mediated cardiomyopathy | 275 |
| Pig | AF model, recapitulates tachycardia-mediated cardiomyopathy | 276 |
| Chronic ventricular pa | acing | |
| Mouse | Requires tethered or ex vivo pacing | 277 |
| Rat | Model tachycardia mediated cardiomyopathy with VF induction with rapid pacing | 278,279 |
| Dog | Recapitulates tachycardia-mediated cardiomyopathy, model of spontaneous AF and VT | 280 |
| Sheep | Recapitulates tachycardia mediated cardiomyopathy with reduced ejection fraction, no studies of arrhythmia | 281 |
| Pig | Studies of affiguillia | 282 |
| Inflammation | | 1 |
| Mouse | Strain-specific susceptibility. C3H/He and DBA/2 mice susceptible to viral myocarditits while C57BL/6 are protected. BALB/c susceptible to immunogen induced myocarditis while C57BL/6 more resistive | 283,284 |
| Rat | Model of AF, but nonphysiological induction of inflammation with talc | 285 |
| Guinea Pig | | 286 |
| Dog | | 287 |
| Sheep | | 288 |
| Metabolic/drug-induc | ned | |
| Mouse | Streptozotocin and DIO models well established, increased susceptibility to AF and VT with PES | 289,290 |
| Rat | Age-dependent fibrosis found in Fisher 344 rat strain, model of AF | 291 |
| Rabbit | Established model of clofilium-induced TdP | 292 |

Representative list of animal models with reference to the method to their development. AF indicates atrial fibrillation; DIO, diet-induced obesity; PES, programmed electrical stimulation; SCD, sudden cardiac death; TdP, Torsades de Pointes; VF, ventricular fibrillation; and VT, ventricular tachycardia.

refractory period, QT prolongation with inducible atrial (50%–60%) and ventricular arrhythmias (40%–50%), but spontaneous arrhythmias in vivo are rare.^{325–328} Redistribution of connexin-43 (Cx43) laterally away from intercalating disk has been purposed to explain, in part, the changes in conduction in this model.³²⁷ Arrhythmia induction can be challenging in this model, which is likely due to variability in the extent of constriction postprocedure,²⁵⁴ the length of time of injury before analysis, sex, and strain.^{329–333} Care must be taken when comparing results from different injury protocols and strains in this model.

Similarly, rabbits, rats, and guinea swine develop ventricular hypertrophy and dilation after TAC, albeit over a more extended period (8 versus 4 weeks in mice)334 and with a higher incidence of sudden cardiac death. Unlike the mouse model, rabbits³³⁵⁻³³⁷ (aortic insufficiency with abdominal aortic constriction), rats,322,326 and guinea pigs^{338–340} develop spontaneous arrhythmias in response to catecholamine challenge. Similarly, these animals develop ventricular hypertrophy, chamber dilation and fibrosis after TAC, albeit over a more extended period (8 versus 4 weeks in mice)334 and with a higher incidence of sudden cardiac death. AP prolongation is a hallmark in these models and is linked to increased $\rm I_{NaL}$ and $\rm I_{NCX}$ with reduced $\rm I_{CaL}$ responsisely. siveness to β-adrenergic stimulus and increased CaMKII activity, which is seen in human heart failure myocytes.341 Overall, these animals seem to represent a better model for arrhythmias than mice, albeit most studies use explanted heart in the Langendorff system.

Unlike small animals, no studies report an increase in arrhythmia in large animals with TAC. Ascending aortic constriction in pig and sheep has been described with polyester band²⁵⁹ or an implanted inflatable cuff.^{309,342} The swine model of TAC differs from other species as they tend to develop heart failure with preserved ejection fraction characterized by LV hypertrophy with diastolic dysfunction.^{343–346} In contrast to pig, sheep develop cardiac dysfunction at 6 to 18 weeks post-TAC with elevated markers of ECM remodeling, chemokine production, and apoptosis,³⁴⁷ but no study reported arrhythmia generation outside procedural effects.

Myocardial Ischemia

Acute and chronic myocardial ischemia is a major cause of ventricular arrhythmias in humans. During acute ischemic, myocytes are exposed to hypoxia, acidosis, increased extracellular K+, and intracellular Ca²+0.348^{,349} Under ischemic conditions, cardiomyocyte mitochondria switch to glycolysis from fatty acid oxidation to maximize ATP production with limited oxygen supply.³⁵⁰ In addition, hypoxia leads to reactive oxygen species production that further damages intracellular proteins and organelles.³⁵¹ Myocardial infarction occurs with myocyte apoptosis and replacement fibrosis if normal blood flow is not restored. This is a highly inflammatory model, with significant infiltration of

CD11b⁺ macrophages and upregulation of inflammatory cytokines (CCL2, TNF-α, and IL-10) after injury, leading to proinflammatory Ly6c^{high}CCR2⁺ macrophages early, then pro-wound healing Ly6c^{low}CXCR1⁺ chronically, which contribute to interstitial fibrosis in the border zone.³⁵² Chronically, myocardial fibrosis leads to conduction heterogeneity, a substrate for reentry arrhythmias.

In mice and rats, myocardial ischemia is induced surgically, either transiently by ischemia/reperfusion (I/R) injury^{260,353} or complete occlusion by coronary artery ligation^{261,354} or cryoinjury.^{262,353,355} Unlike complete occlusion, I/R injury produces reversible ischemia that leads to significant myocardial dysfunction without widespread necrosis in the area at risk. Spontaneous ventricular arrhythmias are observed during the reperfusion phase.²⁶⁰ However, spontaneous ventricular arrhythmias are rare after complete occlusion aside from isolated preventricular contractions (internal data). In addition, LV dysfunction leads to volume overload in the left atrium, causing fibrosis and susceptibility to AF.³⁵⁶

Atrial and ventricular arrhythmias can be induced universally in isolated explanted post-MI hearts, with the occurrence of VT (inducible in up to 90%-100% in mice), VF (inducible in up to 89% of rats), and AF (inducible in 73% of rats and 33% of mice).357-363 In vivo ventricular arrhythmia induction is more difficult, requiring rapid pacing protocols^{364–366} and a catecholamine challenge. Transvenous, pericardial, and transesophageal protocols have been described for AF and ventricular arrhythmia induction, with the occurrence of VT (20%-70% in mice and rats)³⁶⁷ and AF (60%-90% in mice).368 Electrophysiological study of isolated myocytes from acute and chronic infarcted hearts has shown both shortening of the myocardial effective refractory period and slowing of condition time in the left atria, infarct, and border zone.357 The proposed mechanism for AF induction is reduced expression of Cx40, dephosphorylation of Cx40 and Cx43, and redistribution from the intercalated disc to the lateral cell membrane.³⁶²

Established models for rabbit myocardial ischemia focus extensively on ex vivo studies using Langendorf perfusion systems. After infarction, rabbit myocardium shows prolonged APD and Ca²⁺ transients ex vivo.³⁶⁹ Unlike mice and rats, infarction in rabbits leads to delayed AV nodal conduction becaue of fibrosis and reduction in Cx40 expression but did not affect the ventricular effective refractory period.³⁷⁰ No arrhythmia induction protocol was used in these studies.

Dog models focus on atrial ischemia and arrhythmias, with ~40% of animals developing AF. Increased I $_{\rm NCX}$ current, spontaneous Ca $^{2+}$ leak, and conduction heterogeneity were found in myocytes from the border zone of the atrial infarct, supporting both triggered and reentry as the mechanism in this model. The use of beta-blocker (nadolol) and Ca $^{2+}$ channel inhibitor (nifedipine) were more effective at suppressing atrial arrhythmia in this model than class Ic (flecainide) or class III (dofetilide) antiarrhythmic drugs.

Pig myocardial ischemia models are well established as they are of similar size and physiology as humans. Ischemia can be induced by either transient intravascular occlusion of the coronary artery or chronic occlusion with an ameroid constructor.374 Acutely, pig are exquisitely sensitive to ischemia because of lack of functional collaterals at baseline, which leads to significant procedural mortality from VF.375 After chronic ischemia, SCD due to spontaneous ventricular arrhythmias occurs in ≈60% to 70% of animals by 3 months.376,377 Myocytes in the remote zone from have a decreased rapid delayed rectifier K+ current (I_k), altered Na⁺-Ca²⁺ exchange current (I_{NCX}), and increases of late Na⁺ current (I_{Nal}), Ca²⁺-activated K^+ current [$I_{K(Ca)}$], and Ca²⁺activated CI⁻ current [I_{CICa})]. In addition, myocytes in the border zone show the same changes along with a decrease of L-type Ca²⁺ current (I_{Cal}), a decrease of inward rectifier K+ current (I_{K1}), and arrhythmogenic SR Ca²⁺ release-induced EADs and DADs.378 These changes in the current lead to shortening of the APD in the border zone and prolonged APD in the remote region, setting up a substrate for both triggered and reentrant arrhythmias. Gene therapy with dominant-negative K+ channel (KCNH2-G628S)379 or Cx43³⁸⁰ reduced VT induction by prolonging the ADP and effective refractory period in the border zone.

Complete Heart Block and AV Node Ablation

Complete heart block leads to AV dyssynchrony and bradycardia, which acutely causes reduced cardiac output and induces volume overload. Compensatory hemodynamic changes occur to improve cardiac function, including ventricular dilation, hypertrophy, and increased stroke volume but are not able to fully restore cardiac output.³⁸¹ While genetic models are available for primary complete heart block, secondary complete heart block, which is characterized by fibrosis and necrosis of the AV node, is more difficult to reproduce. Outside histological remodeling, little is known about the molecular changes associated with secondary complete heart block.

Given the size, mouse AV node is challenging to identify without immunostaining. Genetic have generated to establish the important transcription factors and ion channels in the AV node (noted in the above section atrioventricular block). Interestingly, disruption of tissue resident macrophages in the AV node, by either knockout of Cx43 in these cells or by genetic ablation of macrophages using diphtheria toxin receptor/diphtheria treatment, lead to progressive AV conduction block. While these resident macrophages were noted in human AV nodes, it unclear if the presence or absence in of these cells contribute to human disease.

Electrical needle AV nodal ablation in rats has been described, as it was noted that alcohol injection leads to either transient block or mortality based on the amount used. This is a technically challenging model with variable success and high early mortality because of bradycardia and

ventricular arrhythmias, as only 6-month-old female rats survived past 3 days.²⁶⁸ Regardless, the surviving rats recapitulated the cardiac remodeling in humans, with 80% showing spontaneous TdP at baseline which could be induced to sustained VT with PES and isoproterenol challenge.²⁶⁸

The dog model of AV ablation by transvenous catheter injection for formaldehyde has been well established and leads to compensated hypertrophy and QT prolongation. This is due to APD prolongation in the setting of bradycardia, with $\approx\!50\%$ of animals developing spontaneous and 90% drug-induced TdP. Chronically, this reduced I current and enhanced Ca²+ influx by NCX, leading to increased SR Ca²+ content, which increases the risk for DADs. The contrast, goats undergoing AV ablation did not show APD prolongation but led to increased PLB (phospholamban), Troponin-I and myosin light chain kinase by PKA and RyR2 by CaMKII, suggesting increased Ca²+ sensitivity in this model.

Chronic Tachypacing

Overriding the normal conduction system has many deleterious effects but is often reversible. Rapid atrial heart rate or pacing can lead to tachycardia-mediated heart failure, and long-term RV pacing in humans can lead to ventricular dyssynchrony, reduced cardiac output, and heart failure.³⁸¹ Long-term pacing has been shown to be detrimental to LV, with significant wall motion abnormalities and perfusion defects in the inferior and apical walls without corresponding coronary disease.³⁸⁸ Conversely, cardiac resynchronization therapy improves HF outcomes for patients with HF and left-bundle branch block.

RV pacing has been used as a model of nonischemic cardiomyopathy, developing significant systolic dysfunction. Long-term RV pacing in AV nodal ablated dogs show the same perfusion mismatches seen in humans and was associated with increased sympathetic innervation of the ventricles. 389 Overdrive RV pacing for 4 weeks in dogs can induce spontaneous ventricular arrhythmias and SCD in 25% of dogs.390 Myocytes isolated from paced ventricles showed prolonged ADP with reduced I_{to} currents, 391 likely due in part to downregulation of Kv4.3³⁹² and increased I_{NaL}³⁹³ in failing heart. In addition, Ca2+ transients showed reduced amplitude, slowed relaxation, and blunted frequency dependence due to reduction in SERCA2a and upregulation of NCX in failing myocytes.³⁹⁴ Cardiac resynchronization therapy in this model was showed to normalization of APD, reduce the I_{Nat} current, and prevent the negative remodeling associated with heart failure in this model. 393,395,396 VF can be induced in pig by applying AC current to the RV, but using arrhythmic drugs to improve resuscitation was not seen.³⁹⁷

Ventricular pacing in dogs also leads to secondary atrial fibrosis, dilation, and reduced function as the LV fails, inducing AF. 398,399 As with rapid ventricular pacing, rapid atrial pacing leads to a reduction in the $\rm I_{to}$, in addition to $\rm I_{Cal.}$ and $\rm I_{Ks}$ currents. 400 If pacing is stopped and

the animal is allowed to recover, the ion currents return to normal, but fibrosis remains, leading to persistent AF. Studying atrial cells from dogs undergoing both rapid atrial and ventricular pacing has shown the atrial ion channel expression in HF, AT, and HF with AT can be significantly different.⁴⁰¹ Upregulation of profibrotic miRNA has also been described in atrial paced dogs, providing a novel target to prevent atrial fibrosis and AF.^{402,403}

In rabbits, rapid atrial pacing increases atrial fibrosis and TGF- β signaling, which can be attenuated with losartan. Atrial pacing leads to decrease KCNE1 KCNB2 expression, reduced I_{Ks}, and shortening the AERP. This was thought to be due to microRNA-1 upregulation in the atria.

In sheep, natriuretic peptide release is found immediately after RV pacing, returning to normal after cessation.²⁸¹ In goats, chronic atrial pacing led to atrial dilation, with reduced PKA phosphorylation of PLB and increased CaMKII phosphorylation of RyR2, leading to reduced SR Ca²⁺ load.²⁷² While structural changes similar to humans with tachycardia mediated cardiomyopathy are seen, increased arrhythmogenesis has not been reported.

Given their small size, in vivo pacing is difficult in mice. Tethered epicardial pacing has been used to study AV dyssynchrony and synchrony in mice after I/R injury. Dyssynchrony leads to further deterioration of cardiac function and activation of p38, ERK1/2, JNK, and MSK1 and inhibition of the GSK3 β pathways. This was reversed by resynchrony.²⁷⁷ Recently, the development of fully implantable epicardial micro pacing technology may allow for longitudinal pacing studies.⁴⁰⁵

In rats, initiation of rapid atrial pacing leads to upregulation of multiple voltage-gated K+ channels (Kv1.5, Kv4.2, and Kv4.3), which contribute to repolarization by $I_{\rm Kr}$ and $I_{\rm To}$. After 2 days of atrial pacing, AF can be induced in $\approx\!20\%$ of animals, with upregulation of associated AF genes (CASQ2, KCNJ2, and TGFB) and activation of the TGF- β and IL-6 pathways. Apply a days are pacing of the LV has been used to reliably induce VF to study the effects of medication for resuscitation.

Inflammation

Myocardial inflammation is a known driver of atrial and ventricular arrhythmias. Postprocedural arrhythmias are common after cardiothoracic surgery but, while they are usually self-limiting, they lead to prolonged hospital stays. While there is usually a preexisting arrhythmogenic substrate due to the underlying disease, surgical scarring and inflammation further exacerbate the system, leading to arrhythmia. Large cohort studies have shown elevated proinflammatory cytokines associated with persistent AF, including CRP, $^{411-413}$ TNF- α , IL-1 β , IL-6, and IL-10. 414,415 After surgery, there is an increase in both macrophages and neutrophils to the surgical site, with reactive oxygen species production from MPO (myeloperoxidase) activity.

Inflammation because of myocarditis is more complex, and the course of the acute and chronic phase of the immune response is dependent on the underlying cause (infectious versus rheumatological). The cytokine profiles are found in viral and autoimmune myocarditis includes more proinflammatory monocyte infiltration and myocyte necrosis than postsurgical injury.⁴¹⁸ This leads to a multitude of ECG changes, including sinus tachycardia, widened QRS patterns, low voltage, prolonged QT, variable AV blocks, and diffuse ST-elevations.⁴¹⁹⁻⁴²¹

Sterile inflammation has been used to induce AF in dogs²⁸⁷ and sheep.²⁸⁸ Pericardial talc treatment of dog atria to induce sterile inflammation induced AF in $\approx\!60\%$ of dogs and was significantly reduced with topical steroids or NSAIDS.²⁸⁷ In sheep, treatment with atorvastatin reduced hs-CRP, IL-6, and TNF- α expression, which improved the atrial effective refractory period at 72 hours.²⁸⁸ While inflammatory myocarditis can be induced in rats²⁸⁵ and guinea pigs,⁴²² no current reports on the arrhythmia potential are available.

Currently, there are 2 models of viral myocarditis due to exposure to coxsackievirus B3 (CVB3)423 and encephalomyocarditis virus A (EMCV).424 There are several issues with the viral myocarditis models in animals. First, of the viruses primarily associated with human myocarditis (parvovirus B19, herpes simplex 9 and coxsackievirus B3), only CVB3 is infectious to animals, with EMCV only rarely causing human disease newborns. Second, the CVB3 mouse model is highly inflammatory and more mimics childhood infections than the milder course in adults. 425 Third, only particular stains of mice are susceptible to viral infection. Regardless, C3H/He mice exposed to develop similar arrhythmias to humans (80% sinus arrest, 30% second or third-degree AV block, 30% PACs, 20% PVCs, and 10% VT).426 Ex vivo electrophysiological studies showed no change to the APD, but mice exposed to CVB3 were hyperpolarized with slightly increased VERP.427 EMCV exposed DBA/2 mice develop AV block in 40% of mice over 2 weeks, with two-thirds of those mice showing mononuclear cell infiltration and edema and another onethird showing necrosis of the conduction system. 428 No further electrophysiological studies were conducted.

Traditionally, myocarditis can be induced in mice by either immunizing with a cardiac structural peptide (myosin heavy chain [MHC- α] or cardiac troponin I)²⁸³ or delivering primed dendritic cells pulsed with MHC- α .²⁸⁴ Importantly, BALB/c mice are susceptible to peptide immunized myocarditis while C57BL/6 strains are resistant, from which a majority of transgenic lines are created.⁴²⁹ MHC- α peptide-induced myocarditis showed significant immune cell infiltration, increased expression of both TNF α and INF γ , fibrosis and prolongation for the APD in ventricular myocytes, which all could be attenuated with atorvastatin.^{285,422,430}

Ctla4+/- Pdcd1-/- mice spontaneously develop myocarditis, modeling immune checkpoint inhibitor-induced myocarditis.⁴³¹ These mice succumb to progressive ventricular hypertrophy and SCD early. Progressive AV block and sinus arrest occurs in ~30% of transgenic mice, similar to arrhythmia seen in patient with immune checkpoint inhibitor–induced myocarditis. Further study using this model would greatly improve treatment for patient with adverse events after immune checkpoint inhibitor therapy.

Metabolic- and Drug-Induced Arrhythmia

Dietary and metabolic considerations also contribute to atrial arrhythmia development as there is a known association with BMI and diabetes in AF, 432,433 and incidence of paroxysmal AF is reduced after gastric bypass surgery. 434 Off-target drug effects lead to adverse clinical outcomes, particularly arrhythmias induction due to block of delayed rectifier K+ channels, I $_{\rm Kr}$, causing drug-induced long QT syndrome and TdP ventricular arrhythmias.

Streptozotocin-induced diabetic models have been developed for both mice and rats435 and consistently shown prolonged APD, increased sympathetic innervations, and inflammation. Studies have shown this is likely due to production of advanced glycation end products (AGEs) in diabetes animals, but they are inconstant in the electrophysiological mechanisms. 436 Initial studies showed a reduction in the I, current as the underlying cause of AF,437-439 while others suggest changes in sinoatrial node connexin channel expression, 440,441 atrial myocyte Ca²⁺ handling proteins (TRPC1/6, RyR3)⁴⁴² and AV nodal ion channels (TRPC1, CASQ2, RYR2, and RYR3)443 are involved. Diabetic mice and rats also have reduced K+ channel expression, leading to overall reduced K+ currents, which was dependent on glycosylation of CaMKII and activation PKC.444 Further studies showed in the setting of hyperglycemia, mice had more diastolic calcium leak through RyR2, prolonged ADP90, ADP alternans, increased DADs, and frequent premature ventricular complex, which could be suppressed by genetic inhibition of CaMKII.445 Blocking IL-1β activation of CaMKII in this model restores the APD to normal and suppresses VT induction in explanted hearts from diabetic mice. 446 These studies all provide a central role for CaMKII in regulating ion channel expression and function in hyperglycemia.

Diet-induced obesity has been shown to affect inflammation and gene expression in multiple disease models. Mice fed 3 months of a high-fat diet had a 15% increase in their QTc and increased I_{Ks} current than mice on a normal diet and developed a 10-fold increase in premature ventricular complex burden.⁴⁴⁷ The increased I_{Ks} current was thought to be due to the reduced expression of voltagegated K⁺ channels. Diet-induced obesity mice were also found to have reduced Na_V1.5 expression and current, leading to reduced ADP and conduction velocity in the atrial and increased incidence of induced AF.⁴⁴⁸ Diet-induced obesity rats over 8 weeks have increased expression of Ca_V1.2, HCN4, Kir2.1, RYR2, NCX, and SERCA2a in the LV, which

may contribute to DADs and triggered PVCs. 449 Rabbits fed a high-fat diet had increased cardiac sympathetic innervation (as seen by increased GAP23 expression), prolonged ADP and increased I_{Ca}, which led to QTc prolongation and repolarization heterogeneity in the ventricle. 450 Isolated hearts were more susceptible to VT induction. While these studies highlight the important changes to ion channel expression animals due to a high-fat Western diet, further study is needed to linking these findings to humans.

A specific line of inbred rats (Fischer F344 at 20–24 months) develop age-dependent adverse cardiac remodeling, with males developing more cardiomyocyte hypertrophy, intestinal fibrosis, and systolic dysfunction and females with more cardiac hypertrophy and diastolic dysfunction. Aged female Fischer F344 rats show enlarged atrial, fibrosis, and CD68+ monocyte infiltration, similar to human disease. Both male and female Fischer 344 rats are more susceptible to AF induction by atrial pacing, with 80% of animals showing atrial arrhythmia. These are the only models of spontaneous AF in aged animals and could provide important insight to mechanism and future therapies for AF in our aging population.

In rabbits treated with clofilium (K+ channel blocker), 70% of animals develop TdP, which can be increased to 100% with the addition of α1 agonist methoxamine. Blockade of CaMK (calmodulin kinase) or PKA in this model reduced pause-dependent VT that was independent of these kinases effect on L-type Ca²+ channel activity. Interestingly, the dose-dependent CaMKII blockade did not change the QT interval, unlike the dose-dependent PKA blockade. Further study showed that blocking CaMKII reduced the ratio of the TU interval, which is likely more critical than the QT interval in the induction of pause-dependent VT. 456

EMERGING ETIOLOGIES OF ARRHYTHMOGENESIS: OPPORTUNITY FOR ANIMAL MODELS

Cardiac electrophysiology is regulated not only by the amino acid sequence at the protein level but also by posttranslational modifications, micropeptides, epigenetics, long noncoding RNAs (lncRNA), micro RNAs (miRs), aging, and environmental factors. Unprecedented advances in sequencing technology, deep mutational scanning, and epigenetic and transcriptome mapping have highlighted several new areas of pathogenesis. A primary challenge with many of the ascribed changes is whether they directly cause disease or, rather, follow disease progression. Animal models have been generated to address this challenge and study new areas of arrhythmia biology, with a select few highlighted below.

In the past decade, micropeptides have been identified that were previously excluded during annotations of open reading frames due to their small size or position in the transcriptome. In the heart, functionally expressed

peptides include examples such as sarcolipin and dwarf open reading frame, which regulate SR Ca reuptake. Sarcolipin protein levels are decreased in humans with AF or heart failure⁴⁵⁷ and the sarcolipin knockout mouse developed AF with age. 169 In mice with DCM, overexpression of dwarf open reading frame attenuated the heart failure phenotype, although arrhythmias were not examined.458 The micropeptides apelin and elabela affect cardiogenesis, fibrosis, hypertrophy, and inotropic responses. Reduced apelin levels predicted major myocardial events and MI scar size in patients with a previous MI event.⁴⁵⁸ Apelin knockout mice developed larger scars and increased mortality following MI, although arrhythmias were not described. 459 These findings establish the pathogenic role and therapeutic potential for micropeptides in the heart, but their impact on arrhythmogenesis remains to be examined.

Posttranslational modifications, primarily phosphorylation, are documented in many arrhythmias. While phosphorylation by PKC, PKA, and CaMKII are the most studied mediators for changes in Na⁺ channels, 137,236,461,462 Ca²⁺ channels, K⁺ channels, Hosphorylation of different phosphorylation sites to ion currents. Phosphomimetic and phosphoablation mouse models have been created to tease out the importance of each site to arrhythmogenesis. 137,463,469,470 In addition, posttranslational modification of signaling kinases themselves, including PKA⁴⁷¹ and CaMKII, Iplay a role in arrhythmia susceptibility in response to oxidative stress. Other posttranslational modifications are less well studied, and it remains to be determined whether these cause disease or are merely epiphenomena.

Epigenetic DNA modifications regulate protein expression. It is now recognized that many cardiovascular diseases are associated with epigenetic changes. For example, cardiac-specific deletion of HDAC1 and HDAC2 in mice led to dilated cardiomyopathy, arrhythmia, and premature death. Are A family with a history of autosomal recessive DCM was later discovered to carry homozygous mutation in GATAD1. Are peigenetics also contribute to risk as it relates to hypertension, obesity, age, and other factors. Are Patients with persistent Are have genome-wide changes in DNA methylation, and animal models could help elucidate the pathophysiology of these changes at the whole genome or protein level.

MiRs regulate gene expression by RNA complementation and silencing. In the heart, miRs control cardiogenesis and pathways to affect gene expression during development. Some miRs are muscle specific and may serve as macroregulators of ion channel expression. In disease, changes in many different miRs have been documented. In dogs with rapid pacing, nicotine administration reduced miR-133 and miR-590 levels and increased AF risk.⁴⁷⁶ Atrial tachypacing in dogs and development of AF was also associated with changes in many miRs compared with control.⁴⁷⁷ Dogs with ventricular tachypacing developed CHF and AF and miR-29b levels declined within the first

24 hours of pacing.⁴⁷⁸ In zebrafish, miR-182 served as a TBX-5 effector and miR-182 upregulation led to block, tachycardia, and arrhythmias.⁴⁷⁹ In mice, it was shown that miR-1 directly binds to the C-terminus of Kir2.1 and represses channel activity.⁴⁸⁰ Moreover, a single nucleotide polymorphism in miR-1 identified from patients with AF did not suppress Kir2.1 channel activity and failed to rescue arrhythmia inducibility in miR-1 knockdown mice (whereas WT miR-1 did). Of most interest is the therapeutic potential for RNA regulation to reverse disease progression.

LncRNA regulates gene expression via several mechanisms and changes in lncRNA expression have been documented in a wide range of diseases. Altered expression of lncRNA has been observed in AF,⁴⁸¹ including altered regulation of PITX2.⁴⁸² Another study in rabbits with AF induced by atrial tachypacing purported a mechanism whereby lncRNA "sponges" up miR-328 to regulate CAC-NA1C.⁴⁸³ Yet another report indicated changes in lncRNA in AF reduce expression of JP2 and RYR2.⁴⁸⁴ These studies demonstrate additional mechanisms of regulation beyond controlling gene expression and open up the field to exciting new opportunities in arrhythmia research.

CONCLUSIONS

For decades, animal models have been an essential tool to study arrhythmia pathophysiology and therapeutic approaches. Recent work with genetic mouse models yielded a new diagnostic tool in CRDS, identified lifesaving drug therapies for CPVT, LQT3, and ARVC, identified pathogenic mechanisms, and advanced our understanding of arrhythmogenesis in ways that other models cannot. The broad availability of transgenic mouse models and the option to generate mice with cell-specific and/or time-dependent regulation of gene expression provides a significant advantage of the mouse over other small animal species. A major limitation is that their fast heart rate, small heart size, and differential ionic currents do not fully recapitulate human cardiac electrophysiology. While guinea pigs and rabbits can overcome some of these electrophysiological limitations, genetic manipulation is limited in these species. Large animal electrophysiological and hemodynamic parameters match humans more closely but can be prohibitively expensive. Regardless, large animals are useful for preclinical studies evaluating new drugs, gene therapy, and devices that necessitate large animal models before moving to humans. Here, we provide an in-depth review of existing animal models to help interpreting published arrhythmia mechanisms as well as planning future experimental studies investigating cardiac arrhythmia diseases.

ARTICLE INFORMATION

Affiliation

Vanderbilt Center for Arrhythmia Research and Therapeutics, Division of Clinical Pharmacology, Vanderbilt University Medical Center, Nashville, TN.

Sources of Funding

This work was supported in part by National Institutes of Health (NIH) grant NHLBI R35 HL144980 (Dr Knollmann), the Leducq Foundation grant 18CVD05 (Dr Knollmann), the American Heart Association grant 19SFRN34830019 (Dr Knollmann), and NIH LRP GSNT8763 (Dr Schmeckpeper).

Disclosures

None.

REFERENCES

- Khurshid S, Choi SH, Weng LC, Wang EY, Trinquart L, Benjamin EJ, Ellinor PT, Lubitz SA. Frequency of cardiac rhythm abnormalities in a half million adults. Circ Arrhythm Electrophysiol. 2018;11:e006273. doi: 10.1161/CIRCEP.118.006273
- Echt DS, Liebson PR, Mitchell LB, Peters RW, Obias-Manno D, Barker AH, Arensberg D, Baker A, Friedman L, Greene HL. Mortality and morbidity in patients receiving encainide, flecainide, or placebo. the cardiac arrhythmia suppression trial. N Engl J Med. 1991;324:781–788. doi: 10.1056/NEJM199103213241201
- Cardiac Arrhythmia Suppression Trial III. Effect of the antiarrhythmic agent moricizine on survival after myocardial infarction. N Engl J Med. 1992;327:227–33. doi: 10.1056/NEJM199207233270403
- Delisle BP, Anson BD, Rajamani S, January CT. Biology of cardiac arrhythmias: ion channel protein trafficking. *Circ Res.* 2004;94:1418–1428. doi: 10.1161/01.RES.0000128561.28701.ea
- Antzelevitch C, Burashnikov A. Overview of basic mechanisms of cardiac arrhythmia. Card Electrophysiol Clin. 2011;3:23–45. doi: 10.1016/j. ccep.2010.10.012
- Gupta DK, Giugliano RP, Ruff CT, Claggett B, Murphy S, Antman E, Mercuri MF, Braunwald E, Solomon SD; Effective Anticoagulation with Factor Xa Next Generation in AFTiMIESI. The prognostic significance of cardiac structure and function in atrial fibrillation: The ENGAGE AF-TIMI 48 echocardiographic substudy. J Am Soc Echocardiogr. 2016;29:537–44. doi: 10.1016/j.echo.2016.03.004
- Morita N, Mandel WJ, Kobayashi Y, Karagueuzian HS. Cardiac fibrosis as a determinant of ventricular tachyarrhythmias. *J Arrhythm.* 2014;30:389–394. doi: 10.1016/j.joa.2013.12.008
- Cao JM, Fishbein MC, Han JB, Lai WW, Lai AC, Wu TJ, Czer L, Wolf PL, Denton TA, Shintaku IP, et al. Relationship between regional cardiac hyperinnervation and ventricular arrhythmia. *Circulation*. 2000;101:1960–1969. doi: 10.1161/01.cir.101.16.1960
- Brachmann J, Kabell G, Scherlag B, Harrison L, Lazarra R. Analysis of interectopic activation patterns during sustained ventricular tachycardia. Circulation. 1983;67:449–456. doi: 10.1161/01.cir.67.2.449
- Kannankeril PJ, Moore JP, Cerrone M, Priori SG, Kertesz NJ, Ro PS, Batra AS, Kaufman ES, Fairbrother DL, Saarel EV, et al. Efficacy of flecainide in the treatment of catecholaminergic polymorphic ventricular tachycardia: a randomized clinical trial. *JAMA Cardiol.* 2017;2:759–766. doi: 10.1001/jamacardio.2017.1320
- Mazzanti A, Maragna R, Faragli A, Monteforte N, Bloise R, Memmi M, Novelli V, Baiardi P, Bagnardi V, Etheridge SP, et al. Gene-specific therapy with mexiletine reduces arrhythmic events in patients with long QT syndrome type 3. J Am Coll Cardiol. 2016;67:1053–1058. doi: 10.1016/j.jacc.2015.12.033
- Zipes DP, Jalife J, Stevenson WG. Cardiac electrophysiology: from cell to bedside. Seventh edition. ed. Elsevier; 2018.
- Varró A, Tomek J, Nagy N, Virág L, Passini E, Rodriguez B, Baczkó I. Cardiac transmembrane ion channels and action potentials: cellular physiology and arrhythmogenic behavior. *Physiol Rev.* 2021;101:1083–1176. doi: 10.1152/physrev.00024.2019
- Varró A, Lathrop DA, Hester SB, Nánási PP, Papp JG. Ionic currents and action potentials in rabbit, rat, and guinea pig ventricular myocytes. *Basic Res Cardiol.* 1993;88:93–102. doi: 10.1007/BF00798257
- Joukar S. A comparative review on heart ion channels, action potentials and electrocardiogram in rodents and human: extrapolation of experimental insights to clinic. Lab Anim Res. 2021;37:25. doi: 10.1186/ s42826-021-00102-3
- Bers DM. Cardiac excitation-contraction coupling. Nature. 2002;415:198– 205. doi: 10.1038/415198a
- 17. Bers DM. Excitation-contraction coupling and cardiac contractile force. 2nd ed. Kluwer Academic Publishers; 2001.
- Mangoni ME, Nargeot J. Genesis and regulation of the heart automaticity. *Physiol Rev.* 2008;88:919–982. doi: 10.1152/physrev.00018.2007

- Haïssaguerre M, Jaïs P, Shah DC, Takahashi A, Hocini M, Quiniou G, Garrigue S, Le Mouroux A, Le Métayer P, Clémenty J. Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. N Engl J Med. 1998;339:659–666. doi: 10.1056/NEJM199809033391003
- Weiss JN, Garfinkel A, Karagueuzian HS, Chen PS, Qu Z. Early afterdepolarizations and cardiac arrhythmias. *Heart Rhythm.* 2010;7:1891–1899. doi: 10.1016/j.hrthm.2010.09.017
- 21. Bers DM. Calcium and cardiac rhythms: physiological and pathophysiological. *Circ Res.* 2002;90:14–17.
- Tse G. Mechanisms of cardiac arrhythmias. J Arrhythm. 2016;32:75–81. doi: 10.1016/j.joa.2015.11.003
- 23. Wellens HJ. Contemporary management of atrial flutter. *Circulation*. 2002;106:649-652. doi: 10.1161/01.cir.0000027683.00417.9a
- Katritsis DG, Camm AJ. Atrioventricular nodal reentrant tachycardia. Circulation. 2010;122:831–840. doi: 10.1161/CIRCULATIONAHA.110.936591
- Killu AM, Stevenson WG. Ventricular tachycardia in the absence of structural heart disease. Heart. 2019;105:645–656. doi: 10.1136/heartjnl-2017-311590
- Wessels A, Sedmera D. Developmental anatomy of the heart: a tale of mice and man. *Physiol Genomics*. 2003;15:165–176. doi: 10.1152/ physiolgenomics.00033.2003
- Kumar D, Hacker TA, Buck J, Whitesell LF, Kaji EH, Douglas PS, Kamp TJ. Distinct mouse coronary anatomy and myocardial infarction consequent to ligation. *Coron Artery Dis.* 2005;16:41–44. doi: 10.1097/00019501-200502000-00008
- Cerrone M, Colombi B, Santoro M, di Barletta MR, Scelsi M, Villani L, Napolitano C, Priori SG. Bidirectional ventricular tachycardia and fibrillation elicited in a knock-in mouse model carrier of a mutation in the cardiac ryanodine receptor. *Circ Res.* 2005;96:e77–e82. doi: 10.1161/01. RES.0000169067.51055.72
- Loaiza R, Benkusky NA, Powers PP, Hacker T, Noujaim S, Ackerman MJ, Jalife J, Valdivia HH. Heterogeneity of ryanodine receptor dysfunction in a mouse model of catecholaminergic polymorphic ventricular tachycardia. Circ Res. 2013;112:298–308. doi: 10.1161/CIRCRESAHA.112.274803
- Lehnart SE, Mongillo M, Bellinger A, Lindegger N, Chen BX, Hsueh W, Reiken S, Wronska A, Drew LJ, Ward CW, et al. Leaky Ca2+ release channel/ryanodine receptor 2 causes seizures and sudden cardiac death in mice. J Clin Invest. 2008;118:2230–2245. doi: 10.1172/JCl35346
- Zhang Y, Fraser JA, Jeevaratnam K, Hao X, Hothi SS, Grace AA, Lei M, Huang CL. Acute atrial arrhythmogenicity and altered Ca(2+) homeostasis in murine RyR2-P2328S hearts. *Cardiovasc Res.* 2011;89:794–804. doi: 10.1093/cyr/cyg229
- Nakamura Y, Yamamoto T, Kobayashi S, Tamitani M, Hamada Y, Fukui G, Xu X, Nishimura S, Kato T, Uchinoumi H, et al. Ryanodine receptor-bound calmodulin is essential to protect against catecholaminergic polymorphic ventricular tachycardia. *JCI Insight*. 2019;4. doi: 10.1172/jci.insight.126112
- Chen W, Wang R, Chen B, Zhong X, Kong H, Bai Y, Zhou Q, Xie C, Zhang J, Guo A, et al. The ryanodine receptor store-sensing gate controls Ca2+ waves and Ca2+-triggered arrhythmias. *Nat Med.* 2014;20:184–192. doi: 10.1038/nm.3440
- 34. Shan J, Xie W, Betzenhauser M, Reiken S, Chen BX, Wronska A, Marks AR. Calcium leak through ryanodine receptors leads to atrial fibrillation in 3 mouse models of catecholaminergic polymorphic ventricular tachycardia. Circ Res. 2012;111:708–717. doi: 10.1161/CIRCRESAHA.112.273342
- Kannankeril PJ, Mitchell BM, Goonasekera SA, Chelu MG, Zhang W, Sood S, Kearney DL, Danila Cl, De Biasi M, Wehrens XH, et al. Mice with the R176Q cardiac ryanodine receptor mutation exhibit catecholamine-induced ventricular tachycardia and cardiomyopathy. *Proc Natl Acad Sci U S A*. 2006;103:12179–12184. doi: 10.1073/pnas.0600268103
- Liu Y, Wang R, Sun B, Mi T, Zhang J, Mu Y, Chen J, Bround MJ, Johnson JD, Gillis AM, Chen SR. Generation and characterization of a mouse model harboring the exon-3 deletion in the cardiac ryanodine receptor. *PLoS One*. 2014;9:e95615. doi: 10.1371/journal.pone.0095615
- Fischer E, Gottschalk A, Schüler C. An optogenetic arrhythmia model to study catecholaminergic polymorphic ventricular tachycardia mutations. Sci Rep. 2017;7:17514. doi: 10.1038/s41598-017-17819-8
- 38. Knollmann BC, Chopra N, Hlaing T, Akin B, Yang T, Ettensohn K, Knollmann BE, Horton KD, Weissman NJ, Holinstat I, et al. Casq2 deletion causes sarcoplasmic reticulum volume increase, premature Ca2+ release, and catecholaminergic polymorphic ventricular tachycardia. *J Clin Invest.* 2006;116:2510–2520. doi: 10.1172/JCl29128
- Song L, Alcalai R, Arad M, Wolf CM, Toka O, Conner DA, Berul CI, Eldar M, Seidman CE, Seidman JG. Calsequestrin 2 (CASO2) mutations increase expression of calreticulin and ryanodine receptors,

- causing catecholaminergic polymorphic ventricular tachycardia. *J Clin Invest.* 2007;117:1814–1823. doi: 10.1172/JCI31080
- Flores DJ, Duong T, Brandenberger LO, Mitra A, Shirali A, Johnson JC, Springer D, Noguchi A, Yu ZX, Ebert SN, et al. Conditional ablation and conditional rescue models for Casq2 elucidate the role of development and of cell-type specific expression of Casq2 in the CPVT2 phenotype. *Hum Mol Genet*. 2018;27:1533–1544. doi: 10.1093/hmg/ddy060
- Blackwell DJ, Faggioni M, Wleklinski MJ, Gomez-Hurtado N, Venkataraman R, Gibbs CE, Baudenbacher FJ, Gong S, Fishman GI, Boyle PM, et al. The Purkinje-myocardial junction is the anatomic origin of ventricular arrhythmia in CPVT. JCI Insight. 2022;7:e151893. doi: 10.1172/jci.insight.151893
- Rizzi N, Liu N, Napolitano C, Nori A, Turcato F, Colombi B, Bicciato S, Arcelli D, Spedito A, Scelsi M, et al. Unexpected structural and functional consequences of the R33Q homozygous mutation in cardiac calsequestrin: a complex arrhythmogenic cascade in a knock in mouse model. *Circ Res.* 2008;103:298–306. doi: 10.1161/CIRCRESAHA.108.171660
- Wleklinski MW, Parikh S, Blackwell DJ, Knollmann BC. Autosomal-dominant CASQ2-K180R causes CPVT by altering intra-SR calcium buffering without reducing Casq2 protein levels. *Biophys J.* 2021;120:238a-238a. doi:10.1016/j.bpj.2020.11.1572
- Søndergaard MT, Sorensen AB, Skov LL, Kjaer-Sorensen K, Bauer MC, Nyegaard M, Linse S, Oxvig C, Overgaard MT. Calmodulin mutations causing catecholaminergic polymorphic ventricular tachycardia confer opposing functional and biophysical molecular changes. FEBS J. 2015;282:803–816. doi: 10.1111/febs.13184
- Da'as SI, Thanassoulas A, Calver BL, Beck K, Salem R, Saleh A, Kontogianni I, Al-Maraghi A, Nasrallah GK, Safieh-Garabedian B, et al. Arrhythmogenic calmodulin E105A mutation alters cardiac RyR2 regulation leading to cardiac dysfunction in zebrafish. *Ann N Y Acad Sci.* 2019;1448:19–29. doi: 10.1111/nyas.14033
- Chopra N, Yang T, Asghari P, Moore ED, Huke S, Akin B, Cattolica RA, Perez CF, Hlaing T, Knollmann-Ritschel BE, et al. Ablation of triadin causes loss of cardiac Ca2+ release units, impaired excitation-contraction coupling, and cardiac arrhythmias. *Proc Natl Acad Sci U S A*. 2009;106:7636–7641. doi: 10.1073/pnas.0902919106
- 47. Reilly L, Alvarado FJ, Lang D, Abozeid S, Van Ert H, Spellman C, Warden J, Makielski JC, Glukhov AV, Eckhardt LL. Genetic loss of IK1 causes adrenergic-induced phase 3 early afterdepolariz ations and polymorphic and bidirectional ventricular tachycardia. *Circ Arrhythm Electrophysiol.* 2020;13:e008638. doi: 10.1161/CIRCEP.120.008638
- Zhao YT, Valdivia CR, Gurrola GB, Powers PP, Willis BC, Moss RL, Jalife J, Valdivia HH. Arrhythmogenesis in a catecholaminergic polymorphic ventricular tachycardia mutation that depresses ryanodine receptor function. *Proc Natl Acad Sci U S A*. 2015;112:E1669–E1677. doi: 10.1073/pnas.1419795112
- Sun B, Yao J, Ni M, Wei J, Zhong X, Guo W, Zhang L, Wang R, Belke D, Chen YX, et al. Cardiac ryanodine receptor calcium release deficiency syndrome. Sci Transl Med. 2021;13. doi: 10.1126/scitranslmed.aba7287
- Casimiro MC, Knollmann BC, Ebert SN, Vary JC Jr, Greene AE, Franz MR, Grinberg A, Huang SP, Pfeifer K. Targeted disruption of the Kcnq1 gene produces a mouse model of Jervell and lange-nielsen syndrome. *Proc Natl Acad Sci U S A*. 2001;98:2526–2531. doi: 10.1073/pnas.041398998
- Demolombe S, Lande G, Charpentier F, van Roon MA, van den Hoff MJ, Toumaniantz G, Baro I, Guihard G, Le Berre N, Corbier A, et al. Transgenic mice overexpressing human KvLQT1 dominant-negative isoform. part I: phenotypic characterisation. *Cardiovasc Res.* 2001;50:314–327. doi: 10.1016/s0008-6363(01)00231-0
- Lubberding AF, Zhang J, Lundh M, Nielsen TS, Søndergaard MS, Villadsen M, Skovhøj EZ, Boer GA, Hansen JB, Thomsen MB, et al. Age-dependent transition from islet insulin hypersecretion to hyposecretion in mice with the long OT-syndrome loss-of-function mutation Kcnq1-A340V. *Sci Rep.* 2021;11:12253. doi: 10.1038/s41598-021-90452-8
- Brunner M, Peng X, Liu GX, Ren XQ, Ziv O, Choi BR, Mathur R, Hajjiri M, Odening KE, Steinberg E, et al. Mechanisms of cardiac arrhythmias and sudden death in transgenic rabbits with long QT syndrome. *J Clin Invest*. 2008;118:2246–2259. doi: 10.1172/JCI33578
- Arnaout R, Ferrer T, Huisken J, Spitzer K, Stainier DY, Tristani-Firouzi M, Chi NC. Zebrafish model for human long QT syndrome. *Proc Natl Acad Sci U S A*. 2007;104:11316–11321. doi: 10.1073/pnas.0702724104
- Langheinrich U, Vacun G, Wagner T. Zebrafish embryos express an orthologue of HERG and are sensitive toward a range of QT-prolonging drugs inducing severe arrhythmia. *Toxicol Appl Pharmacol.* 2003;193:370–382. doi: 10.1016/j.taap.2003.07.012
- 56. Jou CJ, Barnett SM, Bian JT, Weng HC, Sheng X, Tristani-Firouzi M. An in vivo cardiac assay to determine the functional consequences of

- putative long QT syndrome mutations. Circ Res. 2013;112:826-830. doi: 10.1161/CIRCRESAHA.112.300664
- Nuyens D, Stengl M, Dugarmaa S, Rossenbacker T, Compernolle V, Rudy Y, Smits JF, Flameng W, Clancy CE, Moons L, et al. Abrupt rate accelerations or premature beats cause life-threatening arrhythmias in mice with long-QT3 syndrome. Nat Med. 2001;7:1021–1027. doi: 10.1038/nm0901-1021
- Montnach J, Chizelle FF, Belbachir N, Castro C, Li L, Loussouarn G, Toumaniantz G, Carcouët A, Meinzinger AJ, Shmerling D, et al. Arrhythmias precede cardiomyopathy and remodeling of Ca2+ handling proteins in a novel model of long QT syndrome. J Mol Cell Cardiol. 2018;123:13–25. doi: 10.1016/j.yjmcc.2018.08.019
- Tian XL, Yong SL, Wan X, Wu L, Chung MK, Tchou PJ, Rosenbaum DS, Van Wagoner DR, Kirsch GE, Wang Q. Mechanisms by which SCN5A mutation N1325S causes cardiac arrhythmias and sudden death in vivo. Cardiovasc Res. 2004;61:256–267. doi: 10.1016/j.cardiores.2003.11.007
- Remme CA, Verkerk AO, Nuyens D, van Ginneken AC, van Brunschot S, Belterman CN, Wilders R, van Roon MA, Tan HL, Wilde AA, et al. Overlap syndrome of cardiac sodium channel disease in mice carrying the equivalent mutation of human SCN5A-1795insD. Circulation. 2006;114:2584–2594. doi: 10.1161/CIRCULATIONAHA.106.653949
- Priori SG, Napolitano C, Cantù F, Brown AM, Schwartz PJ. Differential response to Na+ channel blockade, beta-adrenergic stimulation, and rapid pacing in a cellular model mimicking the SCN5A and HERG defects present in the long-QT syndrome. Circ Res. 1996;78:1009–1015. doi: 10.1161/01.res.78.6.1009
- Chauhan VS, Tuvia S, Buhusi M, Bennett V, Grant AO. Abnormal cardiac Na(+) channel properties and QT heart rate adaptation in neonatal ankyrin(B) knockout mice. Circ Res. 2000;86:441–447. doi: 10.1161/01.res.86.4.441
- Mohler PJ, Schott JJ, Gramolini AO, Dilly KW, Guatimosim S, duBell WH, Song LS, Haurogné K, Kyndt F, Ali ME, et al. Ankyrin-B mutation causes type 4 long-QT cardiac arrhythmia and sudden cardiac death. *Nature*. 2003;421:634–639. doi: 10.1038/nature01335
- 64. London B, Jeron A, Zhou J, Buckett P, Han X, Mitchell GF, Koren G. Long QT and ventricular arrhythmias in transgenic mice expressing the N terminus and first transmembrane segment of a voltage-gated potassium channel. *Proc Natl Acad Sci U S A.* 1998;95:2926–2931. doi: 10.1073/pnas.95.6.2926
- Li H, Guo W, Yamada KA, Nerbonne JM. Selective elimination of I(K,slow1) in mouse ventricular myocytes expressing a dominant negative Kv1.5alpha subunit. Am J Physiol Heart Circ Physiol. 2004;286:H319-H328. doi: 10.1152/ajpheart.00665.2003
- Xu H, Barry DM, Li H, Brunet S, Guo W, Nerbonne JM. Attenuation of the slow component of delayed rectification, action potential prolongation, and triggered activity in mice expressing a dominant-negative Kv2 alpha subunit. Circ Res. 1999;85:623–633. doi: 10.1161/01.res.85.7.623
- 67. Barry DM, Xu H, Schuessler RB, Nerbonne JM. Functional knockout of the transient outward current, long-QT syndrome, and cardiac remodeling in mice expressing a dominant-negative Kv4 alpha subunit. *Circ Res.* 1998;83:560–567. doi: 10.1161/01.res.83.5.560
- 68. Guo W, Li H, London B, Nerbonne JM. Functional consequences of elimination of i(to,f) and i(to,s): early afterdepolarizations, atrioventricular block, and ventricular arrhythmias in mice lacking Kv1.4 and expressing a dominant-negative Kv4 alpha subunit. *Circ Res.* 2000;87:73–79. doi: 10.1161/01.res.87.1.73
- Major P, Baczkó I, Hiripi L, Odening KE, Juhász V, Kohajda Z, Horváth A, Seprényi G, Kovács M, Virág L, et al. A novel transgenic rabbit model with reduced repolarization reserve: long QT syndrome caused by a dominantnegative mutation of the KCNE1 gene. Br J Pharmacol. 2016;173:2046– 2061. doi: 10.1111/bph.13500
- Drici MD, Arrighi I, Chouabe C, Mann JR, Lazdunski M, Romey G, Barhanin J. Involvement of IsK-associated K+ channel in heart rate control of repolarization in a murine engineered model of Jervell and lange-nielsen syndrome. *Circ Res.* 1998;83:95–102. doi: 10.1161/01.res.83.1.95
- Hu Z, Kant R, Anand M, King EC, Krogh-Madsen T, Christini DJ, Abbott GW. Kcne2 deletion creates a multisystem syndrome predisposing to sudden cardiac death. *Circ Cardiovasc Genet.* 2014;7:33–42. doi: 10.1161/ CIRCGENETICS.113.000315
- Hu Z, Crump SM, Anand M, Kant R, Levi R, Abbott GW. Kcne3 deletion initiates extracardiac arrhythmogenesis in mice. FASEB J. 2014;28:935–945. doi: 10.1096/fj.13-241828
- Zaritsky JJ, Redell JB, Tempel BL, Schwarz TL. The consequences of disrupting cardiac inwardly rectifying K(+) current (I(K1)) as revealed by the targeted deletion of the murine Kir2.1 and Kir2.2 genes. J Physiol. 2001;533(pt 3):697–710. doi: 10.1111/j.1469-7793.2001.t01-1-00697.x

- Lu CW, Lin JH, Rajawat YS, Jerng H, Rami TG, Sanchez X, DeFreitas G, Carabello B, DeMayo F, Kearney DL, et al. Functional and clinical characterization of a mutation in KCNJ2 associated with andersen-tawil syndrome. *J Med Genet*. 2006;43:653–659. doi: 10.1136/jmg.2006.040816
- 75. Kuo HC, Cheng CF, Clark RB, Lin JJ, Lin JL, Hoshijima M, Nguyêñ-Trân VT, Gu Y, Ikeda Y, Chu PH, et al. A defect in the Kv channel-interacting protein 2 (KChIP2) gene leads to a complete loss of I(to) and confers susceptibility to ventricular tachycardia. *Cell.* 2001;107:801–813. doi: 10.1016/s0092-8674(01)00588-8
- Seisenberger C, Specht V, Welling A, Platzer J, Pfeifer A, Kühbandner S, Striessnig J, Klugbauer N, Feil R, Hofmann F. Functional embryonic cardiomyocytes after disruption of the L-type alpha1C (Cav1.2) calcium channel gene in the mouse. *J Biol Chem.* 2000;275:39193–39199. doi: 10.1074/jbc.M006467200
- Cheng EP, Yuan C, Navedo MF, Dixon RE, Nieves-Cintrón M, Scott JD, Santana LF. Restoration of normal L-type Ca2+ channel function during timothy syndrome by ablation of an anchoring protein. *Circ Res.* 2011;109:255-261. doi: 10.1161/CIRCRESAHA.111.248252
- Berchtold MW, Zacharias T, Kulej K, Wang K, Torggler R, Jespersen T, Chen JN, Larsen MR, la Cour JM. The arrhythmogenic calmodulin mutation D129G dysregulates cell growth, calmodulin-dependent kinase II activity, and cardiac function in zebrafish. *J Biol Chem.* 2016;291:26636–26646. doi: 10.1074/jbc.M116.758680
- Lopez-Santiago LF, Meadows LS, Ernst SJ, Chen C, Malhotra JD, McEwen DP, Speelman A, Noebels JL, Maier SK, Lopatin AN, Isom LL. Sodium channel Scn1b null mice exhibit prolonged QT and RR intervals. *J Mol Cell Cardiol*. 2007;43:636–647. doi: 10.1016/j.yjmcc.2007.07.062
- Itokawa K, Sora I, Schindler CW, Itokawa M, Takahashi N, Uhl GR. Heterozygous VMAT2 knockout mice display prolonged OT intervals: possible contributions to sudden death. *Brain Res Mol Brain Res.* 1999;71:354–357. doi: 10.1016/s0169-328x(99)00194-1
- O'Brien SE, Apkon M, Berul CI, Patel HT, Saupe K, Spindler M, Ingwall JS, Zahler R. Phenotypical features of long Q-T syndrome in transgenic mice expressing human Na-K-ATPase alpha(3)-isoform in hearts. Am J Physiol Heart Circ Physiol. 2000;279:H2133-H2142. doi: 10.1152/ajpheart.2000.279.5.H2133
- 82. Odening KE, Bodi I, Franke G, Rieke R, Ryan de Medeiros A, Perez-Feliz S, Fürniss H, Mettke L, Michaelides K, Lang CN, et al. Transgenic short-QT syndrome 1 rabbits mimic the human disease phenotype with QT/action potential duration shortening in the atria and ventricles and increased ventricular tachycardia/ventricular fibrillation inducibility. Eur Heart J. 2019;40:842–853. doi: 10.1093/eurheartj/ehy761
- 83. Hassel D, Scholz EP, Trano N, Friedrich O, Just S, Meder B, Weiss DL, Zitron E, Marquart S, Vogel B, et al. Deficient zebrafish ether-à-go-go-related gene channel gating causes short-QT syndrome in zebrafish reggae mutants. *Circulation*. 2008;117:866–875. doi: 10.1161/CIRCULATIONAHA.107.752220
- 84. Thorsen K, Dam VS, Kjaer-Sorensen K, Pedersen LN, Skeberdis VA, Jurevičius J, Treinys R, Petersen IMBS, Nielsen MS, Oxvig C, et al. Loss-of-activity-mutation in the cardiac chloride-bicarbonate exchanger AE3 causes short QT syndrome. Nat Commun. 2017;8:1696. doi: 10.1038/s41467-017-01630-0
- Pott C, Ren X, Tran DX, Yang MJ, Henderson S, Jordan MC, Roos KP, Garfinkel A, Philipson KD, Goldhaber JI. Mechanism of shortened action potential duration in Na+-Ca2+ exchanger knockout mice. Am J Physiol Cell Physiol. 2007;292:C968-C973. doi: 10.1152/ajpcell. 00177.2006
- Schilling JM, Horikawa YT, Zemljic-Harpf AE, Vincent KP, Tyan L, Yu JK, McCulloch AD, Balijepalli RC, Patel HH, Roth DM. Electrophysiology and metabolism of caveolin-3-overexpressing mice. *Basic Res Cardiol*. 2016;111:28. doi: 10.1007/s00395-016-0542-9
- O'Rourke MF, Avolio AP, Nichols WW. The kangaroo as a model for the study of hypertrophic cardiomyopathy in man. *Cardiovasc Res.* 1986;20:398– 402. doi: 10.1093/cvr/20.6.398
- Papadatos GA, Wallerstein PM, Head CE, Ratcliff R, Brady PA, Benndorf K, Saumarez RC, Trezise AE, Huang CL, Vandenberg JI, et al. Slowed conduction and ventricular tachycardia after targeted disruption of the cardiac sodium channel gene Scn5a. *Proc Natl Acad Sci U S A*. 2002;99:6210–6215. doi: 10.1073/pnas.082121299
- Shy D, Gillet L, Ogrodnik J, Albesa M, Verkerk AO, Wolswinkel R, Rougier JS, Barc J, Essers MC, Syam N, et al. PDZ domain-binding motif regulates cardiomyocyte compartment-specific NaV1.5 channel expression and function. *Circulation*. 2014;130:147–160. doi: 10.1161/ CIRCULATIONAHA.113.007852

- Kirchhof P, Fabritz L, Zwiener M, Witt H, Schäfers M, Zellerhoff S, Paul M, Athai T, Hiller KH, Baba HA, et al. Age- and training-dependent development of arrhythmogenic right ventricular cardiomyopathy in heterozygous plakoglobin-deficient mice. Circulation. 2006;114:1799–1806. doi: 10.1161/CIRCULATIONAHA.106.624502
- Zhang Z, Stroud MJ, Zhang J, Fang X, Ouyang K, Kimura K, Mu Y, Dalton ND, Gu Y, Bradford WH, et al. Normalization of naxos plakoglobin levels restores cardiac function in mice. *J Clin Invest*. 2015;125:1708–1712. doi: 10.1172/JCI80335
- Lombardi R, da Graca Cabreira-Hansen M, Bell A, Fromm RR, Willerson JT, Marian AJ. Nuclear plakoglobin is essential for differentiation of cardiac progenitor cells to adipocytes in arrhythmogenic right ventricular cardiomyopathy. Circ Res. 2011;109:1342–1353. doi: 10.1161/CIRCRESAHA.111.255075
- Martin ED, Moriarty MA, Byrnes L, Grealy M. Plakoglobin has both structural and signalling roles in zebrafish development. *Dev Biol.* 2009;327:83–96. doi: 10.1016/j.ydbio.2008.11.036
- Asimaki A, Kapoor S, Plovie E, Karin Arndt A, Adams E, Liu Z, James CA, Judge DP, Calkins H, Churko J, et al. Identification of a new modulator of the intercalated disc in a zebrafish model of arrhythmogenic cardiomyopathy. Sci Transl Med. 2014;6:240ra74. doi: 10.1126/scitranslmed.3008008
- Garcia-Gras E, Lombardi R, Giocondo MJ, Willerson JT, Schneider MD, Khoury DS, Marian AJ. Suppression of canonical Wnt/beta-catenin signaling by nuclear plakoglobin recapitulates phenotype of arrhythmogenic right ventricular cardiomyopathy. *J Clin Invest*. 2006;116:2012–2021. doi: 10.1172/JCI27751
- Mezzano V, Liang Y, Wright AT, Lyon RC, Pfeiffer E, Song MY, Gu Y, Dalton ND, Scheinman M, Peterson KL, et al. Desmosomal junctions are necessary for adult sinus node function. *Cardiovasc Res.* 2016;111:274–286. doi: 10.1093/cvr/cvw083
- 97. Yang Z, Bowles NE, Scherer SE, Taylor MD, Kearney DL, Ge S, Nadvoretskiy VV, DeFreitas G, Carabello B, Brandon Ll, et al. Desmosomal dysfunction due to mutations in desmoplakin causes arrhythmogenic right ventricular dysplasia/cardiomyopathy. *Circ Res.* 2006;99:646–655. doi: 10.1161/01.RES.0000241482.19382.c6
- Cerrone M, Noorman M, Lin X, Chkourko H, Liang FX, van der Nagel R, Hund T, Birchmeier W, Mohler P, van Veen TA, et al. Sodium current deficit and arrhythmogenesis in a murine model of plakophilin-2 haploinsufficiency. Cardiovasc Res. 2012;95:460–468. doi: 10.1093/cvr/cvs218
- 99. Moncayo-Arlandi J, Guasch E, Sanz-de la Garza M, Casado M, Garcia NA, Mont L, Sitges M, Knöll R, Buyandelger B, Campuzano O, et al. Molecular disturbance underlies to arrhythmogenic cardiomyopathy induced by transgene content, age and exercise in a truncated PKP2 mouse model. *Hum Mol Genet*. 2016;25:3676–3688. doi: 10.1093/hmg/ddw213
- 100. Cerrone M, Montnach J, Lin X, Zhao YT, Zhang M, Agullo-Pascual E, Leo-Macias A, Alvarado FJ, Dolgalev I, Karathanos TV, et al. Plakophilin-2 is required for transcription of genes that control calcium cycling and cardiac rhythm. Nat Commun. 2017;8:106. doi: 10.1038/s41467-017-00127-0
- 101. Cruz FM, Sanz-Rosa D, Roche-Molina M, García-Prieto J, García-Ruiz JM, Pizarro G, Jiménez-Borreguero LJ, Torres M, Bernad A, Ruíz-Cabello J, et al. Exercise triggers ARVC phenotype in mice expressing a disease-causing mutated version of human plakophilin-2. *J Am Coll Cardiol*. 2015;65:1438–1450. doi: 10.1016/j.jacc.2015.01.045
- 102. Moriarty MA, Ryan R, Lalor P, Dockery P, Byrnes L, Grealy M. Loss of plakophilin 2 disrupts heart development in zebrafish. *Int J Dev Biol.* 2012;56:711-718. doi: 10.1387/ijdb.113390mm
- 103. Heuser A, Plovie ER, Ellinor PT, Grossmann KS, Shin JT, Wichter T, Basson CT, Lerman BB, Sasse-Klaassen S, Thierfelder L, et al. Mutant desmocollin-2 causes arrhythmogenic right ventricular cardiomyopathy. Am J Hum Genet. 2006;79:1081–1088. doi: 10.1086/509044
- 104. Hamada Y, Yamamoto T, Nakamura Y, Sufu-Shimizu Y, Nanno T, Fukuda M, Ono M, Oda T, Okuda S, Ueyama T, et al. G790del mutation in DSC2 alone is insufficient to develop the pathogenesis of ARVC in a mouse model. *Biochem Biophys Rep.* 2020;21:100711. doi: 10.1016/j.bbrep.2019.100711
- 105. Brodehl A, Belke DD, Garnett L, Martens K, Abdelfatah N, Rodriguez M, Diao C, Chen YX, Gordon PM, Nygren A, Gerull B. Transgenic mice overexpressing desmocollin-2 (DSC2) develop cardiomyopathy associated with myocardial inflammation and fibrotic remodeling. *PLoS One*. 2017;12:e0174019. doi: 10.1371/journal.pone.0174019
- 106. Eshkind L, Tian Q, Schmidt A, Franke WW, Windoffer R, Leube RE. Loss of desmoglein 2 suggests essential functions for early embryonic development and proliferation of embryonal stem cells. Eur J Cell Biol. 2002;81:592–598. doi: 10.1078/0171-9335-00278

- 107. Kant S, Holthöfer B, Magin TM, Krusche CA, Leube RE. Desmoglein 2-Dependent Arrhythmogenic Cardiomyopathy Is Caused by a Loss of Adhesive Function. Circ Cardiovasc Genet. 2015;8:553-563. doi: 10.1161/CIRCGENETICS.114.000974
- 108. Pilichou K, Remme CA, Basso C, Campian ME, Rizzo S, Barnett P, Scicluna BP, Bauce B, van den Hoff MJ, de Bakker JM, et al. Myocyte necrosis underlies progressive myocardial dystrophy in mouse dsg2related arrhythmogenic right ventricular cardiomyopathy. J Exp Med. 2009;206:1787-1802. doi: 10.1084/jem.20090641
- 109. Calore M, Lorenzon A, Vitiello L, Poloni G, Khan MAF, Beffagna G, Dazzo E, Sacchetto C, Polishchuk R, Sabatelli P, et al. A novel murine model for arrhythmogenic cardiomyopathy points to a pathogenic role of Wnt signalling and miRNA dysregulation. Cardiovasc Res. 2019;115:739-751. doi: 10.1093/cvr/cvv253
- 110. Notari M, Hu Y, Sutendra G, Dedeić Z, Lu M, Dupays L, Yavari A, Carr CA, Zhong S, Opel A, et al. iASPP, a previously unidentified regulator of desmosomes, prevents arrhythmogenic right ventricular cardiomyopathy (ARVC)induced sudden death. Proc Natl Acad Sci U S A. 2015;112:E973-E981. doi: 10.1073/pnas.1408111112
- 111. Ding Y, Yang J, Chen P, Lu T, Jiao K, Tester DJ, Giudicessi JR, Jiang K, Ackerman MJ, Li Y, et al. Knockout of SORBS2 protein disrupts the structural integrity of intercalated disc and manifests features of arrhythmogenic cardiomyopathy. J Am Heart Assoc. 2020;9:e017055. doi: 10.1161/JAHA.119.017055
- 112. Bround MJ, Asghari P, Wambolt RB, Bohunek L, Smits C, Philit M, Kieffer TJ, Lakatta EG, Boheler KR, Moore ED, et al. Cardiac ryanodine receptors control heart rate and rhythmicity in adult mice. Cardiovasc Res. 2012;96:372-380. doi: 10.1093/cvr/cvs260
- 113. Wang Y, Li C, Shi L, Chen X, Cui C, Huang J, Chen B, Hall DD, Pan Z, Lu M, et al. Integrin β1D deficiency-mediated RyR2 dysfunction contributes to catecholamine-sensitive ventricular tachycardia in arrhythmogenic right ventricular cardiomyopathy. Circulation. 2020;141:1477-1493. doi: 10.1161/CIRCULATIONAHA.119.043504
- 114. Zheng G, Jiang C, Li Y, Yang D, Ma Y, Zhang B, Li X, Zhang P, Hu X, Zhao X, et al. TMEM43-S358L mutation enhances NF-κB-TGFβ signal cascade in arrhythmogenic right ventricular dysplasia/cardiomyopathy. Protein Cell. 2019;10:104-119. doi: 10.1007/s13238-018-0563-2
- 115. Quang KL, Maguy A, Qi XY, Naud P, Xiong F, Tadevosyan A, Shi YF, Chartier D, Tardif JC, Dobrev D, Nattel S. Loss of cardiomyocyte integrin-linked kinase produces an arrhythmogenic cardiomyopathy in mice. Circ Arrhythm Electrophysiol. 2015;8:921-932. doi: 10.1161/CIRCEP.115.001668
- 116. Asano Y, Takashima S, Asakura M, Shintani Y, Liao Y, Minamino T, Asanuma H, Sanada S, Kim J, Ogai A, et al. Lamr1 functional retroposon causes right ventricular dysplasia in mice. Nat Genet. 2004;36:123-130. doi: 10.1038/ng1294
- 117. Basso C, Fox PR, Meurs KM, Towbin JA, Spier AW, Calabrese F, Maron BJ, Thiene G. Arrhythmogenic right ventricular cardiomyopathy causing sudden cardiac death in boxer dogs; a new animal model of human disease. Circulation. 2004;109:1180-1185. doi: 10.1161/01. CIR.0000118494.07530.65
- 118. Eason BD, Leach SB, Kuroki K. Arrhythmogenic right ventricular cardiomyopathy in a weimaraner. Can Vet J. 2015;56:1035-1039.
- 119. Cunningham SM, Sweeney JT, MacGregor J, Barton BA, Rush JE. Clinical features of english bulldogs with presumed arrhythmogenic right ventricular cardiomyopathy: 31 cases (2001-2013). J Am Anim Hosp Assoc. 2018;54:95-102. doi: 10.5326/JAAHA-MS-6550
- 120. Fox PR, Maron BJ, Basso C, Liu SK, Thiene G. Spontaneously occurring arrhythmogenic right ventricular cardiomyopathy in the domestic cat: a new animal model similar to the human disease. Circulation. 2000;102:1863-1870. doi: 10.1161/01.cir.102.15.1863
- 121. Wolf CM, Wang L, Alcalai R, Pizard A, Burgon PG, Ahmad F, Sherwood M, Branco DM, Wakimoto H, Fishman GI, et al. Lamin A/C haploinsufficiency causes dilated cardiomyopathy and apoptosis-triggered cardiac conduction system disease. J Mol Cell Cardiol. 2008;44:293-303. doi: 10.1016/j.yjmcc.2007.11.008
- 122. Arimura T, Helbling-Leclerc A, Massart C, Varnous S, Niel F, Lacène E, Fromes Y, Toussaint M, Mura AM, Keller DI, et al. Mouse model carrying H222P-Lmna mutation develops muscular dystrophy and dilated cardiomyopathy similar to human striated muscle laminopathies. Hum Mol Genet. 2005;14:155-169. doi: 10.1093/hmg/ddi017
- 123. Mounkes LC, Kozlov SV, Rottman JN, Stewart CL. Expression of an LMNA-N195K variant of A-type lamins results in cardiac conduction defects and death in mice. Hum Mol Genet. 2005;14:2167-2180. doi: 10.1093/hmg/ddi221

- 124. Osorio FG, Navarro CL, Cadiñanos J, López-Mejía IC, Quirós PM, Bartoli C, Rivera J, Tazi J, Guzmán G, Varela I, et al. Splicing-directed therapy in a new mouse model of human accelerated aging. Sci Transl Med. 2011;3:106ra107. doi: 10.1126/scitranslmed.3002847
- 125. Dorado B, Pløen GG, Barettino A, Macías A, Gonzalo P, Andrés-Manzano MJ, González-Gómez C, Galán-Arriola C, Alfonso JM, Lobo M, et al. Generation and characterization of a novel knockin minipig model of Hutchinson-Gilford progeria syndrome. Cell Discov. 2019;5:16. doi: 10.1038/s41421-019-0084-z
- 126. Ramspacher C, Steed E, Boselli F, Ferreira R, Faggianelli N, Roth S, Spiegelhalter C, Messaddeq N, Trinh L, Liebling M, et al. Developmental alterations in heart biomechanics and skeletal muscle function in desmin mutants suggest an early pathological root for desminopathies. Cell Rep. 2015;11:1564-1576. doi: 10.1016/j.celrep.2015.05.010
- 127. Clemen CS, Stöckigt F, Strucksberg KH, Chevessier F, Winter L, Schütz J, Bauer R, Thorweihe JM, Wenzel D, Schlötzer-Schrehardt U, et al. The toxic effect of R350P mutant desmin in striated muscle of man and mouse. Acta Neuropathol. 2015;129:297-315. doi: 10.1007/s00401-014-1363-2
- 128. Gupta V, Discenza M, Guyon JR, Kunkel LM, Beggs AH. α-Actinin-2 deficiency results in sarcomeric defects in zebrafish that cannot be rescued by α -actinin-3 revealing functional differences between sarcomeric isoforms. FASEB J. 2012;26:1892-1908. doi: 10.1096/fj.11-194548
- 129. Li Z, Ai T, Samani K, Xi Y, Tzeng HP, Xie M, Wu S, Ge S, Taylor MD, Dong JW, et al. A ZASP missense mutation, S196L, leads to cytoskeletal and electrical abnormalities in a mouse model of cardiomyopathy. Circ Arrhythm Electrophysiol. 2010;3:646-656. doi: 10.1161/CIRCEP.109.929240
- 130. Li J, Patel VV, Kostetskii I, Xiong Y, Chu AF, Jacobson JT, Yu C, Morley GE, Molkentin JD. Radice GL. Cardiac-specific loss of N-cadherin leads to alteration in connexins with conduction slowing and arrhythmogenesis. Circ Res. 2005;97:474-481. doi: 10.1161/01.RES.0000181132.11393.18
- 131. Li S, Mo K, Tian H, Chu C, Sun S, Tian L, Ding S, Li TR, Wu X, Liu F, et al. Lmod2 piggyBac mutant mice exhibit dilated cardiomyopathy. Cell Biosci. 2016;6:38. doi: 10.1186/s13578-016-0101-y
- 132. Guo W, Schafer S, Greaser ML, Radke MH, Liss M, Govindarajan T, Maatz H, Schulz H, Li S, Parrish AM, et al. RBM20, a gene for hereditary cardiomyopathy, regulates titin splicing. Nat Med. 2012;18:766-773. doi: 10.1038/nm.2693
- 133. van den Hoogenhof MMG, Beqqali A, Amin AS, van der Made I, Aufiero S, Khan MAF, Schumacher CA, Jansweijer JA, van Spaendonck-Zwarts KY, Remme CA, et al. RBM20 mutations induce an arrhythmogenic dilated cardiomyopathy related to disturbed calcium handling. Circulation. 2018;138:1330-1342. doi: 10.1161/CIRCULATIONAHA.117.031947
- 134. Ihara K, Sasano T, Hiraoka Y, Togo-Ohno M, Soejima Y, Sawabe M, Tsuchiya M, Ogawa H, Furukawa T, Kuroyanagi H. A missense mutation in the RSRSP stretch of Rbm20 causes dilated cardiomyopathy and atrial fibrillation in mice. Sci Rep. 2020;10:17894. doi: 10.1038/s41598-020-74800-8
- 135. Eijgenraam TR, Boukens BJ, Boogerd CJ, Schouten EM, van de Kolk CWA, Stege NM, Te Rijdt WP, Hoorntje ET, van der Zwaag PA, van Rooij E, et al. The phospholamban p.(Arg14del) pathogenic variant leads to cardiomyopathy with heart failure and is unreponsive to standard heart failure therapy. Sci Rep. 2020;10:9819. doi: 10.1038/s41598-020-66656-9
- 136. Hesse M, Kondo CS, Clark RB, Su L, Allen FL, Geary-Joo CT, Kunnathu S, Severson DL, Nygren A, Giles WR, Cross JC. Dilated cardiomyopathy is associated with reduced expression of the cardiac sodium channel Scn5a. Cardiovasc Res. 2007;75:498-509. doi: 10.1016/j.cardiores.2007.04.009
- 137. Glynn P, Musa H, Wu X, Unudurthi SD, Little S, Qian L, Wright PJ, Radwanski PB, Gyorke S, Mohler PJ, Hund TJ. Voltage-Gated Sodium Channel Phosphorylation at Ser571 Regulates Late Current, Arrhythmia, and Cardiac Function In Vivo. Circulation. 2015;132:567-577. doi: 10.1161/CIRCULATIONAHA.114.015218
- 138. Watanabe H, Yang T, Stroud DM, Lowe JS, Harris L, Atack TC, Wang DW, Hipkens SB, Leake B, Hall L, et al. Striking In vivo phenotype of a disease-associated human SCN5A mutation producing minimal changes in vitro. Circulation. 2011;124:1001-1011. doi: 10.1161/ CIRCULATIONAHA.110.987248
- 139. Huttner IG, Trivedi G, Jacoby A, Mann SA, Vandenberg JI, Fatkin D. A transgenic zebrafish model of a human cardiac sodium channel mutation exhibits bradycardia, conduction-system abnormalities and early death. J Mol Cell Cardiol. 2013;61:123-132. doi: 10.1016/j.yjmcc.2013.06.005
- 140. Moretti A, Fonteyne L, Giesert F, Hoppmann P, Meier AB, Bozoglu T, Baehr A, Schneider CM, Sinnecker D, Klett K, et al. Somatic gene editing ameliorates skeletal and cardiac muscle failure in pig and human models of duchenne muscular dystrophy. Nat Med. 2020;26:207-214. doi: 10.1038/s41591-019-0738-2

- 141. Moise NS, Valentine BA, Brown CA, Erb HN, Beck KA, Cooper BJ, Gilmour RF. Duchenne's cardiomyopathy in a canine model: electrocardiographic and echocardiographic studies. J Am Coll Cardiol. 1991;17:812–820. doi: 10.1016/s0735-1097(10)80202-5
- 142. Chu V, Otero JM, Lopez O, Sullivan MF, Morgan JP, Amende I, Hampton TG. Electrocardiographic findings in mdx mice: a cardiac phenotype of Duchenne muscular dystrophy. *Muscle Nerve*. 2002;26:513–519. doi: 10.1002/mus.10223
- 143. Branco DM, Wolf CM, Sherwood M, Hammer PE, Kang PB, Berul CI. Cardiac electrophysiological characteristics of the mdx (5cv) mouse model of Duchenne muscular dystrophy. *J Interv Card Electrophysiol*. 2007;20:1–7. doi: 10.1007/s10840-007-9168-z
- 144. Zemljic-Harpf AE, Miller JC, Henderson SA, Wright AT, Manso AM, Elsherif L, Dalton ND, Thor AK, Perkins GA, McCulloch AD, Ross RS. Cardiac-myocyte-specific excision of the vinculin gene disrupts cellular junctions, causing sudden death or dilated cardiomyopathy. Mol Cell Biol. 2007;27:7522–7537. doi: 10.1128/MCB.00728-07
- 145. Eigenthaler M, Engelhardt S, Schinke B, Kobsar A, Schmitteckert E, Gambaryan S, Engelhardt CM, Krenn V, Eliava M, Jarchau T, et al. Disruption of cardiac Ena-VASP protein localization in intercalated disks causes dilated cardiomyopathy. Am J Physiol Heart Circ Physiol. 2003;285:H2471–H2481. doi: 10.1152/ajpheart.00362.2003
- 146. Kuwahara K, Saito Y, Takano M, Arai Y, Yasuno S, Nakagawa Y, Takahashi N, Adachi Y, Takemura G, Horie M, et al. NRSF regulates the fetal cardiac gene program and maintains normal cardiac structure and function. *EMBO J.* 2003;22:6310–6321. doi: 10.1093/emboj/cdg601
- 147. Wess G, Schulze A, Butz V, Simak J, Killich M, Keller LJ, Maeurer J, Hartmann K. Prevalence of dilated cardiomyopathy in doberman pin-schers in various age groups. J Vet Intern Med. 2010;24:533–538. doi: 10.1111/j.1939-1676.2010.0479.x
- 148. Meurs KM, Miller MW, Wright NA. Clinical features of dilated cardiomyopathy in Great Danes and results of a pedigree analysis: 17 cases (1990-2000). J Am Vet Med Assoc. 2001;218:729–732. doi: 10.2460/iayma.2001.218.729
- 149. Knollmann BC, Kirchhof P, Sirenko SG, Degen H, Greene AE, Schober T, Mackow JC, Fabritz L, Potter JD, Morad M. Familial hypertrophic cardiomyopathy-linked mutant troponin T causes stress-induced ventricular tachycardia and Ca2+-dependent action potential remodeling. *Circ Res.* 2003;92:428–436. doi: 10.1161/01.RES.0000059562:91384.1A
- Du CK, Morimoto S, Nishii K, Minakami R, Ohta M, Tadano N, Lu QW, Wang YY, Zhan DY, Mochizuki M, et al. Knock-in mouse model of dilated cardiomyopathy caused by troponin mutation. *Circ Res.* 2007;101:185–194. doi: 10.1161/CIRCRESAHA.106.146670
- 151. Baudenbacher F, Schober T, Pinto JR, Sidorov VY, Hilliard F, Solaro RJ, Potter JD, Knollmann BC. Myofilament Ca2+ sensitization causes susceptibility to cardiac arrhythmia in mice. *J Clin Invest.* 2008;118:3893–3903. doi: 10.1172/JCl36642
- 152. Frey N, Franz WM, Gloeckner K, Degenhardt M, Müller M, Müller O, Merz H, Katus HA. Transgenic rat hearts expressing a human cardiac troponin T deletion reveal diastolic dysfunction and ventricular arrhythmias. *Cardiovasc Res.* 2000;47:254–264. doi: 10.1016/s0008-6363(00)00114-0
- 153. Tsoutsman T, Chung J, Doolan A, Nguyen L, Williams IA, Tu E, Lam L, Bailey CG, Rasko JE, Allen DG, Semsarian C. Molecular insights from a novel cardiac troponin I mouse model of familial hypertrophic cardiomyopathy. J Mol Cell Cardiol. 2006;41:623–632. doi: 10.1016/j.yjmcc.2006.07.016
- 154. Tsoutsman T, Kelly M, Ng DC, Tan JE, Tu E, Lam L, Bogoyevitch MA, Seidman CE, Seidman JG, Semsarian C. Severe heart failure and early mortality in a double-mutation mouse model of familial hypertrophic cardiomyopathy. *Circulation*. 2008;117:1820–1831. doi: 10.1161/CIRCULATIONAHA.107.755777
- 155. Berul CI, Christe ME, Aronovitz MJ, Maguire CT, Seidman CE, Seidman JG, Mendelsohn ME. Familial hypertrophic cardiomyopathy mice display gender differences in electrophysiological abnormalities. *J Interv Card Electrophysiol*. 1998;2:7–14. doi: 10.1023/a:1009700404218
- 156. Vikstrom KL, Factor SM, Leinwand LA. Mice expressing mutant myosin heavy chains are a model for familial hypertrophic cardiomyopathy. Mol Med. 1996;2:556-567.
- 157. McConnell BK, Jones KA, Fatkin D, Arroyo LH, Lee RT, Aristizabal O, Turnbull DH, Georgakopoulos D, Kass D, Bond M, et al. Dilated cardiomy-opathy in homozygous myosin-binding protein-C mutant mice. *J Clin Invest*. 1999;104:1235–1244. doi: 10.1172/JCI7377
- 158. Toib A, Zhang C, Borghetti G, Zhang X, Wallner M, Yang Y, Troupes CD, Kubo H, Sharp TE, Feldsott E, et al. Remodeling of repolarization and arrhythmia susceptibility in a myosin-binding protein C knockout mouse

- model. *Am J Physiol Heart Circ Physiol.* 2017;313:H620-H630. doi: 10.1152/ajpheart.00167.2017
- 159. Meurs KM, Sanchez X, David RM, Bowles NE, Towbin JA, Reiser PJ, Kittleson JA, Munro MJ, Dryburgh K, Macdonald KA, Kittleson MD. A cardiac myosin binding protein C mutation in the maine coon cat with familial hypertrophic cardiomyopathy. *Hum Mol Genet*. 2005;14:3587–3593. doi: 10.1093/hmg/ddi386
- Meurs KM, Norgard MM, Ederer MM, Hendrix KP, Kittleson MD. A substitution mutation in the myosin binding protein C gene in ragdoll hypertrophic cardiomyopathy. *Genomics*. 2007;90:261–264. doi: 10.1016/j. ygeno.2007.04.007
- Ruan H, Mitchell S, Vainoriene M, Lou Q, Xie LH, Ren S, Goldhaber JI, Wang Y. Gi alpha 1-mediated cardiac electrophysiological remodeling and arrhythmia in hypertrophic cardiomyopathy. *Circulation*. 2007;116:596– 605. doi: 10.1161/CIRCULATIONAHA.106.682773
- 162. Alvarado FJ, Bos JM, Yuchi Z, Valdivia CR, Hernández JJ, Zhao YT, Henderlong DS, Chen Y, Booher TR, Marcou CA, et al. Cardiac hypertrophy and arrhythmia in mice induced by a mutation in ryanodine receptor 2. JCI Insight. 2019;5:126544. doi: 10.1172/jci.insight.126544
- 163. Hu LR, Ackermann MA, Hecker PA, Prosser BL, King B, O'Connell KA, Grogan A, Meyer LC, Berndsen CE, Wright NT, et al. Deregulated Ca2+ cycling underlies the development of arrhythmia and heart disease due to mutant obscurin. Sci Adv. 2017;3:e1603081. doi: 10.1126/sciadv.1603081
- 164. Sood S, Chelu MG, van Oort RJ, Skapura D, Santonastasi M, Dobrev D, Wehrens XH. Intracellular calcium leak due to FKBP12.6 deficiency in mice facilitates the inducibility of atrial fibrillation. *Heart Rhythm.* 2008;5:1047–1054. doi: 10.1016/j.hrthm.2008.03.030
- 165. Faggioni M, Savio-Galimberti E, Venkataraman R, Hwang HS, Kannankeril PJ, Darbar D, Knollmann BC. Suppression of spontaneous ca elevations prevents atrial fibrillation in calsequestrin 2-null hearts. Circ Arrhythm Electrophysiol. 2014;7:313–320. doi: 10.1161/CIRCEP.113.000994
- 166. Beavers DL, Wang W, Ather S, Voigt N, Garbino A, Dixit SS, Landstrom AP, Li N, Wang Q, Olivotto I, et al. Mutation E169K in junctophilin-2 causes atrial fibrillation due to impaired RyR2 stabilization. J Am Coll Cardiol. 2013;62:2010–2019. doi: 10.1016/j.jacc.2013.06.052
- 167. Müller FU, Lewin G, Baba HA, Bokník P, Fabritz L, Kirchhefer U, Kirchhof P, Loser K, Matus M, Neumann J, et al. Heart-directed expression of a human cardiac isoform of cAMP-response element modulator in transgenic mice. J Biol Chem. 2005;280:6906–6914. doi: 10.1074/jbc.M407864200
- 168. Yao C, Veleva T, Scott L Jr, Cao S, Li L, Chen G, Jeyabal P, Pan X, Alsina KM, Abu-Taha I Dr, et al. Enhanced cardiomyocyte NLRP3 inflammasome signaling promotes atrial fibrillation. *Circulation*. 2018;138:2227–2242. doi: 10.1161/CIRCULATIONAHA.118.035202
- 169. Xie LH, Shanmugam M, Park JY, Zhao Z, Wen H, Tian B, Periasamy M, Babu GJ. Ablation of sarcolipin results in atrial remodeling. Am J Physiol Cell Physiol. 2012;302:C1762-C1771. doi: 10.1152/ajpcell.00425.2011
- Wan E, Abrams J, Weinberg RL, Katchman AN, Bayne J, Zakharov SI, Yang L, Morrow JP, Garan H, Marx SO. Aberrant sodium influx causes cardiomyopathy and atrial fibrillation in mice. *J Clin Invest*. 2016;126:112–122. doi: 10.1172/JCI84669
- 171. Blana A, Kaese S, Fortmüller L, Laakmann S, Damke D, van Bragt K, Eckstein J, Piccini I, Kirchhefer U, Nattel S, et al. Knock-in gain-of-function sodium channel mutation prolongs atrial action potentials and alters atrial vulnerability. *Heart Rhythm.* 2010;7:1862–1869. doi: 10.1016/j.hrthm.2010.08.016
- 172. Temple J, Frias P, Rottman J, Yang T, Wu Y, Verheijck EE, Zhang W, Siprachanh C, Kanki H, Atkinson JB, et al. Atrial fibrillation in KCNE1-null mice. Circ Res. 2005;97:62-69. doi: 10.1161/01.RES.0000173047.42236.88
- 173. Glasscock E, Voigt N, McCauley MD, Sun Q, Li N, Chiang DY, Zhou XB, Molina CE, Thomas D, Schmidt C, et al. Expression and function of Kv1.1 potassium channels in human atria from patients with atrial fibrillation. Basic Res Cardiol. 2015;110:505. doi: 10.1007/s00395-015-0505-6
- 174. David JP, Lisewski U, Crump SM, Jepps TA, Bocksteins E, Wilck N, Lossie J, Roepke TK, Schmitt N, Abbott GW. Deletion in mice of X-linked, brugada syndrome- and atrial fibrillation-associated Kcne5 augments ventricular KV currents and predisposes to ventricular arrhythmia. FASEB J. 2019;33:2537–2552. doi: 10.1096/fj.201800502R
- 175. Li N, Timofeyev V, Tuteja D, Xu D, Lu L, Zhang Q, Zhang Z, Singapuri A, Albert TR, Rajagopal AV, et al. Ablation of a Ca2+-activated K+ channel (SK2 channel) results in action potential prolongation in atrial myocytes and atrial fibrillation. *J Physiol.* 2009;587(pt 5):1087-1100. doi: 10.1113/jphysiol.2008.167718
- 176. Kirchhof P, Kahr PC, Kaese S, Piccini I, Vokshi I, Scheld HH, Rotering H, Fortmueller L, Laakmann S, Verheule S, et al. PITX2c is expressed in the

- adult left atrium, and reducing Pitx2c expression promotes atrial fibrillation inducibility and complex changes in gene expression. Circ Cardiovasc Genet. 2011;4:123-133. doi: 10.1161/CIRCGENETICS.110.958058
- 177. Nadadur RD, Broman MT, Boukens B, Mazurek SR, Yang X, van den Boogaard M, Bekeny J, Gadek M, Ward T, Zhang M, et al. Pitx2 modulates a Tbx5-dependent gene regulatory network to maintain atrial rhythm, Sci Transl Med. 2016;8:354ra115. doi: 10.1126/scitranslmed.aaf4891
- 178. Tuomi JM. Tvml K. Jones DL. Atrial tachycardia/fibrillation in the connexin 43 G60S mutant (Oculodentodigital dysplasia) mouse. Am J Physiol Heart Circ Physiol. 2011;300:H1402-H1411. doi: 10.1152/ajpheart.01094.2010
- 179. Ozcan C, Battaglia E, Young R, Suzuki G. LKB1 knockout mouse develops spontaneous atrial fibrillation and provides mechanistic insights into human disease process. J Am Heart Assoc. 2015;4:e001733. doi: 10.1161/JAHA.114.001733
- 180. Moreira LM, Takawale A, Hulsurkar M, Menassa DA, Antanaviciute A, Lahiri SK, Mehta N, Evans N, Psarros C, Robinson P, et al. Paracrine signalling by cardiac calcitonin controls atrial fibrogenesis and arrhythmia. Nature. 2020;587:460-465. doi: 10.1038/s41586-020-2890-8
- 181. Polejaeva IA, Ranjan R, Davies CJ, Regouski M, Hall J, Olsen AL, Meng Q, Rutigliano HM, Dosdall DJ, Angel NA, et al. Increased susceptibility to atrial fibrillation secondary to atrial fibrosis in transgenic goats expressing transforming growth factor-β1. J Cardiovasc Electrophysiol. 2016;27:1220-1229. doi: 10.1111/jce.13049
- 182. Verheule S, Sato T, Everett T 4th, Engle SK, Otten D, Rubart-von der Lohe M, Nakajima HO, Nakajima H, Field LJ, Olgin JE. Increased vulnerability to atrial fibrillation in transgenic mice with selective atrial fibrosis caused by overexpression of TGF-beta1. Circ Res. 2004;94:1458-1465. doi: 10.1161/01.RES.0000129579.59664.9d
- 183. Davies L, Jin J, Shen W, Tsui H, Shi Y, Wang Y, Zhang Y, Hao G, Wu J, Chen S, et al. Mkk4 is a negative regulator of the transforming growth factor beta 1 signaling associated with atrial remodeling and arrhythmogenesis with age. J Am Heart Assoc. 2014;3:e000340. doi: 10.1161/JAHA.113.000340
- 184. Xiao HD, Fuchs S, Campbell DJ, Lewis W, Dudley SC Jr, Kasi VS, Hoit BD, Keshelava G, Zhao H, Capecchi MR, Bernstein KE. Mice with cardiacrestricted angiotensin-converting enzyme (ACE) have atrial enlargement, cardiac arrhythmia, and sudden death. Am J Pathol. 2004;165:1019-1032, doi: 10.1016/S0002-9440(10)63363-9
- 185. Parahuleva MS, Kockskämper J, Heger J, Grimm W, Scherer A, Bühler S, Kreutz J, Schulz R, Euler G. Structural, Pro-Inflammatory and Calcium Handling Remodeling Underlies Spontaneous Onset of Paroxysmal Atrial Fibrillation in JDP2-Overexpressing Mice. Int J Mol Sci. 2020;21:E9095. doi: 10.3390/ijms21239095
- 186. Lei M, Goddard C, Liu J, Léoni AL, Royer A, Fung SS, Xiao G, Ma A, Zhang H, Charpentier F, et al. Sinus node dysfunction following targeted disruption of the murine cardiac sodium channel gene Scn5a. J Physiol. 2005;567(pt 2):387-400. doi: 10.1113/jphysiol.2005.083188
- 187. Hakim P, Brice N, Thresher R, Lawrence J, Zhang Y, Jackson AP, Grace AA, Huang CL, Scn3b knockout mice exhibit abnormal sino-atrial and cardiac conduction properties. Acta Physiol (Oxf). 2010;198:47-59. doi: 10.1111/j.1748-1716.2009.02048.x
- 188. Qiao Y, Lipovsky C, Hicks S, Bhatnagar S, Li G, Khandekar A, Guzy R, Woo KV, Nichols CG, Efimov IR, Rentschler S. Transient notch activation induces long-term gene expression changes leading to sick sinus syndrome in mice. Circ Res. 2017;121:549-563. doi: 10.1161/ CIRCRESAHA.116.310396
- 189. Baker K, Warren KS, Yellen G, Fishman MC. Defective "pacemaker" current (lh) in a zebrafish mutant with a slow heart rate, Proc Natl Acad Sci U S A. 1997;94:4554-4559. doi: 10.1073/pnas.94.9.4554
- 190. Groenke S, Larson ED, Alber S, Zhang R, Lamp ST, Ren X, Nakano H, Jordan MC, Karagueuzian HS, Roos KP, et al. Complete atrial-specific knockout of sodium-calcium exchange eliminates sinoatrial node pacemaker activity. PLoS One. 2013;8:e81633. doi: 10.1371/journal.pone.0081633
- 191. Ebert AM, Hume GL, Warren KS, Cook NP, Burns CG, Mohideen MA, Siegal G. Yelon D. Fishman MC. Garrity DM. Calcium extrusion is critical for cardiac morphogenesis and rhythm in embryonic zebrafish hearts. Proc Natl Acad Sci U S A. 2005;102:17705-17710. doi: 10.1073/pnas.0502683102
- 192. Fenske S, Krause SC, Hassan SI, Becirovic E, Auer F, Bernard R, Kupatt C, Lange P, Ziegler T, Wotjak CT, et al. Sick sinus syndrome in HCN1deficient mice. Circulation. 2013;128:2585-2594. doi: 10.1161/ CIRCULATIONAHA.113.003712
- 193. Ludwig A, Budde T, Stieber J, Moosmang S, Wahl C, Holthoff K, Langebartels A, Wotjak C, Munsch T, Zong X, et al. Absence epilepsy and sinus dysrhythmia in mice lacking the pacemaker channel HCN2. EMBO J. 2003;22:216-224. doi: 10.1093/emboj/cdg032

- 194. Stieber J, Herrmann S, Feil S, Löster J, Feil R, Biel M, Hofmann F, Ludwig A. The hyperpolarization-activated channel HCN4 is required for the generation of pacemaker action potentials in the embryonic heart. Proc Natl Acad Sci U S A. 2003;100:15235-15240. doi: 10.1073/pnas.2434235100
- 195. Harzheim D, Pfeiffer KH, Fabritz L, Kremmer E, Buch T, Waisman A, Kirchhof P. Kaupp UB. Seifert R. Cardiac pacemaker function of HCN4 channels in mice is confined to embryonic development and requires cyclic AMP. EMBO J. 2008;27:692-703. doi: 10.1038/emboj.2008.3
- 196. Alig J, Marger L, Mesirca P, Ehmke H, Mangoni ME, Isbrandt D. Control of heart rate by cAMP sensitivity of HCN channels. Proc Natl Acad Sci U S A. 2009;106:12189-12194. doi: 10.1073/pnas.0810332106
- 197. Baruscotti M, Bucchi A, Viscomi C, Mandelli G, Consalez G, Gnecchi-Rusconi T, Montano N, Casali KR, Micheloni S, Barbuti A, DiFrancesco D. Deep bradycardia and heart block caused by inducible cardiac-specific knockout of the pacemaker channel gene Hcn4. Proc Natl Acad Sci USA. 2011;108:1705-1710. doi: 10.1073/pnas.1010122108
- 198. Herrmann S, Fabritz L, Layh B, Kirchhof P, Ludwig A. Insights into sick sinus syndrome from an inducible mouse model. Cardiovasc Res. 2011;90:38-48. doi: 10.1093/cvr/cva390
- 199. Kasahara H, Wakimoto H, Liu M, Maguire CT, Converso KL, Shioi T, Huang WY, Manning WJ, Paul D, Lawitts J, et al. Progressive atrioventricular conduction defects and heart failure in mice expressing a mutant Csx/Nkx2.5 homeoprotein. J Clin Invest. 2001;108:189-201. doi: 10.1172/JCI12694
- 200. Chowdhury R, Ashraf H, Melanson M, Tanada Y, Nguyen M, Silberbach M, Wakimoto H, Benson DW, Anderson RH, Kasahara H. Mouse model of human congenital heart disease: progressive atrioventricular block induced by a heterozygous Nkx2-5 homeodomain missense mutation. Circ Arrhythm Electrophysiol. 2015;8:1255-1264. doi: 10.1161/CIRCEP.115.002720
- 201. Berul Cl, Maguire CT, Gehrmann J, Reddy S. Progressive atrioventricular conduction block in a mouse myotonic dystrophy model. J Interv Card Electrophysiol. 2000;4:351-358. doi: 10.1023/a:1009842114968
- 202. Simon AM, Goodenough DA, Paul DL. Mice lacking connexin40 have cardiac conduction abnormalities characteristic of atrioventricular block and bundle branch block. Curr Biol. 1998;8:295-298. doi: 10.1016/s0960-9822(98)70113-7
- 203. Demion M, Thireau J, Gueffier M, Finan A, Khoueiry Z, Cassan C, Serafini N, Aimond F, Granier M, Pasquié JL, et al. Trpm4 gene invalidation leads to cardiac hypertrophy and electrophysiological alterations. PLoS One. 2014;9:e115256. doi: 10.1371/journal.pone.0115256
- 204. Sidhu JS, Rajawat YS, Rami TG, Gollob MH, Wang Z, Yuan R, Marian AJ, DeMayo FJ, Weilbacher D, Taffet GE, et al. Transgenic mouse model of ventricular preexcitation and atrioventricular reentrant tachycardia induced by an AMP-activated protein kinase loss-of-function mutation responsible for Wolff-Parkinson-White syndrome. Circulation. 2005;111:21-29. doi: 10.1161/01.CIR.0000151291.32974.D5
- 205. Arad M, Moskowitz IP, Patel VV, Ahmad F, Perez-Atayde AR, Sawyer DB, Walter M, Li GH, Burgon PG, Maguire CT, et al. Transgenic mice overexpressing mutant PRKAG2 define the cause of Wolff-Parkinson-White syndrome in glycogen storage cardiomyopathy. Circulation. 2003;107:2850-2856. doi: 10.1161/01.CIR.0000075270.13497.2B
- 206. Davies JK, Wells DJ, Liu K, Whitrow HR, Daniel TD, Grignani R, Lygate CA, Schneider JE, Noël G, Watkins H, Carling D. Characterization of the role of gamma2 R531G mutation in AMP-activated protein kinase in cardiac hypertrophy and Wolff-Parkinson-White syndrome. Am J Physiol Heart Circ Physiol. 2006;290:H1942-H1951. doi: 10.1152/ ajpheart.01020.2005
- 207. Aanhaanen WT, Boukens BJ, Sizarov A, Wakker V, de Gier-de Vries C, van Ginneken AC, Moorman AF, Coronel R, Christoffels VM. Defective Tbx2dependent patterning of the atrioventricular canal myocardium causes accessory pathway formation in mice. J Clin Invest. 2011;121:534-544. doi: 10.1172/JCI44350
- 208. Lübkemeier I, Requardt RP, Lin X, Sasse P, Andrié R, Schrickel JW, Chkourko H, Bukauskas FF, Kim JS, Frank M, et al. Deletion of the last five C-terminal amino acid residues of connexin43 leads to lethal ventricular arrhythmias in mice without affecting coupling via gap junction channels. Basic Res Cardiol. 2013;108:348. doi: 10.1007/s00395-013-0348-y
- 209. Zi M, Kimura TE, Liu W, Jin J, Higham J, Kharche S, Hao G, Shi Y, Shen W, Prehar S, et al. Mitogen-activated protein kinase kinase 4 deficiency in cardiomyocytes causes connexin 43 reduction and couples hypertrophic signals to ventricular arrhythmogenesis. J Biol Chem. 2011;286:17821-17830. doi: 10.1074/jbc.M111.228791
- 210. Yamada N, Asano Y, Fujita M, Yamazaki S, Inanobe A, Matsuura N, Kobayashi H, Ohno S, Ebana Y, Tsukamoto O, et al. Mutant KCNJ3 and KCNJ5 potassium channels as novel molecular targets in bradyarrhythmias

- and atrial fibrillation. Circulation. 2019;139:2157-2169. doi: 10.1161/CIRCULATIONAHA.118.036761
- 211. Baudot M, Torre E, Bidaud I, Louradour J, Torrente AG, Fossier L, Talssi L, Nargeot J, Barrère-Lemaire S, Mesirca P, Mangoni ME. Concomitant genetic ablation of L-type Cav1.3 (α1D) and T-type Cav3.1 (α1G) Ca2+ channels disrupts heart automaticity. Sci Rep. 2020;10:18906. doi: 10.1038/s41598-020-76049-7
- 212. Mahida S, Mills RW, Tucker NR, Simonson B, Macri V, Lemoine MD, Das S, Milan DJ, Ellinor PT. Overexpression of KCNN3 results in sudden cardiac death. *Cardiovasc Res.* 2014;101:326–334. doi: 10.1093/cvr/cvt269
- 213. Zhang SS, Kim KH, Rosen A, Smyth JW, Sakuma R, Delgado-Olguín P, Davis M, Chi NC, Puviindran V, Gaborit N, et al. Iroquois homeobox gene 3 establishes fast conduction in the cardiac his-purkinje network. Proc Natl Acad Sci U S A. 2011;108:13576–13581. doi: 10.1073/pnas.1106911108
- 214. Hoogaars WM, Engel A, Brons JF, Verkerk AO, de Lange FJ, Wong LY, Bakker ML, Clout DE, Wakker V, Barnett P, et al. Tbx3 controls the sinoatrial node gene program and imposes pacemaker function on the atria. *Genes Dev.* 2007;21:1098–1112. doi: 10.1101/gad.416007
- 215. Bruneau BG, Nemer G, Schmitt JP, Charron F, Robitaille L, Caron S, Conner DA, Gessler M, Nemer M, Seidman CE, Seidman JG. A murine model of Holt-Oram syndrome defines roles of the T-box transcription factor Tbx5 in cardiogenesis and disease. *Cell.* 2001;106:709–721. doi: 10.1016/s0092-8674(01)00493-7
- 216. McCauley MD, Wang T, Mike E, Herrera J, Beavers DL, Huang TW, Ward CS, Skinner S, Percy AK, Glaze DG, et al. Pathogenesis of lethal cardiac arrhythmias in Mecp2 mutant mice: implication for therapy in rett syndrome. Sci Transl Med. 2011;3:113ra125. doi: 10.1126/scitranslmed.3002982
- 217. Karakikes I, Ameen M, Termglinchan V, Wu JC. Human induced pluripotent stem cell-derived cardiomyocytes: insights into molecular, cellular, and functional phenotypes. *Circ Res.* 2015;117:80–88. doi: 10.1161/CIRCRESAHA.117.305365
- 218. Watanabe H, Chopra N, Laver D, Hwang HS, Davies SS, Roach DE, Duff HJ, Roden DM, Wilde AA, Knollmann BC. Flecainide prevents catecholaminergic polymorphic ventricular tachycardia in mice and humans. *Nat Med*. 2009;15:380–383. doi: 10.1038/nm.1942
- 219. Wakula P, Bisping E, Kockskämper J, Post H, Brauer S, Deuter M, Oehlmann R, Besenfelder U, Lai FA, Brem G, Pieske B. CMV promoter is inadequate for expression of mutant human RyR2 in transgenic rabbits. *J Pharmacol Toxicol Methods*. 2011;63:180–185. doi: 10.1016/j.vascn.2010.09.007
- 220. Ng K, Titus EW, Lieve KV, Roston TM, Mazzanti A, Deiter FH, Denjoy I, Ingles J, Till J, Robyns T, et al. An international multicenter evaluation of inheritance patterns, arrhythmic risks, and underlying mechanisms of CASQ2-catecholaminergic polymorphic ventricular tachycardia. Circulation. 2020;142:932–947. doi: 10.1161/CIRCULATIONAHA.120.045723
- Wleklinski MJ, Kannankeril PJ, Knollmann BC. Molecular and tissue mechanisms of catecholaminergic polymorphic ventricular tachycardia. *J Physiol.* 2020;598:2817–2834. doi: 10.1113/JP276757
- 222. Nyegaard M, Overgaard MT, Søndergaard MT, Vranas M, Behr ER, Hildebrandt LL, Lund J, Hedley PL, Camm AJ, Wettrell G, et al. Mutations in calmodulin cause ventricular tachycardia and sudden cardiac death. *Am J Hum Genet*. 2012;91:703–712. doi: 10.1016/j.ajhg.2012.08.015
- 223. Kurtzwald-Josefson E, Yadin D, Harun-Khun S, Waldman M, Aravot D, Shainberg A, Eldar M, Hochhauser E, Arad M. Viral delivered gene therapy to treat catecholaminergic polymorphic ventricular tachycardia (CPVT2) in mouse models. *Heart Rhythm.* 2017;14:1053–1060. doi: 10.1016/j.hrthm.2017.03.025
- 224. Kryshtal DO, Blackwell DJ, Egly CL, Smith AN, Batiste SM, Johnston JN, Laver DR, Knollmann BC. RYR2 channel inhibition is the principal mechanism of flecainide action in CPVT. Circ Res. 2021;128:321–331. doi: 10.1161/CIRCRESAHA.120.316819
- 225. Al-Khatib SM, Stevenson WG, Ackerman MJ, Bryant WJ, Callans DJ, Curtis AB, Deal BJ, Dickfeld T, Field ME, Fonarow GC, et al. 2017 AHA/ACC/HRS guideline for management of patients with ventricular arrhythmias and the prevention of sudden cardiac death: a report of the american college of cardiology/american heart association task force on clinical practice guidelines and the heart rhythm society. Circulation. 2018;138:e272–e391. doi: 10.1161/CIR.00000000000000549
- 226. Priori SG, Wilde AA, Horie M, Cho Y, Behr ER, Berul C, Blom N, Brugada J, Chiang CE, Huikuri H, et al. HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHRS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. Heart Rhythm. 2013;10:1932–1963. doi: 10.1016/j.hrthm.2013.05.014

- 227. Shimizu W, Horie M. Phenotypic manifestations of mutations in genes encoding subunits of cardiac potassium channels. Circ Res. 2011;109:97–109. doi: 10.1161/CIRCRESAHA.110.224600
- 228. Migdalovich D, Moss AJ, Lopes CM, Costa J, Ouellet G, Barsheshet A, McNitt S, Polonsky S, Robinson JL, Zareba W, et al. Mutation and gender-specific risk in type 2 long QT syndrome: implications for risk stratification for life-threatening cardiac events in patients with long QT syndrome. *Heart Rhythm.* 2011;8:1537–1543. doi: 10.1016/j. hrthm.2011.03.049
- 229. Locati EH, Zareba W, Moss AJ, Schwartz PJ, Vincent GM, Lehmann MH, Towbin JA, Priori SG, Napolitano C, Robinson JL, et al. Age- and sex-related differences in clinical manifestations in patients with congenital long-QT syndrome: findings from the International LQTS Registry. Circulation. 1998;97:2237–2244. doi: 10.1161/01.cir.97.22.2237
- Salama G, Bett GC. Sex differences in the mechanisms underlying long QT syndrome. Am J Physiol Heart Circ Physiol. 2014;307:H640-H648. doi: 10.1152/ajpheart.00864.2013
- 231. Odening KE, Koren G. How do sex hormones modify arrhythmogenesis in long QT syndrome? Sex hormone effects on arrhythmogenic substrate and triggered activity. *Heart Rhythm.* 2014;11:2107–2115. doi: 10.1016/j.hrthm.2014.06.023
- Drici MD, Burklow TR, Haridasse V, Glazer RI, Woosley RL. Sex hormones prolong the QT interval and downregulate potassium channel expression in the rabbit heart. *Circulation*. 1996;94:1471–1474. doi: 10.1161/01.cir.94.6.1471
- Drici MD, Knollmann BC, Wang WX, Woosley RL. Cardiac actions of erythromycin: influence of female sex. *JAMA*. 1998;280:1774–1776. doi: 10.1001/jama.280.20.1774
- 234. Wada Y, Yang T, Shaffer CM, Daniel LL, Glazer AM, Davogustto GE, Lowery BD, Farber-Eger EH, Wells QS, Roden DM. Common ancestry-specific ion channel variants predispose to drug-induced arrhythmias. Circulation. 2022;145:299–308. doi: 10.1161/CIRCULATIONAHA.121.054883
- 235. Leoni AL, Gavillet B, Rougier JS, Marionneau C, Probst V, Le Scouarnec S, Schott JJ, Demolombe S, Bruneval P, Huang CL, et al. Variable Na(v)1.5 protein expression from the wild-type allele correlates with the penetrance of cardiac conduction disease in the Scn5a(+/-) mouse model. PLoS One. 2010;5:e9298. doi: 10.1371/journal.pone.0009298
- 236. Wagner S, Dybkova N, Rasenack EC, Jacobshagen C, Fabritz L, Kirchhof P, Maier SK, Zhang T, Hasenfuss G, Brown JH, et al. Ca2+/calmodulin-dependent protein kinase II regulates cardiac Na+ channels. *J Clin Invest*. 2006;116:3127–3138. doi: 10.1172/JCl26620
- 237. Park DS, Cerrone M, Morley G, Vasquez C, Fowler S, Liu N, Bernstein SA, Liu FY, Zhang J, Rogers CS, et al. Genetically engineered SCN5A mutant pig hearts exhibit conduction defects and arrhythmias. *J Clin Invest*. 2015;125:403–412. doi: 10.1172/JCl76919
- 238. van Opbergen CJM, Bagwan N, Maurya SR, Kim JC, Smith AN, Blackwell DJ, Johnston JN, Knollmann BC, Cerrone M, Lundby A, et al. Exercise causes arrhythmogenic remodeling of intracellular calcium dynamics in Plakophilin-2-Deficient Hearts. *Circulation*. 2022;145:1480-1496. doi: 10.1161/CIRCULATIONAHA.121.057757
- 239. Laitinen-Forsblom PJ, Mäkynen P, Mäkynen H, Yli-Mäyry S, Virtanen V, Kontula K, Aalto-Setälä K. SCN5A mutation associated with cardiac conduction defect and atrial arrhythmias. *J Cardiovasc Electrophysiol*. 2006;17:480–485. doi: 10.1111/j.1540-8167.2006.00411.x
- 240. Groenewegen WA, Firouzi M, Bezzina CR, Vliex S, van Langen IM, Sandkuijl L, Smits JP, Hulsbeek M, Rook MB, Jongsma HJ, Wilde AA. A cardiac sodium channel mutation cosegregates with a rare connexin40 genotype in familial atrial standstill. Circ Res. 2003;92:14–22. doi: 10.1161/01.res.0000050585.07097.d7
- 241. Marian AJ, Braunwald E. Hypertrophic Cardiomyopathy: Genetics, Pathogenesis, Clinical Manifestations, Diagnosis, and Therapy. Circ Res. 2017;121:749–770. doi: 10.1161/CIRCRESAHA.117.311059
- 242. Marian AJ, Wu Y, Lim DS, McCluggage M, Youker K, Yu QT, Brugada R, DeMayo F, Quinones M, Roberts R. A transgenic rabbit model for human hypertrophic cardiomyopathy. J Clin Invest. 1999;104:1683–1692. doi: 10.1172/JCl7956
- 243. Schober T, Huke S, Venkataraman R, Gryshchenko O, Kryshtal D, Hwang HS, Baudenbacher FJ, Knollmann BC. Myofilament Ca sensitization increases cytosolic Ca binding affinity, alters intracellular Ca homeostasis, and causes pause-dependent Ca-triggered arrhythmia. Circ Res. 2012;111:170–179. doi: 10.1161/CIRCRESAHA.112.270041
- 244. Huke S, Venkataraman R, Faggioni M, Bennuri S, Hwang HS, Baudenbacher F, Knollmann BC. Focal energy deprivation underlies arrhythmia susceptibility in mice with calcium-sensitized myofilaments. Circ Res. 2013;112:1334–1344. doi: 10.1161/CIRCRESAHA.113.301055

- 245. Venkataraman R, Baldo MP, Hwang HS, Veltri T, Pinto JR, Baudenbacher FJ, Knollmann BC. Myofilament calcium de-sensitization and contractile uncoupling prevent pause-triggered ventricular tachycardia in mouse hearts with chronic myocardial infarction. *J Mol Cell Cardiol*. 2013;60:8–15. doi: 10.1016/j.yjmcc.2013.03.022
- 246. Huke S, Knollmann BC. Increased myofilament Ca2+-sensitivity and arrhythmia susceptibility. *J Mol Cell Cardiol.* 2010;48:824-833. doi: 10.1016/j.yjmcc.2010.01.011
- 247. Schüttler D, Bapat A, Kääb S, Lee K, Tomsits P, Clauss S, Hucker WJ. Animal models of atrial fibrillation. Circ Res. 2020;127:91–110. doi: 10.1161/CIRCRESAHA.120.316366
- 248. Roselli C, Rienstra M, Ellinor PT. Genetics of Atrial Fibrillation in 2020: GWAS, Genome Sequencing, Polygenic Risk, and Beyond. Circ Res. 2020;127:21–33. doi: 10.1161/CIRCRESAHA.120.316575
- 249. Syeda F, Holmes AP, Yu TY, Tull S, Kuhlmann SM, Pavlovic D, Betney D, Riley G, Kucera JP, Jousset F, et al. PITX2 modulates atrial membrane potential and the antiarrhythmic effects of sodium-channel blockers. *J Am Coll Cardiol*. 2016;68:1881–1894. doi: 10.1016/j.jacc.2016.07.766
- 250. Jeevaratnam K, Zhang Y, Guzadhur L, Duehmke RM, Lei M, Grace AA, Huang CL. Differences in sino-atrial and atrio-ventricular function with age and sex attributable to the Scn5a+/- mutation in a murine cardiac model. *Acta Physiol (Oxf)*. 2010;200:23–33. doi: 10.1111/j. 1748-1716.2010.02110.x
- 251. Fenske S, Mader R, Scharr A, Paparizos C, Cao-Ehlker X, Michalakis S, Shaltiel L, Weidinger M, Stieber J, Feil S, et al. HCN3 contributes to the ventricular action potential waveform in the murine heart. *Circ Res.* 2011;109:1015–1023. doi: 10.1161/CIRCRESAHA.111.246173
- 252. Rockman HA, Ross RS, Harris AN, Knowlton KU, Steinhelper ME, Field LJ, Ross J Jr, Chien KR. Segregation of atrial-specific and inducible expression of an atrial natriuretic factor transgene in an in vivo murine model of cardiac hypertrophy. *Proc Natl Acad Sci U S A.* 1991;88:8277–8281. doi: 10.1073/pnas.88.18.8277
- 253. deAlmeida AC, van Oort RJ, Wehrens XH. Transverse aortic constriction in mice. *J Vis Exp.* 2010. doi: 10.3791/1729
- 254. Richards DA, Aronovitz MJ, Calamaras TD, Tam K, Martin GL, Liu P, Bowditch HK, Zhang P, Huggins GS, Blanton RM. Distinct phenotypes induced by three degrees of transverse aortic constriction in mice. *Sci Rep.* 2019;9:5844. doi: 10.1038/s41598-019-42209-7
- 255. Del Monte F, Butler K, Boecker W, Gwathmey JK, Hajjar RJ. Novel technique of aortic banding followed by gene transfer during hypertrophy and heart failure. *Physiol Genomics*. 2002;9:49–56. doi: 10.1152/ physiolgenomics.00035.2001
- 256. McAinsh AM, Turner MA, O'Hare D, Nithythyananthan R, Johnston DG, O'Gorman DJ, Sheridan DJ. Cardiac hypertrophy impairs recovery from ischaemia because there is a reduced reactive hyperaemic response. Cardiovasc Res. 1995;30:113–121.
- 257. Gilson N, el Houda Bouanani N, Corsin A, Crozatier B. Left ventricular function and beta-adrenoceptors in rabbit failing heart. Am J Physiol. 1990;258(3 Pt 2):H634-H641. doi: 10.1152/ajpheart.1990.258.3.H634
- 258. Gyöngyösi M, Pavo N, Lukovic D, Zlabinger K, Spannbauer A, Traxler D, Goliasch G, Mandic L, Bergler-Klein J, Gugerell A, et al. Porcine model of progressive cardiac hypertrophy and fibrosis with secondary post-capillary pulmonary hypertension. *J Transl Med.* 2017;15:202. doi: 10.1186/s12967-017-1299-0
- 259. Walther T, Falk V, Binner C, Löscher N, Schubert A, Rauch T, von Oppel U, Autschbach R, Mohr FW. Experimental aortic stenosis and corresponding left ventricular hypertrophy in sheep. *J Invest Surg.* 2000;13:327–331. doi: 10.1080/089419300750059370
- Sakamoto J, Miura T, Tsuchida A, Fukuma T, Hasegawa T, Shimamoto K. Reperfusion arrhythmias in the murine heart: their characteristics and alteration after ischemic preconditioning. *Basic Res Cardiol*. 1999;94:489–495. doi: 10.1007/s003950050165
- Patten RD, Aronovitz MJ, Deras-Mejia L, Pandian NG, Hanak GG, Smith JJ, Mendelsohn ME, Konstam MA. Ventricular remodeling in a mouse model of myocardial infarction. *Am J Physiol.* 1998;274:H1812–H1820. doi: 10.1152/ajpheart.1998.274.5.H1812
- 262. van den Bos EJ, Mees BM, de Waard MC, de Crom R, Duncker DJ. A novel model of cryoinjury-induced myocardial infarction in the mouse: a comparison with coronary artery ligation. Am J Physiol Heart Circ Physiol. 2005;289:H1291-H1300. doi: 10.1152/ajpheart.00111.2005
- 263. Srikanth G, Prakash P, Tripathy N, Dikshit M, Nityanand S. Establishment of a rat model of myocardial infarction with a high survival rate: a suitable model for evaluation of efficacy of stem cell therapy. J Stem Cells Regen Med. 2009;5:30–36.

- 264. Coker SJ. Anesthetized rabbit as a model for ischemia- and reperfusion-induced arrhythmias: effects of quinidine and bretylium. J Pharmacol Methods. 1989;21:263–279. doi: 10.1016/0160-5402(89)90064-8
- 265. Arimura T, Saku K, Kakino T, Nishikawa T, Tohyama T, Sakamoto T, Sakamoto K, Kishi T, Ide T, Sunagawa K. Intravenous electrical vagal nerve stimulation prior to coronary reperfusion in a canine ischemia-reperfusion model markedly reduces infarct size and prevents subsequent heart failure. Int J Cardiol. 2017;227:704–710. doi: 10.1016/j.ijcard.2016.10.074
- 266. Silvis MJM, van Hout GPJ, Fiolet ATL, Dekker M, Bosch L, van Nieuwburg MMJ, Visser J, Jansen MS, Timmers L, de Kleijn DPV. Experimental parameters and infarct size in closed chest pig LAD ischemia reperfusion models; lessons learned. *BMC Cardiovasc Disord*. 2021;21:171. doi: 10.1186/s12872-021-01995-7
- Charles CJ, Elliott JM, Nicholls MG, Rademaker MT, Richards M. Myocardial infarction with and without reperfusion in sheep: early cardiac and neurohumoral changes. Clin Sci (Lond). 2000;98:703

 –711.
- 268. Kim NK, Wolfson D, Fernandez N, Shin M, Cho HC. A rat model of complete atrioventricular block recapitulates clinical indices of bradycardia and provides a platform to test disease-modifying therapies. *Sci Rep.* 2019;9:6930. doi: 10.1038/s41598-019-43300-9
- Suto F, Cahill SA, Wilson GJ, Hamilton RM, Greenwald I, Gross GJ. A novel rabbit model of variably compensated complete heart block. *J Appl Physiol* (1985). 2002;92:1199–1204. doi: 10.1152/japplphysiol.00714.2001
- 270. Oros A, Beekman JD, Vos MA. The canine model with chronic, complete atrio-ventricular block. *Pharmacol Ther.* 2008;119:168–178. doi: 10.1016/j.pharmthera.2008.03.006
- 271. Steiner C, Kovalik AT. A simple technique for production of chronic complete heart block in dogs. J Appl Physiol. 1968;25:631–632. doi: 10.1152/jappl.1968.25.5.631
- 272. Greiser M, Neuberger HR, Harks E, El-Armouche A, Boknik P, de Haan S, Verheyen F, Verheule S, Schmitz W, Ravens U, Nattel S, Allessie MA, Dobrev D, Schotten U. Distinct contractile and molecular differences between two goat models of atrial dysfunction: AV block-induced atrial dilatation and atrial fibrillation. *J Mol Cell Cardiol.* 2009;46:385–394. doi: 10.1016/j.yjmcc.2008.11.012
- 273. Yamashita T, Murakawa Y, Hayami N, Fukui Ei, Kasaoka Y, Inoue M, Omata M. Short-term effects of rapid pacing on mRNA level of voltage-dependent K(+) channels in rat atrium: electrical remodeling in paroxysmal atrial tachycardia. *Circulation*. 2000;101:2007–2014. doi: 10.1161/01.cir.101.16.2007
- 274. Zhao Y, Gu TX, Zhang GW, Liu HG, Wang C. Losartan affects the substrate for atrial fibrillation maintenance in a rabbit model. *Cardiovasc Pathol.* 2013;22:383–388. doi: 10.1016/j.carpath.2013.02.005
- 275. Morillo CA, Klein GJ, Jones DL, Guiraudon CM. Chronic rapid atrial pacing. Structural, functional, and electrophysiological characteristics of a new model of sustained atrial fibrillation. *Circulation*. 1995;91:1588–1595. doi: 10.1161/01.cir.91.5.1588
- Bauer A, McDonald AD, Donahue JK. Pathophysiological findings in a model of persistent atrial fibrillation and severe congestive heart failure. Cardiovasc Res. 2004;61:764–770. doi: 10.1016/j.cardiores.2003.12.013
- 277. Ståhlberg M, Nakagawa R, Bedja D, Zhu G, Lin BL, Saberi A, Lee DI, Kass DA. Chronic Atrial and Ventricular Pacing in the Mouse. Circ Heart Fail. 2019;12:e005655. doi: 10.1161/CIRCHEARTFAILURE.118.005655
- 278. Mulla W, Etzion S, Elyagon S, Gillis R, Murninkas M, Konstantino Y, Mannhardt I, Eschenhagen T, Liel-Cohen N, Etzion Y. Prominent differences in left ventricular performance and myocardial properties between right ventricular and left ventricular-based pacing modes in rats. *Sci Rep.* 2017;7:5931. doi: 10.1038/s41598-017-06197-w
- Lamoureux L, Radhakrishnan J, Gazmuri RJ. A rat model of ventricular fibrillation and resuscitation by conventional closed-chest technique. J Vis Exp. 2015. doi: 10.3791/52413
- Armstrong PW, Stopps TP, Ford SE, de Bold AJ. Rapid ventricular pacing in the dog: pathophysiologic studies of heart failure. Circulation. 1986;74:1075–1084. doi: 10.1161/01.cir.74.5.1075
- Pemberton CJ, Johnson ML, Yandle TG, Espiner EA. Deconvolution analysis of cardiac natriuretic peptides during acute volume overload. *Hypertension*. 2000;36:355–359. doi: 10.1161/01.hyp.36.3.355
- 282. Amorós-Figueras G, Jorge E, Raga S, Alonso-Martin C, Rodríguez-Font E, Bazan V, Viñolas X, Cinca J, Guerra JM. Comparison between endocardial and epicardial cardiac resynchronization in an experimental model of non-ischaemic cardiomyopathy. *Europace*. 2018;20:1209–1216. doi: 10.1093/europace/eux212
- 283. Pummerer CL, Luze K, Grässl G, Bachmaier K, Offner F, Burrell SK, Lenz DM, Zamborelli TJ, Penninger JM, Neu N. Identification of cardiac myosin

- peptides capable of inducing autoimmune myocarditis in BALB/c mice. J Clin Invest. 1996;97:2057-2062. doi: 10.1172/JCI118642
- 284. Machino-Ohtsuka T, Tajiri K, Kimura T, Sakai S, Sato A, Yoshida T, Hiroe M, Yasutomi Y, Aonuma K, Imanaka-Yoshida K. Tenascin-C aggravates autoimmune myocarditis via dendritic cell activation and Th17 cell differentiation. J Am Heart Assoc. 2014;3:e001052. doi: 10.1161/JAHA.114.001052
- Izumi T, Kodama M, Shibata A. Experimental giant cell myocarditis induced by cardiac myosin immunization. *Eur Heart J.* 1991;12 Suppl D:166–168. doi: 10.1093/eurheartj/12.suppl_d.166
- Hosenpud JD, Campbell SM, Hart MV, Paul MS, Rowles JR, Niles NR. Experimental autoimmune myocarditis in the guinea pig. *Cardiovasc Res.* 1985;19:613–622. doi: 10.1093/cvr/19.10.613
- 287. Tselentakis EV, Woodford E, Chandy J, Gaudette GR, Saltman AE. Inflammation effects on the electrical properties of atrial tissue and inducibility of postoperative atrial fibrillation. *J Surg Res.* 2006;135:68–75. doi: 10.1016/j.jss.2006.03.024
- 288. Zhang Y, Wang YT, Shan ZL, Guo HY, Guan Y, Yuan HT. Role of inflammation in the initiation and maintenance of atrial fibrillation and the protective effect of atorvastatin in a goat model of aseptic pericarditis. *Mol Med Rep.* 2015;11:2615–2623. doi: 10.3892/mmr.2014.3116
- 289. Furman BL. Streptozotocin-induced diabetic models in mice and rats. *Curr Protoc Pharmacol.* 2015;70:5.47.1-5.47.20. doi: 10.1002/0471141755.ph0547s70
- 290. Wang CY, Liao JK. A mouse model of diet-induced obesity and insulin resistance. *Methods Mol Biol.* 2012;821:421–433. doi: 10.1007/978-1-61779-430-8 27
- 291. Holmes A, Coppey LJ, Davidson EP, Yorek MA. Rat models of diet-induced obesity and high fat/low dose streptozotocin type 2 diabetes: effect of reversal of high fat diet compared to treatment with enalapril or menhaden oil on glucose utilization and neuropathic endpoints. *J Diabetes Res.* 2015;2015:307285. doi: 10.1155/2015/307285
- 292. Fenichel RR, Malik M, Antzelevitch C, Sanguinetti M, Roden DM, Priori SG, Ruskin JN, Lipicky RJ, Cantilena LR; Independent Academic Task Force. Drug-induced torsades de pointes and implications for drug development. J Cardiovasc Electrophysiol. 2004;15:475–495. doi: 10.1046/j. 1540-8167.2004.03534.x
- 293. Lip GYH, Coca A, Kahan T, Boriani G, Manolis AS, Olsen MH, Oto A, Potpara TS, Steffel J, Marín F, et al; Reviewers:. Hypertension and cardiac arrhythmias: a consensus document from the european heart rhythm association (EHRA) and esc council on hypertension, endorsed by the heart rhythm society (HRS), asia-pacific heart rhythm society (APHRS) and sociedad latinoamericana de estimulación cardíaca y electrofisiología (SOLEACE). Europace. 2017;19:891–911. doi: 10.1093/europace/eux091
- 294. Narins CR, Aktas MK, Chen AY, McNitt S, Ling FS, Younis A, Zareba W, Daubert JP, Huang DT, Rosero S, et al. Arrhythmic and mortality outcomes among ischemic versus nonischemic cardiomyopathy patients receiving primary ICD therapy. *JACC Clin Electrophysiol.* 2022;8:1–11. doi: 10.1016/j.jacep.2021.06.020
- 295. Thomas KL, Jackson LR 2nd, Shrader P, Ansell J, Fonarow GC, Gersh B, Kowey PR, Mahaffey KW, Singer DE, Thomas L, et al. prevalence, characteristics, and outcomes of valvular heart disease in patients with atrial fibrillation: insights from the ORBIT-AF (outcomes registry for better informed treatment for atrial fibrillation). J Am Heart Assoc. 2017;6:e006475. doi: 10.1161/JAHA.117.006475
- 296. Peretto G, Sala S, Rizzo S, De Luca G, Campochiaro C, Sartorelli S, Benedetti G, Palmisano A, Esposito A, Tresoldi M, et al. Arrhythmias in myocarditis: State of the art. *Heart Rhythm.* 2019;16:793–801. doi: 10.1016/j.hrthm.2018.11.024
- 297. Pinilla-Vera M, Hahn VS, Kass DA. Leveraging signaling pathways to treat heart failure with reduced ejection fraction. *Circ Res.* 2019;124:1618– 1632. doi: 10.1161/CIRCRESAHA.119.313682
- Heineke J, Molkentin JD. Regulation of cardiac hypertrophy by intracellular signalling pathways. Nat Rev Mol Cell Biol. 2006;7:589–600. doi: 10.1038/nrm1983
- 299. Knowlton AA, Chen L, Malik ZA. Heart failure and mitochondrial dysfunction: the role of mitochondrial fission/fusion abnormalities and new therapeutic strategies. *J Cardiovasc Pharmacol.* 2014;63:196–206. doi: 10.1097/01.fjc.0000432861.55968.a6
- Zhou B, Tian R. Mitochondrial dysfunction in pathophysiology of heart failure. J Clin Invest. 2018;128:3716–3726. doi: 10.1172/JCI120849
- Bito V, Heinzel FR, Biesmans L, Antoons G, Sipido KR. Crosstalk between L-type Ca2+ channels and the sarcoplasmic reticulum: alterations during cardiac remodelling. *Cardiovasc Res.* 2008;77:315–324. doi: 10.1093/cvr/cvm063

- 302. Harris DM, Mills GD, Chen X, Kubo H, Berretta RM, Votaw VS, Santana LF, Houser SR. Alterations in early action potential repolarization causes localized failure of sarcoplasmic reticulum Ca2+ release. *Circ Res.* 2005;96:543–550. doi: 10.1161/01.RES.0000158966.58380.37
- 303. Wehrens XH, Lehnart SE, Reiken S, Vest JA, Wronska A, Marks AR. Ryanodine receptor/calcium release channel PKA phosphorylation: a critical mediator of heart failure progression. Proc Natl Acad Sci U S A. 2006;103:511-518. doi: 10.1073/pnas.0510113103
- Wende AR, Brahma MK, McGinnis GR, Young ME. Metabolic Origins of Heart Failure. JACC Basic Transl Sci. 2017;2:297–310. doi: 10.1016/j. iacbts.2016.11.009
- 305. Gloschat CR, Koppel AC, Aras KK, Brennan JA, Holzem KM, Efimov IR. Arrhythmogenic and metabolic remodelling of failing human heart. J Physiol. 2016;594:3963–3980. doi: 10.1113/JP271992
- 306. Koitabashi N, Danner T, Zaiman AL, Pinto YM, Rowell J, Mankowski J, Zhang D, Nakamura T, Takimoto E, Kass DA. Pivotal role of cardiomyocyte TGF-β signaling in the murine pathological response to sustained pressure overload. J Clin Invest. 2011;121:2301–2312. doi: 10.1172/JCI44824
- 307. De Jong AM, Van Gelder IC, Vreeswijk-Baudoin I, Cannon MV, Van Gilst WH, Maass AH. Atrial remodeling is directly related to end-diastolic left ventricular pressure in a mouse model of ventricular pressure overload. PLoS One. 2013;8:e72651. doi: 10.1371/journal.pone.0072651
- Marionneau C, Brunet S, Flagg TP, Pilgram TK, Demolombe S, Nerbonne JM. Distinct cellular and molecular mechanisms underlie functional remodeling of repolarizing K+ currents with left ventricular hypertrophy. Circ Res. 2008;102:1406–1415. doi: 10.1161/CIRCRESAHA.107.170050
- 309. Moorjani N, Catarino P, El-Sayed R, Al-Ahmed S, Meyer B, Al-Mohanna F, Westaby S. A pressure overload model to track the molecular biology of heart failure. Eur J Cardiothorac Surg. 2003;24:920–925. doi: 10.1016/s1010-7940(03)00514-1
- 310. Wang Y, Huang S, Sah VP, Ross J Jr, Brown JH, Han J, Chien KR. Cardiac muscle cell hypertrophy and apoptosis induced by distinct members of the p38 mitogen-activated protein kinase family. *J Biol Chem.* 1998;273:2161–2168. doi: 10.1074/jbc.273.4.2161
- 311. Li XM, Ma YT, Yang YN, Liu F, Chen BD, Han W, Zhang JF, Gao XM. Downregulation of survival signalling pathways and increased apoptosis in the transition of pressure overload-induced cardiac hypertrophy to heart failure. Clin Exp Pharmacol Physiol. 2009;36:1054–1061. doi: 10.1111/j.1440-1681.2009.05243.x
- 312. Patrucco E, Notte A, Barberis L, Selvetella G, Maffei A, Brancaccio M, Marengo S, Russo G, Azzolino O, Rybalkin SD, et al. Pl3Kgamma modulates the cardiac response to chronic pressure overload by distinct kinase-dependent and -independent effects. *Cell.* 2004;118:375–387. doi: 10.1016/j.cell.2004.07.017
- 313. Xia Y, Lee K, Li N, Corbett D, Mendoza L, Frangogiannis NG. Characterization of the inflammatory and fibrotic response in a mouse model of cardiac pressure overload. *Histochem Cell Biol.* 2009;131:471– 481. doi: 10.1007/s00418-008-0541-5
- 314. O'Connell TD, Swigart PM, Rodrigo MC, Ishizaka S, Joho S, Turnbull L, Tecott LH, Baker AJ, Foster E, Grossman W, Simpson PC. Alpha1adrenergic receptors prevent a maladaptive cardiac response to pressure overload. J Clin Invest. 2006;116:1005–1015. doi: 10.1172/JCI22811
- 315. Zhao M, Fajardo G, Urashima T, Spin JM, Poorfarahani S, Rajagopalan V, Huynh D, Connolly A, Quertermous T, Bernstein D. Cardiac pressure overload hypertrophy is differentially regulated by β-adrenergic receptor subtypes. Am J Physiol Heart Circ Physiol. 2011;301:H1461-H1470. doi: 10.1152/ajpheart.00453.2010
- 316. Rockman HA, Wachhorst SP, Mao L, Ross J Jr. ANG II receptor blockade prevents ventricular hypertrophy and ANF gene expression with pressure overload in mice. Am J Physiol. 1994;266(6 Pt 2):H2468–H2475. doi: 10.1152/ajpheart.1994.266.6.H2468
- 317. Qu J, Volpicelli FM, Garcia LI, Sandeep N, Zhang J, Márquez-Rosado L, Lampe PD, Fishman GI. Gap junction remodeling and spironolactone-dependent reverse remodeling in the hypertrophied heart. *Circ Res.* 2009;104:365–371. doi: 10.1161/CIRCRESAHA.108.184044
- 318. Weisheit C, Zhang Y, Faron A, Köpke O, Weisheit G, Steinsträsser A, Frede S, Meyer R, Boehm O, Hoeft A, et al. Ly6C(low) and not Ly6C(high) macrophages accumulate first in the heart in a model of murine pressure-overload. PLoS One. 2014;9:e112710. doi: 10.1371/journal.pone.0112710
- 319. Kent RL, Rozich JD, McCollam PL, McDermott DE, Thacker UF, Menick DR, McDermott PJ, Cooper G 4th. Rapid expression of the Na(+)-Ca2+ exchanger in response to cardiac pressure overload. Am J Physiol. 1993;265(3 Pt 2):H1024-H1029. doi: 10.1152/ajpheart.1993.265.3.H1024

- 320. Ujihara Y, Iwasaki K, Takatsu S, Hashimoto K, Naruse K, Mohri S, Katanosaka Y. Induced NCX1 overexpression attenuates pressure overload-induced pathological cardiac remodelling. Cardiovasc Res. 2016;111:348-361. doi: 10.1093/cvr/cvw113
- 321. del Monte F, Williams E, Lebeche D, Schmidt U, Rosenzweig A, Gwathmey JK, Lewandowski ED, Haijar RJ, Improvement in survival and cardiac metabolism after gene transfer of sarcoplasmic reticulum Ca(2+)-ATPase in a rat model of heart failure. Circulation. 2001;104:1424-1429. doi: 10.1161/hc3601.095574
- 322. Lyon AR, Bannister ML, Collins T, Pearce E, Sepehripour AH, Dubb SS, Garcia E, O'Gara P, Liang L, Kohlbrenner E, et al. SERCA2a gene transfer decreases sarcoplasmic reticulum calcium leak and reduces ventricular arrhythmias in a model of chronic heart failure. Circ Arrhythm Electrophysiol. 2011;4:362-372. doi: 10.1161/CIRCEP.110.961615
- 323. Cutler MJ, Wan X, Plummer BN, Liu H, Deschenes I, Laurita KR, Hajjar RJ, Rosenbaum DS. Targeted sarcoplasmic reticulum Ca2+ ATPase 2a gene delivery to restore electrical stability in the failing heart. Circulation. 2012;126:2095-2104. doi: 10.1161/CIRCULATIONAHA.111.071480
- 324. Greenberg B, Butler J, Felker GM, Ponikowski P, Voors AA, Desai AS, Barnard D, Bouchard A, Jaski B, Lyon AR, et al. Calcium upregulation by percutaneous administration of gene therapy in patients with cardiac disease (CUPID 2): a randomised, multinational, double-blind, placebo-controlled, phase 2b trial. Lancet. 2016;387:1178-1186. doi: 10.1016/S0140-6736(16)00082-9
- 325. Zhang C, Yasuno S, Kuwahara K, Zankov DP, Kobori A, Makiyama T, Horie M. Blockade of angiotensin II type 1 receptor improves the arrhythmia morbidity in mice with left ventricular hypertrophy. Circ J. 2006;70:335-341. doi: 10.1253/circi.70.335
- 326. Jin H, Chemaly ER, Lee A, Kho C, Hadri L, Hajjar RJ, Akar FG. Mechanoelectrical remodeling and arrhythmias during progression of hypertrophy. *FASEB J.* 2010;24:451-463. doi: 10.1096/fj.09-136622
- 327. Boulaksil M, Noorman M, Engelen MA, van Veen TA, Vos MA, de Bakker JM, van Rijen HV. Longitudinal arrhythmogenic remodelling in a mouse model of longstanding pressure overload. Neth Heart J. 2010;18:509-515, doi: 10.1007/BF03091824
- 328. Westphal C, Spallek B, Konkel A, Marko L, Qadri F, DeGraff LM, Schubert C, Bradbury JA, Regitz-Zagrosek V, Falck JR, et al. CYP2J2 overexpression protects against arrhythmia susceptibility in cardiac hypertrophy. PLoS One. 2013;8:e73490. doi: 10.1371/journal.pone.0073490
- 329. Bosch L, de Haan JJ, Bastemeijer M, van der Burg J, van der Worp E, Wesseling M, Viola M, Odille C, El Azzouzi H, Pasterkamp G, et al. The transverse aortic constriction heart failure animal model: a systematic review and meta-analysis. Heart Fail Rev. 2021;26:1515-1524. doi: 10.1007/s10741-020-09960-w
- 330. Zi M, Stafford N, Prehar S, Baudoin F, Oceandy D, Wang X, Bui T, Shaheen M, Neyses L, Cartwright EJ. Cardiac hypertrophy or failure? - a systematic evaluation of the transverse aortic constriction model in C57BL/6NTac and C57BL/6J substrains. Curr Res Physiol. 2019;1:1-10. doi: 10.1016/j.crphys.2019.10.001
- 331. Barrick CJ, Rojas M, Schoonhoven R, Smyth SS, Threadgill DW. Cardiac response to pressure overload in 129S1/SvlmJ and C57BL/6J mice: temporal- and background-dependent development of concentric left ventricular hypertrophy. Am J Physiol Heart Circ Physiol. 2007;292:H2119-H2130. doi: 10.1152/ajpheart.00816.2006
- 332. Koentges C, Pepin ME, Müsse C, Pfeil K, Alvarez SVV, Hoppe N, Hoffmann MM, Odening KE, Sossalla S, Zirlik A, et al. Gene expression analysis to identify mechanisms underlying heart failure susceptibility in mice and humans. Basic Res Cardiol. 2018;113:8. doi: 10.1007/s00395-017-0666-6
- 333. Nickel AG, von Hardenberg A, Hohl M, Löffler JR, Kohlhaas M, Becker J, Reil JC, Kazakov A, Bonnekoh J, Stadelmaier M, et al. Reversal of mitochondrial transhydrogenase causes oxidative stress in heart failure. Cell Metab. 2015;22:472-484. doi: 10.1016/j.cmet.2015.07.008
- 334. Litwin SE, Katz SE, Weinberg EO, Lorell BH, Aurigemma GP, Douglas PS. Serial echocardiographic-doppler assessment of left ventricular geometry and function in rats with pressure-overload hypertrophy, chronic angiotensin-converting enzyme inhibition attenuates the transition to heart failure. Circulation. 1995;91:2642-2654. doi: 10.1161/01.cir.91.10.2642
- 335. Pogwizd SM, Qi M, Yuan W, Samarel AM, Bers DM. Upregulation of Na(+)/Ca(2+) exchanger expression and function in an arrhythmogenic rabbit model of heart failure. Circ Res. 1999;85:1009-1019. doi: 10.1161/01.res.85.11.1009
- 336. Desantiago J, Ai X, Islam M, Acuna G, Ziolo MT, Bers DM, Pogwizd SM. Arrhythmogenic effects of beta2-adrenergic stimulation in the failing heart

- are attributable to enhanced sarcoplasmic reticulum Ca load. Circ Res. 2008;102:1389-1397. doi: 10.1161/CIRCRESAHA.107.169011
- 337. Pogwizd SM, Schlotthauer K, Li L, Yuan W, Bers DM. Arrhythmogenesis and contractile dysfunction in heart failure: roles of sodium-calcium exchange, inward rectifier potassium current, and residual beta-adrenergic responsiveness. Circ Res. 2001;88:1159-1167. doi: 10.1161/ hh1101.091193
- 338. Liu T, Takimoto E, Dimaano VL, DeMazumder D, Kettlewell S, Smith G, Sidor A, Abraham TP, O'Rourke B. Inhibiting mitochondrial Na+/Ca2+ exchange prevents sudden death in a guinea pig model of heart failure. Circ Res. 2014;115:44-54. doi: 10.1161/CIRCRESAHA.115.303062
- 339. Wang Y, Yuan J, Qian Z, Zhang X, Chen Y, Hou X, Zou J. B2 adrenergic receptor activation governs cardiac repolarization and arrhythmogenesis in a guinea pig model of heart failure. Sci Rep. 2015;5:7681. doi: 10.1038/srep07681
- 340. Dey S, DeMazumder D, Sidor A, Foster DB, O'Rourke B. Mitochondrial ROS drive sudden cardiac death and chronic proteome remodeling in heart failure. Circ Res. 2018;123:356-371. doi: 10.1161/ CIRCRESAHA.118.312708
- 341. Hegyi B, Morotti S, Liu C, Ginsburg KS, Bossuyt J, Belardinelli L, Izu LT, Chen-Izu Y, Bányász T, Grandi E, Bers DM. Enhanced depolarization drive in failing rabbit ventricular myocytes: calcium-dependent and βadrenergic effects on late sodium, L-type calcium, and sodium-calcium exchange currents. Circ Arrhythm Electrophysiol. 2019;12:e007061. doi: 10.1161/CIRCEP.118.007061
- 342. Aoyagi T, Fujii AM, Flanagan MF, Arnold LW, Brathwaite KW, Colan SD, Mirsky I. Transition from compensated hypertrophy to intrinsic myocardial dysfunction during development of left ventricular pressureoverload hypertrophy in conscious sheep, systolic dysfunction precedes diastolic dysfunction. Circulation. 1993;88(5 Pt 1):2415-2425. doi: 10.1161/01.cir.88.5.2415
- 343. Ye Y, Gong G, Ochiai K, Liu J, Zhang J. High-energy phosphate metabolism and creatine kinase in failing hearts: a new porcine model. Circulation. 2001;103:1570-1576. doi: 10.1161/01.cir.103.11.1570
- 344. Hiemstra JA, Lee DI, Chakir K, Gutiérrez-Aguilar M, Marshall KD, Zgoda PJ, Cruz Rivera N, Dozier DG, Ferguson BS, Heublein DM, et al. Saxagliptin and tadalafil differentially alter cyclic guanosine monophosphate (cGMP) signaling and left ventricular function in aortic-banded mini-swine. J Am Heart Assoc. 2016;5:e003277. doi: 10.1161/JAHA.116.003277
- 345. Ishikawa K, Aguero J, Oh JG, Hammoudi N, Fish LA, Leonardson L, Picatoste B, Santos-Gallego CG, Fish KM, Hajjar RJ. Increased stiffness is the major early abnormality in a pig model of severe aortic stenosis and predisposes to congestive heart failure in the absence of systolic dysfunction. J Am Heart Assoc. 2015;4:e001925. doi: 10.1161/JAHA.115.001925
- 346. Yarbrough WM, Mukherjee R, Stroud RE, Rivers WT, Oelsen JM, Dixon JA, Eckhouse SR, Ikonomidis JS, Zile MR, Spinale FG. Progressive induction of left ventricular pressure overload in a large animal model elicits myocardial remodeling and a unique matrix signature. J Thorac Cardiovasc Surg. 2012;143:215-223. doi: 10.1016/j.jtcvs.2011.09.032
- 347. Moorjani N, Catarino P, Trabzuni D, Saleh S, Moorji A, Dzimiri N, Al-Mohanna F, Westaby S, Ahmad M. Upregulation of Bcl-2 proteins during the transition to pressure overload-induced heart failure. Int J Cardiol. 2007;116:27-33. doi: 10.1016/j.ijcard.2006.04.037
- 348. Fleet WF, Johnson TA, Graebner CA, Gettes LS. Effect of serial brief ischemic episodes on extracellular K+, pH, and activation in the pig. Circulation. 1985;72:922-932. doi: 10.1161/01.cir.72.4.922
- 349. Johnson TA, Engle CL, Boyd LM, Koch GG, Gwinn M, Gettes LS. Magnitude and time course of extracellular potassium inhomogeneities during acute ischemia in pigs. Effect of verapamil. Circulation. 1991;83:622-634. doi: 10.1161/01.cir.83.2.622
- 350. Huss JM, Kelly DP. Nuclear receptor signaling and cardiac energetics. Circ Res. 2004;95:568-578. doi: 10.1161/01.RES.0000141774.29937.e3
- 351. Cadenas S. ROS and redox signaling in myocardial ischemia-reperfusion injury and cardioprotection. Free Radic Biol Med. 2018;117:76-89. doi: 10.1016/j.freeradbiomed.2018.01.024
- 352. Nahrendorf M, Swirski FK, Aikawa E, Stangenberg L, Wurdinger T, Figueiredo JL, Libby P, Weissleder R, Pittet MJ. The healing myocardium sequentially mobilizes two monocyte subsets with divergent and complementary functions. J Exp Med. 2007;204:3037-3047. doi: 10.1084/jem.20070885
- 353. Xu Z, Alloush J, Beck E, Weisleder N. A murine model of myocardial ischemia-reperfusion injury through ligation of the left anterior descending artery. J Vis Exp. 2014. doi: 10.3791/51329

- Lugrin J, Parapanov R, Krueger T, Liaudet L. Murine myocardial infarction model using permanent ligation of left anterior descending coronary artery. J Vis Exp. 2019. doi: 10.3791/59591
- 355. Wang D, Tediashvili G, Hu X, Gravina A, Marcus SG, Zhang H, Olgin JE, Deuse T, Schrepfer S. A cryoinjury model to study myocardial infarction in the mouse. J Vis Exp. 2019. doi: 10.3791/59958
- 356. Nofi C, Zhang K, Tang YD, Li Y, Migirov A, Ojamaa K, Gerdes AM, Zhang Y. Chronic dantrolene treatment attenuates cardiac dysfunction and reduces atrial fibrillation inducibility in a rat myocardial infarction heart failure model. Heart Rhythm O2. 2020;1:126–135. doi: 10.1016/j.hroo.2020.03.004
- 357. Beiert T, Tiyerili V, Knappe V, Effelsberg V, Linhart M, Stöckigt F, Klein S, Schierwagen R, Trebicka J, Nickenig G, et al. Relaxin reduces susceptibility to post-infarct atrial fibrillation in mice due to anti-fibrotic and anti-inflammatory properties. *Biochem Biophys Res Commun.* 2017;490:643–649. doi: 10.1016/j.bbrc.2017.06.091
- 358. Boixel C, Fontaine V, Rücker-Martin C, Milliez P, Louedec L, Michel JB, Jacob MP, Hatem SN. Fibrosis of the left atria during progression of heart failure is associated with increased matrix metalloproteinases in the rat. J Am Coll Cardiol. 2003;42:336–344. doi: 10.1016/s0735-1097(03)00578-3
- 359. Curtis MJ, Macleod BA, Walker MJ. Models for the study of arrhythmias in myocardial ischaemia and infarction: the use of the rat. *J Mol Cell Cardiol.* 1987;19:399–419. doi: 10.1016/s0022-2828(87)80585-0
- Hundahl LA, Tfelt-Hansen J, Jespersen T. Rat models of ventricular fibrillation following acute myocardial infarction. J Cardiovasc Pharmacol Ther. 2017;22:514–528. doi: 10.1177/1074248417702894
- 361. Kolossov E, Bostani T, Roell W, Breitbach M, Pillekamp F, Nygren JM, Sasse P, Rubenchik O, Fries JW, Wenzel D, et al. Engraftment of engineered ES cell-derived cardiomyocytes but not BM cells restores contractile function to the infarcted myocardium. *J Exp Med.* 2006;203:2315–2327. doi: 10.1084/jem.20061469
- 362. Rucker-Martin C, Milliez P, Tan S, Decrouy X, Recouvreur M, Vranckx R, Delcayre C, Renaud JF, Dunia I, Segretain D, Hatem SN. Chronic hemodynamic overload of the atria is an important factor for gap junction remodeling in human and rat hearts. *Cardiovasc Res.* 2006;72:69–79. doi: 10.1016/j.cardiores.2006.06.016
- 363. Zhang Y, Dedkov El, Lee B 3rd, Li Y, Pun K, Gerdes AM. Thyroid hormone replacement therapy attenuates atrial remodeling and reduces atrial fibrillation inducibility in a rat myocardial infarction-heart failure model. *J Card Fail*. 2014;20:1012–1019. doi: 10.1016/j.cardfail.2014.10.003
- 364. Schrickel JW, Bielik H, Yang A, Schimpf R, Shlevkov N, Burkhardt D, Meyer R, Grohé C, Fink K, Tiemann K, et al. Induction of atrial fibrillation in mice by rapid transesophageal atrial pacing. *Basic Res Cardiol*. 2002;97:452–460. doi: 10.1007/s003950200052
- Berul CI, Aronovitz MJ, Wang PJ, Mendelsohn ME. In vivo cardiac electrophysiology studies in the mouse. *Circulation*. 1996;94:2641–2648. doi: 10.1161/01.cir.94.10.2641
- 366. Clasen L, Eickholt C, Angendohr S, Jungen C, Shin DI, Donner B, Fürnkranz A, Kelm M, Klöcker N, Meyer C, Makimoto H. A modified approach for programmed electrical stimulation in mice: inducibility of ventricular arrhythmias. *PLoS One*. 2018;13:e0201910. doi: 10.1371/journal.pone.0201910
- 367. Gehrmann J, Frantz S, Maguire CT, Vargas M, Ducharme A, Wakimoto H, Lee RT, Berul Cl. Electrophysiological characterization of murine myocardial ischemia and infarction. *Basic Res Cardiol*. 2001;96:237–250. doi: 10.1007/s003950170054
- 368. Lübkemeier I, Andrié R, Lickfett L, Bosen F, Stöckigt F, Dobrowolski R, Draffehn AM, Fregeac J, Schultze JL, Bukauskas FF, et al. The Connexin40A96S mutation from a patient with atrial fibrillation causes decreased atrial conduction velocities and sustained episodes of induced atrial fibrillation in mice. *J Mol Cell Cardiol*. 2013;65:19–32. doi: 10.1016/j.yjmcc.2013.09.008
- 369. Ng GA, Cobbe SM, Smith GL. Non-uniform prolongation of intracellular Ca2+ transients recorded from the epicardial surface of isolated hearts from rabbits with heart failure. *Cardiovasc Res.* 1998;37:489–502. doi: 10.1016/s0008-6363(97)00255-1
- 370. Nisbet AM, Camelliti P, Walker NL, Burton FL, Cobbe SM, Kohl P, Smith GL. Prolongation of atrio-ventricular node conduction in a rabbit model of ischaemic cardiomyopathy: role of fibrosis and connexin remodelling. *J Mol Cell Cardiol*. 2016;94:54–64. doi: 10.1016/j.yjmcc.2016.03.011
- 371. Sinno H, Derakhchan K, Libersan D, Merhi Y, Leung TK, Nattel S. Atrial ischemia promotes atrial fibrillation in dogs. *Circulation*. 2003;107:1930–1936. doi: 10.1161/01.CIR.0000058743.15215.03
- 372. Nishida K, Qi XY, Wakili R, Comtois P, Chartier D, Harada M, Iwasaki YK, Romeo P, Maguy A, Dobrev D, et al. Mechanisms of atrial tachyarrhythmias associated with coronary artery occlusion in a

- chronic canine model. *Circulation*. 2011;123:137-146. doi: 10.1161/CIRCULATIONAHA.110.972778
- Rivard L, Sinno H, Shiroshita-Takeshita A, Schram G, Leung TK, Nattel S. The pharmacological response of ischemia-related atrial fibrillation in dogs: evidence for substrate-specific efficacy. *Cardiovasc Res.* 2007;74:104–113. doi: 10.1016/j.cardiores.2007.01.018
- 374. Keeran KJ, Jeffries KR, Zetts AD, Taylor J, Kozlov S, Hunt TJ. A chronic cardiac ischemia model in swine using an ameroid constrictor. J Vis Exp. 2017. doi: 10.3791/56190
- 375. Millard RW. Induction of functional coronary collaterals in the swine heart. Basic Res Cardiol. 1981;76:468–473. doi: 10.1007/BF01908345
- Fallavollita JA, Riegel BJ, Suzuki G, Valeti U, Canty JM Jr. Mechanism of sudden cardiac death in pigs with viable chronically dysfunctional myocardium and ischemic cardiomyopathy. Am J Physiol Heart Circ Physiol. 2005;289:H2688-H2696. doi: 10.1152/ajpheart.00653.2005
- 377. Canty JM Jr, Suzuki G, Banas MD, Verheyen F, Borgers M, Fallavollita JA. Hibernating myocardium: chronically adapted to ischemia but vulnerable to sudden death. *Circ Res.* 2004;94:1142–1149. doi: 10.1161/01. RES.0000125628.57672.CF
- 378. Hegyi B, Bossuyt J, Griffiths LG, Shimkunas R, Coulibaly Z, Jian Z, Grimsrud KN, Sondergaard CS, Ginsburg KS, Chiamvimonvat N, et al. Complex electrophysiological remodeling in postinfarction ischemic heart failure. Proc Natl Acad Sci U S A. 2018;115:E3036–E3044. doi: 10.1073/pnas.1718211115
- Sasano T, McDonald AD, Kikuchi K, Donahue JK. Molecular ablation of ventricular tachycardia after myocardial infarction. *Nat Med.* 2006;12:1256–1258. doi: 10.1038/nm1503
- Greener ID, Sasano T, Wan X, Igarashi T, Strom M, Rosenbaum DS, Donahue JK. Connexin43 gene transfer reduces ventricular tachycardia susceptibility after myocardial infarction. J Am Coll Cardiol. 2012;60:1103–1110. doi: 10.1016/j.jacc.2012.04.042
- Syed FF, Hayes DL, Friedman PA, Asirvatham, SJ. Hemodynamics of cardiac pacing: optimization and programming to enhance cardiac function. In: Cardiac Pacing, Defibrillation and Resynchronization: A Clinical Approach. Wiley-Blackwell; 2012:41-92.
- 382. Xia R, Vlcek J, Bauer J, Kaab S, Ishikawa-Ankerhold H, van den Heuvel DA, Schulz C, Massberg S, Clauss S. Whole-mount immunofluorescence staining, confocal imaging and 3D reconstruction of the sinoatrial and atrioventricular node in the mouse. J Vis Exp. 2020. doi: 10.3791/62058
- 383. Hulsmans M, Clauss S, Xiao L, Aguirre AD, King KR, Hanley A, Hucker WJ, Wülfers EM, Seemann G, Courties G, et al. Macrophages Facilitate Electrical Conduction in the Heart. Cell. 2017;169:510-522.e20. doi: 10.1016/i.cell.2017.03.050
- 384. Gizurarson S, Lorentzon M, Råmunddal T, Waagstein F, Bergfeldt L, Omerovic E. Effects of complete heart block on myocardial function, morphology, and energy metabolism in the rat. *Europace*. 2007;9:411–416. doi: 10.1093/europace/eum065
- 385. Vos MA, Verduyn SC, Gorgels AP, Lipcsei GC, Wellens HJ. Reproducible induction of early afterdepolarizations and torsade de pointes arrhythmias by d-sotalol and pacing in dogs with chronic atrioventricular block. *Circulation*. 1995;91:864–872. doi: 10.1161/01.cir.91.3.864
- 386. Sipido KR, Volders PG, de Groot SH, Verdonck F, Van de Werf F, Wellens HJ, Vos MA. Enhanced Ca(2+) release and Na/Ca exchange activity in hypertrophied canine ventricular myocytes: potential link between contractile adaptation and arrhythmogenesis. *Circulation*. 2000;102:2137–2144. doi: 10.1161/01.cir.102.17.2137
- 387. de Groot SH, Vos MA, Gorgels AP, Leunissen JD, van der Steld BJ, Wellens HJ. Combining monophasic action potential recordings with pacing to demonstrate delayed afterdepolarizations and triggered arrhythmias in the intact heart value of diastolic slope. *Circulation*. 1995;92:2697–2704. doi: 10.1161/01.cir.92.9.2697
- 388. Tse HF, Lau CP. Long-term effect of right ventricular pacing on myocardial perfusion and function. J Am Coll Cardiol. 1997;29:744-749. doi: 10.1016/s0735-1097(96)00586-4
- 389. Lee MA, Dae MW, Langberg JJ, Griffin JC, Chin MC, Finkbeiner WE, O'Connell JW, Botvinick E, Scheinman MM, Rosenqvist M. Effects of long-term right ventricular apical pacing on left ventricular perfusion, innervation, function and histology. J Am Coll Cardiol. 1994;24:225–232. doi: 10.1016/0735-1097(94)90567-3
- Pak PH, Nuss HB, Tunin RS, Kääb S, Tomaselli GF, Marban E, Kass DA. Repolarization abnormalities, arrhythmia and sudden death in canine tachy-cardia-induced cardiomyopathy. *J Am Coll Cardiol*. 1997;30:576–584. doi: 10.1016/s0735-1097(97)00193-9

- 391. Kääb S, Nuss HB, Chiamvimonvat N, O'Rourke B, Pak PH, Kass DA, Marban E, Tomaselli GF. Ionic mechanism of action potential prolongation in ventricular myocytes from dogs with pacing-induced heart failure. Circ Res. 1996;78:262-273. doi: 10.1161/01.res.78.2.262
- 392. Kääb S, Dixon J, Duc J, Ashen D, Näbauer M, Beuckelmann DJ, Steinbeck G, McKinnon D, Tomaselli GF. Molecular basis of transient outward potassium current downregulation in human heart failure: a decrease in Kv4.3 mRNA correlates with a reduction in current density. Circulation. 1998;98:1383-1393. doi: 10.1161/01.cir.98.14.1383
- 393. Aiba T, Barth AS, Hesketh GG, Hashambhoy YL, Chakir K, Tunin RS, Greenstein JL, Winslow RL, Kass DA, Tomaselli GF. Cardiac resynchronization therapy improves altered Na channel gating in canine model of dyssynchronous heart failure. Circ Arrhythm Electrophysiol. 2013;6:546-554. doi: 10.1161/CIRCEP.113.000400
- 394. Winslow RL, Rice J, Jafri S, Marbán E, O'Rourke B. Mechanisms of altered excitation-contraction coupling in canine tachycardia-induced heart failure, II: model studies. Circ Res. 1999;84:571-586. doi: 10.1161/01. res.84.5.571
- 395. Li H, Lichter JG, Seidel T, Tomaselli GF, Bridge JH, Sachse FB. Cardiac resynchronization therapy reduces subcellular heterogeneity of ryanodine receptors, T-Tubules, and Ca2+ sparks produced by dyssynchronous heart failure. Circ Heart Fail. 2015;8:1105-1114. doi: 10.1161/ CIRCHEARTFAILURE, 115,002352
- 396. DeMazumder D, Kass DA, O'Rourke B, Tomaselli GF. Cardiac resynchronization therapy restores sympathovagal balance in the failing heart by differential remodeling of cholinergic signaling. Circ Res. 2015;116:1691-1699. doi: 10.1161/CIRCRESAHA.116.305268
- 397. Wiesmann T, Freitag D, Dersch W, Eschbach D, Irqsusi M, Steinfeldt T, Wulf H, Feldmann C. Dantrolene versus amiodarone for cardiopulmonary resuscitation: a randomized, double-blinded experimental study. Sci Rep. 2017;7:40875. doi: 10.1038/srep40875
- 398. Shinagawa K, Shi YF, Tardif JC, Leung TK, Nattel S. Dynamic nature of atrial fibrillation substrate during development and reversal of heart failure in dogs. Circulation. 2002;105:2672-2678. doi: 10.1161/01.cir. 0000016826.62813.f5
- 399. Shinagawa K, Li D, Leung TK, Nattel S. Consequences of atrial tachycardia-induced remodeling depend on the preexisting atrial substrate. Circulation. 2002;105:251-257. doi: 10.1161/hc0202.102014
- 400. Cha TJ, Ehrlich JR, Zhang L, Shi YF, Tardif JC, Leung TK, Nattel S. Dissociation between ionic remodeling and ability to sustain atrial fibrillation during recovery from experimental congestive heart failure. Circulation. 2004;109:412-418. doi: 10.1161/01.CIR.0000109501.47603.0C
- 401. Cha TJ, Ehrlich JR, Zhang L, Nattel S. Atrial ionic remodeling induced by atrial tachycardia in the presence of congestive heart failure. Circulation. 2004;110:1520-1526. doi: 10.1161/01.CIR.0000142052.03565.87
- 402. Chen Y, Wakili R, Xiao J, Wu CT, Luo X, Clauss S, Dawson K, Qi X, Naud P, Shi YF, et al. Detailed characterization of microRNA changes in a canine heart failure model: Relationship to arrhythmogenic structural remodeling. J Mol Cell Cardiol. 2014;77:113-124. doi: 10.1016/j.yjmcc.2014.10.001
- 403. Luo X, Pan Z, Shan H, Xiao J, Sun X, Wang N, Lin H, Xiao L, Maguy A, Qi XY, et al. MicroRNA-26 governs profibrillatory inward-rectifier potassium current changes in atrial fibrillation. J Clin Invest. 2013;123:1939-1951. doi: 10.1172/JCI62185
- 404. Jia X, Zheng S, Xie X, Zhang Y, Wang W, Wang Z, Zhang Y, Wang J, Gao M, Hou Y. MicroRNA-1 accelerates the shortening of atrial effective refractory period by regulating KCNE1 and KCNB2 expression: an atrial tachypacing rabbit model. PLoS One. 2013;8:e85639. doi: 10.1371/journal.pone.0085639
- 405. Gutruf P, Yin RT, Lee KB, Ausra J, Brennan JA, Qiao Y, Xie Z, Peralta R, Talarico O, Murillo A, et al. Wireless, battery-free, fully implantable multimodal and multisite pacemakers for applications in small animal models. Nat Commun. 2019;10:5742. doi: 10.1038/s41467-019-13637-w
- 406. Mulla W, Hajaj B, Elyagon S, Mor M, Gillis R, Murninkas M, Klapper-Goldstein H. Plaschkes I. Chalifa-Caspi V. Etzion S. Etzion Y. Rapid atrial pacing promotes atrial fibrillation substrate in unanesthetized instrumented rats. Front Physiol. 2019;10:1218. doi: 10.3389/fphys.2019.01218
- 407. Chen MH, Liu TW, Xie L, Song FQ, He T, Zeng ZY, Mo SR. Ventricular fibrillation induced by transoesophageal cardiac pacing: a new model of cardiac arrest in rats. Resuscitation. 2007;74:546-551. doi: 10.1016/j.resuscitation.2007.01.039
- 408. Bruins P, te Velthuis H, Yazdanbakhsh AP, Jansen PG, van Hardevelt FW, de Beaumont EM, Wildevuur CR, Eijsman L, Trouwborst A, Hack CE. Activation of the complement system during and after cardiopulmonary bypass surgery: postsurgery activation involves C-reactive protein and is

- associated with postoperative arrhythmia. Circulation. 1997;96:3542-3548. doi: 10.1161/01.cir.96.10.3542
- 409. Chung MK, Martin DO, Sprecher D, Wazni O, Kanderian A, Carnes CA, Bauer JA, Tchou PJ, Niebauer MJ, Natale A, Van Wagoner DR. C-reactive protein elevation in patients with atrial arrhythmias: inflammatory mechanisms and persistence of atrial fibrillation. Circulation. 2001:104:2886-2891. doi: 10.1161/hc4901.101760
- 410. Dobrev D, Aguilar M, Heijman J, Guichard JB, Nattel S. Postoperative atrial fibrillation: mechanisms, manifestations and management. Nat Rev Cardiol. 2019;16:417-436. doi: 10.1038/s41569-019-0166-5
- 411. Aviles RJ, Martin DO, Apperson-Hansen C, Houghtaling PL, Rautaharju P, Kronmal RA, Tracy RP, Van Wagoner DR, Psaty BM, Lauer MS, Chung MK. Inflammation as a risk factor for atrial fibrillation. Circulation. 2003;108:3006-3010. doi: 10.1161/01.CIR.0000103131.70301.4F
- 412. Marott SC, Nordestgaard BG, Zacho J, Friberg J, Jensen GB, Tybjaerg-Hansen A, Benn M. Does elevated C-reactive protein increase atrial fibrillation risk? A Mendelian randomization of 47,000 individuals from the general population. J Am Coll Cardiol. 2010;56:789-795. doi: 10.1016/i.iacc,2010.02.066
- 413. Conen D, Ridker PM, Everett BM, Tedrow UB, Rose L, Cook NR, Buring JE, Albert CM. A multimarker approach to assess the influence of inflammation on the incidence of atrial fibrillation in women. Eur Heart J. 2010;31:1730-1736. doi: 10.1093/eurheartj/ehq146
- 414. Hu YF, Chen YJ, Lin YJ, Chen SA. Inflammation and the pathogenesis of atrial fibrillation. Nat Rev Cardiol. 2015;12:230-243. doi: 10.1038/ nrcardio.2015.2
- 415. Guo Y, Lip GY, Apostolakis S. Inflammation in atrial fibrillation. J Am Coll Cardiol. 2012;60:2263-2270. doi: 10.1016/j.jacc.2012.04.063
- 416. Reilly SN, Jayaram R, Nahar K, Antoniades C, Verheule S, Channon KM, Alp NJ, Schotten U, Casadei B. Atrial sources of reactive oxygen species vary with the duration and substrate of atrial fibrillation: implications for the antiarrhythmic effect of statins. Circulation. 2011;124:1107-1117. doi: 10.1161/CIRCULATIONAHA.111.029223
- 417. Antoniades C, Demosthenous M, Reilly S, Margaritis M, Zhang MH, Antonopoulos A, Marinou K, Nahar K, Jayaram R, Tousoulis D, et al. Myocardial redox state predicts in-hospital clinical outcome after cardiac surgery effects of short-term pre-operative statin treatment. J Am Coll Cardiol. 2012;59:60-70. doi: 10.1016/j.jacc.2011.08.062
- 418. Tschöpe C, Ammirati E, Bozkurt B, Caforio ALP, Cooper LT, Felix SB, Hare JM, Heidecker B, Heymans S, Hübner N, et al. Myocarditis and inflammatory cardiomyopathy: current evidence and future directions. Nat Rev Cardiol. 2021;18:169-193. doi: 10.1038/s41569-020-00435-x
- 419. Bailey JR, Loftus A RN, Allan RJC. Myopericarditis: recognition and impact in the military population. J R Army Med Corps. 2019;165:451-453. doi: 10.1136/jramc-2018-001056
- 420. Tselios K, Urowitz MB. Cardiovascular and pulmonary manifestations of systemic lupus erythematosus. Curr Rheumatol Rev. 2017;13:206-218. doi: 10.2174/1573397113666170704102444
- 421. Lazaros G, Oikonomou E, Tousoulis D. Established and novel treatment options in acute myocarditis, with or without heart failure. Expert Rev Cardiovasc Ther. 2017;15:25-34. doi: 10.1080/14779072.2017.1262764
- 422. Radhakrishnan VV. Experimental myocarditis in the guinea-pig. Cardiovasc Res. 1996;31:651-654.
- 423. GRODUMS EI, DEMPSTER G. Myocarditis in experimental Coxsackie B-3 infection. Can J Microbiol. 1959;5:605-615. doi: 10.1139/m59-074
- 424. Kishimoto C, Matsumori A, Ohmae M, Tomioka N, Kawai C. Electrocardiographic findings in experimental myocarditis in DBA/2 mice: complete atrioventricular block in the acute stage, low voltage of the QRS complex in the subacute stage and arrhythmias in the chronic stage. J Am Coll Cardiol. 1984;3:1461-1468. doi: 10.1016/s0735-1097(84)80285-5
- 425. Tracy S, Höfling K, Pirruccello S, Lane PH, Reyna SM, Gauntt CJ. Group B coxsackievirus myocarditis and pancreatitis: connection between viral virulence phenotypes in mice. J Med Virol. 2000;62:70-81. doi: 10.1002/1096-9071(200009)62:1<70::aid-jmv11>3.0.co;2-r
- 426. Terasaki F, Kitaura Y, Hayashi T, Nakayama Y, Deguchi H, Kawamura K. Arrhythmias in Coxsackie B3 virus myocarditis. Continuous electrocardiography in conscious mice and histopathology of the heart with special reference to the conduction system. Heart Vessels Suppl. 1990;5:45-50.
- 427. Kaese S, Larbig R, Rohrbeck M, Frommeyer G, Dechering D, Olligs J, Schönhofer-Merl S, Wessely R, Klingel K, Seebohm G, Eckardt L. Electrophysiological alterations in a murine model of chronic coxsackievirus B3 myocarditis. PLoS One. 2017;12:e0180029. doi: 10.1371/ journal.pone.0180029

- Ohmae M, Kishimoto C, Tomioka N. Complete atrioventricular block in experimental murine myocarditis. *J Electrocardiol.* 2005;38:230–234. doi: 10.1016/j.jelectrocard.2005.02.006
- 429. Błyszczuk P. Myocarditis in Humans and in Experimental Animal Models. Front Cardiovasc Med. 2019;6:64. doi: 10.3389/fcvm.2019.00064
- 430. Tang Q, Huang J, Qian H, Chen L, Wang T, Wang H, Shen D, Wu H, Xiong R. Antiarrhythmic effect of atorvastatin on autoimmune myocarditis is mediated by improving myocardial repolarization. *Life Sci.* 2007;80:601–608. doi: 10.1016/j.lfs.2006.11.045
- 431. Wei SC, Meijers WC, Axelrod ML, Anang NAS, Screever EM, Wescott EC, Johnson DB, Whitley E, Lehmann L, Courand PY, et al. A genetic mouse model recapitulates immune checkpoint inhibitor-associated myocarditis and supports a mechanism-based therapeutic intervention. *Cancer Discov.* 2021;11:614–625. doi: 10.1158/2159-8290.CD-20-0856
- 432. Schnabel RB, Yin X, Gona P, Larson MG, Beiser AS, McManus DD, Newton-Cheh C, Lubitz SA, Magnani JW, Ellinor PT, et al. 50 year trends in atrial fibrillation prevalence, incidence, risk factors, and mortality in the framingham heart study: a cohort study. *Lancet*. 2015;386:154–162. doi: 10.1016/S0140-6736(14)61774-8
- 433. Joseph LC, Subramanyam P, Radlicz C, Trent CM, Iyer V, Colecraft HM, Morrow JP. Mitochondrial oxidative stress during cardiac lipid overload causes intracellular calcium leak and arrhythmia. *Heart Rhythm*. 2016;13:1699–1706. doi: 10.1016/j.hrthm.2016.05.002
- 434. Donnellan E, Wazni OM, Elshazly M, Kanj M, Hussein AA, Baranowski B, Kochar A, Trulock K, Aminian A, Schauer P, et al. Impact of bariatric surgery on atrial fibrillation type. *Circ Arrhythm Electrophysiol*. 2020;13:e007626. doi: 10.1161/CIRCEP.119.007626
- 435. Furman BL. Streptozotocin-induced diabetic models in mice and rats. *Curr Protoc.* 2021;1:e78. doi: 10.1002/cpz1.78
- 436. Grisanti LA. Diabetes and arrhythmias: pathophysiology, mechanisms and therapeutic outcomes. Front Physiol. 2018;9:1669. doi: 10.3389/ fphys.2018.01669
- 437. Shimoni Y, Firek L, Severson D, Giles W. Short-term diabetes alters K+currents in rat ventricular myocytes. Circ Res. 1994;74:620–628. doi: 10.1161/01.res.74.4.620
- 438. Meo M, Meste O, Signore S, Sorrentino A, Cannata A, Zhou Y, Matsuda A, Luciani M, Kannappan R, Goichberg P, et al. Reduction in Kv current enhances the temporal dispersion of the action potential in diabetic myocytes: insights from a novel repolarization algorithm. *J Am Heart Assoc.* 2016;5:e003078. doi: 10.1161/JAHA.115.003078
- 439. Lopez-Izquierdo A, Pereira RO, Wende AR, Punske BB, Abel ED, Tristani-Firouzi M. The absence of insulin signaling in the heart induces changes in potassium channel expression and ventricular repolarization. Am J Physiol Heart Circ Physiol. 2014;306:H747-H754. doi: 10.1152/ajpheart.00849.2013
- 440. Mitašíková M, Lin H, Soukup T, Imanaga I, Tribulová N. Diabetes and thyroid hormones affect connexin-43 and PKC-epsilon expression in rat heart atria. *Physiol Res.* 2009;58:211–217. doi: 10.33549/ physiolres.931425
- 441. Hage C, Michaëlsson E, Linde C, Donal E, Daubert JC, Gan LM, Lund LH. Inflammatory biomarkers predict heart failure severity and prognosis in patients with heart failure with preserved ejection fraction: a holistic proteomic approach. *Circ Cardiovasc Genet.* 2017;10:e001633. doi: 10.1161/CIRCGENETICS.116.001633
- 442. Ferdous Z, Qureshi MA, Jayaprakash P, Parekh K, John A, Oz M, Raza H, Dobrzynski H, Adrian TE, Howarth FC. Different profile of mrna expression in sinoatrial node from streptozotocin-induced diabetic rat. *PLoS One*. 2016;11:e0153934. doi: 10.1371/journal.pone.0153934
- 443. Howarth FC, Parekh K, Jayaprakash P, Inbaraj ES, Oz M, Dobrzynski H, Adrian TE. Altered profile of mRNA expression in atrioventricular node of streptozotocin-induced diabetic rats. *Mol Med Rep.* 2017;16:3720–3730. doi: 10.3892/mmr.2017.7038
- 444. Hegyi B, Borst JM, Bailey LRJ, Shen EY, Lucena AJ, Navedo MF, Bossuyt J, Bers DM. Hyperglycemia regulates cardiac K+ channels via O-GlcNAc-CaMKII and NOX2-ROS-PKC pathways. *Basic Res Cardiol*. 2020;115:71. doi: 10.1007/s00395-020-00834-8
- 445. Hegyi B, Fasoli A, Ko CY, Van BW, Alim CC, Shen EY, Ciccozzi MM, Tapa S, Ripplinger CM, Erickson JR, et al. CaMKII Serine 280 O-GlcNAcylation links diabetic hyperglycemia to proarrhythmia. Circ Res. 2021;129:98–113. doi: 10.1161/CIRCRESAHA.120.318402
- 446. Monnerat G, Alarcón ML, Vasconcellos LR, Hochman-Mendez C, Brasil G, Bassani RA, Casis O, Malan D, Travassos LH, Sepúlveda M, et al. Macrophage-dependent IL-1β production induces cardiac arrhythmias in diabetic mice. Nat Commun. 2016;7:13344. doi: 10.1038/ncomms13344

- 447. Huang H, Amin V, Gurin M, Wan E, Thorp E, Homma S, Morrow JP. Diet-induced obesity causes long QT and reduces transcription of voltage-gated potassium channels. *J Mol Cell Cardiol*. 2013;59:151–158. doi: 10.1016/j.yjmcc.2013.03.007
- 448. McCauley MD, Hong L, Sridhar A, Menon A, Perike S, Zhang M, da Silva IB, Yan J, Bonini MG, Ai X, et al. Ion channel and structural remodeling in obesity-mediated atrial fibrillation. Circ Arrhythm Electrophysiol. 2020;13:e008296. doi: 10.1161/CIRCEP.120.008296
- 449. Ashrafi R, Yon M, Pickavance L, Yanni Gerges J, Davis G, Wilding J, Jian K, Zhang H, Hart G, Boyett M. Altered left ventricular ion channel transcriptome in a high-fat-fed rat model of obesity: insight into obesity-induced arrhythmogenesis. J Obes. 2016;2016;7127898. doi: 10.1155/2016/7127898
- 450. Liu YB, Wu CC, Lu LS, Su MJ, Lin CW, Lin SF, Chen LS, Fishbein MC, Chen PS, Lee YT. Sympathetic nerve sprouting, electrical remodeling, and increased vulnerability to ventricular fibrillation in hypercholesterolemic rabbits. *Circ Res.* 2003;92:1145–1152. doi: 10.1161/01. RES.0000072999.51484.92
- 451. Forman DE, Cittadini A, Azhar G, Douglas PS, Wei JY. Cardiac morphology and function in senescent rats: gender-related differences. J Am Coll Cardiol. 1997;30:1872–1877. doi: 10.1016/s0735-1097(97)00411-7
- 452. Mesquita TRR, Zhang R, de Couto G, Valle J, Sanchez L, Rogers RG, Holm K, Liu W, Marbán E, Cingolani E. Mechanisms of atrial fibrillation in aged rats with heart failure with preserved ejection fraction. *Heart Rhythm.* 2020;17:1025–1033. doi: 10.1016/j.hrthm.2020.02.007
- 453. Hayashi H, Wang C, Miyauchi Y, Omichi C, Pak HN, Zhou S, Ohara T, Mandel WJ, Lin SF, Fishbein MC, et al. Aging-related increase to inducible atrial fibrillation in the rat model. *J Cardiovasc Electrophysiol*. 2002;13:801–808. doi: 10.1046/j.1540-8167.2002.00801.x
- 454. Carlsson L, Almgren O, Duker G. QTU-prolongation and torsades de pointes induced by putative class III antiarrhythmic agents in the rabbit: etiology and interventions. *J Cardiovasc Pharmacol.* 1990;16:276–285. doi: 10.1097/00005344-199008000-00014
- 455. Mazur A, Roden DM, Anderson ME. Systemic administration of calmodulin antagonist W-7 or protein kinase A inhibitor H-8 prevents torsade de pointes in rabbits. *Circulation*. 1999;100:2437–2442. doi: 10.1161/ 01.cir.100.24.2437
- 456. Gbadebo TD, Trimble RW, Khoo MS, Temple J, Roden DM, Anderson ME. Calmodulin inhibitor W-7 unmasks a novel electrocardiographic parameter that predicts initiation of torsade de pointes. *Circulation*. 2002;105:770– 774. doi: 10.1161/hc0602.103724
- 457. Shanmugam M, Molina CE, Gao S, Severac-Bastide R, Fischmeister R, Babu GJ. Decreased sarcolipin protein expression and enhanced sarco(endo) plasmic reticulum Ca2+ uptake in human atrial fibrillation. *Biochem Biophys Res Commun.* 2011;410:97–101. doi: 10.1016/j.bbrc.2011.05.113
- 458. Makarewich CA, Munir AZ, Schiattarella GG, Bezprozvannaya S, Raguimova ON, Cho EE, Vidal AH, Robia SL, Bassel-Duby R, Olson EN. The DWORF micropeptide enhances contractility and prevents heart failure in a mouse model of dilated cardiomyopathy. *Elife*. 2018;7:e38319. doi: 10.7554/eLife.38319
- 459. Wang W, McKinnie SM, Patel VB, Haddad G, Wang Z, Zhabyeyev P, Das SK, Basu R, McLean B, Kandalam V, et al. Loss of Apelin exacerbates myocardial infarction adverse remodeling and ischemia-reperfusion injury: therapeutic potential of synthetic Apelin analogues. *J Am Heart Assoc.* 2013;2:e000249. doi: 10.1161/JAHA.113.000249
- Marks AR. Calcium cycling proteins and heart failure: mechanisms and therapeutics. J Clin Invest. 2013;123:46–52. doi: 10.1172/JCI62834
- 461. Herren AW, Bers DM, Grandi E. Post-translational modifications of the cardiac Na channel: contribution of CaMKII-dependent phosphorylation to acquired arrhythmias. *Am J Physiol Heart Circ Physiol.* 2013;305:H431–H445. doi: 10.1152/ajpheart.00306.2013
- 462. Galleano I, Harms H, Choudhury K, Khoo K, Delemotte L, Pless SA. Functional cross-talk between phosphorylation and disease-causing mutations in the cardiac sodium channel Nav1.5. Proc Natl Acad Sci U S A. 2021;118:e2025320118. doi: 10.1073/pnas.2025320118
- Kamp TJ, Hell JW. Regulation of cardiac L-type calcium channels by protein kinase A and protein kinase C. Circ Res. 2000;87:1095–1102. doi: 10.1161/01.res.87.12.1095
- 464. Hell JW. Beta-adrenergic regulation of the L-type Ca2+ channel Ca(V)1.2 by PKA rekindles excitement. Sci Signal. 2010;3:pe33. doi: 10.1126/scisignal.3141pe33
- 465. Fu Y, Westenbroek RE, Scheuer T, Catterall WA. Phosphorylation sites required for regulation of cardiac calcium channels in the fight-or-flight response. *Proc Natl Acad Sci U S A.* 2013;110:19621–19626. doi: 10.1073/pnas.1319421110

- Park KS, Yang JW, Seikel E, Trimmer JS. Potassium channel phosphorylation in excitable cells: providing dynamic functional variability to a diverse family of ion channels. *Physiology (Bethesda)*. 2008;23:49–57. doi: 10.1152/physiol.00031.2007
- 467. Zhang YH, Hancox JC. Regulation of cardiac Na+-Ca2+ exchanger activity by protein kinase phosphorylation-still a paradox? *Cell Calcium*. 2009;45:1-10. doi: 10.1016/j.ceca.2008.05.005
- 468. Dobrev D, Wehrens XH. Role of RyR2 phosphorylation in heart failure and arrhythmias: controversies around ryanodine receptor phosphorylation in cardiac disease. *Circ Res.* 2014;114:1311–9; discussion 1319. doi: 10.1161/CIRCRESAHA.114.300568
- 469. van Oort RJ, McCauley MD, Dixit SS, Pereira L, Yang Y, Respress JL, Wang Q, De Almeida AC, Skapura DG, Anderson ME, et al. Ryanodine receptor phosphorylation by calcium/calmodulin-dependent protein kinase II promotes life-threatening ventricular arrhythmias in mice with heart failure. Circulation. 2010;122:2669–2679. doi: 10.1161/CIRCULATIONAHA.110.982298
- 470. Campbell HM, Quick AP, Abu-Taha I, Chiang DY, Kramm CF, Word TA, Brandenburg S, Hulsurkar M, Alsina KM, Liu HB, et al. Loss of SPEG inhibitory phosphorylation of ryanodine receptor Type-2 promotes atrial fibrillation. *Circulation*. 2020;142:1159–1172. doi: 10.1161/CIRCULATIONAHA.120.045791
- 471. Trum M, Islam MMT, Lebek S, Baier M, Hegner P, Eaton P, Maier LS, Wagner S. Inhibition of cardiac potassium currents by oxidation-activated protein kinase a contributes to early afterdepolarizations in the heart. Am J Physiol Heart Circ Physiol. 2020;319:H1347-H1357. doi: 10.1152/ajpheart.00182.2020
- 472. Montgomery RL, Davis CA, Potthoff MJ, Haberland M, Fielitz J, Qi X, Hill JA, Richardson JA, Olson EN. Histone deacetylases 1 and 2 redundantly regulate cardiac morphogenesis, growth, and contractility. *Genes Dev.* 2007;21:1790–1802. doi: 10.1101/gad.1563807
- 473. Theis JL, Sharpe KM, Matsumoto ME, Chai HS, Nair AA, Theis JD, de Andrade M, Wieben ED, Michels VV, Olson TM. Homozygosity mapping and exome sequencing reveal GATAD1 mutation in autosomal recessive dilated cardiomyopathy. Circ Cardiovasc Genet. 2011;4:585–594. doi: 10.1161/CIRCGENETICS.111.961052
- 474. Zhang W, Song M, Qu J, Liu GH. Epigenetic Modifications in Cardiovascular Aging and Diseases. *Circ Res.* 2018;123:773–786. doi: 10.1161/CIRCRESAHA.118.312497

- 475. Zhao G, Zhou J, Gao J, Liu Y, Gu S, Zhang X, Su P. Genome-wide DNA methylation analysis in permanent atrial fibrillation. *Mol Med Rep.* 2017;16:5505–5514. doi: 10.3892/mmr.2017.7221
- 476. Shan H, Zhang Y, Lu Y, Zhang Y, Pan Z, Cai B, Wang N, Li X, Feng T, Hong Y, Yang B. Downregulation of miR-133 and miR-590 contributes to nicotine-induced atrial remodelling in canines. *Cardiovasc Res.* 2009;83:465–472. doi: 10.1093/cvr/cvp130
- 477. Lu Y, Zhang Y, Wang N, Pan Z, Gao X, Zhang F, Zhang Y, Shan H, Luo X, Bai Y, et al. MicroRNA-328 contributes to adverse electrical remodeling in atrial fibrillation. *Circulation*. 2010;122:2378–2387. doi: 10.1161/CIRCULATIONAHA.110.958967
- 478. Dawson K, Wakili R, Ordög B, Clauss S, Chen Y, Iwasaki Y, Voigt N, Qi XY, Sinner MF, Dobrev D, et al. MicroRNA29: a mechanistic contributor and potential biomarker in atrial fibrillation. *Circulation*. 2013;127:1466–75, 1475e1. doi: 10.1161/CIRCULATIONAHA.112.001207
- 479. Guzzolino E, Pellegrino M, Ahuja N, Garrity D, D'Aurizio R, Groth M, Baumgart M, Hatcher CJ, Mercatanti A, Evangelista M, et al. miR-182-5p is an evolutionarily conserved Tbx5 effector that impacts cardiac development and electrical activity in zebrafish. *Cell Mol Life Sci.* 2020;77:3215–3229. doi: 10.1007/s00018-019-03343-7
- 480. Yang D, Wan X, Dennis AT, Bektik E, Wang Z, Costa MGS, Fagnen C, Vénien-Bryan C, Xu X, Gratz DH, Hund TJ, et al. MicroRNA biophysically modulates cardiac action potential by direct binding to ion channel. *Circulation*. 2021;143:1597–1613. doi: 10.1161/CIRCULATIONAHA.120.050098
- 481. Ruan Z, Sun X, Sheng H, Zhu L. Long non-coding RNA expression profile in atrial fibrillation. *Int J Clin Exp Pathol.* 2015;8:8402–8410.
- 482. Gore-Panter SR, Hsu J, Barnard J, Moravec CS, Van Wagoner DR, Chung MK, Smith JD. PANCR, the PITX2 adjacent noncoding RNA, is expressed in human left atria and regulates PITX2c expression. Circ Arrhythm Electrophysiol. 2016;9:e003197. doi: 10.1161/CIRCEP.115.003197
- 483. Li Z, Wang X, Wang W, Du J, Wei J, Zhang Y, Wang J, Hou Y. Altered long non-coding RNA expression profile in rabbit atria with atrial fibrillation: TCONS_00075467 modulates atrial electrical remodeling by sponging miR-328 to regulate CACNA1C. *J Mol Cell Cardiol*. 2017;108:73–85. doi: 10.1016/j.yjmcc.2017.05.009
- 484. Wang LY, Shen H, Yang Q, Min J, Wang Q, Xi W, Yin L, Le SG, Zhang YF, Xiao J, et al. LncRNA-LINC00472 contributes to the pathogenesis of atrial fibrillation (Af) by reducing expression of JP2 and RyR2 via miR-24. Biomed Pharmacother. 2019;120:109364. doi: 10.1016/j.biopha.2019.109364