

Depression and coronary heart disease

Robert M. Carney and Kenneth E. Freedland

Abstract | Depression is a highly prevalent risk factor for incident coronary heart disease (CHD) and for cardiovascular morbidity and mortality in patients with established CHD. Several biological and behavioural mechanisms have been hypothesized to underlie the relationship between depression and CHD, but none has been shown to account for more than a small proportion of the risk. Only a few clinical trials have examined whether treating depression decreases the risk of cardiac events in patients with established CHD. None of these trials has shown that treatment results in improved cardiac outcomes, but the differences in depression outcomes between the intervention and control groups have been small and not clinically significant. Nevertheless, secondary analyses of these trials suggest that prognosis improves when depression improves. Concerted efforts to develop more potent interventions for depression, identification of high-risk subtypes of depression, and further research on the biobehavioural mechanisms linking depression to CHD are needed to pave the way for definitive clinical trials.

Sir William Harvey observed over 350 years ago that negative emotions adversely affect the heart¹, but scant scientific evidence was available to support this claim until the 1930s, when two longitudinal studies of psychiatric patients found that depression might be a risk factor for early death, especially death owing to cardiovascular disease^{2,3}. These observations were largely ignored until the late 1980s when interest in the role of depression in coronary heart disease (CHD) surged. Most of the studies during this period focused not on psychiatric patients with depression, but on clinically depressed patients with established CHD^{4–7} or on community-dwelling populations at risk of CHD^{8,9}.

Since then, hundreds of studies from around the world have evaluated depression as a risk factor for incident CHD or for cardiovascular morbidity and mortality in patients with established CHD. Researchers have also investigated the relevance of other psychiatric disorders in relation to heart disease^{10–13}, but these disorders are beyond the scope of this article. In this Review, we consider the evidence for depression as a cardiac risk factor and summarize the research that has been conducted on selected biological and behavioural mechanisms that might link depression to CHD. We also consider whether treatment of depression can prevent cardiac morbidity and mortality in patients with CHD.

Defining and measuring depression

Efforts to develop a biological test for major depression are ongoing¹⁴, but structured clinical interviews are still the only well-validated method for its diagnosis. In some studies, however, cut-off scores on symptom

questionnaires are used to define ‘clinically significant’ depression. Many, but not all, individuals who are classified as having clinically significant depression also meet the standard criteria for major depression, and *vice versa*. Despite efforts to promote specific instruments to assess depression in studies of patients with cardiac disease in order to facilitate the comparison of these studies¹⁵, many different interviews and questionnaires have been used in this area of research. This variation can make comparing studies and interpreting divergent findings difficult.

The criteria for a major depressive episode, as outlined in the latest edition of the Diagnostic and Statistical Manual of Mental Disorders (DSM-5)¹⁶, require the presence of depressed mood or loss of interest or pleasure in daily activities (anhedonia) nearly every day for ≥ 2 weeks. In addition, five or more of the following symptoms must be present nearly every day: depressed or irritable mood, loss of interest, significant change in weight or appetite (increase or decrease), change in sleep (insomnia or hypersomnia), change in activity level (psychomotor agitation or retardation), fatigue, poor concentration, excessive feelings of guilt or worthlessness, and thoughts of death or suicide¹⁶. The symptoms cannot be caused by the direct physiological effects of a drug or a medical condition, and they must cause substantial distress or impairment.

Some of the most widely-used depression questionnaires include symptoms that were omitted from the DSM criteria, such as feelings of hopelessness, helplessness, and social isolation. Both DSM and non-DSM symptoms count when questionnaires are used to identify

Washington University
School of Medicine,
4320 Forest Park Avenue,
Suite 301, St. Louis,
Missouri 63108, USA.

Correspondence to R.M.C.
carneyr@wustl.edu

doi:10.1038/nrcardio.2016.181
Published online 17 Nov 2016

Key points

- Depression is prevalent in the general population, and even more common among patients with coronary heart disease (CHD)
- Depression is a risk factor for incident CHD, and for cardiovascular morbidity and mortality in patients with established CHD
- Potential biological and behavioural mechanisms for this risk have been identified, but no single factor has been shown to account for more than a fraction of the total risk
- Very few clinical trials have been conducted to determine whether treating depression decreases the risk of cardiac events in patients with established CHD
- Each of the trials has substantial shortcomings, and more trials are needed
- Effective treatments for depression, identification of 'high-risk' depression subtypes, and an improved understanding of the biobehavioural pathways linking depression to cardiac morbidity and mortality are needed

clinically significant depression. Evidence exists for a dose–response relationship between depression severity and cardiac risk, such that the presence of even a few mild symptoms carries some risk^{17,18}. However, patients with only a few symptoms of depression might be categorized as 'nondepressed' according to the scoring rules of widely used depression questionnaires. Consequently, artificial categorization of depression severity scores (for example, 'nondepressed' versus 'depressed') might attenuate the strength of the association between depression measures and cardiac outcomes.

Depression screening. A number of depression screening instruments have been recommended for patients with CHD¹⁵. Two of the most popular and easiest to use are the Patient Health Questionnaire (PHQ)-2 and the PHQ-9 (REFS 19,20). The PHQ-2 is a two-item questionnaire that screens for depressed mood and loss of interest — the two cardinal symptoms of major depression. The PHQ-9 includes these two cardinal symptoms along with the other seven DSM-5 criterion symptoms of major depression. In some studies, the PHQ-9 has been used only with those patients with a positive PHQ-2. In other studies, this two-step screening process has been replaced by single use of the PHQ-9. Both forms of the PHQ are sensitive and fairly specific for major depression as well as for clinically significant depression in patients with cardiac disease²¹. However, a diagnostic interview is advised before beginning treatment to confirm a diagnosis of major depression and rule out other psychiatric disorders.

Depression and incident CHD

More than 100 studies and at least six meta-analyses of depression as a risk factor for incident heart disease in individuals without evidence of CHD at baseline have been published. Two limitations have made conducting meta-analyses of these studies difficult. First, the populations have varied widely between studies. Some studies have focused on elderly people, on men or women only, on community residents, ethnic groups, or on specific medical populations (such as patients with hypertension)^{22–27}. Second, many different diagnostic interviews and questionnaires have been used to assess and define depression in these studies.

Another challenge in this area of research is that depression is associated with several other cardiac risk factors, including smoking, diabetes mellitus, sedentary lifestyle, and obesity²⁸. These relationships can make it difficult to determine whether depression is an independent risk factor. Although including these other risk factors as covariates in predictive models is possible, some of them might mediate rather than confound the association between depression and CHD. For example, if depression leads to a sedentary lifestyle, which in turn leads to the early development of atherosclerosis, covariate adjustment for sedentary lifestyle would be inappropriate because this covariate would be on the causal pathway between depression and CHD.

Six meta-analyses of this literature have been published^{29–34} (TABLE 1). Various differences exist between these studies, including the study selection criteria. Nevertheless, all six meta-analyses reported a significant relationship between depression and incident CHD. Five studies reported a large (60% to 80%) increased risk of CHD associated with depression^{29–33}, and one found a more modest (30%) increased risk³⁴.

Some of the studies that were included in these meta-analyses, especially those with small sample sizes, adjusted for only a few potential confounders, and others did not adjust for any. The risks attributable to depression tend to be reduced, but not eliminated, in the covariate-adjusted models compared with the univariate models. Nicholson and colleagues identified 11 studies that reported both unadjusted and adjusted risk estimates for depression³⁰. These investigators found a 12% lower risk of CHD in an adjusted compared with an unadjusted risk estimate, suggesting that some confounding by other cardiac risk factors existed²⁸. Nevertheless, the adjusted risk was still considerable. Similarly, Gan and colleagues found that studies that adjusted for established cardiac risk factors, including smoking, BMI, hypertension, diabetes, physical activity, and socioeconomic status, did not greatly influence the risk estimates for depression that included studies without adjustment for these covariates³⁴. In one of the larger studies of depression and cardiovascular mortality, after adjusting for nearly all the major cardiac risk factors, including age, smoking, sex, systolic blood pressure, BMI, diabetes, social class, heavy alcohol use, and antidepressant medications, Surtees and colleagues found a 2.7-fold increased risk of cardiac death in patients with depression over a median follow-up period of 8 years³⁵. Thus, although depression is associated with other cardiac risk factors, the evidence suggests that depression is an independent risk factor for incident CHD. However, residual confounding must be considered as a possible explanation for at least part of the association, because no study has adjusted for all known potential confounders, and because unknown confounders might exist.

Publication bias can lead to inflated meta-analytical risk estimates. Gan and colleagues conducted a sensitivity analysis by use of the trim-and-fill method to correct for potential publication bias in their meta-analysis, but this adjustment had little effect on the

risk estimates³⁴. No evidence indicates that publication bias explains the documented effect of depression on incident CHD.

Three of the meta-analyses found that studies based on clinical diagnoses of depression reported higher risk estimates than studies based on depression symptom questionnaires^{30–32}. By contrast, one meta-analysis found no difference between the two methods²⁹, and one found that clinical diagnoses were associated with lower risk estimates than symptom questionnaires³⁴. Although patients with major depression tend to have more severe depression than those patients classified as ‘clinically depressed’ on symptom questionnaires, patients with major depression might also be more likely to receive treatment, which can attenuate the risks associated with major depression in some studies. Also, the composition of the ‘nondepressed’ comparison groups might differ between studies of major depression and studies of clinically significant depression. Comparison groups in studies of major depression often include patients with mild or subthreshold depression, which can make detecting an effect of depression more difficult, because the presence of even a few symptoms of depression increases the risk of cardiac events^{17,18}. Nevertheless, the results of these meta-analyses suggest that clinically significant depression, whether defined by questionnaire scores or interview-based diagnoses of major depression, increases the risk of developing CHD.

These meta-analyses were limited by significant heterogeneity of the risk estimates. The estimates have varied between studies as a result of differences in the definition and assessment of depression, the populations, the length of follow-up, and the definition of incident CHD. In addition, depression was assessed only once in most of the studies³², despite the knowledge that depression can follow a variety of course patterns. Depression can remit and never recur, remit and recur one or more times, or become chronic with or without a fluctuating pattern of severity. Consequently, repeated measures would provide more-valid estimates of exposure to depression than a single assessment, and would make it possible to determine how much exposure to depression is needed to increase the risk of developing CHD.

Most of the literature in this area has focused on adults, but evidence also exists that depressed children can be at risk of developing early atherosclerosis. A 2015 scientific statement from the AHA concluded that major depressive disorder and bipolar disorder in children and adolescents are associated with a moderate risk of accelerated atherosclerosis and early cardiovascular disease, independent of traditional CVD risk factors³⁶.

Depression and established CHD

Prevalence of depression in patients with CHD.

Numerous studies have reported the prevalence of major depression or clinically significant depression in patients with established CHD²⁸. In a comprehensive review of the literature, Thombs and colleagues identified eight studies with a total of 10,785 patients with a recent acute myocardial infarction (MI) that used a structured clinical interview to diagnose major depression³⁷. Major depression was present in 19.8% (95% CI 19.1–20.6%) of these patients. To estimate the proportion of patients with previous MI and clinically significant depression symptoms, the investigators analysed studies that used the Beck Depression Inventory (BDI), a widely used questionnaire. The analysis showed that 31.1% (95% CI 29.2–33.0%) of these patients had clinically significant depression (defined as a BDI score >10)³⁷. An estimated 15–18% of patients with medically stable CHD have major depression, and the 1-year prevalence of clinically significant depression is about 30% in these patients²⁸. By comparison, the National Comorbidity Study³⁸ estimated the point prevalence of major depression in the general population to be about 5%.

Depression as a risk factor in established CHD. More than 200 studies have evaluated depression as a risk factor for cardiac events in patients with established CHD. Between 2004 and 2013, five meta-analyses of this literature were published^{30,39–42} (TABLE 2). Three meta-analyses evaluated studies of depression as a risk factor for all-cause or cardiac-related mortality after an acute MI or acute coronary syndrome (ACS)^{39–41}, and two included both post-MI and post-ACS studies as well as studies of patients undergoing revascularization or cardiac catheterization and angiography^{30,42}. All the meta-analyses found that depression was predictive of all-cause mortality, cardiac-related mortality, and/or a combined end point of all-cause mortality and cardiac morbidity.

The largest meta-analysis comparing patients with or without depression after an MI included 29 studies⁴⁰. Depression was associated with a 2.7-fold increased risk of cardiac-related death, a 2.3-fold increased risk of all-cause death, and a 1.6-fold increased risk of cardiovascular events in the 2 years after an acute MI⁴⁰.

An individual patient-level meta-analysis by the same group examined depression severity scores as a continuous variable instead of classifying patients as depressed or nondepressed³⁹. The hazard ratios and odds ratios in this report represent one standard deviation of the mean depression score. For the three studies that included data

Table 1 | Depression as a risk factor for incident CHD

Meta-analysis	Number of studies	Number of participants	Odds ratio or relative risk (95% CI) of CHD
Rugulies (2002) ³¹	11	36,549	1.64 (1.29–2.08)
Cuijpers & Smit (2002) ²⁹	25	106,628	1.81 (1.58–2.07)
Wulsin & Singal (2003) ³³	10	NR	1.64 (1.41–1.90)
Nicholson <i>et al.</i> (2006) ³⁰	21	124,509	1.81 (1.53–2.15)
	11*	NR	• 2.08 (1.69–2.55) unadjusted • 1.90 (1.48–2.42) adjusted
Van der Kooy <i>et al.</i> (2007) ³²	16†	659,991	1.57 (1.36–1.81)
Gan <i>et al.</i> (2014) ³⁴	30	893,850	1.30 (1.22–1.40)

CHD, coronary heart disease. *Studies that included unadjusted and adjusted analyses.

†Includes only those studies of participants without CHD at baseline.

for all the covariates, the analysis found an unadjusted hazard ratio of 1.33 (95% CI 1.23–1.44, $P < 0.001$) per standard deviation, and 1.23 (95% CI 1.15–1.31, $P < 0.001$) after adjusting for age, sex, history of MI, left ventricular ejection fraction, Killip class, smoking, diabetes, and BMI³⁹. For the five studies that did not have time-to-event data, the odds ratio adjusted for age and sex was 1.41 (95% CI 1.34–1.49, $P < 0.001$) per standard deviation³⁹.

As in studies of depression as a risk factor for incident CHD, the potential for residual confounding is a concern in studies of depression in established CHD. Although efforts were made to adjust for major cardiac risk factors in many of these studies, none adjusted for every possible risk factor. Because the risk associated with depression usually decreases after adjustment for potential confounders³⁰, some researchers have speculated that some of the depression symptoms that are present in patients with heart disease might be expressions of severe heart disease rather than of a comorbid depressive disorder. These researchers further speculate that depression predicts worse outcomes, because patients with depression have worse heart disease than patients without depression-like symptoms. This speculation is essentially a reverse causality hypothesis, but the observation that the risks associated with depression continue to be significant after adjustment for multiple indices of CHD severity^{5,43} casts doubt on it. Nevertheless, severe heart disease increasing the risk or severity of comorbid depression does seem plausible. This situation has led some researchers to argue that the relationship between depression and CHD is bidirectional⁴⁴, and this model does seem to fit the findings. However, this hypothesis poses some additional challenges for understanding this common form of comorbidity. In addition to providing careful attention to potential confounding, long-term studies are needed to track the interrelationships over time between CHD, mood, depression, and psychosocial functioning.

Recognizing how difficult it can be to adjust for multiple cardiac risk factors, especially in smaller studies in which statistical overfitting is a concern⁴⁵, two research teams chose instead to use the Global Registry of Acute

Coronary Events (GRACE) score to evaluate the independent effect of depression on recurrent cardiac events after an ACS^{46,47}. The GRACE score was developed to estimate the risk of death in patients with a recent ACS⁴⁸. This score is highly predictive of cardiac outcomes even 5 years after a cardiac event⁴⁹. Adjustment for the GRACE score reduced the estimated risk, but depression was still an independent predictor of all-cause mortality⁴⁶ and of a composite index of fatal and nonfatal cardiac events⁴⁷. The GRACE score, and perhaps other validated prediction models, might offer a useful alternative to adjusting for the large number of risk factors that could be considered as confounders in a study of depression as a risk factor for cardiac mortality after ACS.

A scientific advisory statement from the AHA published in 2014 critically reviewed the evidence for depression as a risk factor for morbidity and mortality after an ACS⁵⁰. A total of 53 studies met the review panel's criteria for study quality. The statement concluded that “despite the heterogeneity of published studies included in this review, the preponderance of evidence supports the recommendation that the American Heart Association should elevate depression to the status of a risk factor for adverse medical outcomes in patients with acute coronary syndrome” (REF. 50). Nevertheless, until the biobehavioural mechanisms by which depression affects the course and outcome of heart disease have been identified, and until treating depression has been shown to improve cardiac outcomes, the question of whether depression is a causal risk factor for cardiac events or merely an epiphenomenon will remain controversial.

Biobehavioural mechanisms

This section provides a brief overview of some of the potential mechanisms that have been proposed to explain the relationship between depression and CHD^{51–54} (FIG. 1). For a more comprehensive evaluation of the literature on these and other potential mechanisms, see the Review published previously in this journal⁵⁵.

Since the 1970s, studies of psychiatric patients with depression have shown evidence of dysregulation of the

Table 2 | Depression as a predictor of morbidity or mortality in patients with CHD

Meta-analysis	Number of studies	Number of participants	Outcomes	Odds ratio or relative risk (95% CI) of CHD	P value
Van Melle <i>et al.</i> (2004) ⁴¹	16	6,367	All-cause mortality	2.38 (1.76–3.22)	<0.00001
			Cardiac mortality	2.59 (1.77–3.77)	<0.00001
			Cardiovascular events	1.95 (1.33–2.85)	<0.0006
Barth <i>et al.</i> (2004) ⁴²	20*	NR	All-cause mortality	2.24 (1.37–3.60)	NR
Nicholson <i>et al.</i> (2006) ³⁰	34	17,842	All-cause mortality	1.80 (1.50–2.15)	<0.00001
Meijer <i>et al.</i> (2011) ⁴⁰	29	16,889	All-cause mortality	2.25 (1.73–2.93)	<0.001
			Cardiac mortality	2.71 (1.68–4.36)	<0.001
			Cardiovascular events	1.59 (1.37–1.85)	<0.001
Meijer <i>et al.</i> (2013) ³⁹	3†	10,175	All-cause mortality	1.33 (1.23–1.44) [§]	<0.001
			Cardiovascular events	1.19 (1.14–1.24)	<0.001

*Includes five studies of patients with documented CHD, but no recent myocardial infarction. †Data from studies included in the 2011 meta-analysis, but combined into one database. §Hazard ratios indicate increased risk per standard deviation above the mean depression score.




Figure 1 | **Potential mechanisms linking depression and coronary heart disease.** Several biological and behavioural mechanisms linking depression and adverse cardiac outcomes have been hypothesized.

autonomic nervous system including higher levels of plasma and urinary catecholamines and higher resting heart rates compared with control individuals without depression^{56–59}. Veith and colleagues showed that elevated circulating levels of plasma catecholamines were caused by systemic increases in sympathetic activity in these patients⁶⁰. Other studies found elevated cortisol levels or that dexamethasone did not suppress the rising morning levels of cortisol^{61,62}. The potential effect of altered autonomic nervous system or hypothalamic–pituitary–adrenal axis functioning on cardiac outcomes in patients with CHD and depression was clear, leading to studies in patients with CHD and depression.

Cardiac autonomic dysfunction. Studies of patients with CHD and depression have found many indications of cardiovascular dysregulation, including elevated 24-h and resting heart rates^{52,63–65}, increased heart-rate responses to physical stressors⁶⁶, reduced heart-rate variability (HRV) and decreased baroreceptor sensitivity^{52,65,67–69}, increased QT variability reflecting abnormal ventricular repolarization⁷⁰, and heart-rate turbulence (that is, abnormal heart rate responses to ventricular arrhythmias)⁷¹. All these indicators have been associated with an elevated mortality and cardiac morbidity, especially in vulnerable populations such as patients with previous MI⁷².

Measures of HRV are widely used to study cardiac autonomic activity in humans⁷³. Beat-to-beat variability in the heart's rhythm is determined primarily by autonomic nervous system modulation of the intrinsic cardiac pacemakers. HRV reflects the balance between sympathetic and parasympathetic regulation of heart rate, with low HRV reflecting inadequate cardiac parasympathetic and/or excessive cardiac sympathetic modulation⁷³. Low HRV is a strong predictor of mortality in patients with a recent MI^{74–77}, and of morbidity and mortality even in patients with stable CHD⁷⁸.

Most studies of patients with CHD have found lower HRV and higher heart rate in those patients with depression compared with those without depression^{52,65–67,79,80}. One exception, however, found no association between HRV and high scores on a depression questionnaire in patients with stable CHD⁸¹. However, a secondary analysis of these data found that somatic (for example, insomnia) but not cognitive (for example, sense of guilt) symptoms of depression were associated with lower HRV⁸². This observation raises the possibility that low HRV might be more common in some subtypes of depression than in others.

In an attempt to determine whether low HRV accounts for the effect of depression on mortality, we studied 311 patients with depression and a recent acute MI who were enrolled in the ENRICH clinical trial⁸³, and 367 patients who met the ENRICH medical inclusion criteria but who were not depressed⁸⁴. Very low frequency power (VLF; 0.0033–0.04 Hz) was selected as the primary index of HRV for this study because of the prognostic importance of this index in patients with previous MI^{74,75}. Although the physiological determinants of HRV are not completely understood, VLF is unaffected by β -blockade, but greatly reduced by atropine, suggesting that the parasympathetic nervous system is an important determinant⁸⁵. VLF was significantly lower in patients with depression compared with patients who were non-depressed⁸⁴. During a median follow-up of 24 months, a total of 47 deaths occurred (6.1% of the cohort). The patients with depression were found to be at higher risk of all-cause death, even after adjusting for potential confounders (HR 2.8, 95% CI 1.4–5.4, $P < 0.003$)⁸⁴. When VLF was entered into the model, the hazard ratio for depression dropped to 2.1 (95% CI 1.1–4.2, $P = 0.03$), indicating that VLF accounted for about one-quarter of the overall mortality. When a measure of the heart's response to premature ventricular contraction (heart-rate turbulence) was added to the model, the adjusted hazard ratio decreased to 1.6 (95% CI 0.8–3.4, $P = 0.18$)⁷¹. Thus, the combination of VLF and heart-rate turbulence explained nearly half of the effect of depression on survival in these patients. These results do not prove that a causal connection between depression, autonomic nervous system dysfunction, and mortality after acute MI exists, but these findings are consistent with this possibility.

Conversely, Whooley and colleagues found that adding VLF to their model did not reduce the effect of depression on cardiac events in a sample of medically stable patients with CHD⁸⁶. Their previous study showed that somatic, but not cognitive, symptoms of depression were associated with low HRV⁸¹, but their subsequent study examined only the relationship between VLF and a depression scale score that counted both cognitive and somatic symptoms.

Inflammation. Inflammatory processes have been associated with the progression of coronary artery disease and with cardiac events, including MI. Depression is associated with elevated levels of inflammatory biomarkers, including proinflammatory cytokines, acute

phase proteins, chemokines, and adhesions molecules⁸⁷. Meta-analyses have found increased levels of C-reactive protein (CRP) and interleukin-6 (IL-6) to be associated with depression⁸⁸. These markers have also been found to be elevated in patients with depression and comorbid CHD^{88–90}. Another meta-analysis showed that levels of tumour necrosis factor are increased in major depression⁹¹. All three of these markers are associated with an increased risk of cardiac events in patients with established CHD^{92–96}. Evidence also indicates that the relationship between inflammation and depression might be bidirectional^{97,98}, so that inflammation might lead to symptoms of depression, and depression might be proinflammatory.

Several studies have investigated the extent to which inflammation mediates the association between depression and the risk of incident CHD. Although most studies found that CRP, IL-6, and soluble intercellular adhesion molecule 1 tend to be elevated in patients with depression, these inflammatory markers explained $\leq 20\%$ of the estimated risk^{99–104}. Thus, although inflammation might contribute to the effect of depression on incident CHD, this mechanism accounts for a fairly small part of the total effect.

Little evidence indicates that inflammation accounts for much of the morbidity and mortality associated with depression in patients with established CHD. Whooley and colleagues found that CRP reduced the effect of depression on cardiac events by 11.3% in their adjusted model in a follow-up of 1,017 outpatients with stable CHD⁸⁶. Two studies of patients with previous MI also found no evidence of higher levels of inflammatory markers in patients with depression compared with those who were nondepressed^{105,106}. These data suggest that inflammation assessed shortly after an acute MI might not explain the effect of depression on cardiac outcomes. However, as more markers of inflammation are studied in patients with depression and CHD, including the activity of anti-inflammatory cytokines, inflammation might turn out to have a more prominent role in the link between depression and cardiac outcomes.

Levels of inflammatory markers are not elevated in everyone with depression^{98,107}. For example, the relationship between inflammation and depression might even depend on race or ethnicity⁹⁸, as well as other factors. Thus, inflammation might have a role in the progression of CHD in some individuals with depression, but not others. Moreover, the way in which depression was assessed in many of these studies might not have been optimal. For example, CRP has been shown to be more strongly associated with persistent than with transient depression¹⁰⁸. However, few studies have included repeated measures of depression. When cases of transient and persistent depression are combined in a single analysis, the former might attenuate the overall association with CRP.

Endothelial dysfunction. Endothelial dysfunction is associated with most of the traditional cardiac risk factors, and can be detected in the earliest preclinical stages

of the atherosclerotic process. Normally, the vascular endothelium produces nitric oxide to maintain vascular tone and inhibit smooth muscle cell growth, leukocyte adhesion, and platelet aggregation. Endothelial dysfunction occurs when endothelial nitric oxide availability is reduced, thereby leaving the atherosclerotic process unopposed¹⁰⁹. Evidence indicates that depression is associated with endothelial dysfunction, even in the absence of other cardiac risk factors^{110–114}.

Endothelial dysfunction can be identified by measuring flow-mediated dilatation (FMD). During FMD testing, endothelium-dependent vasodilatation is evoked by a reactive hyperaemia-induced increase in endothelial shear stress¹¹⁵. In a comprehensive review and meta-analysis of the relationship between depression and FMD, Cooper and colleagues identified 12 studies with a total of 1,491 adults¹¹¹. All studies included in the review were either cross-sectional or retrospective analyses of the correlation between FMD and depressed mood. Across diverse populations including healthy adults and patients with CHD, the meta-analysis found a combined effect size correlation of $r = 0.19$ (95% CI 0.08–0.29, $P = 0.001$)¹¹¹. The relationship was stronger in patients with other cardiac risk factors or comorbidities ($r = 0.29$), and in studies in which maximum vasodilatation was used to quantify FMD ($r = 0.27$)¹¹¹.

Platelet dysfunction. Factors that promote coagulation, platelet activation, and platelet aggregation have a role in ACS. Available evidence, albeit mixed, shows that both psychiatric patients who are depressed but medically well, and patients who are depressed and have CHD, have elevated levels of markers of coagulation and platelet activity, especially β -thromboglobulin and platelet factor 4 (REFS 116–119). To date, these markers have not been evaluated as potential mediators of the relationship between depression and ACS. However, any attempt to measure platelet function and correlate this parameter with clinical outcomes is problematic, because no single measure or marker adequately reflects platelet biology or function¹²⁰. Therefore, although platelet dysfunction might contribute to the relationship between depression and cardiac outcomes, evaluating this pathway might be challenging.

Adherence to medical treatment regimens. Depression is associated with poor adherence to prescribed treatments for chronic medical illnesses¹²¹. In patients with cardiac disease, depression predicts decreased adherence to medication regimens, risk-factor modification interventions, and cardiac rehabilitation^{122–125}. Using electronic medication monitors to obtain objective data on adherence, our group found that patients with CHD and depression were significantly less adherent to a daily regimen of aspirin than patients with CHD without depression¹²². Rieckmann and colleagues reported similar findings in a sample of 172 patients who were studied for 3 months after an ACS¹²⁴. These observations are potentially very important because nonadherence to medications such as aspirin, statins, antiplatelet drugs, and angiotensin-converting-enzyme inhibitors

is associated with cardiac morbidity and mortality in patients with CHD¹²⁶. However, to our knowledge, only one study has examined whether poor adherence mediates the effect of depression on cardiac outcomes. Surprisingly, the study found that self-reported cardiac medication adherence accounted for only about 5% of the effect of the relationship between depression and cardiac events in patients with stable CHD⁸⁶. Whether objective measures of adherence can explain a larger proportion of the effect of depression, or whether adherence mediates more of the effect in patients with recent cardiac events, is currently unknown.

Physical inactivity. Many studies have found that depression is associated with physical inactivity in patients with CHD^{127,128}. Most^{86,129–132}, but not all¹³³, studies on the role of sedentary behaviour as a mediator between depression and cardiac outcomes have found that sedentary behaviour accounts for a substantial proportion of the risk of cardiac events associated with depression. In the Cardiovascular Health Study¹³¹, depression symptoms were assessed annually and self-reported physical activity was assessed at baseline and at 3 and 7 years in a cohort of 5,888 older adults (mean age 72.8 ± 5.6 years). After an average of 10.3 years, both physical activity and depressive symptoms were independent predictors of cardiovascular mortality and were strongly associated with each other. When physical inactivity was added to the model for depression, the risk associated with depression decreased by 26%. The reduction in risk associated with depression was similar for subgroups with (25%) or without (23%) established CHD¹³¹. Most studies have used self-report questionnaires to measure physical activity, even though depression can bias patients' responses. However, Whooley and colleagues found a similar effect when they substituted an objective measure of exercise capacity for a self-report assessment in their model with depression and cardiac events⁸⁶.

Multiple factors model. In 2013, Burg and colleagues argued that linear models of cardiac outcomes, in which one or more risk factors are assumed to act independently, oversimplify the process¹³⁴. As an alternative, the investigators proposed a 'perfect storm' conceptual model in which cardiac events result from a confluence of multiple factors and environmental events that activate the critical pathophysiological processes. In this model, depression might be associated with any or all of the factors described above, but no single factor invariably leads to a cardiac event¹³⁴. This model is very compelling, but underscores the complexity of the processes through which depression can lead to cardiac morbidity and mortality.

Treatment

A number of treatments for depression in patients with CHD have been tested in randomized clinical trials, including antidepressants (such as citalopram, fluoxetine, mirtazapine, and sertraline), cognitive behavioural therapy (CBT), interpersonal psychotherapy, exercise,

and stepped care^{83,135–142}. Most trials have found that the intervention under study was superior to a control condition. Although the effect sizes reported in antidepressant monotherapy trials for patients with depression and CHD have been modest ($d = 0.20–0.38$)¹⁴³, the effect sizes are similar to those reported in published and unpublished trials of antidepressants in psychiatric patients ($d = 0.24–0.35$)^{144,145}. Little evidence indicates that any form of treatment for depression is less effective in patients with CHD than in medically well patients. However, monoamine oxidase inhibitors and tricyclic antidepressants have cardiotoxic adverse effects and are seldom used to treat depression in patients with cardiac disease.

Several large, randomized, controlled trials have examined whether treating depression can improve medical outcomes in patients with CHD^{83,136,140}. The largest of these, the ENRICHD study⁸³, was designed to determine whether treating depression and inadequate social support after acute MI reduced the risk of recurrent infarction and death. A sample of 2,481 patients (1,084 women, 1,397 men) with major or minor depression and/or low perceived social support were randomly assigned to receive either usual care alone or CBT in addition to usual care⁸³. Patients in the intervention group who had severe depression or who did not respond to CBT were also given sertraline. The intervention group showed a significantly greater improvement in depression after 6 months than the usual-care group⁸³, but the difference between the groups was small and not clinically relevant. The other two trials also found modest differences in depression^{136,140}, and none of the trials (including ENRICHD) found a between-group difference in cardiac end points.

A smaller ($n = 157$) study, the COPES trial¹⁴⁶, was designed to determine the acceptability and efficacy of stepped-care depression intervention after ACS. The investigators found a significant difference between treated and control groups in depression outcomes and, despite the small sample size, a trend for a reduced risk of death or hospitalization in the intervention group during the trial period¹⁴⁶. However, this trend was reversed at the 12-month follow-up, resulting in no overall difference between the groups¹⁴⁷.

A meta-analysis of studies on the use of selective serotonin reuptake inhibitor antidepressants to treat depression in patients with CHD reported an overall positive effect on depression, but no difference in mortality or CHD hospitalizations between intervention and control groups¹⁴⁸. However, the study showed modest reductions in hospitalizations and mortality when nonrandomized studies were included in the analysis¹⁴⁸.

The small differences in depression outcomes between treated and control groups, and the small number of end points in the larger trials, have made the detection of an effect of depression treatment on cardiac morbidity or mortality difficult. However, secondary analyses of the three largest trials found that patients whose depression symptoms significantly improved with treatment tended to survive longer than those whose symptoms showed only minimal or

no improvement^{82,149,150}. In the intervention group of ENRICHD¹⁴⁹, for example, a linear relationship was observed between improvement in depression symptoms from baseline to 6 months and survival beyond 6 months, which remained significant after adjustment for all major demographic and medical predictors of mortality (HR 1.05 per point on the BDI, 95% CI 1.01–1.09, $P < 0.003$). Improvement in the intervention group was not related to the presence of medical or logistical problems that might have interfered with participation in treatment.

Secondary analyses of ENRICHD¹⁴⁹ and the other large trials^{150,151} suggest, therefore, that survival might improve if depression improves. Similar findings have been reported in a nonrandomized trial of exercise training and cardiac rehabilitation in patients with depression and previous MI¹⁵², and in a nonrandomized¹⁵³ and a randomized¹⁵⁴ clinical trial of depression interventions for patients with heart failure. Of course, demonstrating a relationship between improvement in depression and survival is not as convincing as showing that an intervention for depression improves survival compared with a control condition, but this evidence might be the best that can be obtained. To our knowledge, no large clinical trials are either ongoing or planned to address the question of whether treating depression can improve survival in patients with CHD.

Future research

Substantial evidence shows that depression is a risk factor for cardiac morbidity and mortality, both in patients without clinical evidence of CHD and in patients with established CHD. Although residual confounding remains a possible explanation for some of these effects, the preponderance of evidence supports an independent effect of depression. The lack of positive findings among the few clinical trials that examined the effect of treating depression on cardiac outcomes, however, has led some researchers to conclude that depression might not be an appropriate therapeutic target for these patients¹⁵⁵.

Compared with most trials in cardiology, the sample sizes in the existing clinical trials of treating depression in patients with CHD have been very small. As cardiovascular mortality continues to decline^{156,157}, detecting a survival benefit for any treatment becomes increasingly difficult. Unfortunately, depression treatments that are deemed safe in patients with CHD are only modestly effective. Ongoing research is directed towards developing more effective treatments for depression in these patients. However, additional research on factors that mediate the effect of depression on cardiac outcomes, and on the identification of high-risk subtypes or symptoms of depression, might help to provide better targets for intervention.

As discussed earlier, depression has been associated with plausible biological pathways, including elevated inflammation and altered autonomic nervous system functioning, and with behavioural pathways, including poor adherence to diet and medication treatments and a sedentary lifestyle. However, none of these

mediators has accounted for more than a small proportion of the associated risk in mediation studies. Few studies of potential mechanisms have simultaneously evaluated multiple candidates. Consequently, little is known about the relationships between putative mediators. These mediators might be additive or synergistic, or they might represent different expressions of a common aetiology (such as hypothalamic–pituitary–adrenal axis dysfunction). Some putative mechanisms, such as increased inflammation, might be largely the result of other risk factors associated with depression, such as obesity, physical inactivity, or smoking^{158–160}. Furthermore, not all patients with depression have low HRV, elevated levels of inflammatory markers, or any of the other possible mediators considered in this Review. Distinct pathways might mediate cardiac risk in different subgroups of patients with depression. Indeed, perhaps some, but not all, cases of depression are associated with elevated cardiac risk¹⁶¹. These questions should be explored in future studies.

Behavioural risk factors, especially sedentary behaviour and poor adherence to medical treatment regimens, might be targeted in addition to depression. Medications commonly administered to patients with cardiac disease might improve many of the putative biological mediators, but only if patients adhere to the treatment regimen. Methods to improve adherence in these patients have been studied, but research is needed to determine how best to improve adherence in patients with CHD and depression.

In 2012, we concluded that the research to identify depression subtypes or specific symptoms that confer a particularly high risk of cardiac events was promising, but methodological problems and contradictory findings made it premature to draw any definite conclusions^{161,162}. Since then, studies have continued to find that the presence of somatic symptoms of depression, such as fatigue or problems sleeping, are better predictors of cardiac risk than cognitive symptoms, such as feelings of guilt or thoughts of death¹⁶³. Other studies have found that fatigue and sleep problems are among the most common symptoms of depression before treatment, and two of the most likely symptoms to remain in patients meeting the standard criteria for remission^{164–167}. In a 3-year study of primary-care patients, for example, loss of energy and sleep problems were reported in 90% and 85% of patients, respectively, before treatment¹⁶⁸. These symptoms continued to be present in 35% and 39% of the patients, respectively, after remission of depression, whereas the two core symptoms of depression — sadness and loss of interest — were present in only 21% of the patients¹⁶⁸. If somatic symptoms are more strongly associated with cardiac morbidity and mortality, these findings might help to explain the difficulty of improving survival even with ‘successful’ treatment of depression.

Research is needed to determine how best to target somatic symptoms of depression. Although little evidence exists that any traditional treatment has a greater effect on a specific symptom or group of depression symptoms^{169–172}, this area has not been well studied in

patients with CHD and depression. Exercise, which has been shown to reduce fatigue, improve sleep, and to be modestly effective in improving depression^{173–176}, might be considered as a primary intervention or as an augmentation strategy when these somatic symptoms remain after other depression treatments. Aerobic exercise can have the additional benefits of lowering heart rate, increasing HRV^{177–179}, and increasing anti-inflammatory and decreasing proinflammatory cytokine levels^{180–183}. CBT for disordered sleep might also be considered as an adjunctive therapy when sleep problems do not improve after standard depression treatments^{184–186}. Studies testing these interventions and others to address somatic symptoms at baseline or after traditional treatments are needed.

Conclusions

Since the late 1980s, hundreds of studies have been conducted on depression as a risk factor for the onset or progression of cardiovascular disease, together with studies of the biobehavioural pathways that might explain this risk, and several large treatment trials. Sir William Harvey would probably have been amazed if he had known how much work would eventually be devoted to this area, and perhaps at how long it took after his initial observation before this work began. How depression increases the risk of cardiac events, and whether successful treatment of depression can reduce this risk, are questions that remain to be answered. Further research will, we hope, give us definitive answers to these questions in much less time than it took to confirm Harvey's initial observation.

- Willis, R. *The Works of William Harvey* (Univ. of Pennsylvania Press, 1989).
- Fuller, R. G. What happens to mental patients after discharge from the hospital? *Psychiatr. Quart.* **9**, 95–104 (1935).
- Malzberg, B. Mortality among patients with involution melancholia. *Am. J. Psychiatry* **93**, 1231–1238 (1937).
- Carney, R. M. *et al.* Major depressive disorder predicts cardiac events in patients with coronary artery disease. *Psychosom. Med.* **50**, 627–633 (1988).
- Frasere-Smith, N., Lesperance, F. & Talajic, M. Depression following myocardial infarction: impact on 6-month survival. *JAMA* **270**, 1819–1825 (1993).
- Ladwig, K. H., Kieser, M., Konig, J., Breithardt, G. & Borggrefe, M. Affective disorders and survival after acute myocardial infarction. Results from the post-infarction late potential study. *Eur. Heart J.* **12**, 959–964 (1991).
- Silverstone, P. H. Depression and outcome in acute myocardial infarction. *Br. Med. J. (Clin. Res. Ed.)* **294**, 219–220 (1987).
- Bruce, M. L., Leaf, P. J., Rozal, G. P., Florio, L. & Hoff, R. A. Psychiatric status and 9-year mortality data in the New Haven Epidemiologic Catchment Area Study. *Am. J. Psychiatry* **151**, 716–721 (1994).
- Murphy, J. M., Monson, R. R., Olivier, D. C., Sobol, A. M. & Leighton, A. H. Affective disorders and mortality: a general population study. *Arch. Gen. Psychiatry* **44**, 473–480 (1987).
- Gale, C. R., Batty, G. D., Osborn, D. P., Tynelius, P. & Rasmussen, F. Mental disorders across the adult life course and future coronary heart disease: evidence for general susceptibility. *Circulation* **129**, 186–193 (2014).
- Roest, A. M., Martens, E. J., de Jonge, P. & Denollet, J. Anxiety and risk of incident coronary heart disease: a meta-analysis. *J. Am. Coll. Cardiol.* **56**, 38–46 (2010).
- Sowden, G. L. & Huffman, J. C. The impact of mental illness on cardiac outcomes: a review for the cardiologist. *Int. J. Cardiol.* **132**, 30–37 (2009).
- Vaccarino, V. *et al.* Post-traumatic stress disorder and incidence of coronary heart disease: a twin study. *J. Am. Coll. Cardiol.* **62**, 970–978 (2013).
- Bilello, J. A. *et al.* MDDScore: confirmation of a blood test to aid in the diagnosis of major depressive disorder. *J. Clin. Psychiatry* **76**, e199–e206 (2015).
- Davidson, K. W. *et al.* Assessment and treatment of depression in patients with cardiovascular disease: National Heart, Lung, and Blood Institute Working Group Report. *Psychosom. Med.* **68**, 645–650 (2006).
- American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders (DSM-5)* 5th edn (American Psychiatric Association, 2013).
- Bush, D. E. *et al.* Even minimal symptoms of depression increase mortality risk after acute myocardial infarction. *Am. J. Cardiol.* **88**, 337–341 (2001).
- Lesperance, F., Frasure-Smith, N., Talajic, M. & Bourassa, M. G. Five-year risk of cardiac mortality in relation to initial severity and one-year changes in depression symptoms after myocardial infarction. *Circulation* **105**, 1049–1053 (2002).
- Kroenke, K., Spitzer, R. L. & Williams, J. B. The PHQ-9: validity of a brief depression severity measure. *J. Gen. Intern. Med.* **16**, 606–613 (2001).
- Kroenke, K., Spitzer, R. L. & Williams, J. B. The Patient Health Questionnaire-2: validity of a two-item depression screener. *Med. Care* **41**, 1284–1292 (2003).
- Thombs, B. D., Ziegelstein, R. C. & Whooley, M. A. Optimizing detection of major depression among patients with coronary artery disease using the patient health questionnaire: data from the Heart and Soul Study. *J. Gen. Intern. Med.* **23**, 2014–2017 (2008).
- O'Brien, E. C. *et al.* Depressive symptoms and risk of cardiovascular events in blacks: findings from the Jackson Heart Study. *Circ. Cardiovasc. Qual. Outcomes* **8**, 552–559 (2015).
- O'Neil, A. *et al.* Depression is a risk factor for incident coronary heart disease in women: an 18-year longitudinal study. *J. Affect. Disord.* **196**, 117–124 (2016).
- Penninx, B. W. *et al.* Cardiovascular events and mortality in newly and chronically depressed persons > 70 years of age. *Am. J. Cardiol.* **81**, 988–994 (1998).
- Pequignot, R. *et al.* High level of depressive symptoms at repeated study visits and risk of coronary heart disease and stroke over 10 years in older adults: the Three-City Study. *J. Am. Geriatr. Soc.* **64**, 118–125 (2016).
- Schulz, R. *et al.* Association between depression and mortality in older adults: the Cardiovascular Health Study. *Arch. Intern. Med.* **160**, 1761–1768 (2000).
- Wassertheil-Smoller, S. *et al.* Depression and cardiovascular sequelae in postmenopausal women: the Women's Health Initiative (WHI). *Arch. Intern. Med.* **164**, 289–298 (2004).
- Carney, R. M. & Freedland, K. E. Depression in patients with coronary heart disease. *Am. J. Med.* **121**, S20–S27 (2008).
- Cuijpers, P. & Smit, F. Excess mortality in depression: a meta-analysis of community studies. *J. Affect. Disord.* **72**, 227–236 (2002).
- Nicholson, A., Kuper, H. & Hemingway, H. Depression as an aetiological and prognostic factor in coronary heart disease: a meta-analysis of 6,362 events among 146,538 participants in 54 observational studies. *Eur. Heart J.* **27**, 2763–2774 (2006).
- Rugulies, R. Depression as a predictor for coronary heart disease: a review and meta-analysis. *Am. J. Prev. Med.* **23**, 51–61 (2002).
- Van der Kooy, K. *et al.* Depression and the risk for cardiovascular diseases: systematic review and meta analysis. *Int. J. Geriatr. Psychiatry* **22**, 613–626 (2007).
- Wulsin, L. R. & Singal, B. M. Do depressive symptoms increase the risk for the onset of coronary disease? A systematic quantitative review. *Psychosom. Med.* **65**, 201–210 (2003).
- Gan, Y. *et al.* Depression and the risk of coronary heart disease: a meta-analysis of prospective cohort studies. *BMC Psychiatry* **14**, 371 (2014).
- Surtees, P. G. *et al.* Depression and ischemic heart disease mortality: evidence from the EPIC-Norfolk United Kingdom prospective cohort study. *Am. J. Psychiatry* **165**, 515–523 (2008).
- Goldstein, B. I. *et al.* Major depressive disorder and bipolar disorder predispose youth to accelerated atherosclerosis and early cardiovascular disease: a scientific statement from the American Heart Association. *Circulation* **132**, 965–986 (2015).
- Thombs, B. D. *et al.* Prevalence of depression in survivors of acute myocardial infarction. *J. Gen. Intern. Med.* **21**, 30–38 (2006).
- Blazer, D. G., Kessler, R. C., McGonagle, K. A. & Swartz, M. S. The prevalence and distribution of major depression in a national community sample: the National Comorbidity Survey. *Am. J. Psychiatry* **151**, 979–986 (1994).
- Meijer, A. *et al.* Adjusted prognostic association of depression following myocardial infarction with mortality and cardiovascular events: individual patient data meta-analysis. *Br. J. Psychiatry* **203**, 90–102 (2013).
- Meijer, A. *et al.* Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis of 25 years of research. *Gen. Hosp. Psychiatry* **33**, 203–216 (2011).
- van Melle, J. P. *et al.* Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis. *Psychosom. Med.* **66**, 814–822 (2004).
- Barth, J., Schumacher, M. & Herrmann-Lingen, C. Depression as a risk factor for mortality in patients with coronary heart disease: a meta-analysis. *Psychosom. Med.* **66**, 802–813 (2004).
- Carney, R. M. *et al.* Depression as a risk factor for mortality after acute myocardial infarction. *Am. J. Cardiol.* **92**, 1277–1281 (2003).
- de Jonge, P. & Roest, A. M. Depression and cardiovascular disease: the end of simple models. *Br. J. Psychiatry* **201**, 337–338 (2012).
- Babiyak, M. A. What you see may not be what you get: a brief, nontechnical introduction to overfitting in regression-type models. *Psychosom. Med.* **66**, 411–421 (2004).
- Kronish, I. M., Rieckmann, N., Schwartz, J. E., Schwartz, D. R. & Davidson, K. W. Is depression after an acute coronary syndrome simply a marker of known prognostic factors for mortality? *Psychosom. Med.* **71**, 697–703 (2009).
- Meurs, M., Zuidersma, M., Dickens, C. & de Jonge, P. Examining the relation between post myocardial infarction depression and cardiovascular prognosis using a validated prediction model for post myocardial mortality. *Int. J. Cardiol.* **167**, 2533–2538 (2013).
- Eagle, K. A. *et al.* A validated prediction model for all forms of acute coronary syndrome: estimating the risk of 6-month postdischarge death in an international registry. *JAMA* **291**, 2727–2733 (2004).
- Fox, K. A. *et al.* The Global Registry of Acute Coronary Events, 1999 to 2009—GRACE. *Heart* **96**, 1095–1101 (2010).
- Lichtman, J. H. *et al.* Depression as a risk factor for poor prognosis among patients with acute coronary syndrome: systematic review and recommendations: a scientific statement from the American Heart Association. *Circulation* **129**, 1350–1369 (2014).
- Carney, R. M., Freedland, K. E., Miller, G. E. & Jaffe, A. S. Depression as a risk factor for cardiac mortality and morbidity: a review of potential mechanisms. *J. Psychosom. Res.* **53**, 897–902 (2002).
- Carney, R. M., Freedland, K. E., Rich, M. W. & Jaffe, A. S. Depression as a risk factor for cardiac events in established coronary heart disease: a review of possible mechanisms. *Ann. Behav. Med.* **17**, 142–149 (1995).

53. Lett, H. S. *et al.* Depression as a risk factor for coronary artery disease: evidence, mechanisms, and treatment. *Psychosom. Med.* **66**, 305–315 (2004).
54. Skala, J. A., Freedland, K. E. & Carney, R. M. Coronary heart disease and depression: a review of recent mechanistic research. *Can. J. Psychiatry* **51**, 738–745 (2006).
55. Nemeroff, C. B. & Goldschmidt-Clermont, P. J. Heartache and heartbreak — the link between depression and cardiovascular disease. *Nat. Rev. Cardiol.* **9**, 526–539 (2012).
56. Esler, M. *et al.* The peripheral kinetics of norepinephrine in depressive illness. *Arch. Gen. Psychiatry* **39**, 295–300 (1982).
57. Lake, C. R. *et al.* High plasma norepinephrine levels in patients with major affective disorder. *Am. J. Psychiatry* **139**, 1315–1318 (1982).
58. Roy, A., Pickar, D., De Jong, J., Karoum, F. & Linnoila, M. Norepinephrine and its metabolites in cerebrospinal fluid, plasma, and urine: relationship to hypothalamic–pituitary–adrenal axis function in depression. *Arch. Gen. Psychiatry* **45**, 849–857 (1988).
59. Siever, L. J. & Davis, K. L. Overview: toward a dysregulation hypothesis of depression. *Am. J. Psychiatry* **142**, 1017–1031 (1985).
60. Veith, R. C. *et al.* Sympathetic nervous system activity in major depression: basal and desipramine-induced alterations in plasma norepinephrine kinetics. *Arch. Gen. Psychiatry* **51**, 411–422 (1994).
61. Charlton, B. G., Leake, A., Wright, C., Griffiths, H. W. & Ferrier, I. N. A combined study of cortisol, ACTH and dexamethasone concentrations in major depression. Multiple time-point sampling. *Br. J. Psychiatry* **150**, 791–796 (1987).
62. Ribeiro, S. C., Tandon, R., Grunhaus, L. & Greden, J. F. The DST as a predictor of outcome in depression: a meta-analysis. *Am. J. Psychiatry* **150**, 1618–1629 (1993).
63. Carney, R. M. *et al.* Change in heart rate and heart rate variability during treatment for depression in patients with coronary heart disease. *Psychosom. Med.* **62**, 639–647 (2000).
64. Carney, R. M. *et al.* The relationship between heart rate, heart rate variability and depression in patients with coronary artery disease. *J. Psychosom. Res.* **32**, 159–164 (1988).
65. Stein, P. K. *et al.* Severe depression is associated with markedly reduced heart rate variability in patients with stable coronary heart disease. *J. Psychosom. Res.* **48**, 493–500 (2000).
66. Carney, R. M. *et al.* Major depression, heart rate, and plasma norepinephrine in patients with coronary heart disease. *Biol. Psychiatry* **45**, 458–463 (1999).
67. Carney, R. M. *et al.* Depression, heart rate variability, and acute myocardial infarction. *Circulation* **104**, 2024–2028 (2001).
68. Pitzalis, M. V. *et al.* Depression but not anxiety influences the autonomic control of heart rate after myocardial infarction. *Am. Heart J.* **141**, 765–771 (2001).
69. Watkins, L. L. & Grossman, P. Association of depressive symptoms with reduced baroreflex cardiac control in coronary artery disease. *Am. Heart J.* **137**, 453–457 (1999).
70. Carney, R. M. *et al.* Effects of depression on QT interval variability after myocardial infarction. *Psychosom. Med.* **65**, 177–180 (2003).
71. Carney, R. M. *et al.* Heart rate turbulence, depression, and survival after acute myocardial infarction. *Psychosom. Med.* **69**, 4–9 (2007).
72. Carney, R. M., Freedland, K. E. & Veith, R. C. Depression, the autonomic nervous system, and coronary heart disease. *Psychosom. Med.* **67** (Suppl. 1), S29–S33 (2005).
73. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. Heart rate variability: standards of measurement, physiological interpretation and clinical use. *Circulation* **93**, 1043–1065 (1996).
74. Bigger, J. T. Jr, Fleiss, J. L., Rolnitzky, L. M. & Steinman, R. C. Frequency domain measures of heart period variability to assess risk late after myocardial infarction. *J. Am. Coll. Cardiol.* **21**, 729–736 (1993).
75. Bigger, J. T. Jr *et al.* Frequency domain measures of heart period variability and mortality after myocardial infarction. *Circulation* **85**, 164–171 (1992).
76. Kleiger, R. E., Miller, J. P., Bigger, J. T. Jr & Moss, A. J. Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *Am. J. Cardiol.* **59**, 256–262 (1987).
77. Vaishnav, S. *et al.* Relation between heart rate variability early after acute myocardial infarction and long-term mortality. *Am. J. Cardiol.* **73**, 653–657 (1994).
78. Rich, M. W. *et al.* Correlation of heart rate variability with clinical and angiographic variables and late mortality after coronary angiography. *Am. J. Cardiol.* **62**, 714–717 (1988).
79. Kang, H. J. *et al.* Predictors of depressive disorder following acute coronary syndrome: results from K-DEPACS and EsDEPACS. *J. Affect. Disord.* **181**, 1–8 (2015).
80. Krittayaphong, R. *et al.* Heart rate variability in patients with coronary artery disease: differences in patients with higher and lower depression scores. *Psychosom. Med.* **59**, 231–235 (1997).
81. Gehi, A., Mangano, D., Pipkin, S., Browner, W. S. & Whooley, M. A. Depression and heart rate variability in patients with stable coronary heart disease: findings from the Heart and Soul Study. *Arch. Gen. Psychiatry* **62**, 661–666 (2005).
82. de Jonge, P., Mangano, D. & Whooley, M. A. Differential association of cognitive and somatic depressive symptoms with heart rate variability in patients with stable coronary heart disease: findings from the Heart and Soul Study. *Psychosom. Med.* **69**, 735–739 (2007).
83. Berkman, L. F. *et al.* Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICH) randomized trial. *JAMA* **289**, 3106–3116 (2003).
84. Carney, R. M. *et al.* Low heart rate variability and the effect of depression on post-myocardial infarction mortality. *Arch. Intern. Med.* **165**, 1486–1491 (2005).
85. Taylor, J. A., Carr, D. L., Myers, C. W. & Eckberg, D. L. Mechanisms underlying very-low-frequency RR-interval oscillations in humans. *Circulation* **98**, 547–555 (1998).
86. Whooley, M. A. *et al.* Depressive symptoms, health behaviors, and risk of cardiovascular events in patients with coronary heart disease. *JAMA* **300**, 2379–2388 (2008).
87. Miller, A. H., Maletic, V. & Raison, C. L. Inflammation and its discontents: the role of cytokines in the pathophysiology of major depression. *Biol. Psychiatry* **65**, 732–741 (2009).
88. Howren, M. B., Lamkin, D. M. & Suls, J. Associations of depression with C-reactive protein, IL-1, and IL-6: a meta-analysis. *Psychosom. Med.* **71**, 171–186 (2009).
89. Miller, G. E., Freedland, K. E., Carney, R. M., Stetler, C. A. & Banks, W. A. Cynical hostility, depressive symptoms, and the expression of inflammatory risk markers for coronary heart disease. *J. Behav. Med.* **26**, 501–515 (2003).
90. Miller, G. E., Freedland, K. E., Duntley, S. & Carney, R. M. Relation of depressive symptoms to C-reactive protein and pathogen burden (cytomegalovirus, herpes simplex virus, Epstein-Barr virus) in patients with earlier acute coronary syndromes. *Am. J. Cardiol.* **95**, 317–321 (2005).
91. Dowlati, Y. *et al.* A meta-analysis of cytokines in major depression. *Biol. Psychiatry* **67**, 446–457 (2010).
92. Danesh, J., Collins, R., Appleby, P. & Peto, R. Association of fibrinogen, C-reactive protein, albumin, or leukocyte count with coronary heart disease: meta-analyses of prospective studies. *JAMA* **279**, 1477–1482 (1998).
93. Ridker, P. M., Glynn, R. J. & Hennekens, C. H. C-reactive protein adds to the predictive value of total and HDL cholesterol in determining risk of first myocardial infarction. *Circulation* **97**, 2007–2011 (1998).
94. Ridker, P. M., Hennekens, C. H., Buring, J. E. & Rifai, N. C-reactive protein and other markers of inflammation in the prediction of cardiovascular disease in women. *N. Engl. J. Med.* **342**, 836–843 (2000).
95. Ridker, P. M. *et al.* Elevation of tumor necrosis factor- α and increased risk of recurrent coronary events after myocardial infarction. *Circulation* **101**, 2149–2153 (2000).
96. Ridker, P. M., Rifai, N., Stampfer, M. J. & Hennekens, C. H. Plasma concentration of interleukin-6 and the risk of future myocardial infarction among apparently healthy men. *Circulation* **101**, 1767–1772 (2000).
97. Gimeno, D. *et al.* Associations of C-reactive protein and interleukin-6 with cognitive symptoms of depression: 12-year follow-up of the Whitehall II study. *Psychol. Med.* **39**, 413–423 (2009).
98. Stewart, J. C. One effect size does not fit all — is the depression-inflammation link missing in racial/ethnic minority individuals? *JAMA Psychiatry* **73**, 301–302 (2016).
99. Davidson, K. W. *et al.* Relation of inflammation to depression and incident coronary heart disease (from the Canadian Nova Scotia Health Survey [NSHS95] Prospective Population Study). *Am. J. Cardiol.* **103**, 755–761 (2009).
100. Empana, J. P. *et al.* Contributions of depressive mood and circulating inflammatory markers to coronary heart disease in healthy European men: the Prospective Epidemiological Study of Myocardial Infarction (PRIME). *Circulation* **111**, 2299–2305 (2005).
101. Hiles, S. A. *et al.* The role of inflammatory markers in explaining the association between depression and cardiovascular hospitalisations. *J. Behav. Med.* **38**, 609–619 (2015).
102. Kop, W. J. *et al.* Autonomic nervous system dysfunction and inflammation contribute to the increased cardiovascular mortality risk associated with depression. *Psychosom. Med.* **72**, 626–635 (2010).
103. Vaccarino, V. *et al.* Depression, inflammation, and incident cardiovascular disease in women with suspected coronary ischemia: the National Heart, Lung, and Blood Institute-sponsored WISE study. *J. Am. Coll. Cardiol.* **50**, 2044–2050 (2007).
104. Surtees, P. G. *et al.* Major depression, C-reactive protein, and incident ischemic heart disease in healthy men and women. *Psychosom. Med.* **70**, 850–855 (2008).
105. Schins, A. *et al.* Inflammatory markers in depressed post-myocardial infarction patients. *J. Psychiatr. Res.* **39**, 137–144 (2005).
106. Smolderen, K. G. *et al.* Association of somatic and cognitive depressive symptoms and biomarkers in acute myocardial infarction: insights from the translational research investigating underlying disparities in acute myocardial infarction patients' health status registry. *Biol. Psychiatry* **71**, 22–29 (2012).
107. Glassman, A. H. & Miller, G. E. Where there is depression, there is inflammation... sometimes! *Biol. Psychiatry* **62**, 280–281 (2007).
108. Shimbo, D., Rieckmann, N., Paulino, R. & Davidson, K. W. Relation between C reactive protein and depression remission status in patients presenting with acute coronary syndrome. *Heart* **92**, 1316–1318 (2006).
109. Shimokawa, H. Primary endothelial dysfunction: atherosclerosis. *J. Mol. Cell. Cardiol.* **31**, 23–37 (1999).
110. Broadley, A. J., Korszun, A., Jones, C. J. & Frenneaux, M. P. Arterial endothelial function is impaired in treated depression. *Heart* **88**, 521–523 (2002).
111. Cooper, D. C. *et al.* Depressed mood and flow-mediated dilation: a systematic review and meta-analysis. *Psychosom. Med.* **73**, 360–369 (2011).
112. Rajagopalan, S. *et al.* Abnormal brachial artery flow-mediated vasodilation in young adults with major depression. *Am. J. Cardiol.* **88**, 196–198 (2001).
113. Rybakowski, J. K., Wykretowicz, A., Heymann-Szlachcinska, A. & Wysocki, H. Impairment of endothelial function in unipolar and bipolar depression. *Biol. Psychiatry* **60**, 889–891 (2006).
114. Sherwood, A., Hinderliter, A. L., Watkins, L. L., Waugh, R. A. & Blumenthal, J. A. Impaired endothelial function in coronary heart disease patients with depressive symptomatology. *J. Am. Coll. Cardiol.* **46**, 656–659 (2005).
115. Corretti, M. C. *et al.* Guidelines for the ultrasound assessment of endothelial-dependent flow-mediated vasodilation of the brachial artery: a report of the International Brachial Artery Reactivity Task Force. *J. Am. Coll. Cardiol.* **39**, 257–265 (2002).
116. Kuijpers, P. M., Hamulyak, K., Strik, J. J., Wellens, H. J. & Honig, A. Beta-thromboglobulin and platelet factor 4 levels in post-myocardial infarction patients with major depression. *Psychiatry Res.* **109**, 207–210 (2002).
117. Serebruany, V. L. *et al.* Enhanced platelet/endothelial activation in depressed patients with acute coronary syndromes: evidence from recent clinical trials. *Blood Coagul. Fibrinolysis* **14**, 563–567 (2003).
118. Shimbo, D. *et al.* Exaggerated serotonin-mediated platelet reactivity as a possible link in depression and acute coronary syndromes. *Am. J. Cardiol.* **89**, 331–333 (2002).
119. von Kanel, R. Platelet hyperactivity in clinical depression and the beneficial effect of antidepressant drug treatment: how strong is the evidence? *Acta Psychiatr. Scand.* **110**, 163–177 (2004).
120. Gurbel, P. A., Becker, R. C., Mann, K. G., Steinhilber, S. R. & Michelson, A. D. Platelet function monitoring in patients with coronary artery disease. *J. Am. Coll. Cardiol.* **50**, 1822–1834 (2007).
121. DiMatteo, M. R., Lepper, H. S. & Croghan, T. W. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch. Intern. Med.* **160**, 2101–2107 (2000).

122. Carney, R. M., Freedland, K. E., Eisen, S. A., Rich, M. W. & Jaffe, A. S. Major depression and medication adherence in elderly patients with coronary artery disease. *Health Psychol.* **14**, 88–90 (1995).
123. Gehl, A., Haas, D., Pipkin, S. & Whoolley, M. A. Depression and medication adherence in outpatients with coronary heart disease: findings from the Heart and Soul Study. *Arch. Intern. Med.* **165**, 2508–2513 (2005).
124. Rieckmann, N. *et al.* Course of depressive symptoms and medication adherence after acute coronary syndromes: an electronic medication monitoring study. *J. Am. Coll. Cardiol.* **48**, 2218–2222 (2006).
125. Ziegelstein, R. C. *et al.* Patients with depression are less likely to follow recommendations to reduce cardiac risk during recovery from a myocardial infarction. *Arch. Intern. Med.* **160**, 1818–1823 (2000).
126. Kumbhani, D. J. *et al.* Adherence to secondary prevention medications and four-year outcomes in outpatients with atherosclerosis. *Am. J. Med.* **126**, 693–700.e1 (2013).
127. Boschloo, L., Rieuwijk, K. G., Schoevers, R. A. & Penninx, B. W. J. H. The impact of lifestyle factors on the 2-year course of depressive and/or anxiety disorders. *J. Affect. Disord.* **159**, 73–79 (2014).
128. Roshanaei-Moghaddam, B., Katon, W. J. & Russo, J. The longitudinal effects of depression on physical activity. *Gen. Hosp. Psychiatry* **31**, 306–315 (2009).
129. Brummett, B. H. *et al.* Effect of smoking and sedentary behavior on the association between depressive symptoms and mortality from coronary heart disease. *Am. J. Cardiol.* **92**, 529–532 (2003).
130. Kurdyak, P. A., Chong, A., Gnam, W. H., Goering, P. & Alter, D. A. Depression and self-reported functional status: impact on mortality following acute myocardial infarction. *J. Eval. Clin. Pract.* **17**, 444–451 (2011).
131. Win, S. *et al.* Depressive symptoms, physical inactivity and risk of cardiovascular mortality in older adults: the Cardiovascular Health Study. *Heart* **97**, 500–505 (2011).
132. Larsen, K. K., Christensen, B., Sondergaard, J. & Vestergaard, M. Depressive symptoms and risk of new cardiovascular events or death in patients with myocardial infarction: a population-based longitudinal study examining health behaviors and health care interventions. *PLoS ONE* **8**, e74393 (2013).
133. Kamphuis, M. H. *et al.* Physical inactivity, depression, and risk of cardiovascular mortality. *Med. Sci. Sports Exerc.* **39**, 1693–1699 (2007).
134. Burg, M. M. *et al.* The ‘perfect storm’ and acute coronary syndrome onset: do psychosocial factors play a role? *Prog. Cardiovasc. Dis.* **55**, 601–610 (2013).
135. Freedland, K. E. *et al.* Treatment of depression after coronary artery bypass surgery. *Arch. Gen. Psychiatry* **66**, 387–396 (2009).
136. Glassman, A. H. *et al.* Sertraline treatment of major depression in patients with acute MI or unstable angina. *JAMA* **288**, 701–709 (2002).
137. Honig, A. *et al.* Treatment of post-myocardial infarction depressive disorder: a randomized, placebo-controlled trial with mirtazapine. *Psychosom. Med.* **69**, 606–613 (2007).
138. Lesperance, F. *et al.* Effects of citalopram and interpersonal psychotherapy on depression in patients with coronary artery disease: the Canadian Cardiac Randomized Evaluation of Antidepressant and Psychotherapy Efficacy (CREATE) trial. *JAMA* **297**, 367–379 (2007).
139. Strik, J. J. *et al.* Efficacy and safety of fluoxetine in the treatment of patients with major depression after first myocardial infarction: findings from a double-blind, placebo-controlled trial. *Psychosom. Med.* **62**, 783–789 (2000).
140. van Melle, J. P. *et al.* Effects of antidepressant treatment following myocardial infarction. *Br. J. Psychiatry* **190**, 460–466 (2007).
141. Davidson, K. W. *et al.* Centralized, stepped, patient preference-based treatment for patients with post-acute coronary syndrome depression: CODIACS vanguard randomized controlled trial. *JAMA Intern. Med.* **173**, 997–1004 (2013).
142. Herrmann-Lingen, C. *et al.* A stepwise psychotherapy intervention for reducing risk in coronary artery disease (SPIRR-CAD): results of an observer-blinded, multicenter, randomized trial in depressed patients with coronary artery disease. *Psychosom. Med.* **78**, 704–715 (2016).
143. Thombs, B. D. *et al.* Depression screening and patient outcomes in cardiovascular care. *JAMA* **300**, 2161–2171 (2008).
144. Khan, A., Leventhal, R. M., Khan, S. R. & Brown, W. A. Severity of depression and response to antidepressants and placebo: an analysis of the Food and Drug Administration database. *J. Clin. Psychopharmacol.* **22**, 40–45 (2002).
145. Kirsch, I. *et al.* Initial severity and antidepressant benefits: a meta-analysis of data submitted to the Food and Drug Administration. *PLoS Med.* **5**, e45 (2008).
146. Davidson, K. W. *et al.* Enhanced depression care for patients with acute coronary syndrome and persistent depressive symptoms: coronary psychosocial evaluation studies randomized controlled trial. *Arch. Intern. Med.* **170**, 600–608 (2010).
147. Ye, S. *et al.* Long-term outcomes of enhanced depression treatment in patients with acute coronary syndromes. *Am. J. Med.* **127**, 1012–1016 (2014).
148. Pizzi, C. *et al.* Meta-analysis of selective serotonin reuptake inhibitors in patients with depression and coronary heart disease. *Am. J. Cardiol.* **107**, 972–979 (2011).
149. Carney, R. M. *et al.* Depression and late mortality after myocardial infarction in the Enhancing Recovery in Coronary Heart Disease (ENRICHD). *Psychosom. Med.* **66**, 466–474 (2004).
150. Glassman, A. H., Bigger, J. T. Jr & Gaffney, M. Psychiatric characteristics associated with long-term mortality among 361 patients having an acute coronary syndrome and major depression. *Arch. Gen. Psychiatry* **66**, 1022–1029 (2009).
151. de Jonge, P. *et al.* Nonresponse to treatment for depression following myocardial infarction: association with subsequent cardiac events. *Am. J. Psychiatry* **164**, 1371–1378 (2007).
152. Milani, R. V. & Lavie, C. J. Impact of cardiac rehabilitation on depression and its associated mortality. *Am. J. Med.* **120**, 799–806 (2007).
153. Chung, M. L., Dekker, R. L., Lennie, T. A. & Moser, D. K. Antidepressants do not improve event-free survival in patients with heart failure when depressive symptoms remain. *Heart Lung* **42**, 85–91 (2013).
154. Jiang, W. *et al.* Characteristics of depression remission and its relation with cardiovascular outcome among patients with chronic heart failure. *Am. J. Cardiol.* **107**, 545–551 (2011).
155. Rafanelli, C., Sirri, L., Grandi, S. & Fava, G. A. Is depression the wrong treatment target for improving outcome in coronary artery disease? *Psychother. Psychosom.* **82**, 285–291 (2013).
156. Hartley, A. *et al.* Trends in mortality from ischemic heart disease and cerebrovascular disease in Europe: 1980 to 2009. *Circulation* **133**, 1916–1926 (2016).
157. Rodriguez, T. *et al.* Trends in mortality from coronary heart and cerebrovascular diseases in the Americas: 1970–2000. *Heart* **92**, 453–460 (2006).
158. Duvis, H. E. *et al.* Depressive symptoms, health behaviors, and subsequent inflammation in patients with coronary heart disease: prospective findings from the Heart and Soul Study. *Am. J. Psychiatry* **168**, 913–920 (2011).
159. Morris, A. A. *et al.* Association between depression and inflammation — differences by race and sex: the META-Health study. *Psychosom. Med.* **73**, 462–468 (2011).
160. Whoolley, M. A. *et al.* Depression and inflammation in patients with coronary heart disease: findings from the Heart and Soul Study. *Biol. Psychiatry* **62**, 314–320 (2007).
161. Carney, R. M. & Freedland, K. E. Is there a high-risk subtype of depression in patients with coronary heart disease? *Curr. Psychiatry Rep.* **14**, 1–7 (2012).
162. Carney, R. M. & Freedland, K. E. Are somatic symptoms of depression better predictors of cardiac events than cognitive symptoms in coronary heart disease? *Psychosom. Med.* **74**, 33–38 (2012).
163. de Miranda Azevedo, R., Roest, A. M., Hoen, P. W. & de Jonge, P. Cognitive/affective and somatic/affective symptoms of depression in patients with heart disease and their association with cardiovascular prognosis: a meta-analysis. *Psychol. Med.* **44**, 2689–2703 (2014).
164. Fava, M. *et al.* A cross-sectional study of the prevalence of cognitive and physical symptoms during long-term antidepressant treatment. *J. Clin. Psychiatry* **67**, 1754–1759 (2006).
165. McClintock, S. M. *et al.* Residual symptoms in depressed outpatients who respond by 50% but do not remit to antidepressant medication. *J. Clin. Psychopharmacol.* **31**, 180–186 (2011).
166. Nierenberg, A. A. *et al.* Residual symptoms in depressed patients who respond acutely to fluoxetine. *J. Clin. Psychiatry* **60**, 221–225 (1999).
167. Zajecka, J. M. Residual symptoms and relapse: mood, cognitive symptoms, and sleep disturbances. *J. Clin. Psychiatry* **74** (Suppl. 2), 9–13 (2013).
168. Conradi, H. J., Ormel, J. & de Jonge, P. Presence of individual (residual) symptoms during depressive episodes and periods of remission: a 3-year prospective study. *Psychol. Med.* **41**, 1165–1174 (2011).
169. Carney, C. E., Segal, Z. V., Edinger, J. D. & Krystal, A. D. A comparison of rates of residual insomnia symptoms following pharmacotherapy or cognitive-behavioral therapy for major depressive disorder. *J. Clin. Psychiatry* **68**, 254–260 (2007).
170. Kang, E. H. *et al.* Mirtazapine versus venlafaxine for the treatment of somatic symptoms associated with major depressive disorder: a randomized, open-labeled trial. *Psychiatry Res.* **169**, 118–123 (2009).
171. Nelson, J. C., Portera, L. & Leon, A. C. Are there differences in the symptoms that respond to a selective serotonin or norepinephrine reuptake inhibitor? *Biol. Psychiatry* **57**, 1535–1542 (2005).
172. Stewart, J. G. & Harkness, K. L. Symptom specificity in the acute treatment of major depressive disorder: a re-analysis of the treatment of depression collaborative research program. *J. Affect. Disord.* **137**, 87–97 (2012).
173. Blumenthal, J. A. *et al.* Exercise, depression, and mortality after myocardial infarction in the ENRICHD trial. *Med. Sci. Sports Exerc.* **36**, 746–755 (2004).
174. Daley, A. Exercise and depression: a review of reviews. *J. Clin. Psychol. Med. Settings* **15**, 140–147 (2008).
175. Mead, G. E. *et al.* Exercise for depression. *Cochrane Database of Systematic Reviews*, Issue 3. Art. No.: CD004366. <http://dx.doi.org/10.1002/14651858.CD004366.pub4> (2009).
176. Rethorst, C. D., Wipfli, B. M. & Landers, D. M. The antidepressive effects of exercise: a meta-analysis of randomized trials. *Sports Med.* **39**, 491–511 (2009).
177. Poirier, P. Exercise, heart rate variability, and longevity: the cocoon mystery? *Circulation* **129**, 2085–2087 (2014).
178. Rutledge, T., Redwine, L. S., Linke, S. E. & Mills, P. J. A meta-analysis of mental health treatments and cardiac rehabilitation for improving clinical outcomes and depression among patients with coronary heart disease. *Psychosom. Med.* **75**, 335–349 (2013).
179. Sandercock, G. R., Bromley, P. D. & Brodie, D. A. Effects of exercise on heart rate variability: inferences from meta-analysis. *Med. Sci. Sports Exerc.* **37**, 433–439 (2005).
180. Beavers, K. M., Brinkley, T. E. & Nicklas, B. J. Effect of exercise training on chronic inflammation. *Clin. Chim. Acta* **411**, 785–793 (2010).
181. Pedersen, B. K. & Fischer, C. P. Physiological roles of muscle-derived interleukin-6 in response to exercise. *Curr. Opin. Clin. Nutr. Metab. Care* **10**, 265–271 (2007).
182. Pedersen, M. *et al.* Circulating levels of TNF- α and IL-6 relation to truncal fat mass and muscle mass in healthy elderly individuals and in patients with type-2 diabetes. *Mech. Ageing Dev.* **124**, 495–502 (2003).
183. Ploeger, H. E., Takken, T., de Greef, M. H. & Timmons, B. W. The effects of acute and chronic exercise on inflammatory markers in children and adults with a chronic inflammatory disease: a systematic review. *Exerc. Immunol. Rev.* **15**, 6–41 (2009).
184. Edinger, J. D., Wohlgemuth, W. K., Radtke, R. A., Marsh, G. R. & Quillian, R. E. Cognitive behavioral therapy for treatment of chronic primary insomnia: a randomized controlled trial. *JAMA* **285**, 1856–1864 (2001).
185. Morin, C. M. *et al.* Cognitive behavioral therapy, singly and combined with medication, for persistent insomnia: a randomized controlled trial. *JAMA* **301**, 2005–2015 (2009).
186. Siversen, B. *et al.* Cognitive behavioral therapy versus zopiclone for treatment of chronic primary insomnia in older adults: a randomized controlled trial. *JAMA* **295**, 2851–2858 (2006).

Acknowledgements

The authors are supported by grant number R01HL089336 from the National Heart, Lung, and Blood Institute of the National Institutes of Health, Bethesda, Maryland, USA.

Author contributions

Both authors researched data for the article, discussed the content, wrote the manuscript, and revised and edited the article before submission.

Completing interests statement

R.M.C. or a member of his family owns stock in Pfizer. K.E.F. declares no competing interests.