

JACC REVIEW TOPIC OF THE WEEK

Ischemic Stroke Risk in Patients With Nonvalvular Atrial Fibrillation

JACC Review Topic of the Week

Mohamad Alkhouli, MD, Paul A. Friedman, MD



JACC JOURNAL CME/MOC/ECME

This article has been selected as the month's JACC CME/MOC/ECME activity, available online at <http://www.acc.org/jacc-journals-cme> by selecting the JACC Journals CME/MOC/ECME tab.

Accreditation and Designation Statement

The American College of Cardiology Foundation (ACCF) is accredited by the Accreditation Council for Continuing Medical Education to provide continuing medical education for physicians.

The ACCF designates this Journal-based CME activity for a maximum of 1 AMA PRA Category 1 Credit(s)[™]. Physicians should claim only the credit commensurate with the extent of their participation in the activity.

Successful completion of this CME activity, which includes participation in the evaluation component, enables the participant to earn up to 1 Medical Knowledge MOC point in the American Board of Internal Medicine's (ABIM) Maintenance of Certification (MOC) program. Participants will earn MOC points equivalent to the amount of CME credits claimed for the activity. It is the CME activity provider's responsibility to submit participant completion information to ACCME for the purpose of granting ABIM MOC credit.

Ischemic Stroke Risk in Patients With Nonvalvular Atrial Fibrillation: JACC Review Topic of the Week will be accredited by the European Board for Accreditation in Cardiology (EBAC) for 1 hour of External CME credits. Each participant should claim only those hours of credit that have actually been spent in the educational activity. The Accreditation Council for Continuing Medical Education (ACCME) and the European Board for Accreditation in Cardiology (EBAC) have recognized each other's accreditation systems as substantially equivalent. Apply for credit through the post-course evaluation. While offering the credits noted above, this program is not intended to provide extensive training or certification in the field.

Method of Participation and Receipt of CME/MOC/ECME Certificate

To obtain credit for JACC CME/MOC/ECME, you must:

1. Be an ACC member or JACC subscriber.
2. Carefully read the CME/MOC/ECME-designated article available online and in this issue of the *Journal*.
3. Answer the post-test questions. A passing score of at least 70% must be achieved to obtain credit.
4. Complete a brief evaluation.
5. Claim your CME/MOC/ECME credit and receive your certificate electronically by following the instructions given at the conclusion of the activity.

CME/MOC/ECME Objective for This Article: Upon completion of this activity, the learner should be able to: 1) relate to the evolution of stroke risk prediction in patients with nonvalvular AF; 2) identify the advantages and limitations of the commonly used CHA₂DS₂-VASC score; and 3) discuss the emerging tools for ischemic stroke risk determination in patients with nonvalvular AF.

CME/MOC/ECME Editor Disclosure: JACC CME/MOC/ECME Editor Ragavendra R. Baliga, MD, FACC, has reported that he has no financial relationships or interests to disclose.

Author Disclosures: Both authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Medium of Participation: Print (article only); online (article and quiz).

CME/MOC/ECME Term of Approval

Issue Date: December 17, 2019

Expiration Date: December 16, 2020



Listen to this manuscript's
audio summary by
Editor-in-Chief
Dr. Valentin Fuster on
JACC.org.

From the Department of Cardiovascular Medicine, Mayo Clinic, Rochester, Minnesota. Both authors have reported that they have no relationships relevant to the contents of this paper to disclose.

Manuscript received July 30, 2019; revised manuscript received October 16, 2019, accepted October 21, 2019.

Ischemic Stroke Risk in Patients With Nonvalvular Atrial Fibrillation

JACC Review Topic of the Week

Mohamad Alkhouli, MD, Paul A. Friedman, MD

ABSTRACT

The last decade has witnessed remarkable advances in pharmacological and nonpharmacological strategies for stroke prevention in patients with atrial fibrillation. However, the currently available clinical stroke risk prediction models do not account for key nonclinical factors (arrhythmia burden, left atrial physiology and anatomy, chemical and electrocardiographic markers) and other competing clinical risks. Hence, their ability to identify patients who will derive the most benefit from pharmacological and mechanical risk prevention strategies remain limited. In this paper, the authors review the current and evolving ischemic stroke risk prediction schemes in patients with nonvalvular atrial fibrillation, highlight the strengths and weaknesses of the models, and discuss the unmet needs in this field.

(J Am Coll Cardiol 2019;74:3050-65) © 2019 by the American College of Cardiology Foundation.

Atrial fibrillation (AF) is a growing epidemic, with >12 million patients projected to have AF in the United States by 2030 (1-3). Stroke is the most debilitating complication of AF, and its prevention is hence considered the cornerstone of AF management (1,4). Numerous stroke prevention strategies have been introduced, including the use of anticoagulation centers of excellence, direct oral anticoagulants, and percutaneous left atrial (LA) appendage occlusion (LAAO) devices (1,5). However, stroke risk prediction models have not kept pace with these advances. Current societal guidelines remain, for the most part, depended on a single clinical risk score (CHA₂DS₂-VASc) to determine an individual's candidacy for pharmacological or mechanical stroke prevention (6-8). Unfortunately, despite its utility and ease of calculation, the CHA₂DS₂-VASc score has demonstrated modest performance in predicting stroke or other cerebrovascular ischemic events in real-world cohorts (9). In addition, the score does not consider several key anatomic, physiological, and other factors that have been shown to impact the risk of stroke in patients with AF (10). The aim of this paper is to review current and emerging tools for ischemic stroke risk determination in patients with nonvalvular AF (NVAF), and to propose a framework for further investigation to achieve optimal stroke prevention in these patients. Data provided in this paper were gathered through a focused review of the published studies on risk prediction of ischemic stroke in patients with AF (nonsystematic review).

HISTORICAL PERSPECTIVES

William Osler characterized medicine as a science of uncertainty and an art of probability (11). This is clearly illustrated in the management of NVAF, where the prevention of AF-associated ischemic stroke is based on balancing the probability of benefit versus harm from a specific strategy using probability calculation tools that are known to have considerable limitations. Herein, we review the origin and evolution of the prior and currently utilized stroke risk prediction scores.

In 1978, Wolf et al. (12) established the association between NVAF and stroke by demonstrating a 5-fold increase in the incidence of ischemic stroke among patients with NVAF compared with patients without NVAF in the landmark Framingham study. This observation prompted several randomized clinical trials (RCTs) aiming to evaluate the efficacy of oral anticoagulation in reducing stroke risk in patients with NVAF (13). A patient-level meta-analysis of the first 5 RCTs demonstrated that warfarin was associated with a 68% relative risk (RR) reduction in stroke compared with placebo (14). This substantial benefit was confirmed in further studies and additional meta-analyses, granting warfarin a central role in the management of NVAF for several decades (1,15-18). However, the necessity of objectively estimating stroke risk in NVAF patients became quickly evident, as the risk/benefit of oral anticoagulation (OAC) varied substantially between individuals on the basis of the underlying risk of stroke (19-21). Hence,

ABBREVIATIONS AND ACRONYMS

AF	= atrial fibrillation
AI	= artificial intelligence
CI	= confidence interval
HR	= hazard ratio
LA	= left atrium
LAA	= left atrial appendage
LAAO	= left atrial appendage occlusion
NT-proBNP	= N-terminal pro-B-type natriuretic peptide
NVAF	= nonvalvular atrial fibrillation
OAC	= oral anticoagulation
RCT	= randomized clinical trial
RR	= relative risk
SE	= systemic embolization

considerable efforts were made to devise an individualized risk classification score for ischemic stroke in patients with NVAF (22,23).

CLINICAL RISK SCORES

THE CHADS₂ SCORE. The AFI (Atrial Fibrillation Investigators) and the SPAF (Stroke Prevention and Atrial Fibrillation) trial classification schemes were the first to provide objective evidence of the association between clinical features and a heightened risk of stroke in patients with NVAF (14,17,24). In the AFI classification, age, hypertension, prior cerebral ischemia, and diabetes mellitus were identified as independent predictors of stroke (17). In the SPAF classification, blood pressure >160 mm Hg, prior cerebral ischemia, recent clinical or echocardiography-proven heart failure, or the combination of age ≥75 years and female sex were associated with higher odds of ischemic stroke (24). However, these classifications were viewed as ambiguous, were often conflicting, and had modest performance when validated outside of the clinical trial setting (19). Hence, Gage et al. (19) combined elements from the 2 schemes into a new risk score (CHADS₂: congestive heart failure, hypertension, age ≥75, diabetes, prior stroke/transient ischemic attack) and validated its predictive value in a large cohort of Medicare-age patients. Due to its C-statistic of 0.82 (95% confidence interval [CI]: 0.80 to 0.84) and ease of use, the CHADS₂ score emerged as the most accurate predictor of stroke, and was promulgated widely in societal guidelines and in clinical practice.

With the wide adoption of CHADS₂, some of its limitations have become apparent. First, the score was found to have a weak discriminatory value in identifying low-risk patients, as patients with a CHADS₂ score of 0 still had an annual stroke risk of ~2%. Second, the score was noninclusive of key risk factors (e.g., female sex, vascular disease) that were increasingly recognized as independently predisposing towards thromboembolism in NVAF patients (25). Third, age was treated as a binary variable in CHADS₂, whereas emerging data had shown that increasing age among NVAF patients >65 years of age was associated with an incremental increase in stroke risk (19). Fourth, several studies have shown modest-to-poor validity of the CHADS₂ score in real-world cohorts. Hence, attempts were made to further modify/supplement the CHADS₂ score.

HIGHLIGHTS

- Ischemic stroke risk prediction is a cornerstone in the management of patients with atrial fibrillation.
- The paper reviews the evolution of these risk scores, discusses their strengths and limitations, and appraises the emerging risk assessment tools and their incremental utility.
- There is an unmet need for a comprehensive study incorporating various clinical, anatomic, and biophysiological risk factors to optimize our stroke prevention practices in patients with atrial fibrillation.

CHA₂DS₂-VASc SCORE. To address the shortcomings of the CHADS₂ score, Lip et al. (25) added additional risk factors (vascular disease [coronary, peripheral arterial and venous], and female sex) and age categories (<60 years, 60 to 74 years, and ≥75 years) to create a new risk scoring system: CHA₂DS₂-VASc. In their study, the investigators found that the CHA₂DS₂-VASc score outperformed the CHADS₂ score in identifying NVAF patients at risk for stroke. The incremental benefit of the CHA₂DS₂-VASc score documented by Lip et al. (25) was initially questioned due to its modest discriminatory power (C-statistic = 0.606), its limited follow-up (1 year), and its incomplete surveillance (no follow-up data in 31% of patients) (26). However, several subsequent validation studies confirmed the superiority of CHA₂DS₂-VASc over CHADS₂ in quantifying stroke risk, especially in its ability to identify low-risk patients who do not benefit from stroke prevention therapy (27-30). Hence, the European Society of Cardiology and the American College of Cardiology/American Heart Association guidelines were updated in 2012 and 2014, respectively, recommending the use of CHA₂DS₂-VASc instead of the CHADS₂ score to guide OAC in NVAF patients.

OTHER CLINICAL RISK SCORES. In the 2010s, the CHA₂DS₂-VASc score became the leading stroke risk prediction tool for patients with NVAF, both in the guidelines and in clinical practice. Indeed, the enthusiasm for the CHA₂DS₂-VASc score has fueled dozens of investigations demonstrating its potential role, not only for predicting ischemic stroke, but also for predicting other outcomes in patients with and without NVAF (13,31-34). However, the score is not without limitations, and its suboptimal performance

TABLE 1 Characteristics of the Main Clinical Stroke Risk Stratifications Schemes in NVAF Patients

Risk Score First Author (Ref. #)	Data Source	Score Definition	Supportive Data
CHADS ₂ Gage et al. (19)	Medicare Database* N = 1,733	CHF, hypertension, age (≥65 yrs = 1 point, ≥75 yrs = 2 points), diabetes, and stroke/TIA (2 points)	CHADS ₂ was superior to existing risk classification schemes AFI scheme: C-statistic = 0.68 (0.65–0.71) SPAF-III scheme: C-statistic = 0.74 (0.71–0.76) CHADS ₂ score: C-statistic = 0.82 (0.80–0.84)
R ₂ CHADS ₂ Piccini et al. (35)	ROCKET-AF trial, ATRIA trial N = 14,264	Added +2 points for creatinine clearance <60 ml/min	In the derivation cohort (ROCKET-AF): C-statistic was 0.587 vs. 0.575 for CHADS ₂ , NRI = 8.2% (2.5%–14%) In the validation cohort (ATRIA): C-statistic was 0.704 vs. 0.696 for CHADS ₂ , NRI = 22.6% (14.5%–30.7%)
CHA ₂ DS ₂ -VASC Lip et al. (25)	Euro Heart Survey for AF N = 1,084	CHF, hypertension, age ≥75 yrs, diabetes, stroke, or TIA, vascular disease, age 65–74 yrs, sex	C-statistic was 0.606 (0.513–0.699) for CHA ₂ DS ₂ -VAs vs. 0.561 (0.450–0.672) for CHADS ₂ . Patients were classified as low (9.2% vs. 20.4%), intermediate (15.1% vs. 61.9%), or high (75.7% vs. 17.7%) risk with CHA ₂ DS ₂ -VAs vs. CHADS ₂ .
CHA ₂ DS ₂ -VASC-R Kabra et al. (40)	Medicare Database N = 460,417	Added +1 point to African-American race	CHA ₂ DS ₂ -VASC-R superior to CHA ₂ DS ₂ -VASC: HR: 1.36 (1.29–1.43; p < 0.001), NRI = 7.6% (p < 0.001)
CHA ₂ DS ₂ -VA Tomita et al. (42)	J-RHYTHM registry N = 997	Eliminated sex category	CHA ₂ DS ₂ -VA superior to CHA ₂ DS ₂ -VASC overall [C-statistic = 0.029; p = 0.02, NRI = 11% (1%–20%)] and in low-risk patients: [C-statistic = 0.053; p < 0.001, NRI = 11% (7%–14%)]
mCHA ₂ DS ₂ -VASC Chao et al. (37)	Taiwan National Insurance Database N = 224,866	Expanded lower threshold for age to 50 yrs (1 point for age 50–74 yrs)	mCHA ₂ DS ₂ -VASC was superior to CHA ₂ DS ₂ -VAs C-statistics = 0.708 (0.703–0.712) vs. 0.689 (0.684–0.694) NRI = 3.39% (2.16%–4.59%); p < 0.001
CHA ₂ DS ₂ -VAK Cha et al. (36)	Korean single-center cohort N = 12,876	Replaced sex category with kidney disease (GFR <60 ml/min/1.73 m ²)	CHA ₂ DS ₂ -VAK was superior to CHA ₂ DS ₂ -VAs C-statistics = 0.650 (0.64–0.66) vs. 0.639 (0.62–0.65)
ATRIA Singer et al. (41)	ATRIA, ATRIA-CVRN cohort N = 10,927	Age (65–74 yrs = 3 points, 75–84 yrs = 5 points, ≥85 yrs = 6 points), hypertension, diabetes, CHF, proteinuria, GFR <45 ml/min/1.73 m ² , sex	ATRIA was superior to CHA ₂ DS ₂ -VASC: C-statistics = 0.708 (0.704–0.713) vs. 0.694 (0.690–0.700), NRI 16% (14%–17%).
ABC Hijazi et al. (39)	ARISTOTLE trial, STABILITY trial N = 18,201	Age, biomarkers (hs-troponin T, NT-proBNP), prior history of stroke	The ABC score yielded higher C-statistics than CHA ₂ DS ₂ -VAs in the derivation cohort (0.68 vs. 0.62; p < 0.001) and the external validation cohort (0.66 vs. 0.58; p < 0.001)
GARFIELD-AF Fox et al. (38)	GARFIELD-AF, ORBIT-AF registry N = 39,898	Computed machine learning model (web-based)	The GARFIELD-AF score yielded higher C-statistics than CHA ₂ DS ₂ -VAs 0.69 (0.67–0.71) vs. 0.64 (0.61–0.66) in the derivation cohort. In the validation cohort, GARFIELD-AF C-statistic was 0.68 (0.62–0.74)

*NRAF (National Registry of Atrial Fibrillation) data.


ABC = age, biomarker, clinical history; AFI = Atrial Fibrillation Investigators; ARISTOTLE = Apixaban for Reduction in Stroke and Other Thromboembolic Events in Atrial Fibrillation; ATRIA = Anticoagulation and Risk Factors in Atrial Fibrillation; BNP = B-type natriuretic peptide; CHF = congestive heart failure; CVRN = Cardiovascular Research Network; GARFIELD = Global Anticoagulant Registry in the FIELD; GFR = glomerular filtration rate; hs = high-sensitivity; NRI = net reclassification index; NT-proBNP = N-terminal pro-B-type natriuretic peptide; NVAF = nonvalvular atrial fibrillation; ROCKET-AF = Rivaroxaban versus Warfarin in Nonvalvular Atrial Fibrillation trial; SPAF III = Stroke Prevention in Atrial Fibrillation III trial; STABILITY = The Stabilization of Atherosclerotic Plaque by Initiation of Darapladib Therapy Trial; TIA = transient ischemic attack.

in selected populations (e.g., patients with renal insufficiency, Asian patients) stimulated a quest for a superior stroke prediction scheme. Between 2010 and 2018, novel clinical risk scores were derived and validated in large cohorts of patients including: the ATRIA (Anticoagulation and Risk Factors in Atrial Fibrillation), ABC (Age, Biomarkers, Clinical History), and the GARFIELD-AF (Global Anticoagulant Registry in the Field) (Table 1, Figure 1) (35–42). However, none of these scores has gained the wide spread adoption that the CHA₂DS₂-VASC score has enjoyed.

Despite the proven utility of the CHA₂DS₂-VASC score, and the data supporting an incremental value of the newer risk scores in selected populations, these scores all share a common theme: they are solely based on clinical risk factors (e.g., age, hypertension,

diabetes, vascular disease) that also increase the risk of stroke in patients without AF. Hence, concerns have been raised about the lack of specificity of the CHA₂DS₂-VASC score for predicting AF-related versus non-AF-related ischemic stroke. These concerns were substantiated by several studies that documented a similar predictive value for stroke with the application of the CHA₂DS₂-VASC score in patients with and without AF (13,31–34,43). Hence, investigators sought to identify AF-specific risk factors for ischemic stroke, and tested their additive value to the CHA₂DS₂-VASC and other clinical scores. Here, we summarize the published reports on anatomic, physiological, and electrocardiographic factors that have been shown to have an impact on the risk of stroke in patients with NVAF. It should be noted, nonetheless, that all of the established and emerging risk scores have been

FIGURE 1 Temporal Evolution of the Current Stroke Reduction Scores



AF = atrial fibrillation; OAC = oral anticoagulation.

evaluated in terms of their ability to predict clinical ischemic strokes in patients with NVAf. Future studies should consider silent ischemic strokes, which have been shown to be prevalent and to carry a negative prognostic impact in this population (44).

PREDICTORS OF STROKE RISK BEYOND CLINICAL RISK SCORES


1. BURDEN OF AF AND STROKE. Current guidelines recommend the use of clinical risk scores (e.g., CHA₂DS₂-VASc) to determine eligibility for anticoagulation in patients with NVAf and do not account for AF burden. These recommendations may be based on older studies showing no differential impact of AF burden on stroke risk. In the ACTIVE W trial (Atrial Fibrillation Clopidogrel Trial With Irbesartan for Prevention of Vascular Events) (45), the risk of stroke/systemic embolization (SE) was 2.0% in patients with paroxysmal AF versus 2.2% in sustained AF (RR: 0.87; 95% CI: 0.59 to 1.30; $p = 0.50$) (45). Similarly, in the SPAF trial, the annualized rate of ischemic stroke of 3.2% in patients with intermittent AF was 3.3% in those with sustained AF (46). However, there is growing evidence suggesting a strong association between the burden of AF and the risk of subsequent ischemic events.

The ENGAGE AF-TIMI 48 (Effective Anticoagulation With Factor Xa Next Generation in Atrial Fibrillation-Thrombolysis In Myocardial Infarction 48) trial showed a linear association between AF burden and risk of stroke (47). In this trial, paroxysmal AF (<7 days) was associated with lower stroke/SE event rates compared with persistent AF

(≥ 7 days, but <1 year) (1.83%/year; hazard ratio [HR]: 0.79; 95% CI: 0.66 to 0.90; $p = 0.015$), or permanent AF (≥ 1 year) (1.95%/year; HR: 0.78; 95% CI: 0.67 to 0.93; $p = 0.004$). These results were replicated in post hoc analyses of major RCTs assessing anticoagulation (ARISTOTLE [Apixaban for Reduction in Stroke and Other Thromboembolic Event], ROCKET-AF [Rivaroxaban Once-Daily, Oral, Direct Factor Xa Inhibition Compared With Vitamin K Antagonism for Prevention of Stroke and Embolism Trial in Atrial Fibrillation]), or antiplatelet therapy (ACTIVE A [Atrial Fibrillation Clopidogrel Trial With Irbesartan for Prevention of Vascular Events], AVERROES [A Phase III Study of Apixaban in Patients With Atrial Fibrillation]) for AF. Two study-level meta-analyses including RCTs, population-based studies, and reports from AF registries found a strong relationship between AF burden and stroke risk. In the first meta-analysis (12 studies; 99,996 patients), the unadjusted RR ratio of stroke/SE in patients with nonparoxysmal versus paroxysmal AF was 1.35 (95% CI: 1.17 to 1.57; $p < 0.001$), and this persisted after risk adjustment: 1.38 (95% CI: 1.19 to 1.61; $p < 0.001$) (Figure 2) (48). In the second meta-analysis (18 studies; 239,528 patient-years), the incidence rate of stroke or SE was 1.6% versus 2.3% in patients with in paroxysmal versus nonparoxysmal AF, respectively (RR: 0.72; 95% CI: 0.65 to 0.80; $p < 0.001$) (49).

It remained unclear, however, whether the risk of stroke in paroxysmal AF increases continuously or whether a threshold exists. Analyses of patients with pre-existing cardiac implantable electronic were confounded by several issues: AF burden cutoffs were arbitrarily pre-specified, stroke events were not

FIGURE 2 Meta-Analysis Comparing Stroke Rates Between Patients With PAF and NPAF



(A) Pooled hazard risk ratio for stroke or systemic embolization in paroxysmal versus nonparoxysmal atrial fibrillation (12 studies; n = 99,996). (B) Pooled adjusted hazard risk ratio for stroke or systemic embolization in paroxysmal versus non-paroxysmal atrial fibrillation (7 studies; n = 58,421). Multivariable risk adjustment was performed at the study level and was not uniform across the studies. Reprinted with permission from Ganesan and Chew (48). CI = confidence interval; NPAF = nonparoxysmal atrial fibrillation; PAF = paroxysmal atrial fibrillation.

TABLE 2 Knowledge Gaps in the Association Between the Burden of NVAF and Stroke Risk

Unified definitions AF burden
Monitoring technologies that accurately measure the burden of AF
Threshold of AF that increases stroke risk in AF patients
Differential impact of AF burden on patients with various CHA ₂ DS ₂ -VASc scores
Association between AF burden and stroke severity and outcomes
Association between AF burden and cognitive function
Relationship between AF burden and quality of life
Reasons for the weak temporal relationship between AF episodes and stroke events
Impact of AF burden reduction therapies on stroke risk

AF = atrial fibrillation; NVAF = nonvalvular atrial fibrillation.

found to be temporally related to AF episodes, and data from patients with implantable devices were not deemed generalizable given the high prevalence of advanced comorbidities in the implantable electronic devices' population. Nonetheless, post hoc analysis of prospective randomized trial data suggests a dose effect for AF burden and stroke, with various studies proposing increased risk above 6 min, 6 h, and 24 h (50). A recent study by Go et al. (51) attempted to mitigate these issues by assessing the impact of various durations of paroxysmal AF on stroke/SE rates in patients not on anticoagulation, using 14-day continuous ambulatory electrocardiographic monitoring. In this study, the median burden of AF was 4.4% (interquartile range: 1.1% to 17.2%). After adjusting for the CHA₂DS₂-VASc score, the highest tertile of AF burden ($\geq 11.4\%$) was associated with a >3-fold higher adjusted rate of stroke/SE (adjusted HR: 3.16, 95% CI: 1.5 to 6.6) compared with the combined lower 2 tertiles of AF burden. A recent statement from the American Heart Association acknowledged the limitations of the current guidelines that classify AF in a binary fashion (AF vs. no AF), and called for further research to bridge the knowledge gaps in our understanding of the association between AF burden and stroke (Table 2) (52). Furthermore, the lack of temporal association between AF and stroke has raised the question as to whether AF is pathogenic for stroke or simply a marker of the extent of atrial fibrosis and dysfunction necessary for stroke. This has important implications for some stroke prevention strategies such as a "pill in the pocket approach" for paroxysmal AF (53).

2. LA ANATOMY AND FUNCTION. The size, function, and structure of the LA have been shown to modify the risk of stroke in patients with NVAF (Figure 3).


Left atrial size and stroke risk. Benjamin et al. (54) documented an ~40% and >100% increase in

adjusted stroke rates for each 10-mm increase in LA size in women and men enrolled in the Framingham Heart Study, respectively. Several other studies have corroborated a similar relationship between LA size and stroke risk specifically in patients with NVAF (55-57). Moreover, the LA size has also been found predictive of ischemic stroke recurrence in patients with NVAF (58,59).

LA function and stroke risk. The pathogenesis of AF is complex. However, certain LA functional and structural abnormalities are common in patients with AF including: loss of LA reservoir function, LA fibrosis, reduced LA flow, and LA electrical remodeling. The prognostic implications of these abnormalities have been an area of interest. The association between LA abnormalities and AF recurrence after transcatheter AF ablation has been documented in several studies (60). However, data on the association between LA structural abnormalities and stroke risk remain limited. In a study by Leong et al. (61), each 1% reduction in LA reservoir function (as measured by speckle tracking longitudinal strain imaging) was associated with 7% increase in stroke ($p < 0.001$). In another study by Daccarett et al. (62), patients with AF with a history of stroke had significantly more LA fibrosis on cardiac magnetic resonance imaging (CMR) compared with those without a history of stroke ($24.4 \pm 12.4\%$ vs. $16.2 \pm 9.9\%$, respectively; $p < 0.001$). Incorporating the extent of LA fibrosis into the CHADS₂ score improved the predictive value of the score (C-statistics = 0.58 to 0.72) (62). In the MESA (Multi-Ethnic Study of Atherosclerosis) study, LA phasic function (LA ejection fraction) as measured by CMR was associated with incident ischemic stroke (HR: 0.85/SD; 95% CI: 0.74 to 0.98; $p = 0.027$) (63). However, technical issues related to LA imaging in patients with NVAF have tempered the enthusiasm for these approaches, given the relatively simple and readily available clinical risk scores.

3. LA APPENDAGE ANATOMY AND FUNCTION. The role of the LA appendage (LAA) in the causation of stroke in patients with NVAF has been an area of increasing interest since Blackshear first showed that the LAA is the nidus for thrombus in 91% of patients with NVAF (64). This unique observation led to the development of the concept of mechanical exclusion of the LAA as an alternative strategy to OAC for stroke prevention. The proliferation of innovative technologies in this field has shed more light on the enormous variation in the size, shape, and function of the LAA (Figure 4). The impact of these variations on stroke risk is now supported by a large body of evidence (65).

FIGURE 3 Assessment of LA Fibrosis by CMR in Patients With AF




(Reprinted with permission from Siebermair J, Kholmovski EG, Marrouche N. Assessment of left atrial fibrosis by late gadolinium enhancement magnetic resonance imaging: methodology and clinical implications. *J Am Coll Cardiol EP* 2017;3:791-802). 3D = 3-dimensional; AF = atrial fibrillation; CMR = cardiac magnetic resonance; LA = left atrial.

LAA size/function and stroke risk. The plausible hypothesis that large and/or less mobile LAAs may be associated with higher odds of clot formation and subsequent embolic events compared with small contractile LAAs has been confirmed in several studies. In a cross-sectional study of 218 patients, Lee et al. (66) found that large LAA orifice area confers a 6-fold increase in the adjusted risk of stroke (OR: 6.2; 95% CI: 2.67 to 14.18; $p < 0.001$). In another study using CMR, LAA volume $>34 \text{ cm}^3$ was associated with a 7-fold increase in adjusted stroke rate (OR: 7.1; $p = 0.003$). Fatkin et al. (67) documented a substantial correlation between low LAA emptying velocity ($<35 \text{ cm/s}$) and LAA thrombus/

spontaneous echo contrast (OR: 28; 95% CI: 9.1 to 84.4). In the SPAF-III (Stroke Prevention in Atrial Fibrillation) study, peak antegrade flow velocity $<20 \text{ cm/s}$ was independently associated with LAA thrombus (RR: 2.6; $p = 0.02$) (68). Similarly, Kamp et al. (69) found that LAA peak velocity $<20 \text{ cm/s}$ was an independent predictor of subsequent thromboembolic events (OR: 4.1; 95% CI: 1.4 to 11.6; $p < 0.01$). Extent of LAA fibrosis may also play a role in the assessment of stroke risk in patients with NVAF (70).

LAA morphology and stroke risk. The role of the LAA anatomy in stroke prediction has also been assessed in a number of studies. In a multicenter

FIGURE 4 LAA Angiography Illustrating the Variation in the LAA Size and Shape in 4 High-Risk Patients With Paroxysmal AF



(A) Small nontrabeculated appendage, (B) medium-size heavily trabeculated appendage, (C) large anterior chicken wing-shaped appendage, (D) very large windsock shape appendage. All 4 patients had a CHA₂DS₂-VASc score of 4. AF = atrial fibrillation; LAA = left atrial appendage.


study including 433 patients with drug-refractory AF who underwent computed tomographic imaging before AF ablation, the rates of ischemic stroke were 4.08-fold, 4.5-fold, and 8.0-fold higher in patients with cactus, windsock, or cauliflower LAA anatomy compared with those with chicken wing anatomy ($p = 0.046$, 0.038 , and 0.056 , respectively) (71). In a meta-analysis of 8 studies including 2,596 patients with NVAF, thromboembolism risk was lower in patients with chicken wing morphology versus those with non-chicken wing morphology (OR: 0.46; 95% CI: 0.36 to 0.58) (72). A recent proof-of-concept study showed that a simplified classification of LAA anatomy into low and high risk on the basis of an acute angle bend or fold in the proximal/middle portion of the appendage is useful, not only in

identifying high-risk patients, but also in predicting subtypes of ischemic strokes (73).

4. BIOMARKERS. AF is a chronic condition with a variable disease burden. Cardiac and inflammatory biomarkers in patients with AF might be surrogates for the chronicity and/or severity of the disease (74,75). Hence, the incorporation of such biomarkers into the current clinically based stroke risk prediction models may improve their performance (74).

Cardiac biomarkers. The incremental value of troponin-I and N-terminal pro-B-type natriuretic peptide (NT-proBNP) in predicting stroke risk in patients with NVAF was tested in a substudy of the RE-LY (Randomized Evaluation of Long-Term Anticoagulation Therapy) trial (76). In this trial, rates of

FIGURE 5 LA Abnormalities Associated With Stroke on 12-Lead Electrocardiograms in Patients With Paroxysmal AF




Reprinted with permission from Maheshwari et al. (84). AF = atrial fibrillation; LA = left atrial.

stroke were 2.1% versus 0.8% per year in the highest versus the lowest troponin-I quartile groups (HR: 1.99; 95% CI: 1.17 to 3.39; $p = 0.004$), and to NT-proBNP with 2.3% versus 0.9% per year in the highest versus the lowest NT-proBNP quartile groups (HR: 2.40; 95% CI: 1.41 to 4.07; $p = 0.001$). Similarly, in the ARISTOTLE trial, stroke rates were 2.1% versus 0.9% per year in the highest versus the lowest high-sensitivity troponin-T quartile groups (HR: 1.94; 95% CI: 1.35 to 2.78; $p = 0.001$), 2.2% versus 0.7% per

year in the highest versus the lowest NT-proBNP quartile groups (HR: 2.35; 95% CI: 1.62 to 3.40; $p < 0.0001$) (77,78). Other observational studies have shown similar findings (79,80). These observations formed the foundation for the first hybrid score (ABC score), which included risk factors (age, prior stroke) and cardiac biomarker levels (39).

Inflammatory biomarkers. Inflammatory biomarkers are elevated in patients with AF compared with those without AF (75). However, studies assessing their




values in predicting stroke in patients with AF yielded mixed results. In the SPAF-III study, C-reactive protein levels were associated with all-cause mortality, but not with stroke events (81). In the RE-LY study, C-reactive protein and fibrinogen were not associated with stroke, whereas interleukin-6 showed a modest association with stroke/SE (C-statistics for CHA₂DS₂-VASC increased from 0.615 to 0.642 [p = 0.002]) (82). In the ARISTOTLE trial, interleukin-6 was not associated with stroke after adjusting for clinical risk factors and other cardiac biomarkers (83). Although some biomarkers do correlate with stroke risk, none have reached a discriminatory power and ease of use sufficient for widespread adoption.

5. ELECTROCARDIOGRAPHIC MARKERS. The presence of a P wave during a period of sinus rhythm in patients with paroxysmal AF allows measurements of

several P-wave indices that have been shown to be surrogates for atrial cardiopathy (84). This has led to several investigations seeking to assess the potential prognostic value of these indices in predicting subsequent ischemic events (34,84-86). The largest relevant study assessed the association between 4 P-wave indices (abnormal P-wave axis, PR duration, advanced intra-atrial block, abnormal P-wave terminal force in lead V₁) and stroke events in 2,229 patients enrolled in the ARIC (Atherosclerosis Risk In Communities) study and 700 patients enrolled in the MESA study (Figure 5) (84). The study found that an abnormal P-wave axis was the only P-wave parameter that was associated with higher odds of stroke independent of the CHA₂DS₂-VASC score (HR: 1.84; 95% CI: 1.33 to 2.55). Incorporating the P-wave axis in the CHA₂DS₂-VASC score improved its C-statistics from 0.68 (95% CI: 0.52

FIGURE 6 A Proposed Outline for a Prospective Study to Comprehensively Assess Stroke Risk in Patients With NVAF



AI = artificial intelligence; BNP = B-type natriuretic peptide; CRP = C-reactive protein; EF = ejection fraction; EKG = electrocardiogram; hs = high-sensitivity; LAAO = left atrial appendage occlusion; MRI = magnetic resonance imaging; TIA = transient ischemic attack; other abbreviations as in [Figures 3 and 4](#).

to 0.84) to 0.75 (95% CI: 0.60 to 0.91). These observations, albeit intriguing, remain to be confirmed in additional studies, especially in light of the emergence of other data suggesting that surrogates of abnormal atrial substrates (e.g., P-wave terminal force in V₁, intra-atrial block) are indeed predictive of incident stroke independent of the presence of AF (87,88).

The advances in artificial intelligence (AI) algorithms might also provide an additional opportunity to optimize stroke prevention in patients with AF. Recent studies have shown that AI permits identification of occult or recent AF on a simple 12-lead electrocardiogram in patients with sinus rhythm (89). This may prove useful for determining therapy in patients with embolic stroke of undetermined source. In patients with embolic stroke of undetermined source, or with an elevated CHA₂DS₂-VASc score, it is possible that a positive AI electrocardiogram indicating recent or impending AF may obviate the need for prolonged monitoring, and identify individuals who may benefit from stroke prevention therapies. This strategy has not been clinically tested and will require validation.

6. COMPETING RISKS. Patients with AF have a large burden of cardiovascular diseases that can cause

stroke regardless of AF. Those include: complex aortic plaque, carotid or intracranial arterial stenosis, uncontrolled hypertension, advanced left ventricular dysfunction, or structural heart defects (e.g., patent foramen ovale, atrial septal aneurysm) (90-93). Unfortunately, data on the relative contribution of these concomitant pathologies to stroke risk in patients with NVAF are scarce (87). In a prospective stroke registry, a concomitant potential source of cardiovascular atheroembolic events was present in 38.8% of patients (94). In another study, the presence of carotid stenosis in patients with NVAF was not uncommon and was associated with doubling in the risk of stroke after adjusting for risk factors and antithrombotic therapy (8.1 vs. 3.6 events/100 follow-up years; p = 0.005) (95).

CURRENT CHALLENGES AND FUTURE DIRECTIONS

It is evident that the currently used clinical risk scores for ischemic stroke prediction in NVAF (e.g., CHA₂DS₂-VASc) suffer from marked limitations. Yet, these scores continue to be the main determinant of eligibility for various stroke prevention strategies in current practice, highlighting the need for better stroke prediction tools. The weak performance of

the available risk scores might partially stem from their exclusion of key anatomic, physiological, and electrical factors that are now proven to have an independent value in identifying patients at risk for AF-related strokes. However, studies attempting to address this issue have mostly focused on assessing the association of a single variable and the odds of stroke (e.g., duration of AF, non-chicken wing LAA anatomy, abnormal P-wave axis, etc.). Hence, these studies are unlikely to have an impact on the current practice.

A study that provides a holistic assessment of clinical, anatomic, and other measurable factors that are associated with excess stroke risk in patients with NVAF is direly needed (**Central Illustration**). Nonetheless, conducting such a study faces significant logistical challenges. First, the absolute annual event rate in patients with NVAF who are not on anticoagulation is small (e.g., a CHA₂DS₂-VASc of 4 corresponds with an annual stroke risk of 4.8%, and an annual risk of SE of 6.7%). This suggests that a large number of patients and/or long-term follow-up are needed to avoid the proposed study being underpowered. Second, the majority of patients with NVAF are on various anticoagulation regimens with marked variations in compliance, which further complicates the risk assessment. Third, a comprehensive risk assessment will require multiple tests (transesophageal echo, computed tomography, CMR, loop recorders, etc.). This might not be feasible (e.g., cost) nor practical (e.g., presence of a pacemaker, intolerance of transesophageal echo) in many patients. Indeed, the strongest argument for the adoption of CHA₂DS₂-VASc is that it is simple and costless, which would not be the case with any risk models that incorporate imaging, electrocardiographic, or laboratory data. In addition, given the wide adoption of the simple CHA₂DS₂-VASc in current practice, concerns arise about the potential for more complicated scores to sway some clinicians (e.g., general practitioners) from prescribing appropriate anticoagulation (96). Fourth, some of the proven and suggested clinical (age, hypertension, diabetes, congestive heart failure) and nonclinical (LAA velocity, AF burden, biomarkers, etc.) risk factors for AF-associated stroke are dynamic (97). A single recording/measurement of those factors might underestimate or overestimate the actual risk of an individual patient. For example, well controlled and poorly controlled hypertension or diabetes are viewed equally in the current risk prediction models, although their association with a heightened risk of stroke might be remarkably different.

Despite these challenges, the unmet needs for a proper stroke risk assessment scheme in the growing population with NVAF mandates practical solutions. Machine learning techniques might provide an opportunity to better understand the interplay between various risk factors in large populations of patients with NVAF. Potential sources of data for such analyses may be the completed randomized trials of catheter ablation or LAAO, as those trials would have recorded multiple clinical, imaging, electrocardiographic, and laboratory variables and have adequate follow-up and credible adjudications of events. Nonetheless, the modifications of stroke risk introduced by those interventions (catheter ablation, LAAO) will be difficult to control for. Another source would be large single or multicenter databases of patients who were referred for cardioversion and underwent LAA imaging, although data completion, variability of follow-up, and ascertainment of clinical events may lead to major limitations. A prospective long-term observational study will ultimately be needed to arrive at meaningful conclusions. A proposed outline for such study is shown in **Figure 6**. The availability of novel tools such as smartphone-based echocardiography and the application of AI to interpretation of the images acquired may significantly lower the cost and improve the availability of additional important parameters useful for risk stratification at the bedside. Integrated electronic medical records may provide additional assistance, although to date, these have been wanting.

CONCLUSIONS

The prevalence of AF and the burden of its associated morbidities continue to rise worldwide. Despite the advances in ischemic stroke prevention strategies in these patients, our current stroke risk prediction tools remain basic and imprecise. There is an unmet need to incorporate various elements to the current risk factor-based stroke prediction schemes to optimize our stroke prevention practices. The time has come for a comprehensive study that provides a holistic approach to risk assessment in this high-risk population.

ADDRESS FOR CORRESPONDENCE: Dr. Mohamad Alkhouli, Department of Cardiovascular Medicine, Mayo Clinic School of Medicine, Mayo Clinic, 200 First Street SW, Rochester, Minnesota 55905. E-mail: Alkhouli.Mohamad@mayo.edu. Twitter: [@adnanalkhouli](https://twitter.com/adnanalkhouli), [@drpaulfriedman](https://twitter.com/drpaulfriedman).

REFERENCES

- Alkhouli M, Noseworthy PA, Rihal CS, Holmes DR Jr. Stroke prevention in nonvalvular atrial fibrillation: a stakeholder perspective. *J Am Coll Cardiol* 2018;71:2790-801.
- Schnabel RB, Yin X, Gona P, et al. 50 year trends in atrial fibrillation prevalence, incidence, risk factors, and mortality in the Framingham Heart Study: a cohort study. *Lancet* 2015;386:154-62.
- Chugh SS, Havmoeller R, Narayanan K, et al. Worldwide epidemiology of atrial fibrillation: a Global Burden of Disease 2010 Study. *Circulation* 2014;129:837-47.
- Alkhouli M, Alqahtani F, Aljohani S, Alvi M, Holmes DR. Burden of atrial fibrillation-associated ischemic stroke in the United States. *J Am Coll Cardiol EP* 2018;4:618-25.
- Ruff CT, Giugliano RP, Braunwald E, et al. Comparison of the efficacy and safety of new oral anticoagulants with warfarin in patients with atrial fibrillation: a meta-analysis of randomised trials. *Lancet* 2014;383:955-62.
- January CT, Wann LS, Calkins H, et al. 2019 AHA/ACC/HRS focused update of the 2014 AHA/ACC/HRS guideline for the management of patients with atrial fibrillation: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines and the Heart Rhythm Society. *J Am Coll Cardiol* 2019;74:104-32.
- Andrade JG, Verma A, Mitchell LB, et al. 2018 focused update of the Canadian Cardiovascular Society guidelines for the management of atrial fibrillation. *Can J Cardiol* 2018;34:1371-92.
- Kirchhof P, Benussi S, Kotecha D, et al. 2016 ESC guidelines for the management of atrial fibrillation developed in collaboration with EACTS. *Eur Heart J* 2016;37:2893-962.
- Quinn GR, Severdija ON, Chang Y, Singer DE. Wide variation in reported rates of stroke across cohorts of patients with atrial fibrillation. *Circulation* 2017;135:208-19.
- Yaghi S, Kamel H. Stratifying stroke risk in atrial fibrillation: beyond clinical risk scores. *Stroke* 2017;48:2665-70.
- Saunders J. The practice of clinical medicine as an art and as a science. *Med Humanit* 2000;26:18-22.
- Wolf PA, Dawber TR, Thomas HE Jr., Kannel WB. Epidemiologic assessment of chronic atrial fibrillation and risk of stroke: the Framingham study. *Neurology* 1978;28:973-7.
- Mitchell LB, Southern DA, Galbraith D, et al. Prediction of stroke or TIA in patients without atrial fibrillation using CHADS2 and CHA2DS2-VASc scores. *Heart* 2014;100:1524-30.
- Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation. Analysis of pooled data from five randomized controlled trials. *Arch Intern Med* 1994;154:1449-57.
- Hart RG, Pearce LA, Aguilar MI. Meta-analysis: antithrombotic therapy to prevent stroke in patients who have nonvalvular atrial fibrillation. *Ann Intern Med* 2007;146:857-67.
- van Walraven C, Hart RG, Singer DE, et al. Oral anticoagulants vs aspirin in nonvalvular atrial fibrillation: an individual patient meta-analysis. *JAMA* 2002;288:2441-8.
- The Atrial Fibrillation Investigators. The efficacy of aspirin in patients with atrial fibrillation. Analysis of pooled data from 3 randomized trials. *Arch Intern Med* 1997;157:1237-40.
- The Stroke Prevention in Atrial Fibrillation Investigators. Predictors of thromboembolism in atrial fibrillation: I. clinical features of patients at risk. *Ann Intern Med* 1992;116:1-5.
- Gage BF, Waterman AD, Shannon W, Boechler M, Rich MW, Radford MJ. Validation of clinical classification schemes for predicting stroke: results from the National Registry of Atrial Fibrillation. *JAMA* 2001;285:2864-70.
- Laupacis A, Albers G, Dalen J, Dunn MI, Jacobson AK, Singer DE. Antithrombotic therapy in atrial fibrillation. *Chest* 1998;114:579S-89S.
- Gage BF, Cardinali AB, Albers GW, Owens DK. Cost-effectiveness of warfarin and aspirin for prophylaxis of stroke in patients with nonvalvular atrial fibrillation. *JAMA* 1995;274:1839-45.
- Gullov AL, Koefoed BG, Petersen P. Bleeding during warfarin and aspirin therapy in patients with atrial fibrillation: the AFASAK 2 study. *Arch Intern Med* 1999;159:1322-8.
- Gullov AL, Koefoed BG, Petersen P. Bleeding complications to long-term oral anticoagulant therapy. *J Thromb Thrombolysis* 1994;1:17-25.
- Stroke Prevention in Atrial Fibrillation Investigators. Risk factors for thromboembolism during aspirin therapy in patients with atrial fibrillation: the stroke prevention in atrial fibrillation study. *J Stroke Cerebrovasc Dis* 1995;5:147-57.
- Lip GY, Nieuwlaet R, Pisters R, Lane DA, Crijns HJ. Refining clinical risk stratification for predicting stroke and thromboembolism in atrial fibrillation using a novel risk factor-based approach: the Euro Heart Survey on Atrial Fibrillation. *Chest* 2010;137:263-72.
- Singer DE, Fang MC, Go AS. CHA2DS2-VASc risk scheme: not ready for clinical use. *Chest* 2010;138:1020. author reply 1020-1.
- Chen JY, Zhang AD, Lu HY, Guo J, Wang FF, Li ZC. CHADS2 versus CHA2DS2-VASc score in assessing the stroke and thromboembolism risk stratification in patients with atrial fibrillation: a systematic review and meta-analysis. *J Geriatr Cardiol* 2013;10:258-66.
- Olesen JB, Torp-Pedersen C, Hansen ML, Lip GY. The value of the CHA2DS2-VASc score for refining stroke risk stratification in patients with atrial fibrillation with a CHADS2 score 0-1: a nationwide cohort study. *Thromb Haemost* 2012;107:1172-9.
- Olesen JB, Lip GY, Hansen ML, et al. Validation of risk stratification schemes for predicting stroke and thromboembolism in patients with atrial fibrillation: nationwide cohort study. *BMJ* 2011;342:d124.
- Van Staa TP, Setakis E, Di Tanna GL, Lane DA, Lip GY. A comparison of risk stratification schemes for stroke in 79,884 atrial fibrillation patients in general practice. *J Thromb Haemost* 2011;9:39-48.
- Wei-Syun H, Lin CL. CHA2DS2-VASc score for prediction of ischemic stroke in patients with systemic lupus erythematosus without atrial fibrillation. *Lupus* 2018;27:1240-6.
- Hu WS, Lin CL. CHA2DS2-VASc score for ischaemic stroke risk stratification in patients with chronic obstructive pulmonary disease with and without atrial fibrillation: a nationwide cohort study. *Europace* 2018;20:575-81.
- Wu JT, Wang SL, Chu YJ, et al. CHADS2 and CHA2DS2-VASc scores predict the risk of ischemic stroke outcome in patients with interatrial block without atrial fibrillation. *J Atheroscler Thromb* 2017;24:176-84.
- Chan YH, Yiu KH, Lau KK, et al. The CHADS2 and CHA2DS2-VASc scores predict adverse vascular function, ischemic stroke and cardiovascular death in high-risk patients without atrial fibrillation: role of incorporating PR prolongation. *Atherosclerosis* 2014;237:504-13.
- Piccini JP, Stevens SR, Chang Y, et al. Renal dysfunction as a predictor of stroke and systemic embolism in patients with nonvalvular atrial fibrillation: validation of the R(2)CHADS(2) index in the ROCKET AF (Rivaroxaban Once-daily, oral, direct factor Xa inhibition Compared with vitamin K antagonism for prevention of stroke and Embolism Trial in Atrial Fibrillation) and ATRIA (AnTicoagulation and Risk factors In Atrial fibrillation) study cohorts. *Circulation* 2013;127:224-32.
- Cha MJ, Cho Y, Oh IY, Choi EK, Oh S. Validation of conventional thromboembolic risk factors in a Korean atrial fibrillation population: suggestion for a novel scoring system, CHA2DS2-VAK. *Circ J* 2018;82:2970-5.
- Chao TF, Lip GY, Liu CJ, et al. Validation of a modified CHA2DS2-VASc score for stroke risk stratification in Asian Patients with atrial fibrillation: a nationwide cohort study. *Stroke* 2016;47:2462-9.
- Fox KAA, Lucas JE, Pieper KS, et al. Improved risk stratification of patients with atrial fibrillation: an integrated GARFIELD-AF tool for the prediction of mortality, stroke and bleed in patients with and without anticoagulation. *BMJ Open* 2017;7:e017157.
- Hijazi Z, Lindback J, Alexander JH, et al. The ABC (age, biomarkers, clinical history) stroke risk score: a biomarker-based risk score for predicting stroke in atrial fibrillation. *Eur Heart J* 2016;37:1582-90.
- Kabra R, Girotra S, Vaughan Sarrazin M. Refining stroke prediction in atrial fibrillation patients by addition of African-American ethnicity to CHA2DS2-VASc score. *J Am Coll Cardiol* 2016;68:461-70.

41. Singer DE, Chang Y, Borowsky LH, et al. A new risk scheme to predict ischemic stroke and other thromboembolism in atrial fibrillation: the ATRIA study stroke risk score. *J Am Heart Assoc* 2013;2:e000250.
42. Tomita H, Okumura K, Inoue H, et al. Validation of risk scoring system excluding female sex from CHA2DS2-VASc in Japanese patients with nonvalvular atrial fibrillation: subanalysis of the J-RHYTHM registry. *Circ J* 2015;79:1719-26.
43. Parsons C, Patel SJ, Cha S, et al. CHA2DS2-VASc score: a predictor of thromboembolic events and mortality in patients with an implantable monitoring device without atrial fibrillation. *Mayo Clin Proc* 2017;92:360-9.
44. Cha MJ, Park HE, Lee MH, Cho Y, Choi EK, Oh S. Prevalence of and risk factors for silent ischemic stroke in patients with atrial fibrillation as determined by brain magnetic resonance imaging. *Am J Cardiol* 2014;113:655-61.
45. Hohnloser SH, Pajitnev D, Pogue J, et al. Incidence of stroke in paroxysmal versus sustained atrial fibrillation in patients taking oral anticoagulation or combined antiplatelet therapy: an ACTIVE W substudy. *J Am Coll Cardiol* 2007;50:2156-61.
46. Hart RG, Pearce LA, Rothbart RM, McAnulty JH, Asinger RW, Halperin JL, Stroke Prevention in Atrial Fibrillation Investigators. Stroke with intermittent atrial fibrillation: incidence and predictors during aspirin therapy. *J Am Coll Cardiol* 2000;35:183-7.
47. Link MS, Giugliano RP, Ruff CT, et al., ENGAGE AF-TIMI 48 Investigators. Stroke and mortality risk in patients with various patterns of atrial fibrillation: results from the ENGAGE AF-TIMI 48 trial (Effective Anticoagulation With Factor Xa Next Generation in Atrial Fibrillation-Thrombolysis in Myocardial Infarction 48). *Circ Arrhythm Electrophysiol* 2017;10:e004267.
48. Ganesan AN, Chew DP, Hartshorne T, et al. The impact of atrial fibrillation type on the risk of thromboembolism, mortality, and bleeding: a systematic review and meta-analysis. *Eur Heart J* 2016;37:1591-602.
49. Lilli A, Di Cori A, Zaca V. Thromboembolic risk and effect of oral anticoagulation according to atrial fibrillation patterns: a systematic review and meta-analysis. *Clin Cardiol* 2017;40:641-7.
50. Van Gelder IC, Healey JS, Crijns H, et al. Duration of device-detected subclinical atrial fibrillation and occurrence of stroke in ASSERT. *Eur Heart J* 2017;38:1339-44.
51. Go AS, Reynolds K, Yang J, et al. Association of burden of atrial fibrillation with risk of ischemic stroke in adults with paroxysmal atrial fibrillation: the KP-RHYTHM study. *JAMA Cardiol* 2018;3:601-8.
52. Chen LY, Chung MK, Allen LA, et al. Atrial fibrillation burden: moving beyond atrial fibrillation as a binary entity: a scientific statement from the American Heart Association. *Circulation* 2018;137:e623-44.
53. Zimetbaum P, Ellis ER, Waks JW, Passman RS. The importance of atrial fibrillation burden and the origin of device-tailored anticoagulation. *Pacing Clin Electrophysiol* 2013;36:1319-24.
54. Benjamin EJ, D'Agostino RB, Belanger AJ, Wolf PA, Levy D. Left atrial size and the risk of stroke and death. The Framingham Heart Study. *Circulation* 1995;92:835-41.
55. Tiwari S, Lochen ML, Jacobsen BK, et al. CHA2DS2-VASc score, left atrial size and atrial fibrillation as stroke risk factors in the Tromsø Study. *Open Heart* 2016;3:e000439.
56. Hamatani Y, Ogawa H, Takabayashi K, et al. Left atrial enlargement is an independent predictor of stroke and systemic embolism in patients with non-valvular atrial fibrillation. *Sci Rep* 2016;6:31042.
57. The Stroke Prevention in Atrial Fibrillation Investigators. Predictors of thromboembolism in atrial fibrillation: II. echocardiographic features of patients at risk. *Ann Intern Med* 1992;116:6-12.
58. Ogata T, Matsuo R, Kiyuna F, et al. Left atrial size and long-term risk of recurrent stroke after acute ischemic stroke in patients with nonvalvular atrial fibrillation. *J Am Heart Assoc* 2017;6:e006402.
59. Yaghi S, Moon YP, Mora-McLaughlin C, et al. Left atrial enlargement and stroke recurrence: the Northern Manhattan Stroke Study. *Stroke* 2015;46:1488-93.
60. Marrouche NF, Wilber D, Hindricks G, et al. Association of atrial tissue fibrosis identified by delayed enhancement MRI and atrial fibrillation catheter ablation: the DECAAF study. *JAMA* 2014;311:498-506.
61. Leong DP, Joyce E, Debonnaire P, et al. Left atrial dysfunction in the pathogenesis of cryptogenic stroke: novel insights from speckle-tracking echocardiography. *J Am Soc Echocardiogr* 2017;30:71-9.e1.
62. Daccarett M, Badger TJ, Akoum N, et al. Association of left atrial fibrosis detected by delayed-enhancement magnetic resonance imaging and the risk of stroke in patients with atrial fibrillation. *J Am Coll Cardiol* 2011;57:831-8.
63. Habibi M, Zareian M, Ambale Venkatesh B, et al. Left atrial mechanical function and incident ischemic cerebrovascular events independent of AF: insights from the MESA study. *J Am Coll Cardiol* 2019. Apr 17 [E-pub ahead of print].
64. Blackshear JL, Odell JA. Appendage obliteration to reduce stroke in cardiac surgical patients with atrial fibrillation. *Ann Thorac Surg* 1996;61:755-9.
65. Yaghi S, Song C, Gray WA, Furie KL, Elkind MS, Kamel H. Left atrial appendage function and stroke risk. *Stroke* 2015;46:3554-9.
66. Lee JM, Shim J, Uhm JS, et al. Impact of increased orifice size and decreased flow velocity of left atrial appendage on stroke in nonvalvular atrial fibrillation. *Am J Cardiol* 2014;113:963-9.
67. Fatkin D, Kelly RP, Feneley MP. Relations between left atrial appendage blood flow velocity, spontaneous echocardiographic contrast and thromboembolic risk in vivo. *J Am Coll Cardiol* 1994;23:961-9.
68. Goldman ME, Pearce LA, Hart RG, et al. Pathophysiologic correlates of thromboembolism in nonvalvular atrial fibrillation: I. Reduced flow velocity in the left atrial appendage (The Stroke Prevention in Atrial Fibrillation [SPAF-III] study). *J Am Soc Echocardiogr* 1999;12:1080-7.
69. Kamp O, Verhorst PM, Welling RC, Visser CA. Importance of left atrial appendage flow as a predictor of thromboembolic events in patients with atrial fibrillation. *Eur Heart J* 1999;20:979-85.
70. Akoum N, Marrouche N. Assessment and impact of cardiac fibrosis on atrial fibrillation. *Curr Cardiol Rep* 2014;16:518.
71. Di Biase L, Santangeli P, Anselmino M, et al. Does the left atrial appendage morphology correlate with the risk of stroke in patients with atrial fibrillation? Results from a multicenter study. *J Am Coll Cardiol* 2012;60:531-8.
72. Lupercio F, Carlos Ruiz J, Briceno DF, et al. Left atrial appendage morphology assessment for risk stratification of embolic stroke in patients with atrial fibrillation: a meta-analysis. *Heart Rhythm* 2016;13:1402-9.
73. Yaghi S, Chang AD, Akiki R, et al. The left atrial appendage morphology is associated with embolic stroke subtypes using a simple classification system: a proof of concept study. *J Cardiovasc Comput Tomogr* 2019 Apr 16. Apr 16 [E-pub ahead of print].
74. Hylek EM. Biomarkers for prediction of stroke and bleeds in atrial fibrillation. *Circulation* 2019;139:772-4.
75. Koenig W. High-sensitivity C-reactive protein and atherosclerotic disease: from improved risk prediction to risk-guided therapy. *Int J Cardiol* 2013;168:5126-34.
76. Hijazi Z, Oldgren J, Andersson U, et al. Cardiac biomarkers are associated with an increased risk of stroke and death in patients with atrial fibrillation: a Randomized Evaluation of Long-term Anticoagulation Therapy (RE-LY) substudy. *Circulation* 2012;125:1605-16.
77. Hijazi Z, Wallentin L, Siegbahn A, et al. High-sensitivity troponin T and risk stratification in patients with atrial fibrillation during treatment with apixaban or warfarin. *J Am Coll Cardiol* 2014;63:52-61.
78. Hijazi Z, Wallentin L, Siegbahn A, et al. N-terminal pro-B-type natriuretic peptide for risk assessment in patients with atrial fibrillation: insights from the ARISTOTLE Trial (Apixaban for the Prevention of Stroke in Subjects With Atrial Fibrillation). *J Am Coll Cardiol* 2013;61:2274-84.
79. Vafaie M, Giannitsis E, Mueller-Hennessen M, et al. High-sensitivity cardiac troponin T as an independent predictor of stroke in patients admitted to an emergency department with atrial fibrillation. *PLoS One* 2019;14:e0212278.
80. Cushman M, Judd SE, Howard VJ, et al. N-terminal pro-B-type natriuretic peptide and stroke risk: the reasons for geographic and racial differences in stroke cohort. *Stroke* 2014;45:1646-50.
81. Lip GY, Patel JV, Hughes E, Hart RG. High-sensitivity C-reactive protein and soluble CD40 ligand as indices of inflammation and platelet

activation in 880 patients with nonvalvular atrial fibrillation: relationship to stroke risk factors, stroke risk stratification schema, and prognosis. *Stroke* 2007;38:1229-37.

82. Aulin J, Siegbahn A, Hijazi Z, et al. Interleukin-6 and C-reactive protein and risk for death and cardiovascular events in patients with atrial fibrillation. *Am Heart J* 2015;170:1151-60.

83. Hijazi Z, Aulin J, Andersson U, et al. Biomarkers of inflammation and risk of cardiovascular events in anticoagulated patients with atrial fibrillation. *Heart* 2016;102:508-17.

84. Maheshwari A, Norby FL, Roetker NS, et al. Refining prediction of atrial fibrillation-related stroke using the P2-CHA2DS2-VASc score. *Circulation* 2019;139:180-91.

85. He J, Tse G, Korantzopoulos P, et al. P-wave indices and risk of ischemic stroke: a systematic review and meta-analysis. *Stroke* 2017;48:2066-72.

86. Martinez-Selles M, Baranchuk A, Elosua R, de Luna AB. Rationale and design of the BAYES (Interatrial Block and Yearly Events) registry. *Clin Cardiol* 2017;40:196-9.

87. Kamel H, Okin PM, Elkind MS, Iadecola C. Atrial fibrillation and mechanisms of stroke: time for a new model. *Stroke* 2016;47:895-900.

88. Skov MW, Ghouse J, Kuhl JT, et al. Risk prediction of atrial fibrillation based on electrocardiographic interatrial block. *J Am Heart Assoc* 2018;7:e008247.

89. Attia ZI, Kapa S, Lopez-Jimenez F, et al. Screening for cardiac contractile dysfunction using an artificial intelligence-enabled electrocardiogram. *Nat Med* 2019;25:70-4.

90. Hart RG, Pearce LA, Miller VT, et al. Cardioembolic vs. noncardioembolic strokes in atrial fibrillation: frequency and effect of antithrombotic agents in the stroke prevention in atrial fibrillation studies. *Cerebrovasc Dis* 2000;10:39-43.

91. Katsi V, Georgiopoulos G, Skafida A, et al. Noncardioembolic stroke in patients with atrial fibrillation. *Angiology* 2019;70:299-304.

92. Yang PS, Pak HN, Park DH, et al. Non-cardioembolic risk factors in atrial fibrillation-associated ischemic stroke. *PLoS One* 2018;13:e0201062.

93. Paciaroni M, Agnelli G, Caso V, et al. Causes and risk factors of cerebral ischemic events in patients with atrial fibrillation treated with non-vitamin K antagonist oral anticoagulants for stroke prevention. *Stroke* 2019;50:2168-74.

94. Cha MJ, Kim YD, Nam HS, Kim J, Lee DH, Heo JH. Stroke mechanism in patients with non-valvular atrial fibrillation according to the CHADS2 and CHA2 DS2-VASc scores. *Eur J Neurol* 2012;19:473-9.

95. Lehtola H, Airaksinen KEJ, Hartikainen P, et al. Stroke recurrence in patients with atrial fibrillation: concomitant carotid artery stenosis doubles the risk. *Eur J Neurol* 2017;24:719-25.

96. Gage BF. Stroke prediction rules in atrial fibrillation. *J Am Coll Cardiol* 2018;71:133-4.

97. Chao TF, Lip GYH, Liu CJ, et al. Relationship of aging and incident comorbidities to stroke risk in patients with atrial fibrillation. *J Am Coll Cardiol* 2018;71:122-32.

KEY WORDS atrial fibrillation, oral anticoagulation, risk assessment, stroke



Go to <http://www.acc.org/jacc-journals-cme> to take the CME/MOC/ECME quiz for this article.