

Depressive Symptoms Predict Heart Rate Recovery After Exercise Treadmill Testing in Patients With Coronary Artery Disease: Results From the Psychophysiological Investigations of Myocardial Ischemia Study

JOEL W. HUGHES, PhD, KAKI M. YORK, PhD, QIN LI, MA, KENNETH E. FREEDLAND, PhD, ROBERT M. CARNEY, PhD, AND DAVID S. SHEPS, MD, MSPH

Background: Depression is associated with increased risk of death among patients with coronary disease. Cardiovascular autonomic dysregulation may be one of the mechanisms by which depression exerts its effects on cardiovascular function. The purpose of this study was to determine whether depressive symptoms are associated with low heart rate variability (HRV) and prolonged HR recovery after exercise testing in patients with coronary artery disease (CAD). **Methods:** The Psychophysiological Investigation of Myocardial Ischemia (PIMI) was a large, multicenter study designed to assess psychological and physiological correlates of stress in patients with CAD. One hundred and eighty-eight patients with CAD as evidenced by at least 50% blockage of one major artery and a previous positive exercise stress test were included in this study. Patients included in this report were not taking beta blockers. Cardiovascular functioning was assessed by a modified Bruce protocol treadmill stress test. Measures of psychological functioning, including the Beck Depression Inventory (BDI), were also obtained. **Results:** BDI scores were negatively correlated with HR recovery ($r = -0.15, p = .04$). Depression scores accounted for 3.5% of the variance in HR recovery when controlling for participant age ($p < .01$). Depressive symptoms were related to two HRV indices (ultra-low frequency, high frequency). **Conclusions:** Depressive symptoms are associated with cardiovascular autonomic nervous system dysfunction as assessed by HR recovery. This relationship is not merely due to an association of depression severity with beta blocker usage or a failure of depressed patients to achieve an adequate chronotropic response. **Key words:** coronary disease, depression, heart rate recovery, autonomic nervous system.

ANS = autonomic nervous system; CAD = coronary artery disease; HR = heart rate; HRV = heart rate variability; PIMI = Psychophysiological Investigation of Myocardial Ischemia

INTRODUCTION

Depression is associated with increased risk of death among patients with coronary disease (1–10). For example, the presence of depression during or shortly after hospitalization for myocardial infarction confers a twofold to threefold increase in risk for mortality or nonfatal cardiac events. In the Enhancing Recovery in Coronary Heart Disease (ENRICHD) trial, however, psychological treatment had modest effects on depression and social support but did not reduce risk of mortality (11).

The results of the ENRICHD trial underscore the importance of identifying the mechanisms whereby depression is associated with increased risk. A number of possible mechanisms have been investigated, including associations with other cardiac risk factors (e.g., alcohol consumption, smoking) and poor adherence to treatment recommendations and lifestyle modification (e.g., medication adherence, regular exercise) (12–15). One that has received considerable empirical support is autonomic nervous system (ANS) dysregulation. A

growing body of literature documents compromised ANS functioning among cardiac patients. Heart rate variability (HRV) during ambulatory monitoring has been used to assess cardiovascular autonomic dysregulation in a number of studies (16–19). HRV has been shown to predict mortality among cardiac patients (20–22), and it was recently shown to partially mediate the relationship between depression and mortality in an ENRICHD ancillary study (23).

The ANS also modulates the rate at which HR returns to resting levels after the cessation of exercise (HR recovery). The rapid decrease in HR after exercise is predominantly accomplished by parasympathetic reactivation, making HR recovery a marker of parasympathetic control of the heart (24–28). Prolonged HR recovery during exercise treadmill testing (ETT) has been shown to predict mortality (25,26,29–37). In one study, for example, prolonged HR recovery was associated with increased risk of sudden cardiac death over a 23-year follow-up period among healthy men after controlling for several potential confounders (38). Recently, symptoms of depression were shown to be associated with slower HR recovery from ETT among 260 patients starting cardiac rehabilitation (39). After controlling for age, sex, and beta-blocker usage, Beck Depression Inventory (BDI) (40) scores were associated with slower HR recovery after maximal exercise on an ETT. Furthermore, estimated METS from the ETT helped to explain the association of depressive symptoms and HR recovery. This is consistent with the hypothesis that depression leads to physical inactivity and low aerobic capacity, which then contributes to ANS dysregulation. Although the results of this study were promising, the fact that over 80% of the sample was taking beta-blockers was a serious limitation.

Therefore, the purpose of this investigation was to replicate the association of depressive symptoms and HR recovery in patients who were not taking beta blockers or who were temporarily withdrawn from beta blockers before testing. In keeping with previous findings, we hypothesized that depres-

From the Kent State University, Kent, Ohio (J.W.H.); University of Florida, and North Florida/South Georgia VAMC, Gainesville, Florida (K.M.Y., Q.L., D.S.S.); and Washington University School of Medicine, St. Louis, Missouri (K.E.F., R.M.C.).

Address correspondence and reprint requests to Joel W. Hughes, PhD, Department of Psychology, P.O. Box 5190, Kent, OH 44242. E-mail: jhughes1@kent.edu.

Received for publication January 5, 2007; revision received August 8, 2007.

Supported by contracts HV 18114, HV 18119, HV 18120, HV 18121, and HV 28127 of the National Heart Lung and Blood Institute.

Because co-authors Kenneth E. Freedland and David S. Sheps are editors of this journal, the review of this paper was overseen by a guest editor. Drs. Freedland and Sheps were not involved in the decision-making process and, like all authors, were blinded to the identity of the peer reviewers.

DOI: 10.1097/PSY.0b013e31816fcab3

DEPRESSION AND HR RECOVERY

sive symptoms are inversely related to HR recovery from ETT. We also hypothesized that physical fitness, estimated from total exercise time during the ETT, help to account for the relationship between depression and HR recovery.

As a secondary aim, we evaluated the relationship between depressive symptoms and HRV measures derived from 48 hours of Holter monitoring. HRV measures are complementary to HR recovery, and there are far more reports of a relationship between depression symptoms and HRV. However, few studies have incorporated both of these measures of ANS regulation. Employing both measures provided a basis for evaluating whether similar relationships with depression would be observed. Finally, we explored the relationship between HR recovery, and HRV measures from 48 hours of Holter monitoring.

METHODS

These data are from the Psychophysiological Investigations of Myocardial Ischemia (PIMI) study and were collected during 1993 through 1994. The PIMI methodology is described more fully in an earlier report (41).

Participants

Participants in the PIMI study were originally recruited from clinical sites of the Asymptomatic Cardiac Ischemia Pilot (ACIP) study (42). Eligibility criteria included having verified coronary artery disease, a history of myocardial infarction, or evidence of myocardial ischemia on an exercise treadmill test while off medications (43). Exclusion criteria for the PIMI study included pregnancy, myocardial infarction within 3 months of the ETT, percutaneous transluminal coronary angioplasty (PTCA) within 6 months of the qualifying ETT, cardiac surgery requiring thoracotomy, unstable angina within 4 weeks of the qualifying ETT, serious noncardiac illness, inability to discontinue medications before assessments, and neurologic disease (e.g., stroke, dementia). For these analyses, three participants taking beta-blockers were also excluded. Four participants were taking antidepressants and were included in the analyses. The protocol was approved by the institutional review board at each participating medical center. Informed consent was obtained from all participants.

The current report includes data from 188 patients (162 men and 26 women). Participants were predominantly white (87%). Additional patient characteristics are reported in Table 1.

TABLE 1. Clinical and Demographic Characteristics of Study Population

	N	Percent
Age (years)		
41–50	17	9
51–60	44	24
61–70	87	46
71–80	39	21
Gender (male)	162	86
Race (Caucasian)	164	87
Current smoker	32	17
History of diabetes	28	15
History of MI	80	43
Medications		
Beta blockers	0	0
Calcium channel blocker	3	2
Digitalis	0	0
Nitrates	0	0

Procedures

The participants were asked to stop antianginal medications before all study assessments, for a period ranging in duration from 1 hour (e.g., sublingual nitroglycerin) to 3 days or five half-lives, whichever was longer (e.g., beta blockers). They were asked to discontinue sedatives and analgesics for 24 hours, and other medications (digitalis, theophylline, steroids, ACE inhibitors, antipsychotic medications) for 2 to 4 weeks. They were also asked to fast and to abstain from nicotine products for 12 hours before testing.

Exercise Stress Testing

An ETT was performed during the morning hours with patients off antianginal medications except for sublingual nitroglycerin, if needed. Resting HR, maximum HR, exercise capacity, and HR recovery measures were derived from the ETT results. The treadmill exercise test was conducted according to a modified ACIP protocol (43) using standard criteria for stopping. Heart rate and BP were recorded during each exercise stage and for each minute of recovery. HR recovery was defined as the difference in HR from maximum HR (HR at exercise termination) to HR 1 minute later (25,29). Exercise capacity in METS was not recorded in the PIMI database, and so total exercise time was used as an estimate of physical fitness. Exercise duration has been used as an estimate of physical fitness in other studies that also employed oxygen consumption at peak exercise (44,45).

Uniform interpretation and execution of this and other aspects of the PIMI study protocol were facilitated by training, pilot testing, and certification of study staff based on performance criteria. All data were centrally reviewed and entered by the PIMI Clinical Coordinating Center. Results of ETT were sent to their respective core laboratories for blinded analysis.

HRV From Ambulatory ECG Monitoring

After the ETT, the patients were asked to wear an Applied Cardiac Systems AM cassette AMBULATORY ECG (AECG) monitor (Laguna Beach, California) for 48 hours, producing two 24-hour tapes. The recordings were analyzed at the PIMI AECG Core Laboratory at the Brigham and Women's Hospital, Harvard Medical School, Boston, Massachusetts. Standard 24-hour time and frequency domain HRV indices were obtained. The frequency domain indices are reported as the natural log of milliseconds squared. Specifically, the ultra-low frequency (ULF; 0.0001–0.003 Hz), very-low frequency (VLF; 0.003–0.04 Hz), low frequency (LF; 0.04–0.15 Hz), and high-frequency (HF; 0.15–0.4 Hz) components of HRV are reported. The values from tape 1 and tape 2 were averaged to increase reliability. HRV data were available for 163 of the 188 participants in this study. Missing data were related to poor recording quality, and the 25 patients with missing HRV data were not different from the larger sample with respect to study variables (i.e., age, gender, race, smoking status, history of diabetes, resting HR, peak HR, HR recovery, BDI score, and total exercise time).

Depressive Symptoms

The BDI (40) is a 21-item questionnaire used to assess symptoms of depression. Each item consists of four statements representing increasing degrees of severity with scores ranging from 0 to 3. Patients select the statement that best described themselves in the previous 2 weeks. Scores can range from 0 to 63, and a cutoff score of 10 or higher is widely used to screen for clinically significant depression (40).

Data Analysis

All statistical analyses were performed using SAS 9.1 software (SAS, Cary, NC). The criterion for statistical significance was set at 0.05 per analysis. The distribution of HR recovery values was skewed, so they were square-root transformed to normalize the distribution. The primary analyses, evaluating the relationship between depression and HR recovery, consisted of bivariate correlations and hierarchical multiple linear regression analyses. Age was included in the regression models as a control variable because increasing age is associated with reductions in parasympathetic cardiac control (46). Resting and maximum HR were also included as covariates. The potential contribution of total exercise time to HR recovery was evaluated as an estimate of physical fitness. The relationships among depression, HR

TABLE 2. Beck Depression Inventory Score, Resting Heart Rate, Maximum HR, HR Recovery, and Total Exercise Time(s): Correlations and Descriptive Statistics (*N* = 188)

	BDI	Resting HR		Peak HR		HR Recovery		Exercise Time(s)	
		<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>
BDI	—	0.07	.38	−0.02	.78	−0.15	.04	−0.11	.12
Resting HR				0.40	.001	−0.32	.001	−0.14	.05
Peak HR				—		0.00	.96	0.38	.001
HR recovery						—		0.22	.002
M	6.67	79.3		138.0		24.0		464.26	
SD	5.7	13.7		18.3		3.0		178.6	
Range	0–34	43–123		89–187		2–71		157–1178	

HR recovery is transformed in correlation analyses.

BDI = Beck Depression Inventory; HR = heart rate; HRR = heart rate recovery; M = mean; SD = 1 standard deviation.

recovery, and HRV from AECG were explored in correlation and regression analyses.

RESULTS

Sample characteristics are presented in Table 1 for the sample as a whole. Table 2 summarizes BDI scores and HR variables. BDI scores ranged from 0 to 34 ($M = 6.36$, interquartile range = 2.5–8.5) and 18% had scores ≥ 10 , suggesting the possibility of clinically significant depression. Total exercise time ranged from 157 to 1178 seconds ($M = 4635.2$, $SD = 179.8$). HR recovery at 1-minute postexercise averaged 23.9 seconds ($SD = 13.0$).

Correlations

Univariate correlation analyses are presented in Table 2. Depression scores were not associated with total exercise time ($r = -0.12$, $p = .12$), nor were they associated with resting or peak HR. Depression scores were negatively correlated with HR recovery ($r = -0.15$, $p = .04$).

Regression Analyses

Hierarchical multiple linear regression analyses were performed to further evaluate the relationship between depressive symptoms and HR recovery. The first step regressed HR recovery on age, resting HR, and maximum HR. The resulting equation accounted for 15.6% of the variance in HR recovery (adjusted $R^2 = 0.14$, $F(4,182) = 8.46$, $p < .001$). Significant effects were found for age ($\beta = -0.19$) and resting HR ($\beta = -0.39$). When BDI scores were added in the second step, the model accounted for 18.3% of the variance in HR recovery

(adjusted $R^2 = 0.16$, $F(5,181) = 8.1$, $p < .001$), and BDI scores explained 2.6% of the variance in HR recovery ($\Delta R^2 = 0.026$, $F(1,181) = 5.18$, $p = .02$). In an alternate model, when total exercise time was added to the model including age, resting HR, and peak HR, total exercise time did not explain additional variability in HR recovery ($\Delta R^2 = 0.01$, $F(1,181) = 2.14$, $p = .15$). When BDI score was added to this model, it explained an additional 2.3% of the variability in HR recovery ($\Delta R^2 = 0.023$, $F(1,180) = 5.15$, $p = .02$). Thus, the prior inclusion of total exercise time did not eliminate the relationship between depressive symptoms and HR recovery. The final model including age, resting HR, peak HR, total exercise time, and BDI score explained a total of 19% of the variability in HR recovery (adjusted $R^2 = 0.16$, $F(6,180) = 7.02$, $p < .001$). These results were not appreciably altered by controlling for current smoking status, diabetes, or history of myocardial infarction, nor were they altered by excluding the individuals taking antidepressants.

Relationship of Depressive Symptoms and HR Recovery With HRV

Table 3 reports correlations in the 163 participants with complete data for both HRV and HR recovery measures. Regression analyses followed the same procedures as above. In the first step, age was entered as a control variable. In the second step, depression score (or HR recovery value) was added to evaluate the additional variance explained. After controlling for age, depression symptoms were related to ULF ($\Delta R^2 = 0.038$, $F(1,160) = 6.4$, $p = .01$) and HF ($\Delta R^2 =$

TABLE 3. HR Recovery, Total Exercise Time(s), and HRV: Correlations and Descriptive Statistics (*N* = 163)

HRV (ln ms ²)	ULF		VLF		LF		HF		Ratio	
	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>	<i>r</i>	<i>p</i>	<i>R</i>	<i>p</i>	<i>R</i>	<i>p</i>
BDI	−0.17	.025	−0.10	.20	−0.09	.23	−0.15	.053	0.09	.27
HR recovery	0.31	.001	0.34	.001	0.24	.002	0.18	.02	−0.03	.67
Exercise time(s)	0.21	.006	0.23	.003	0.14	.08	0.04	.58	0.05	.53

HR recovery is transformed in correlation analyses. HRV is log transformed, with the exception of the HRV ratio.

BDI = Beck Depression Inventory; HR = heart rate; HRR = heart rate recovery; M = mean; SD = 1 standard deviation.

DEPRESSION AND HR RECOVERY

0.024, $F(1,159) = 3.93$, $p = .04$). Depression scores were not related to the VLF or LF components of HRV or the LF/HF ratio. After controlling for age, HR recovery predicted 9.4% of the variability in ULF, 11.1% of the variability in VLF, 4.9% of the variability in LF and 3.3% of the variability in HF (F values >4.0 , p values $<.05$). In contrast, HR recovery was not related to the LF/HF ratio ($F = 0.4$, $p = .52$).

DISCUSSION

Here, we report that depressive symptoms are associated with HR recovery, after controlling for age, resting HR, and peak HR. Although these associations were not large, these results are similar to those reported by Hughes et al. (39), and extend the findings to a group of patients who were not taking beta-blockers. Specifically, beta blockers had been withdrawn for five half-lives from those patients prescribed beta blockers. However, it is possible that were residual effects of beta blocker usage or possibly HR rebound due to withdrawal. Nevertheless, these findings suggest that the relationship between depressive symptoms and HR recovery is not merely due to an association of depression severity with beta blocker usage or a failure of patients with more depressive symptoms to achieve an adequate chronotropic response. Although the magnitude of the relationship was not very strong, with depression scores accounting for about 2.6% of the variability in HR recovery, depressive symptoms were as powerful a predictor of HR recovery as were age and total exercise time.

Controlling for the length of time that patients exercised during the ETT did not eliminate the relationship between depressive symptoms and HR recovery. It was expected that total exercise time, as an estimate of physical fitness (44,45), would help to explain the relationship between depressive symptoms and HR recovery. However, in this sample, depressive symptoms were not associated with total exercise time, which contrasts with other reports that higher depression scores were associated with lower estimated METS (39,47). These results suggest that mechanisms besides reduced physical activity (48,49) and poor physical fitness are involved in the relationship between depressive symptoms and HR recovery.

We also found that depressive symptoms were associated with two HRV indices from two 24-hour Holter monitoring periods, which is similar to the typical finding that depressive symptoms are associated with lower HRV (16–19). For example, Carney et al. (23) recently reported that clinical depression was associated with lower HRV, and also that HRV accounts for 27% of the variance in risk of mortality associated with depression. Our findings are not completely consistent, as depression was associated with ULF and HF, but not VLF or LF. This may be due to the restricted range of depressive symptoms in our sample and the relatively small sample. Specifically, among the 166 individuals for whom HRV data were available, only 26 had BDI scores of 10 or greater. Because the study employed a relatively small sample, it may have been underpowered to detect a relationship between depression and HRV.

Considered together with our finding that depressive symptoms are related to HR recovery, it is also possible that HR recovery is more a more sensitive indicator of the relationship between depression and ANS functioning than VLF and HF in some contexts (e.g., in patients who show evidence of myocardial ischemia on an exercise treadmill test while off medications). Studies with larger samples and a broader range of depressive symptoms are needed to evaluate the relative strengths of association among depression, HRV, and HR recovery.

We found that HR recovery is associated with the ULF, VLF, LF, and HF components of HRV but not with the ratio of LF to HF. One other study found that HR recovery at 1 minute was positively correlated with LF and uncorrelated with HF among elderly men (50), but to our knowledge, there are no other published reports on the relationship between HR recovery and HRV indices from 24-hour monitoring. Although more research is necessary, these results imply that HR recovery and HRV are complementary measures. HR recovery from ETT is easily obtained from routine assessments of cardiac patients (e.g., at enrollment in cardiac rehabilitation), and thus may be a convenient, inexpensive, and valuable measure to employ in studies of depression and autonomic functioning among cardiac patients.

CONCLUSION

The results of this study suggest that depressive symptoms are associated with ANS dysfunction as indicated by prolonged HR recovery after ETT, when controlling for age and gender. HR recovery may be a particularly sensitive marker of ANS dysregulation among patients with depressive symptoms.

This material is the result of work supported with resources and the use of facilities at the Department of Veterans Affairs Medical Center, Gainesville, FL.

REFERENCES

1. Barefoot JC, Helms MJ, Mark DB, Blumenthal JA, Califf RM, Haney TL, O'Connor CM, Siegler IC, Williams RB. Depression and long-term mortality risk in patients with coronary artery disease. *Am J Cardiol* 1996;78:613–7.
2. Carney RM, Rich MW, Freedland KE, Saini J, teVelde A, Simeone C, Clark K. Major depressive disorder predicts cardiac events in patients with coronary artery disease. *Psychosom Med* 1988;50:627–33.
3. Ahern DK, Gorkin L, Anderson JL, Tierney C, Hallstrom A, Ewart C, Capone RJ, Schron E, Kornfeld D, Herd JA. Biobehavioral variables and mortality or cardiac arrest in the Cardiac Arrhythmia Pilot Study (CAPS). *Am J Cardiol* 1990;66:59–62.
4. Frasure-Smith N, Lesperance F. Reflections on depression as a cardiac risk factor. *Psychosom Med* 2005;67(Suppl 1):S19–S25.
5. Frasure-Smith N, Lesperance F, Talajic M. Depression and 18-month prognosis after myocardial infarction. *Circulation* 1995;91:999–1005.
6. Frasure-Smith N, Lesperance F, Talajic M. Depression following myocardial infarction. Impact on 6-month survival. *JAMA* 1993;270:1819–25.
7. Lesperance F, Frasure-Smith N. Negative emotions and coronary heart disease: getting to the heart of the matter. [Letter; comment.]. *Lancet* 1996;347:414–5.
8. Lesperance F, Frasure-Smith N, Talajic M, Bourassa MG. Five-year risk of cardiac mortality in relation to initial severity and one-year changes in depression symptoms after myocardial infarction. *Circulation* 2002;105:1049–53.
9. Welin C, Lappas G, Wilhelmsen L. Independent importance of psycho-

- social factors for prognosis after myocardial infarction. *J Intern Med* 2000;247:629.
10. Grace SL, Abbey SE, Kapral MK, Fang J, Nolan RP, Stewart DE. Effect of depression on five-year mortality after an acute coronary syndrome. *Am J Cardiol* 2005;96:1179–85.
 11. Writing Committee for the ENRICHD Investigators. Effects of treating depression and low perceived social support on clinical events after myocardial infarction: The Enhancing Recovery in Coronary Heart Disease Patients (ENRICHD) Randomized Trial. *JAMA* 2003;289:3106–16.
 12. Carney RM, Freedland KE, Rich MW, Jaffe AS. Depression as a risk factor for cardiac events in established coronary heart disease: a review of possible mechanisms. *Ann Behav Med* 1995;17:142–9.
 13. Dalack GW, Roose SP. Perspectives on the relationship between cardiovascular disease and affective disorder. *J Clin Psychiatry* 1990;51:4–11.
 14. Carney RM, Freedland KE, Miller GE, Jaffe AS. Depression as a risk factor for cardiac mortality and morbidity: a review of potential mechanisms. *J Psychosom Res* 2002;53:897–902.
 15. Carney RM, Freedland KE, Veith RC. Depression, the autonomic nervous system, and coronary heart disease. *Psychosom Med* 2005;67(Suppl 1):S29–S33.
 16. Carney RM, Saunders RD, Freedland KE, Stein P, Rich MW, Jaffe AS. Association of depression with reduced heart rate variability in coronary artery disease. *Am J Cardiol* 1995;76:562–4.
 17. Carney RM, Blumenthal JA, Stein PK, Watkins L, Catellier D, Berkman LF, Czajkowski SM, O'Connor C, Stone PH, Freedland KE. Depression, heart rate variability, and acute myocardial infarction. *Circulation* 2001;104:2024–8.
 18. Hughes JW, Stoney CM. Depressed mood is related to high-frequency heart rate variability during stressors. *Psychosom Med* 2000;62:796–803.
 19. Kittayaphong R, Cascio WE, Light KC, Sheffield D, Golden RN, Finkel JB, Glekas G, Koch GG, Sheps DS. Heart rate variability in patients with coronary artery disease: differences in patients with higher and lower depression scores. *Psychosom Med* 1997;59:231–5.
 20. Bigger JT, Fleiss JL, Steinman RC, Rolnitzky LM, Klieger RE, Rottman JN. Frequency domain measures of heart period variability and mortality after myocardial infarction. *Circulation* 1992;85:164–71.
 21. Klieger RE, Miller JP, Bigger JT, Moss AJ. The Multicenter Post-infarction Research Group. Decreased heart rate variability and its association with increased mortality after acute myocardial infarction. *Am J Cardiol* 1987;59:256–62.
 22. La Rovere MT, Bigger JTJ, Marcus FI, Mortara A, Schwartz PJ. Baroreflex sensitivity and heart-rate variability in prediction of total cardiac mortality after myocardial infarction. ATRAMI (Autonomic Tone and Reflexes After Myocardial Infarction) Investigators. *Lancet* 1998;351:478–84.
 23. Carney RM, Blumenthal JA, Freedland KE, Stein P, Howells WB, Berkman LF, Watkins L, Czajkowski SM, Hayano J, Domitrovich PP, Jaffe AS. Low heart rate variability and the effect of depression on post-myocardial infarction mortality. *Arch Intern Med* 2005;165:1486–91.
 24. Arai Y, Saul JP, Albrecht P, Hartley LH, Lilly LS, Cohen RJ, Colucci WS. Modulation of cardiac autonomic activity during and immediately after exercise. *Am J Physiol* 1989;256(1 Pt 2):H132–H141.
 25. Cole CR, Foody JM, Blackstone EH, Lauer MS. Heart rate recovery after submaximal exercise testing as a predictor of mortality in a cardiovascularly healthy cohort. *Ann Intern Med* 2000;132:552–5.
 26. Nishime EO, Cole CR, Blackstone EH, Pashkow FJ, Lauer MS. Heart rate recovery and treadmill exercise score as predictors of mortality in patients referred for exercise ECG. *JAMA* 2000;284:1392–8.
 27. Pierpont GL, Stolpman DR, Gornick CC. Heart rate recovery post-exercise as an index of parasympathetic activity. *J Auton Nerv Syst* 2000;80:169–74.
 28. Pierpont GL, Voth EJ. Assessing autonomic function by analysis of heart rate recovery from exercise in healthy subjects. *Am J Cardiol* 2004;94:64–8.
 29. Lipinski MJ, Vetrovec GW, Froelicher VF. Importance of the first two minutes of heart rate recovery after exercise treadmill testing in predicting mortality and the presence of coronary artery disease in men. *Am J Cardiol* 2004;93:445–9.
 30. Vivekananthan DP, Blackstone EH, Pothier CE, Lauer MS. Heart rate recovery after exercise is a predictor of mortality, independent of the angiographic severity of coronary disease. *J Am Coll Cardiol* 2003;42:831–8.
 31. Cole CR, Blackstone EH, Pashkow FJ, Snader CE, Lauer MS. Heart-rate recovery immediately after exercise as a predictor of mortality. *N Engl J Med* 1999;341:1351–7.
 32. Camerini A. [Heart rate recovery and treadmill exercise score as predictors of mortality in patients referred for exercise ECG]. *Ital Heart J Suppl* 2001;2:200–2.
 33. Cheng YJ, Lauer MS, Earnest CP, Church TS, Kampert JB, Gibbons LW, Blair SN. Heart rate recovery following maximal exercise testing as a predictor of cardiovascular disease and all-cause mortality in men with diabetes. *Diabetes Care* 2003;26:2052–7.
 34. Desai MY, Pena-Almaguer E, Mannting F. Abnormal heart rate recovery after exercise: a comparison with known indicators of increased mortality. *Cardiology* 2001;96:38–44.
 35. Diaz LA, Brunken RC, Blackstone EH, Snader CE, Lauer MS. Independent contribution of myocardial perfusion defects to exercise capacity and heart rate recovery for prediction of all-cause mortality in patients with known or suspected coronary heart disease. *J Am Coll Cardiol* 2001;37:1558–64.
 36. Nissinen SI, Makikallio TH, Seppanen T, Tapanainen JM, Salo M, Tulppo MP, Huikuri HV. Heart rate recovery after exercise as a predictor of mortality among survivors of acute myocardial infarction. *Am J Cardiol* 2003;91:711–4.
 37. Watanabe J, Thamilarasan M, Blackstone EH, Thomas JD, Lauer MS. Heart rate recovery immediately after treadmill exercise and left ventricular systolic dysfunction as predictors of mortality: the case of stress echocardiography. *Circulation* 2001;104:1911–6.
 38. Jouven X, Empana JP, Schwartz PJ, Desnos M, Courbon D, Ducimetiere P. Heart-rate profile during exercise as a predictor of sudden death. *N Engl J Med* 2005;352:1951–8.
 39. Hughes JW, Casey E, Luyster F, Doe VH, Waechter D, Rosneck J, Josephson R. Depression symptoms predict heart rate recovery after treadmill stress testing. *Am Heart J* 2006;151:1122.
 40. Beck AT, Ward CH, Mendelson M, Mock J, Erbaugh J. An inventory for measuring depression. *Arch Gen Psychiatry* 1961;4:561–71.
 41. Kaufmann PG, McMahon RP, Becker LC, Bertolet B, Bonsall R, Chaitman B, Cohen JD, Forman S, Goldberg AD, Freedland K, Ketterer MW, Krantz DS, Pepine CJ, Raczynski J, Stone PH, Taylor H, Knatterud GL, Sheps DS. The Psychophysiological Investigations of Myocardial Ischemia (PIMI) study: objective, methods, and variability of measures. *Psychosom Med* 1998;60:56–63.
 42. Pepine CJ, Geller NL, Knatterud GL, Bourassa MG, Chaitman BR, Davies RF, Day P, Deanfield JE, Goldberg AD, McMahon RP. The Asymptomatic Cardiac Ischemia Pilot (ACIP) study: design of a randomized clinical trial, baseline data and implications for a long-term outcome trial. *J Am Coll Cardiol* 1994;24:1–10.
 43. Tamesis B, Stelken A, Byers S, Shaw L, Younis L, Miller DD, Chaitman BR. Comparison of the Asymptomatic Cardiac Ischemia Pilot and modified Asymptomatic Cardiac Ischemia Pilot versus Bruce and Cornell exercise protocols. *Am J Cardiol* 1993;72:715–20.
 44. Hollenberg M, Haight T, Tager IB. Depression decreases cardiorespiratory fitness in older women. *J Clin Epidemiol* 2003;56:1111–7.
 45. Tager IB, Hollenberg M, Satariano WA. Association between self-reported leisure-time physical activity and measures of cardiorespiratory fitness in an elderly population. *Am J Epidemiol* 1998;147:921–31.
 46. Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation* 1996;93:1043–65.
 47. Ruo B, Rumsfeld JS, Pipkin S, Whooley MA. Relation between depressive symptoms and treadmill exercise capacity in the Heart and Soul Study. *Am J Cardiol* 2004;94:96–9.
 48. Bonnet F, Irving K, Terra JL, Nony P, Berthezene F, Moulin P. Anxiety and depression are associated with unhealthy lifestyle in patients at risk of cardiovascular disease. *Atherosclerosis* 2005;178:339–44.
 49. Carnethon MR, Jacobs DRJ, Sidney S, Sternfeld B, Gidding SS, Shoushtari C, Liu K. A longitudinal study of physical activity and heart rate recovery: CARDIA, 1987–1993. *Med Sci Sports Exerc* 2005;37:606–12.
 50. Lind L, Andren B. Heart rate recovery after exercise is related to the insulin resistance syndrome and heart rate variability in elderly men. *Am Heart J* 2002;144:666–72.