

## Secular Trends in Deaths From Cardiovascular Diseases A 25-Year Community Study

Yariv Gerber, PhD; Steven J. Jacobsen, MD, PhD; Robert L. Frye, MD; Susan A. Weston, MS;  
Jill M. Killian, BS; Véronique L. Roger, MD, MPH

**Background**—Although age-adjusted cardiovascular disease (CVD) mortality has declined over the past decades, controversies remain about whether this trend was similar across locations of death and disease categories and about the existence of age and sex disparities.

**Methods and Results**—We examined CVD mortality trends in Olmsted County, Minnesota, between 1979 and 2003 using the categories defined by the American Heart Association, including coronary heart disease (CHD), non-CHD diseases of the heart, and noncardiac circulatory diseases. Data on demographics, cause, and location of death of all 6378 residents who died of CVD were analyzed. Although decreases in the age-adjusted rates occurred in all groups, the magnitude of the decline varied widely. Lesser annual declines were noted in out-of-hospital than in-hospital deaths (1.8% versus 4.8%;  $P<0.001$ ), in older than in younger persons (1.5% at age  $\geq 85$  years versus 3.9% for those  $\leq 74$  years of age;  $P<0.001$ ), and in women relative to men (2.5% versus 3.3%;  $P=0.007$ ). Furthermore, although CHD showed a marked annual decrease (3.3%), more modest decrements were found for non-CHD diseases of the heart (2.1%) and noncardiac circulatory diseases (2.4%) ( $P=0.02$  and  $P=0.04$  for the comparison with CHD decline, respectively).

**Conclusions**—Over the past 25 years, CVD mortality declined markedly in the community, but there were large disparities in the magnitude of the decline, resulting in a shift in the distribution toward out-of-hospital and non-CHD deaths. Further reduction in CVD mortality will require strategies directed at elderly persons and women, in whom out-of-hospital rates have improved only minimally. (*Circulation*. 2006;113:2285-2292.)

**Key Words:** cardiovascular diseases ■ epidemiology ■ mortality ■ population ■ prevention

Although age-adjusted cardiovascular disease (CVD) mortality has declined over the past decades,<sup>1–3</sup> several important controversies remain with regard to this trend. First, conflicting results have been obtained on the decline according to location of death, with some<sup>4–8</sup> but not all<sup>9,10</sup> studies reporting a steeper decline in in-hospital than in out-of-hospital mortality. Nevertheless, because most studies did not include persons  $>74$  years of age, the age past which most cardiac deaths occur,<sup>7,8</sup> the interpretation of these findings is uncertain.

### Clinical Perspective p 2292

Second, conflicting evidence has been reported with regard to age and sex disparities in the magnitude of the decline. A few community studies have noted slower decreases in women<sup>10,11</sup> and the elderly,<sup>7</sup> but these findings have not been supported by others.<sup>4–6,9</sup> Again, because many of these studies do not include elderly persons<sup>4,5,9,10</sup> and others lack statistical power to detect age and sex interactions,<sup>6,10</sup> an important variation in trends could have been missed.

Third, recent classification used by the American Heart Association divides CVD into 3 categories: coronary heart

disease (CHD), diseases of the heart, and diseases of the circulatory system.<sup>1</sup> Whereas trends in CHD mortality have been the focus of many studies,<sup>4–10,12,13</sup> the other categories have rarely been investigated. Because these include heart failure and stroke, for which attenuations in the decline have been documented recently,<sup>3,14–16</sup> it is important to examine whether the magnitude of the decline applied equally to all categories of CVD.

Prior studies in Olmsted County, Minnesota, have suggested that improved case fatality rates,<sup>17</sup> reduced myocardial infarction (MI) severity,<sup>18</sup> and decreasing MI recurrence rates<sup>19</sup> may have played a greater role in overall mortality decline than did changing CHD incidence rates.<sup>17,20</sup> However, because diverging trends across age groups and sex for MI incidence and survival,<sup>17</sup> anatomic coronary disease prevalence,<sup>21</sup> and cardiac death<sup>7,11</sup> had been noted by the mid-1990s, it is unknown to what extent these patterns affected subsequent mortality in an aging population.

Thus, the present study was undertaken to examine trends in CVD mortality in Olmsted County over 25 years,

Received September 22, 2005; revision received March 15, 2006; accepted March 21, 2006.

From the Division of Cardiovascular Diseases (Y.G., R.L.F., V.L.R.) and Department of Health Sciences Research (Y.G., S.J.J., S.A.W., J.M.K., V.L.R.), Mayo Clinic College of Medicine, Rochester, Minn.

Correspondence to Dr Véronique L. Roger, Division of Cardiovascular Diseases, Mayo Clinic, 200 First St SW, Rochester, MN 55905. E-mail roger.veronique@mayo.edu

© 2006 American Heart Association, Inc.

*Circulation* is available at <http://www.circulationaha.org>

DOI: 10.1161/CIRCULATIONAHA.105.590463

**TABLE 1. CVD Deaths According to AHA Classification**

CVD Categories*	ICD-9	ICD-10	ICD Blocks Included
Total CVD	390–459	I00–I99	All circulatory diseases
CHD	410–414	I20–I25	Ischemic heart diseases
Non-CHD diseases of the heart	390–398, 402, 404, 405, 415–429	I00–I09, I11, I13, I26–I51	Acute rheumatic fever, chronic rheumatic heart diseases, hypertensive heart diseases, pulmonary circulatory diseases, other forms of heart disease
Noncardiac circulatory diseases	401, 403, 430–459	I10, I12, I15, I52–I99	Cerebrovascular diseases; other hypertensive diseases (not included elsewhere); diseases of arteries, arterioles, and capillaries; diseases of veins, lymphatic vessels, and lymph nodes; other and unspecified disorders of the circulatory system

\*CVD categories are adapted from those used by the AHA<sup>1</sup> to produce a set of mutually exclusive groups containing all CVD deaths.

between 1979 and 2003. Specifically, we assessed variations according to location of death, age, sex, and specific CVD categories.

## Methods

### Study Setting

The population of Olmsted County is served by a largely unified medical care system that has accumulated comprehensive clinical records over an extensive period.<sup>22</sup> Olmsted County (2000 census population, 124 277) is 144 km southeast of Minneapolis and St Paul, with ≈70% of its population residing in Rochester, the centrally located county seat. In 2000, ≈90% of all residents were white and 11% were ≥65 years of age. The Olmsted population is largely middle class, with 91% of adults having graduated from high school. Except for a higher proportion being employed in the healthcare industry, the population characteristics are similar to those of US whites.

Epidemiological research in Olmsted County is feasible because the city and county are relatively isolated from other urban centers, and nearly all medical care is delivered to local residents by a handful of providers. The Mayo Medical Center and the Olmsted Medical Group and its affiliated Olmsted Community Hospital provide comprehensive care for the region in every clinical discipline. The epidemiological potential of this situation is enhanced by the fact that each provider uses a unit medical record system through which all data collected for an individual are assembled in one place. The unit records of each provider in the county are available for use. These medical records are easily retrievable because the Mayo Clinic has maintained since the early 1900s extensive indexes based on clinical and histological diagnoses and surgical and billable procedures. The Rochester Epidemiology Project<sup>22</sup> has developed a similar index for the records of other providers of medical care to local residents. This indexing system includes death certificates for Olmsted County residents. The result is a linkage of medical records from essentially all sources of medical care available to and used by the Olmsted County population. All aspects of the study were approved by the appropriate institutional review boards.

### Death Enumeration and Coding

We searched death certificate data obtained from the Minnesota Department of Health to identify individuals ≥25 years of age who were listed as residents of Olmsted County at the time of death. Information on date of birth, date of death, sex, underlying cause of death, and site of death was collected. The *International Classification of Diseases, Clinical Modification* (ICD-CM), was used to classify deaths. The 9th version was used for the years 1979 through 1998 and the 10th thereafter. Since the late 1960s, all death certificates issued in Minnesota have described the location of death as coded by the Minnesota Department of Health. We defined out-of-hospital deaths as those taking place outside of short-term-

care or long-term-care hospitals, including deaths occurring in emergency departments, private homes, public places, nursing or boarding-care homes, and infirmaries, as well as deaths among persons declared dead on arrival at a hospital. In-hospital deaths were those coded as occurring in short-term-care or long-term-care hospitals.

The procedures used in Olmsted County for completion of death certificates are different from those used in most other places. The coroner (chief medical examiner) or a pathologist member of the staff of the Mayo Clinic completes the death certificate for almost every Olmsted County resident who dies under the care of a Mayo Clinic physician, regardless of whether an autopsy is performed. When an autopsy is performed, its findings take precedence over clinical information. The autopsy rate is 30%, higher than the national average of 15%.<sup>23</sup>

### Statistical Analyses

We adapted the classification used by the AHA<sup>1</sup> to produce a set of mutually exclusive CVD categories, including CHD, non-CHD diseases of the heart, and noncardiac circulatory diseases (Table 1). The underlying cause of death was used for all analyses.

Age-, sex-, and year-specific mortality rates were calculated for each CVD category. The denominators were determined by Olmsted County population census data for 1970, 1980, 1990, and 2000, with linear interpolation for the intercensal years and extrapolation after 2000.<sup>24</sup> These rates were directly standardized to the age and sex distributions of the 2000 US population.

Poisson regression models were used to assess temporal trends in CVD mortality rates overall and by selected categories. Specific counts for each calendar year, age (using midpoints of 10-year intervals), sex, and location of death (ie, in and out of hospital) were used as the units of observation. Calendar year and successive age intervals were entered as continuous variables, with their quadratic terms tested and rejected. The logarithm of the count was modeled with the logarithm of the year-, age-, and sex-specific population size as the offset using the SAS GENMOD procedure. Annual percentage change and an overall relative risk (RR) for the entire study period in specific groups and categories were estimated through stratified analyses. Comparisons of temporal trends across locations of death, age, sex, and CVD categories were performed by including the 2-way interaction term of each variable with year, separately, after adjustment for all main effects. The 3-way interaction terms year by location by age and year by location by sex were used to examine differential age and sex trends according to location and were evaluated on the basis of fully interacted 2-way models. Model goodness of fit was assessed, and there was no evidence of overdispersion or autocorrelation.

The contribution (in percentage) of each specific CVD category to the total age- and sex-adjusted decline was estimated from this formula:  $100 \times (dM_s/dM_t)$ , where  $dM_s$  is specific adjusted mortality rate in 1979 through 1983 minus specific adjusted mortality

**TABLE 2. Frequency of CVD Deaths by ICD-Defined Blocks**

ICD Blocks	Deaths, n	Percent Within CVD Categories	Percent of Total CVD
CHD (n=3622)			
Ischemic heart diseases	3622	100.0	56.8
Non-CHD diseases of the heart (n=1138)			
Acute rheumatic fever	1	0.1	0.0
Chronic rheumatic heart diseases	81	7.1	1.3
Hypertensive heart diseases	65	5.7	1.0
Pulmonary circulatory diseases	142	12.5	2.2
Other forms of heart disease	849	74.6	13.3
Noncardiac circulatory diseases (n=1618)			
Cerebrovascular diseases	1185	73.2	18.6
Other hypertensive diseases	63	3.9	1.0
Diseases of arteries, arterioles, and capillaries	330	20.4	5.2
Diseases of veins, lymphatic vessels, and lymph nodes	35	2.2	0.5
Other and unspecified disorders of the circulatory system	5	0.3	0.1
Total CVD deaths (n=6378)			

rate in 1999 through 2003 and  $dM_t$  is total adjusted mortality rate in 1979 through 1983 minus total adjusted mortality rate in 1999 through 2003.<sup>9</sup>

The authors had full access to the data and take full responsibility for their integrity. All authors have read and agree to the manuscript as written.

## Results

Over the 25-year period covered by the study, 6378 CVD deaths were recorded, including 57% CHD, 18% non-CHD diseases of the heart, and 25% noncardiac circulatory diseases. Of all CVD deaths, 54% occurred in women and 69% in persons  $\geq 75$  years of age (39% in persons  $\geq 85$  years of age). The most frequent causes of death were CHD and stroke, accounting for 57% and 19% of all CVD deaths, respectively (Table 2). Heart failure, which is included in the "other forms of heart disease" group, was recorded as the underlying cause of death in an additional 4% of the cases.

From the period 1979 through 1983 to the period 1999 through 2003, the average  $\pm$  SD age at CVD death increased from  $76 \pm 13$  to  $80 \pm 14$  years ( $P < 0.001$ ), as did the proportion of decedents  $\geq 85$  years of age (30% to 45%;  $P < 0.001$ ). The age- and sex-adjusted CVD mortality declined substantially over this period, with an average annual decrease of 2.9% (95% CI, 2.6 to 3.3). This equates to an age- and sex-adjusted RR of 0.50 (95% CI, 0.46 to 0.54) for CVD death in 2003 versus 1979.

## Trends by Location of Death


The temporal decline in CVD mortality differed markedly by location of death ( $P < 0.001$  for the year-by-location interaction term). The rate reduction, adjusted for age and sex, was greater for in-hospital deaths (annual decline, 4.8%; 95% CI, 4.2 to 5.3%) than for out-of-hospital deaths (annual decline, 1.8%; 95% CI, 1.4 to 2.2) (Figure 1). In the most recent years (1999 through 2003), 72% of all CVD deaths were categorized as out of hospital, as compared with 55% in the initial

5 years (1979 through 1983). Despite the fact that most CVD deaths occurred outside the hospital, the decline in CVD mortality was primarily the result of an in-hospital decrease (66%), with only 34% attributable to an out-of-hospital decrease.


Within out-of-hospital deaths, similar trends were observed in nursing homes (the site of death for 45% of out-of-hospital decedents) and in other sites (respective annual declines, 2.0% and 1.6%;  $P = 0.39$ ).

## Trends by Age and Sex

The decline in CVD mortality differed markedly according to age. Although an overall reduction occurred for all ages, its magnitude was greater in younger than in older persons ( $P < 0.001$  for the year-by-age interaction). Indeed, the aver-



**Figure 1.** CVD mortality rates (SE) by location of death from 1979 to 2003 in Olmsted County, Minnesota. Yearly rates (per 100 000 persons) are shown on a logarithmic scale and have been standardized by the direct method to the age and sex distribution of the US population in 2000.



**Figure 2.** CVD mortality rates (SE) for selected age groups from 1979 to 2003 in Olmsted County, Minnesota. Yearly rates are presented on a logarithmic scale and reported per 100 000 persons.

age yearly decline in CVD mortality was 3.9% (95% CI, 3.3 to 4.5) for persons <75 years of age, 3.4% (95% CI, 2.8 to 4.0) for those 75 to 84 years of age, and 1.5% (95% CI, 0.9 to 2.0) for those  $\geq 85$  years of age (Figure 2). Over the 25 years of the study, this equates to RRs for fatal CVD of 0.38 (95% CI, 0.33 to 0.44), 0.44 (95% CI, 0.38 to 0.50), and 0.70 (95% CI, 0.61 to 0.80) for the 3 age groups, respectively, in 2003 versus 1979 (Figure 3).

Furthermore, as indicated by the significance of the 3-way interaction year by location by age ( $P<0.001$ ), trends across age groups differed between in-hospital deaths (annual declines, 5.6% [95% CI, 4.7 to 6.5] in those <75 years of age, 4.7% [95% CI, 3.8 to 5.6] in those 75 to

84 years of age, and 3.6% [95% CI, 2.5 to 4.6] in those  $\geq 85$  years of age) and out-of-hospital rates (annual declines, 2.8% [95% CI, 2.0 to 3.5], 2.5% [95% CI, 1.7 to 3.3], and 0.8% [95% CI, 0.1 to 1.4], for the respective 3 age groups). This corresponds to RRs in 2003, as compared with 1979, of 0.25 (95% CI, 0.20 to 0.32), 0.32 (95% CI, 0.25 to 0.40), and 0.42 (95% CI, 0.32 to 0.54) in in-hospital and 0.51 (95% CI, 0.42 to 0.62), 0.55 (95% CI, 0.45 to 0.66), and 0.84 (95% CI, 0.72 to 0.98) in out-of-hospital CVD deaths, respectively (Figure 3).

Large sex disparities also were observed. Although the age-adjusted CVD mortality was higher among men, this gap diminished over time as a result of a steeper downward trend in men than in women ( $P=0.007$  for the year-by-sex interaction). Indeed, the annual decline in men was 3.3% (95% CI, 2.8 to 3.8), as compared with 2.5% (95% CI, 2.0 to 3.0) in women, and the RRs of CVD mortality in 2003 versus 1979 were 0.45 (95% CI, 0.40 to 0.50) in men and 0.54 (95% CI, 0.49 to 0.61) in women (Figure 3).

Furthermore, although trends in the age-adjusted in-hospital deaths were similar across sexes (annual decline, 4.8%; 95% CI, 4.0 to 5.6), the decline in out-of-hospital deaths was greater for men (annual decline, 2.4%; 95% CI, 1.7 to 3.0) than for women (annual decline, 1.4%; 95% CI, 0.8 to 2.0) ( $P<0.001$  for the year-by-location-by-sex interaction). As shown in Figure 3, over the 25 years of the study, this corresponds to an RR of 0.31 (95% CI, 0.25 to 0.38) in both sexes for in-hospital death, as compared with 0.56 (95% CI, 0.49 to 0.66) among men and 0.71 (95% CI, 0.62 to 0.82) among women for out-of-hospital CVD death.


### Trends Across CVD Categories

Trends have differed across CVD categories. The age- and sex-adjusted CHD mortality decreased markedly over time (annual decline, 3.3%; 95% CI, 2.8 to 3.7), contrasting with more modest decreases in non-CHD diseases of the heart (annual decline, 2.1%; 95% CI, 1.3 to 2.9;  $P=0.02$  for the comparison with CHD trend) and noncardiac circulatory diseases (annual decline, 2.4%; 95% CI, 1.8 to 3.1;  $P=0.04$  for the comparison with CHD trend). In the latest 5 years, the distribution of CVD deaths was 55% CHD, 18% non-CHD diseases of the heart, and 27% noncardiac circulatory diseases, as compared with 59%, 16%, and 25%, respectively, in the initial 5 years (Figure 4). Of the total decline in CVD mortality over the study period, 64% was attributable to decreasing CHD, 22% to decreasing noncardiac circulatory diseases, and 14% to declining non-CHD diseases of the heart.

## Discussion


### Summary of Findings

Over the past 25 years in the geographically defined population of Olmsted County, the age- and sex-adjusted mortality from CVD declined steadily. This favorable trend was observed in all CVD categories, for both in- and out-of-hospital locations, and across all demographic groups examined.



**Figure 3.** Age- and sex-adjusted RR estimates and their 95% CIs for fatal CVD in 2003 vs 1979. The sex-specific RRs are adjusted for age; the age-specific RRs are adjusted for sex.





**Figure 4.** Decline in CVD mortality rates (per 100 000 persons) from 1979 to 2003 partitioned into specific AHA-defined categories. Rates have been standardized by the direct method to the age and sex distribution of the US 2000 population and are shown at 5-year intervals.

Although these findings are encouraging, sizable variations in the magnitude of these declines were found: Out-of-hospital declined less than in-hospital; slower decrements occurred in the elderly and in women; and CHD mortality decreased more rapidly than did other CVD categories. The sex disparities were driven by diverging trends in out-of-hospital deaths. Overall, despite a high proportion of CVD deaths occurring outside the hospital, the contribution of the decline in out-of-hospital rates to the total decline in CVD mortality was relatively modest, whereas much greater contribution was made through in-hospital reduction. These data indicate that declining trends were clustered mostly in the hospital setting and among younger subjects and men. Furthermore, the distribution of CVD deaths has shifted toward non-CHD causes.

### Overall Decline in CVD Mortality

Mortality trends in the Olmsted County population are similar to those measured in other cohorts. Indeed, we report here a 50% decline in CVD mortality between 1979 and 2003, which is commensurate with the 52% decline in heart disease observed in the total US population between 1970 and 2002<sup>3</sup> and similarly congruent with the 64% and 70% decline in CHD mortality reported in the Framingham Heart Study<sup>6</sup> and in the Minnesota Heart Study<sup>4</sup> over similar time periods.

In the context of this decline, a controversy still exists as to the relative contributions of primary prevention, acute cardiac care, and secondary prevention. Although reports from the Framingham Heart Study have emphasized the role of temporal reduction in CVD risk factors,<sup>6,25</sup> community surveillance data from the Minnesota Heart Study<sup>4</sup> and the Atherosclerosis Risk in Communities Study<sup>12</sup> have attributed the decline largely to improvements in acute cardiac care and secondary prevention after MI. However, neither of these studies included persons >74 years of age, which limits their ability to detect age- and sex-specific trends.

In accord with the other community surveillance studies, reports from Olmsted County have indicated secular improvements in several aspects related to the management and survival of coronary patients,<sup>17-19,26</sup> whereas no major decline has been shown in the overall incidence of CHD.<sup>17,20</sup> However, all of the trends examined in this community, without exception, have differed by age and sex, indicating the need for specific assessment.

### Specific Trends in Olmsted County

#### *Trends by Location of Death*

A marked discrepancy in the magnitude of CVD decline was found with regard to location of death in that in-hospital deaths declined more rapidly than out-of-hospital deaths. Within the latter setting, trends found in nursing homes were similar to those in other sites. Previous reports on this issue yielded discrepant results, in part because of differences in study populations<sup>5,9</sup> and the inclusion or exclusion of elderly persons.<sup>4-10</sup> As demonstrated by the present data, inclusion of all age groups in such analyses is crucial because data indicated a differential rate reduction by age.

Reduction in length of hospital stay for MI<sup>27</sup> and improved survival of patients with CHD who formerly might have died in the hospital<sup>17</sup> could have contributed to this shift in the location of deaths. On the other hand, a more rapid admission to hospital after symptom onset has been shown over time,<sup>28</sup> which may balance the effect of the shortened hospital stay.

It has been argued that time trends in out-of-hospital mortality likely reflect the success of prevention strategies, whereas trends in in-hospital mortality are more closely related to acute medical care and early treatment.<sup>4,8,13</sup> The encouraging decline in the proportion of CVD deaths occurring in hospital may therefore reflect the improvements in emergency services and in patient treatment.<sup>19,26,28</sup> However, the increased proportion of out-of-

hospital deaths indicates a need for public health initiatives toward prevention, early recognition of cardiac symptoms and signs, and access to emergency cardiac care.<sup>8</sup>

### **Trends by Age and Sex**

Conflicting findings have been published with regard to the existence of age and sex disparities in CVD mortality trends,<sup>4–7,9–11</sup> but many of these studies did not include elderly persons, which hinders their ability to detect such variations. The present data indicate that older persons and women benefit the least from the decline in CVD mortality and that the disparities in trends apply mainly to out-of-hospital deaths. The driving forces of these diverging declines are complex, likely multifactorial, and may themselves have evolved over time. Prior studies in Olmsted County have shown that, over time, the incidence of MI and the prevalence of atherosclerosis decreased in men but remained stable or even increased slightly in women and elderly persons.<sup>17,20,21</sup> Among patients with MI, survival benefits were clustered in younger ages and men.<sup>17</sup> The increase in MI incidence among women and older persons suggests that primary prevention has been less effective in these groups. This is supported by data on differential trends in risk factor prevalence with increasing age.<sup>29</sup> However, because most of the age and sex disparities pertain to out-of-hospital deaths and given that more than half of out-of-hospital decedents have antecedent coronary disease,<sup>6,7,9</sup> secondary prevention (or lack of) also might contribute largely to the disparities in trends. Indeed, according to recent studies of the Olmsted County population, traditional risk factors at the time of MI are more prevalent and confer a higher risk for recurrent ischemic events in women than in men.<sup>30</sup> Furthermore, women and older patients are less likely to participate in secondary prevention programs after MI.<sup>31</sup>

Studies have shown that women with MI wait longer than men after the onset of symptoms to seek care.<sup>32,33</sup> Lack of awareness, more ambiguous MI signs and symptoms, and social isolation may contribute to delays.<sup>32,34</sup> These factors may be involved in the lesser temporal decline in out-of-hospital CVD death rates among women.

Finally, although studies have suggested that men and younger MI patients tend to be treated more aggressively, particularly with reperfusion therapy and cardiac catheterization,<sup>26,35–37</sup> this may not be the primary explanation for the diverging trends shown in our study because in-hospital death trends have been relatively homogeneous across age groups and similar in men and women. Furthermore, sex disparities in the management of MI appear to have remained constant over time and thus are unlikely to confound the observed temporal trends.<sup>37</sup>

### **Trends Across CVD Categories**

CHD has declined more rapidly than non-CHD diseases of the heart and noncardiac circulatory diseases. Improved survival after MI may have contributed to the diverging trends across CVD categories because the risk of stroke and heart failure is substantially increased in MI survivors.<sup>38,39</sup> However, this hypothesis was not fully supported by a recent study that failed to explain the observed slowing of the stroke

mortality rate decline during the 1990s by the prolonged survival of CHD patients.<sup>15</sup> Thus, other possible explanations, including less successful prevention and treatment for the non-CHD causes, should be considered.

### **Implications**

The disparities in the magnitude of the decline in CVD mortality presented in this study delineate opportunities for prevention and interventions. First, the lesser decline in non-CHD categories underscores the fact that the healthcare community should be increasingly prepared to address entities other than acute coronary disease. Second, because persons dying of CVD do so increasingly out of the hospital, clinical and public health efforts should be directed at prevention. To this end, it is important to emphasize that prevention should apply not only to primary but also to secondary prevention because about half of out-of-hospital CVD decedents have a history of coronary disease.<sup>6,7,9</sup> Finally, the burden of fatal CVD has shifted to elderly persons and women, which defines the population toward whom these initiatives should be primarily targeted. It should be underscored that in the very elderly, interventions may revolve around end-of-life issues in the setting of chronic CVD rather than aggressive acute care.

### **Potential Limitations**

Although the population of Olmsted County is becoming more diverse, during the study period, it consisted primarily of US whites. Nevertheless, trends in CVD mortality in this population seem to parallel national data,<sup>1,3</sup> which suggests that these data may be generalized outside of this community.

The reliance on death certificates may introduce bias. On the other hand, death certificates are the only data source currently available to assess population trends in CVD mortality, provided that the accuracy remains relatively constant over time. To this end, a validation study performed in Olmsted County has found a similar accuracy of death certificate data for out-of-hospital CHD, typically considered more challenging than in-hospital events,<sup>40</sup> throughout the 1980s and 1990s.<sup>23</sup> In addition, our analyses are restricted to broad disease categories for which death certificate data are more accurate than for specific diseases.<sup>41</sup>

### **Conclusions**

In this community over the past 25 years, CVD mortality declined steadily, but the magnitude of the decline differed greatly, with less improvement among elderly persons and women. Furthermore, the proportion of persons dying out of hospital and from non-CHD causes has increased, thus changing the face of CVD deaths. Prevention strategies should be directed toward the out-of-hospital setting, with increased focus on non-CHD causes, while targeting elderly persons and women.

### **Acknowledgments**

This study was supported by grants from the Public Health Service and the National Institutes of Health (AR30582, R01 HL 59205, and R01 HL 72435) and a Postdoctoral Fellowship Award from the AHA, Greater Midwest Affiliate (0525753Z). Dr Roger is an

established investigator of the AHA. The funding sources for this study played no role in the design or conduct of the study; data management and analysis; or manuscript preparation, review, and authorization for submission.

## Disclosures

None.

## References

1. *Heart Disease and Stroke Statistics: 2005 Update*. Dallas, Tex: American Heart Association; 2006.
2. Decline in deaths from heart disease and stroke: United States, 1900–1999. *MMWR*. 1999;48:649–656.
3. Jemal A, Ward E, Hao Y, Thun M. Trends in the leading causes of death in the United States, 1970–2002. *JAMA*. 2005;294:1255–1259.
4. McGovern PG, Jacobs DR Jr, Shahar E, Arnett DK, Folsom AR, Blackburn H, Luepker RV. Trends in acute coronary heart disease mortality, morbidity, and medical care from 1985 through 1997: the Minnesota Heart Survey. *Circulation*. 2001;104:19–24.
5. Rosamond WD, Folsom AR, Chambless LE, Wang CH. Coronary heart disease trends in four United States communities: the Atherosclerosis Risk in Communities (ARIC) study 1987–1996. *Int J Epidemiol*. 2001;30(suppl 1):S17–S22.
6. Fox CS, Evans JC, Larson MG, Kannel WB, Levy D. Temporal trends in coronary heart disease mortality and sudden cardiac death from 1950 to 1999: the Framingham Heart Study. *Circulation*. 2004;110:522–527.
7. Goraya TY, Jacobsen SJ, Kottke TE, Frye RL, Weston SA, Roger VL. Coronary heart disease death and sudden cardiac death: a 20-year population-based study. *Am J Epidemiol*. 2003;157:763–770.
8. Zheng ZJ, Croft JB, Giles WH, Mensah GA. Sudden cardiac death in the United States, 1989 to 1998. *Circulation*. 2001;104:2158–2163.
9. Salomaa V, Ketonen M, Koukkuinen H, Immonen-Raiha P, Jerkkola T, Karja-Koskenkari P, Mahonen M, Niemela M, Kuulasmaa K, Palomaki P, Mustonen J, Arstila M, Vuorenmaa T, Lehtonen A, Lehto S, Miettinen H, Torppa J, Tuomilehto J, Kesaniemi YA, Pyorala K. Decline in out-of-hospital coronary heart disease deaths has contributed the main part to the overall decline in coronary heart disease mortality rates among persons 35 to 64 years of age in Finland: the FINAMI Study. *Circulation*. 2003;108:691–696.
10. Derby CA, Lapane KL, Feldman HA, Carleton RA. Sex-specific trends in validated coronary heart disease rates in southeastern New England, 1980–1991. *Am J Epidemiol*. 2000;151:417–429.
11. Roger VL, Jacobsen SJ, Weston SA, Bailey KR, Kottke TE, Frye RL. Trends in heart disease deaths in Olmsted County, Minnesota, 1979–1994. *Mayo Clin Proc*. 1999;74:651–657.
12. Rosamond WD, Chambless LE, Folsom AR, Cooper LS, Conwill DE, Clegg L, Wang CH, Heiss G. Trends in the incidence of myocardial infarction and in mortality due to coronary heart disease, 1987 to 1994. *N Engl J Med*. 1998;339:861–867.
13. McGovern PG, Pankow JS, Shahar E, Doliszny KM, Folsom AR, Blackburn H, Luepker RV. Recent trends in acute coronary heart disease: mortality, morbidity, medical care, and risk factors: the Minnesota Heart Survey Investigators. *N Engl J Med*. 1996;334:884–890.
14. McGovern PG, Burke GL, Sprafka JM, Xue S, Folsom AR, Blackburn H. Trends in mortality, morbidity, and risk factor levels for stroke from 1960 through 1990: the Minnesota Heart Survey. *JAMA*. 1992;268:753–759.
15. Peeters A, Bonneux L, Barendregt JJ, Mackenbach JP. Improvements in treatment of coronary heart disease and cessation of stroke mortality rate decline. *Stroke*. 2003;34:1610–1614.
16. Goldberg RJ, Glatfelter K, Burbank-Schmidt E, Farmer C, Spencer FA, Meyer T. Trends in mortality attributed to heart failure in Worcester, Massachusetts, 1992 to 2001. *Am J Cardiol*. 2005;95:1324–1328.
17. Roger VL, Jacobsen SJ, Weston SA, Goraya TY, Killian J, Reeder GS, Kottke TE, Yawn BP, Frye RL. Trends in the incidence and survival of patients with hospitalized myocardial infarction, Olmsted County, Minnesota, 1979 to 1994. *Ann Intern Med*. 2002;136:341–348.
18. Hellermann JP, Reeder GS, Jacobsen SJ, Weston SA, Killian JM, Roger VL. Longitudinal trends in the severity of acute myocardial infarction: a population study in Olmsted County, Minnesota. *Am J Epidemiol*. 2002;156:246–253.
19. Jokhadar M, Jacobsen SJ, Reeder GS, Weston SA, Roger VL. Sudden death and recurrent ischemic events after myocardial infarction in the community. *Am J Epidemiol*. 2004;159:1040–1046.
20. Arciero TJ, Jacobsen SJ, Reeder GS, Frye RL, Weston SA, Killian JM, Roger VL. Temporal trends in the incidence of coronary disease. *Am J Med*. 2004;117:228–233.
21. Roger VL, Weston SA, Killian JM, Pfeifer EA, Belau PG, Kottke TE, Frye RL, Bailey KR, Jacobsen SJ. Time trends in the prevalence of atherosclerosis: a population-based autopsy study. *Am J Med*. 2001;110:267–273.
22. Melton LJ 3rd. History of the Rochester Epidemiology Project. *Mayo Clin Proc*. 1996;71:266–274.
23. Goraya TY, Jacobsen SJ, Belau PG, Weston SA, Kottke TE, Roger VL. Validation of death certificate diagnosis of out-of-hospital coronary heart disease deaths in Olmsted County, Minnesota. *Mayo Clin Proc*. 2000;75:681–687.
24. Bergstralh EJ, Offord KP, Chu C-P, Beard CM, O'Fallon WM. Calculating incidence, prevalence, and mortality rates for Olmsted County, Minnesota residents: an update. Rochester, Minn: Section of Biostatistics, Mayo Clinic; 1992. Technical Report Series 49.
25. Sytkowski PA, Kannel WB, D'Agostino RB. Changes in risk factors and the decline in mortality from cardiovascular disease: the Framingham Heart Study. *N Engl J Med*. 1990;322:1635–1641.
26. Perschbacher JM, Reeder GS, Jacobsen SJ, Weston SA, Killian JM, Slobodova A, Roger VL. Evidence-based therapies for myocardial infarction: secular trends and determinants of practice in the community. *Mayo Clin Proc*. 2004;79:983–991.
27. Spencer FA, Lessard D, Gore JM, Yarzebski J, Goldberg RJ. Declining length of hospital stay for acute myocardial infarction and postdischarge outcomes: a community-wide perspective. *Arch Intern Med*. 2004;164:733–740.
28. Rogers WJ, Canto JG, Lambrew CT, Tiefenbrunn AJ, Kinkaid B, Shoultz DA, Frederick PD, Every N. Temporal trends in the treatment of over 1.5 million patients with myocardial infarction in the US from 1990 through 1999: the National Registry of Myocardial Infarction 1, 2 and 3. *J Am Coll Cardiol*. 2000;36:2056–2063.
29. Greenlund KJ, Zheng ZJ, Keenan NL, Giles WH, Casper ML, Mensah GA, Croft JB. Trends in self-reported multiple cardiovascular disease risk factors among adults in the United States, 1991–1999. *Arch Intern Med*. 2004;164:181–188.
30. Gerber Y, Weston SA, Killian JM, Jacobsen SJ, Roger VL. Sex and classic risk factors after myocardial infarction: a community study. *Am Heart J*. In press.
31. Witt BJ, Jacobsen SJ, Weston SA, Killian JM, Meverden RA, Allison TG, Reeder GS, Roger VL. Cardiac rehabilitation after myocardial infarction in the community. *J Am Coll Cardiol*. 2004;44:988–996.
32. Marrugat J, Sala J, Masia R, Pavesi M, Sanz G, Valle V, Molina L, Seres L, Elosua R. Mortality differences between men and women following first myocardial infarction: RESCATE Investigators: Recursos Empleados en el Síndrome Coronario Agudo y Tiempo de Espera. *JAMA*. 1998;280:1405–1409.
33. Vaccarino V, Parsons L, Every NR, Barron HV, Krumholz HM. Sex-based differences in early mortality after myocardial infarction: National Registry of Myocardial Infarction 2 Participants. *N Engl J Med*. 1999;341:217–225.
34. Sonke GS, Beaglehole R, Stewart AW, Jackson R, Stewart FM. Sex differences in case fatality before and after admission to hospital after acute cardiac events: analysis of community based coronary heart disease register. *BMJ*. 1996;313:853–855.
35. Qureshi AI, Suri MF, Guterman LR, Hopkins LN. Ineffective secondary prevention in survivors of cardiovascular events in the US population: report from the Third National Health and Nutrition Examination Survey. *Arch Intern Med*. 2001;161:1621–1628.
36. Chandra NC, Ziegelstein RC, Rogers WJ, Tiefenbrunn AJ, Gore JM, French WJ, Rubison M. Observations of the treatment of women in the United States with myocardial infarction: a report from the National Registry of Myocardial Infarction-I. *Arch Intern Med*. 1998;158:981–988.
37. Vaccarino V, Rathore SS, Wenger NK, Frederick PD, Abramson JL, Barron HV, Manhapra A, Mallik S, Krumholz HM. Sex and racial differences in the management of acute myocardial infarction, 1994 through 2002. *N Engl J Med*. 2005;353:671–682.

38. Hellermann JP, Jacobsen SJ, Gersh BJ, Rodeheffer RJ, Reeder GS, Roger VL. Heart failure after myocardial infarction: a review. *Am J Med.* 2002;113:324–330.
39. Witt BJ, Brown RD Jr, Jacobsen SJ, Weston SA, Yawn BP, Roger VL. A community-based study of stroke incidence after myocardial infarction. *Ann Intern Med.* 2005;143:785–792.
40. Folsom AR, Gomez-Marin O, Gillum RF, Kottke TE, Lohman W, Jacobs DR Jr. Out-of-hospital coronary death in an urban population: validation of death certificate diagnosis: the Minnesota Heart Survey. *Am J Epidemiol.* 1987;125:1012–1018.
41. Gittelsohn A, Royston PN. Annotated bibliography of cause-of-death validation studies. *Vital Health Stat 2.* 1982;2:1–42.

### CLINICAL PERSPECTIVE

Despite the dramatic decline in age-adjusted mortality from cardiovascular disease (CVD) in the United States over the past few decades, it is uncertain whether this decline was distributed uniformly across locations of death, age, sex, and CVD categories as defined by the American Heart Association (ie, coronary heart disease [CHD], diseases of the heart, and diseases of the circulatory system). We examined mortality trends in the population of Olmsted County (Minnesota) between 1979 and 2003, during which time 6378 residents died of CVD. Over the 25-year period, CVD mortality rates declined steadily overall and across all locations of death, age groups, sex, and CVD categories examined. Although overall CVD mortality declined by 50%, large variations in the magnitude of the decline were detected. Out-of-hospital deaths declined less than in-hospital deaths; slower decrements occurred in the elderly and in women; and CHD decreased more rapidly than did other CVD categories. Thus, declining trends were clustered mostly in the hospital setting, among younger subjects, and among men, and the distribution of CVD deaths has shifted toward non-CHD causes. To maintain the decline in CVD mortality, prevention strategies should be directed toward the out-of-hospital setting, with increased focus on non-CHD causes while targeting elderly persons and women.