WHITE PAPER

Clinical neurocardiology defining the value of neuroscience-based cardiovascular therapeutics

Kalyanam Shivkumar¹, Olujimi A. Ajijola¹, Inder Anand², J. Andrew Armour¹, Peng-Sheng Chen³, Murray Esler⁴, Gaetano M. De Ferrari⁵, Michael C. Fishbein⁶, Jeffrey J. Goldberger⁷, Ronald M. Harper⁸, Michael J. Joyner⁹, Sahib S. Khalsa¹⁰, Rajesh Kumar¹¹, Richard Lane¹², Aman Mahajan¹³, Sunny Po^{14,15}, Peter J. Schwartz¹⁶, Virend K. Somers⁹, Miguel Valderrabano¹⁷, Marmar Vaseghi¹ and Douglas P. Zipes¹⁸

Abstract The autonomic nervous system regulates all aspects of normal cardiac function, and is recognized to play a critical role in the pathophysiology of many cardiovascular diseases. As such, the value of neuroscience-based cardiovascular therapeutics is increasingly evident. This White Paper reviews the current state of understanding of human cardiac neuroanatomy, neurophysiology, pathophysiology in specific disease conditions, autonomic testing, risk stratification, and neuromodulatory strategies to mitigate the progression of cardiovascular diseases.

(Received 6 February 2016; accepted after revision 8 April 2016; first published online 26 April 2016)

Corresponding author K. Shivkumar: UCLA Cardiac Arrhythmia Center & Neurocardiology Research Center of Excellence Molecular, Cellular & Integrative Physiology Interdepartmental Program, David Geffen School of Medicine at UCLA, 100 UCLA Medical Plaza, suite 660, Los Angeles, CA, USA. Email: kshivkumar@mednet.ucla.edu

Abbreviations ABVN, auricular branch of the vagus nerve; AE, adverse event; AF, atrial fibrillation; ANS, autonomic nervous system; ARI, activation recovery interval; ART, autonomic regulation therapy; ASV, adaptive servo-ventilation; AV, atrioventricular; CHF, clinical heart failure; CPVT, catecholaminergic polymorphic ventricular tachycardia; CSA, central sleep apnoea; CSD, cardiac sympathetic denervation; CSR, Cheyne–Stokes respiration; GP, ganglionated plexi; HF, heart failure; HRV, heart rate variability; ICD, implantable cardioverter defibrillator; ICM, ischaemic cardiomyopathy; ICN, intrinsic cardiac nervous system; LBNP, lower body negative pressure; LCSD, left cardiac sympathetic denervation; LLTS, low-level tragus stimulation; LL-VNS, low-level vagus nerve stimulation; LOM, ligament of Marshall; LQTS, long QT syndrome; LSG, left stellate ganglion; LVEF, left ventricular ejection fraction; MI, myocardial infarction; MSNA, muscle sympathetic nerve activity; NICM, non-ischaemic cardiomyopathy; NGF, nerve growth factor; PV, pulmonary vein; PVI, pulmonary vein isolation; SCD, sudden cardiac death; SG, stellate ganglion; SCS, spinal cord stimulation; TEA, thoracic epidural anaesthesia; OSA, obstructive sleep apnoea; RSNA, renal sympathetic nerve activity; VA, ventricular arrhythmia; VNS, vagus nerve stimulation; VOM, vein of Marshall; VT, ventricular tachycardia.

¹ UCLA Cardiac Arrhythmia Center and Neurocardiology Research Center of Excellence, Los Angeles, CA, USA

²Department of Cardiology, University of Minnesota Medical School, Minneapolis, MN, USA

³Krannert Institute of Cardiology, Indiana University School of Medicine, Indianapolis, IN, USA

⁴Baker IDI Heart and Diabetes Institute, Melbourne, Victoria, Australia

⁵Department of Molecular Medicine, University of Pavia, Pavia, Italy

⁶Department of Pathology and Laboratory Medicine, David Geffen School of Medicine at UCLA, Los Angeles, California, USA

⁷Division of Cardiology, University of Miami Miller School of Medicine, Miami, FL, USA

⁸ Department of Neurobiology and the Brain Research Institute, University of California, Los Angeles, CA, USA

⁹Division of Cardiovascular Diseases, Mayo Clinic and Mayo Foundation, Rochester, MN, USA

¹⁰Laureate Institute for Brain Research, Tulsa, OK, USA

¹¹Departments of Anesthesiology and Radiological Sciences, David Geffen School of Medicine at UCLA, Los Angeles, CA, USA

¹²Department of Psychiatry, University of Arizona College of Medicine, Tucson, AZ, USA

¹³Department of Anesthesia, UCLA, Los Angeles, CA, USA

¹⁴Heart Rhythm Institute, University of Oklahoma Health Sciences Center, Oklahoma City, OK, USA

¹⁵University of Tulsa Oxley College of Health Sciences, Tulsa, OK, USA

¹⁶ Center for Cardiac Arrhythmias of Genetic Origin, IRCCS Instituto Auxologico Italiano, c/o Centro Diagnostico e di Ricerrca San Carlo, Milan, Italy

¹⁷Methodist DeBakey Heart and Vascular Center and Methodist Hospital Research Institute, Houston Methodist Hospital, Houston, TX, USA

¹⁸Indiana University School of Medicine, Indianapolis, IN, USA

As such, although often using some reductionistic models, I do my best to remain a pupil at the school of complexity Alberto Malliani (1935–2006)

(Principles of Cardiovascular Neural Regulation in Health and Disease, pp. xiv–xv, Springer Science + Business Media, LLC, 2000).

Introduction

Understanding cardiac autonomic regulation is a prerequisite to comprehending the development and progression of most cardiovascular diseases (hypertension, heart failure, arrhythmias and myocardial infarction) and cardio-respiratory diseases such as sleep apnoea. Enhanced cardiac sympathetic activity and attenuated parasympathetic central drive are noted in several cardiac disorders, portend poor prognosis, and are associated with arrhythmias and sudden death (Chugh et al. 2008; Shen & Zipes, 2014). The autonomic nervous system (ANS) especially plays an intricate role in the pathophysiology of arrhythmias leading to sudden cardiac death (SCD), and as such, neuraxial modulation is emerging as an important avenue of scientific inquiry and therapeutic intervention (Vaseghi & Shivkumar, 2008; Schwartz, 2014) (Figs 1 and 2).

Anatomy and physiology

To understand the relevance of the cardiac nervous system in the evolution of cardiac disease (Armour, 2008; Kember et al. 2013b; Florea & Cohn, 2014; Fukuda et al. 2015), one must appreciate the anatomical foundations of its functional organization for cardiac control. As a corollary, to develop targeted neuromodulation therapies to treat specific cardiac pathologies, one must comprehend the anatomical basis of neuraxial coordination of regional cardiac function (Buckley et al. 2016). Targeted neuromodulatory therapies have been recently devised for sustained treatment of cardiac arrhythmias as well as heart failure. Bioelectric therapies, for example, have proven to be efficacious following myocardial infarction (Beaumont et al. 2015), as well as in modulating autonomic imbalances that increase the incidence of ventricular arrhythmias (VAs) (Ardell et al. 2015). It is by gaining a thorough understanding of the anatomical basis of neuronal interactivity that one can intelligently devise therapeutic targets for neuromodulation therapy in the management of such disease entities as ventricular arrhythmias (bilateral partial stellectomy) or heart failure (vagus nerve stimulation (VNS) therapy).

Anatomical basis of functional control. A considerable literature exists outlining the gross anatomy of intrathoracic neurons and their ganglia and nerves that

innervate the human heart (Kuntz & Morehouse, 1930; Saccomanno, 1943; Mizeres, 1963; Randall et al. 1972; Hopkins & Armour, 1984). With respect to the entire cardiac neuraxis, focusing at the level of the heart, human intrinsic cardiac ganglia are known to contain a complex neural network involving different neuronal anatomical subtypes that include unipolar neurons (putative afferent neurons), along with cholinergic and adrenergic neurons that presumably represent the two major motor phenotypes (Hoover et al. 2009; Pauza et al. 2014). Intrinsic cardiac ganglia also possess relatively large diameter neuronal somata that form clusters of rosettes within single ganglia. These somata frequently project axons centrally to interconnect primarily with dendrites of similar somata within that ganglion, an anatomical arrangement that forms an anatomical substrate for somatal interactions within or among adjacent intrinsic cardiac ganglia, which are second order neuronal interactions. Based on this anatomy, intrinsic cardiac ganglionated plexus neurons may function not only as an efferent neuronal relay station under medullary and




Figure 1. Cardiac neurotransmission DRG, dorsal root ganglion; ICN, intrinsic nervous system of the heart. Figure modified from W. Jänig (Janig, 2006) with permission.

spinal cord (adrenergic) motor control, but also as an important element in cardiocentric local reflex control (Armour, 2004).

Mammalian intrathoracic extra-cardiac ganglia contain efferent sympathetic postganglionic, afferent and local circuit neuronal somata. In fact, their varied neuronal types and synaptology represents the morphological substrate for various cardiocentric reflexes that depend on peripheral neuronal interactions involved in coordinating regional cardiac indices. These cardiocentric reflexes depend on intrathoracic afferent neuronal transduction of the cardiovascular milieu via intrathoracic local circuit neurons to cardiomotor neurons depicted above.

Cardiovascular afferent neurons. The somata of afferent neurons associated with cardiac or great thoracic vascular mechano- and/or chemo-receptors are located in nodose

ganglia (Thoren, 1977; Hopkins & Armour, 1989; Armour et al. 1994), as well as through the C6-T6 dorsal root ganglia (Vance & Bowker, 1983; Hopkins & Armour, 1989) bilaterally. The cardiac sensory neurites associated with these afferent somata are located throughout both atria and ventricles, being concentrated in the sinoatrial nodal region as well as in the outflow tracts of each ventricle (Armour & Ardell, 2004). They transduce (1) regional mechanical deformation, (2) the local chemical milieu, or (3) both - thereby displaying multimodal transduction properties, depending in large part in which of the above ganglia they are located (Armour & Ardell, 2004). Most cardiac afferent neurons in intrinsic cardiac ganglia also display multimodal transduction properties. A major exception to such multimodal transduction is the population of sensory neurons in middle cervical and stellate ganglia, associated with mechanosensory neurites




Figure 2. Simplified construct of neurohumoral control and functional organization of cardiac innervation

Aff, afferent; β , β -adrenergic receptor; C, cervical; DRG, dorsal root ganglion; G_i , inhibitory G-protein; G_s , stimulatory G-protein; L, lumbar; LCN, local circuit neuron; M, muscarinic receptor; T, thoracic. Additional pathways, for example nitrergic neurotransmission, are not shown.

located particularly along the inner arch of the aorta (Kember *et al.* 2004),(Armour & Ardell, 2004). Their inputs initiate central (nodose and dorsal root ganglion) and peripheral (intrathoracic extra-cardiac and intrinsic cardiac) reflexes (Armour & Ardell, 2004).

Cardiac motor neurons. Cardiac motor neurons have long been classified as belonging to the cholinergic *versus* adrenergic categories of the ANS. These two motor neuronal populations generally exert opposing effects with respect to the electrical and mechanical indices of the atria and ventricles.

Cholinergic efferent neurons. The somata of medullary parasympathetic efferent preganglionic neurons innervate cardiac parasympathetic efferent postganglionic neurons located in every major intrinsic cardiac ganglionated plexus. Their preganglionic components are located primarily in the ventral lateral region of the nucleus ambiguus, with fewer being located in the medullary dorsal motor nucleus and a scattering of somata located in the intermediate zone between these two medullary nuclei (Hopkins *et al.* 1996).

Parasympathetic efferent postganglionic neurons found in individual atrial or ventricular ganglionated plexi receive input from both vagal trunks. In turn, parasympathetic efferent postganglionic neurons located in individual atrial and ventricular ganglionated plexi influence both atrial and ventricular tissues. Therefore, cholinergic motor control of the heart is widely distributed (Ardell *et al.* 2015).

Adrenergic motor neurons. Cardiac-related sympathetic efferent pre-ganglionic neurons located in the intermediolateral cell column of the spinal cord project axons via C7-T6 rami (Norris et al. 1974, 1977) to cardiac post-ganglionic neuronal somata located in superior cervical, middle cervical, mediastinal ganglia and stellate ganglia (Hopkins & Armour, 1984). Given that direct innervation from T3-T4 ganglia to the heart has not been confirmed, it is possible that fibres from these cardiac neurons travel up to the stellate ganglion, and join cardiopulmonary nerves as they travel to the heart (Janes et al. 1986). This is an area that requires more study. These postganglionic neurons from sympathetic ganglia synapse in atrial and ventricular myocardium, releasing noradrenaline at their nerve terminals, in addition to other neurotransmitters, such as neuropeptide Y and galanin, which can inhibit acetylcholine release from parasympathetic terminals (Herring et al. 2008, 2012). The middle cervical ganglia, which reside superior to the stellate ganglia, contain many cardiac sympathetic neurons that send direct projections to the heart as well as interneurons that mediate cardiac reflexes (Armour, 1985; Kostreva et al. 1985; Tomney et al. 1985). In addition, there are multiple afferent mechano-sensory fibres that travel from the heart, and either synapse within the middle cervical or stellate ganglia or transit through these ganglia and synapse in the dorsal root ganglia of the spinal cord, transmitting information from the heart to the central nervous system (Malliani et al. 1973; Bosnjak & Kampine, 1989). Efferent postganglionic neurons throughout any peripheral sympathetic ganglia target all cardiac regions. Thus, no anatomical topology exists within peripheral autonomic ganglia with respect to the clustering of sympathetic efferent postganglionic neurons which innervate specific cardiac regions (Armour & Ardell, 2004). Sympathetic efferent postganglionic neurons in any major atrial or ventricular ganglionic plexus exert control over widely divergent regions of the heart (Cardinal et al. 2009). It should be recalled that such divergence of control resides within its local circuit neuronal populations.

Anatomical basis of peripheral neuronal interactions – fundamental importance of intrathoracic local circuit neurons. Intrathoracic ganglia have long been considered to act as simple efferent monosynaptic relay stations. In such a thesis, they represent waystations that distribute centrifugal efferent sympathetic vs. parasympathetic information from spinal cord and medullary neurons to cardiac tissues, in a reciprocal fashion (Mitchell, 1956; Levy & Zieske, 1969; Henning et al. 1990; Carlson et al. 1992). The intrinsic cardiac nervous system – as a parasympathetic efferent neuronal way station – acts to suppress cardiac indices all under central neuronal control (Henning et al. 1990).

The mammalian intrinsic cardiac nervous system processes both centrifugal and centripetal information in the ongoing coordination of regional cardiac indices (Armour & Ardell, 2004). In that regard, it is relevant to understand that intrathoracic ganglia, including those on the heart, contain a sizable population of neurons which do not project axons out of their respective ganglion or ganglionated plexi to neuronal somata restricted to other intrathoracic ganglia (Murphy et al. 1994; Sica et al. 1994; Yuan et al. 1994; Armour & Ardell, 2004). Neurons projecting axons solely to adjacent neurons are called interneurons. Those which project axons to neurons in other intrathoracic ganglia have also been termed local circuit in nature (Armour & Ardell, 2004). In a number of instances their anatomical arrangement is such that their somata form rosettes in intrathoracic ganglia, projecting centrally in their respective ganglion to solely interact with other somata within one ganglion. In this arrangement - individual neurons only interacting with adjacent somata - presumably is one anatomical substrate for peripheral interactivity (Armour, 2004). Presumably such an anatomical arrangement forms part of the basis whereby significant populations of intrinsic

cardiac neurons, for instance convergent local circuit neurons (Beaumont *et al.* 2013), initiate interneuronal interactions in the periphery (Armour & Ardell, 2004). Such local control persists when the intrinsic cardiac nervous system is chronically disconnected from neurons in the intrathoracic extra-cardiac ganglia and the central nervous system (Ardell *et al.* 1991).

The complexity of peripheral anatomy, upon which functional interactions occur among intrinsic cardiac neurons, has been described. Additional interactions with intrathoracic and central neurons in normal cardiac control add even more complexity (Kember et al. 2013b). As such, intrathoracic neurons interact not only with neurons in their same ganglion, but also with neuronal somata in other intrinsic cardiac ganglia via axo-axonal and axo-dendritic synapses (Gray et al. 2004). They also do so with neuronal somata in intrathoracic extra-cardiac ganglia, providing overall cardiocentric control (Ardell et al. 2015; Beaumont et al. 2015). The synaptology of such interactivity employs various neurotransmitters, including a variety of peptides, excitatory and inhibitory amino acids, nitric oxide donors, etc., all under the control of central neurons (Yuan et al. 1994; Armour, 2004).

Peripheral (intrathoracic) control. While our understanding of the anatomical basis of the multitude of central and peripheral reflexes in which such neurons are engaged remains incomplete, a tentative organization of the intrathoracic cardiac nervous system has been proposed in which cardiac afferent neurons influence, primarily via local circuit neurons, autonomic motor neurons via multiple feedback loops (Armour & Ardell, 2004). For instance, the varied peripheral (intrathoracic) reflexes so engendered represent cardio-centric short-loop reflexes that regulate cardiodynamics on a beat-to-beat basis. They continue to do so even when the intrathoracic nervous system is disconnected from the central nervous system (Arora et al. 2000; Murphy et al. 2000). In addition, it should be appreciated that a number of neurochemicals modify the neurons involved in intrathoracic extra-cardiac and intrinsic cardiac reflexes (Murphy et al. 1994; Yuan et al. 1994; Hoover et al. 2009). It is known that both excitatory and inhibitory synapses are present in intrathoracic reflexes. How a given population of intrathoracic extra-cardiac or intrinsic cardiac neurons interact in the coordination of regional cardiac indices remains very much dependent on the nature and content of sensory neuronal inputs arising from adjacent cardiovascular structures and from distant tissues (Kember et al. 2004). The latter reflect inputs from central neurons that are involved in whole body transduction. Thus, the transduction of the immediate past, as reflective of ganglionic capacity to exhibit memory, is very much dependent upon local circuit neuronal population function (Kember et al. 2013a).

Perspectives. The anatomical evidence provided above, as related to function, indicates that the peripheral (cervical and intrathoracic) nervous system acts as a distributive processor under the control of the CNS. Its multiple nested feedback control loops, each with different latencies of function, modulates regional cardiac indices throughout each cardiac cycle to assure adequacy of cardiac output in physiological states. What is now becoming evident is the fact that such complex control depends upon the behaviour of its various neuronal elements given the fact that its peripheral reflexes exert considerable influence on regional cardiac rate and force. Thorough understanding of the nerves controlling the heart is a pre-requisite for therapeutic neuromodulation.

Questions and controversies

- What are the functional capabilities of peripheral vs. central reflexes in control of cardiac function?
- What are the roles of support cells (glia) and small intensely fluorescent cells in modulating ganglia function?
- How do sex specific hormonal differences impact autonomic control of the heart?

Cardiorespiratory interactions. There is compelling evidence for strong interactions between cardiac and respiratory control, with such interactions playing an important role in an efficient and effective coordination between the lungs and the circulatory system. This interaction is manifest physiologically by phenomena such as sinus arrhythmia, which consists of cardiac acceleration during inspiration due to the vagolytic effects of lung expansion. Other manifestations are evident using approaches such as frequency domain analyses of breathing, heart rate, blood pressure and muscle sympathetic nerve activity (MSNA) (Pagani et al. 1997). Here breathing frequency entrains rhythmic changes in the other three variables. Cardiorespiratory interactions are especially evident in the responses to stressors such as hypoxaemia and hypercapnia. Hypoxaemia affects primarily peripheral chemoreceptors in the carotid bodies and hypercapnia acts mainly through central chemoreceptors in the brainstem. Both hypoxaemia and hypercapnia result in hyperventilation, modest increases in sympathetic nerve traffic to the muscle vascular bed and consequent increases in blood pressure (Somers et al. 1989; Somers et al. 1992). The overriding importance of hyperventilation in defining the neural circulatory response to hypoxaemia and hypercapnia is evident during apnoea. Lung inflation during hyperventilation acts to inhibit central sympathetic outflow. Therefore during apnoea, absence of lung inflation in the setting of hypoxaemia and/or hypercapnia results in disinhibition of sympathetic outflow, with consequent marked increases in sympathetic vasoconstrictor activity, and hence in blood pressure (Somers *et al.* 1989, 1995).

In contrast to hypercapnia, hypoxaemia has the added effect of increasing cardiac vagal drive (Daly et al. 1979). Again, the cardiovagal response to hypoxaemia is inhibited by lung inflation that occurs as a consequence of hyperventilation during hypoxaemia. Therefore, when apnoea is imposed on a hypoxaemic stress, the absence of lung inflation, with resulting vagal disinhibition, can manifest as significant bradyarrhythmias including prolonged asystole (Fig. 3) (Somers et al. 1992). This is part of the diving reflex which allows diving mammals such as sea lions to breathe atmospheric air but still continue to function for extended periods under water. The widespread sympathetic vasoconstriction during submersion and apnoea limits and 'redirects' use of the finite oxygen reservoir preferentially to the heart and brain, which undergo auto-regulatory vasodilatation, at the expense of more 'dispensable' tissues such as the muscle and kidney. Vagally mediated bradycardia enables less myocardial oxygen demand and improved myocardial perfusion due to the prolonged diastole. Evolutionary preservation of the diving reflex in humans, particularly in young children, helps explain the ability of humans to survive prolonged underwater immersion, for periods of up to 30 min (Thalmann et al. 2001). With subsequent circulatory, respiratory and renal support, and intensive care, it is possible that these individuals can eventually recover and be physically and neurologically intact despite the extended period of hypoxaemia and apnoea during immersion (Thalmann *et al.* 2001).

While respiratory-mediated chemoreflex activation can significantly influence neural circulatory control, circulatory mechanisms such as the baroreflex can in turn modulate chemoreflex sensitivity. Specifically, baroreflex activation, whether by raising blood pressure using intravenous phenylephrine or by neck suction, can attenuate the ventilatory, sympathetic and vagal responses to activation of the chemoreflex (Heistad *et al.* 1974; Somers *et al.* 1991, 1992). This appears particularly true for the response to hypoxaemia rather than hypercapnia (Somers *et al.* 1991), suggesting a specific interaction between the peripheral chemoreceptors in the carotid bodies and the arterial baroreflex in the carotid sinus, occurring probably at the level of the nucleus tractus solitarii.

Cardiac interoception. Cardiac interoception refers to the process of sensing, storing and representing information about the state of the cardiovascular system (Sherrington, 1961; Cameron, 2002). This information is represented at non-conscious as well as conscious levels (Fig. 4A), and a majority of early studies examined non-conscious mechanisms (Ádám, 1998; Dworkin, 2007). Cardiac interoceptive information is relayed predominantly via baroreceptors (Dworkin et al. 2000), chemoreceptors (O'Regan & Majcherczyk, 1982), the renin–angiotensin–aldosterone system (Re, 2004), the ANS (Janig, 1996) and the intrinsic cardiac nervous system (Rajendran et al. 2016), and most cardiovascular diseases




Figure 3. Autonomic influences on cardio-respiratory interactions
Recordings of intra-arterial blood pressure (BP) central venous pressure (CVP), electrocardiogram (ECG), sympathetic nerve activity (SNA) and breathing (RESP) in a healthy young subject during voluntary end expiratory apnoea while awake. Chemoreflex activation during apnoea results in progressively increasing sympathetic outflow to muscle blood vessels, simultaneous with vagally mediated sinus bradycardia with AV block (note the P-wave with absent QRS). The simultaneous activation of vascular sympathetic outflow and cardiac vagal drive during apnoea represents components of the diving reflex. Reproduced with permission from Somers et al. (1992).

A Cardiovascular neural representation Prefrontal cortex **VMPFC** Orbitofrontal Cingulate cortex pACC/dACC/rACC/pCC Amygdala/BNST Insular mid/posterior Somatosensory: Hippocampus anterior Primary and secondary Thalamus Cerebellum VM/VPL Spino/cerebral Hypothalamus PVN/LHA Periaqueductal gray Parabrachial nucleus NTS **RVLM** DVN/NA IML column/ICNS Parasympathetic Parasympathetic inhibition activation Cardiovascular tone: HR, HRV, BP, baroreceptor, RAAS ${\it B}$ Cardiac interoceptive prediction coding Outcome Illness state Symptom misinterpretation Autonomic hypervigilance Interoceptive Bodily preoccupation prediction error large Avoidance behaviours Anxiety Depression Predicted cardiac Incoming cardiac = Different? VS signals (expected) signals (actual)

The Journal of **Physiology**

Healthy state

Recovery

Resilience

Intact learning

Interoceptive

prediction

error small

Figure 4. Models of Cardiac Interoception

!!!

???

Prior

experience

A, model of cardiovascular neural representation. Cardiovascular information is relayed in ascending fashion via visceral and somatosensory afferent pathways, which project through brainstem and thalamic relay stations, ultimately reaching the insula and somatosensory cortical regions. At each level this sensory information is hierarchically processed, which results in increasingly conceptual representations that are polymodal (i.e. heterogeneous) and can be integrated with emotional states and conscious visceral perceptions. The primary sites

are expressed via abnormalities in these systems. However, cardiac symptoms can reflect very different forms of pathology, or none at all. For example, palpitations can indicate symptoms of a cardiac arrhythmia, a panic attack, or physical exertion. Cardiac patients also occasionally develop comorbid anxiety or depressive disorders, which further complicates the role of interoception and symptom reporting in these individuals.

Measuring cardiac interoception. The methods used to assess human cardiac interoception vary widely. Most tasks assess heartbeat perception under physiological resting conditions. Popular tasks include silently counting heartbeats during pre-specified time periods ('heartbeat counting; Schandry, 1981), tapping a button to indicate each felt heartbeat ('heartbeat tapping'; Ludwick-Rosenthal & Neufeld, 1985), or comparing the heartbeat sensation with tones presented simultaneously or non-simultaneously with the heartbeat ('heartbeat detection'; Whitehead et al. 1977; Katkin et al. 1982; Brener & Kluvitse, 1988). Each task has drawbacks. Heartbeat counting is confounded by the influence of prior knowledge of one's own heart rate (Ring et al. 2015) and reduced sensitivity to heart rate changes (Windmann et al. 1999). Heartbeat tapping provides temporal detail about each felt heartbeat (unlike heartbeat counting), but requires a sophisticated analysis approach (Davidson et al. 1981; Canales-Johnson et al. 2015). Heartbeat detection provides statistical measures of individual accuracy (Schneider et al. 1998; Katkin et al. 2001; Khalsa et al. 2008), but is complex to implement. However, when performed under physiological resting conditions, all of these tasks fail to measure changes in perceived cardiac intensity. This is a key limitation as dynamic disruptions in cardiovascular homeostasis are frequently associated with clinically and/or emotionally significant events. Furthermore, all of these tasks exhibit the finding that cardiac interoception is quite low at baseline. Less than 35% of individuals accurately detect cardiac sensations under resting conditions, suggesting that the majority of individuals do not perceive cardiac sensations at baseline (Jones, 1994). It is possible to more accurately assess individual differences in interoceptive intensity by increasing sympathetic tone. For example, bolus infusions of isoproterenol reliably induce brief increases in cardiac interoception (Khalsa *et al.* 2009*a*). During this 'infusion task', participants concurrently rate the intensity of perceived cardiac sensations while receiving isoproterenol in a double-blinded placebo-controlled manner. At sufficiently high doses cardiac intensity changes are correctly perceived by 100% of individuals. Although this task is invasive, it provides good experimental control over the cardiovascular system and enables accurate measurement of all facets of cardiac interoception, in every subject (Khalsa *et al.* 2015*a*).

Factors modulating cardiac interoception. Cardiac interoception decreases with age (Khalsa et al. 2009a) and increasing body mass index (Rouse et al. 1988; Kleckner et al. 2015). It increases acutely during exercise (Jones & Hollandsworth, 1981; Montgomery et al. 1984; Schandry et al. 1993) and after caffeine ingestion (Zoellner & Craske, 1999). Sex differences have sometimes been reported (Jones & Hollandsworth, 1981; Katkin et al. 1981), but there is debate whether these reflect true differences (Rouse et al. 1988; Ring & Brener, 1992) and there is currently no clear consensus on this question. Similarly, patients with panic disorder frequently report increased sensitivity to heartbeat sensations, but whether that perception of the heartbeat is accurate is unclear. Numerous studies under resting conditions fail to clearly demonstrate whether panic disorder patients perceive heartbeat sensations differently or have a bias towards reporting such feelings (Ehlers & Breuer, 1996; Van der Does et al. 2000; Domschke et al. 2010). This contrasts with many studies showing that panic disorder patients perceive cardiovascular sensations more intensely when stimulated pharmacologically via caffeine (Charney et al. 1985; Zoellner & Craske, 1999), carbon dioxide (Rassovsky & Kushner, 2003), sodium lactate (Dillon et al. 1987), yohimbine (Gurguis et al.

where conscious integration typically begins is at the level of cortical regions, including the insula, somatosensory areas, hippocampus and amygdala. These representations are modulated predominantly by the medial, orbital and cingulate regions of the prefrontal cortex, which issues visceromotor commands that propagate back through the system resulting in alterations in cardiovascular tone as well as changes in cardiac prediction signals. *B*, cardiac interoceptive prediction coding. The brain's active ongoing comparison between incoming and predicted signals results in calculation of an interoceptive prediction error. Disruptions in cardiovascular homeostasis result in large interoceptive prediction error signals, and this can produce illness states if the system is unable to adequately adapt. Boxes with grey outlines indicate neuroanatomical structures, and black arrows indicate anatomical connections. Boxes with blue outlines indicate conceptual processes, and blue arrows indicate information transfer between anatomo-functional systems and conceptual systems. BNST, bed nucleus of the stria terminalis; BP, blood pressure; dACC, dorsal anterior cingulate cortex; DVN, dorsal motor nucleus of the vagus; HR, heart rate; HRV, heart rate variability; ICNS, intrinsic cardiac nervous system; IML, intermediolateral column; LHA, lateral hypothalamic area; NA, nucleus ambiguous; pACC, pregenual anterior cingulate cortex; PVN, paraventricular nucleus; pACC, pregenual anterior cingulate cortex; PCC, posterior cingulate cortex; VMPFC, ventromedial prefrontal cortex.

1997), cholecystokinin (Schunck *et al.* 2006) or isoproterenol (Pohl *et al.* 1988). Another consideration is whether there is an inherent threshold for cardiac sensation. This threshold is likely to differ from person to person, but might be modulated by organic or psychiatric diseases, and/or past experiences.

Cardiac interoception in cardiovascular disorders. Cardiac interoception is possible when a failing heart is mechanically assisted with external devices (Couto et al. 2013), and even when another person's heart is transplanted inside the chest (Barsky et al. 1998). This suggests cardiac interoception can include non-viscerosensory pathways. At rest there appears to be no difference in the ability to detect the heartbeat in the chronic period following myocardial infarction (Jones et al. 1985) or hypertension (O'Brien et al. 1998; Koroboki et al. 2010), or following diagnosis with cardiac or non-cardiac chest pain (Schroeder et al. 2015). However, several studies have observed heightened cardiac interoception in palpitation patients with abnormal ECGs (Barsky et al. 1993; Ehlers et al. 2000). The explanatory mechanism for this finding is unknown, though it is possible that palpitations are atypical autonomic events that more clearly differentiate the heartbeat signal from other bodily sensations (Zamir et al. 2012). It is well known that some patients with arrhythmias, such as atrial fibrillation (AF), frequently mistake normal heart rhythms for recurrence of AF, or fail to correctly detect the presence of AF (Flaker et al. 2005; Hindricks et al. 2005). On the other hand, AF patients with anxiety and depression are more likely to incorrectly overestimate the presence of AF (Garimella et al. 2015). Whether such patients reflect a unique subpopulation, and whether interventions targeting anxiety and depression can improve AF symptom burden are currently unknown.

Predictive coding and neurovisceral integration. Human studies of cardiovascular sensation support a role for the insular cortex, but they also identify a broader network of cortical brain regions including the somatosensory, anterior cingulate and dorsomedial prefrontal cortices, as well as thalamus and cerebellum (Cameron & Minoshima, 2002; Critchley et al. 2004; Khalsa et al. 2009b; Kern et al. 2013). New perspectives also suggest that interoceptive perceptions are constructed by the brain, indicating that interoceptive experiences may substantially reflect predictions about the state of the body (Seth & Critchley, 2013; Barrett & Simmons, 2015). When the actual state of the body differs from the prediction, this results in a so-called 'prediction error' that is followed by active attempts to modulate the relative value of incoming inputs in order to restore balance (Paulus & Stein, 2010) (Fig. 4B). The brain may thus try to reduce interoceptive prediction error by attenuating the processing of incoming sensory input, by triggering changes in the body that resemble the expected (i.e. predicted) input, or by altering perceptual inferences about bodily states. These observations have contributed to a model of neurovisceral integration, which postulates that a series of structures in the brain are hierarchically organized to mediate emotional experience and vagal control (Thayer & Lane, 2000; Lane et al. 2009). Greater complexity and differentiation of experience is mediated by intact medial prefrontal mechanisms and greater vagal regulation, which are electrically stabilizing for the myocardium. By contrast, emotional disorders characterized by anxiety and depression are associated with more persistent and unchanging dysfunctional emotional states and reductions in vagal tone (Thayer & Lane, 2000, 2009). The latter is not altered by serotonin-specific reuptake inhibitors (Kemp et al. 2010), which are the most frequently prescribed medications for anxiety and depressive disorders. This highlights the need for alternative treatments that are potentially more cardioprotective.

Questions and controversies

- What governs differences in inherent ability to sense the heart? Some patients fail to detect life-threatening pathological alterations in cardiovascular tone (e.g. arrhythmia or myocardial ischaemia), whereas others misinterpret cardiovascular signals and are convinced of impending doom despite a normal workup.
- Does the onset of a cardiac arrhythmia or anxiety disorder enhance the ability to sense heartbeat signals, or conversely, does it increase the brain's prediction bias towards detecting such signals?
- To what extent is the insular cortex responsible for the ability to feel the heartbeat relative to other somatosensory brain regions?
- Is there actually a difference between the sexes in the ability to feel the heartbeat?

Pathophysiology

Myocardial remodelling. The concept and importance of cardiac remodelling are well established. The recognition that cardiac neural structures remodel in a variety of disorders is a more recent observation. Neural remodelling may be positive if it improves organ function, but it may be negative if it leads to electrical instability and arrhythmogenesis. This section will focus on recent developments in our understanding of neural remodelling in human diseases.




Figure 5. Intramyocardial neural remodelling following cardiac injury

A, nerve sprouting in explanted, transplanted heart, with regions of replacement fibrosis due to transplant vasculopathy (\$100 immunohistochemical staining, \times 400). B, autopsy specimen from a patient dying from coronary artery disease. Note nerve sprouting at junction of healed myocardial infarction (MI) and normal myocardium (NM) (\$100 immunohistochemical stain, \times 200). C, left atrial appendage from patient with coronary artery disease undergoing coronary artery bypass surgery. Note marked nerve sprouting in atrial tissue (\$100 immunohistochemical staining, \times 400). D, pulmonary vein from experimental animal with AF. Note pale cell (arrow) that has morphological features of cells of the conduction system (toluidine blue stain, \times 400) (the scale bar for A–D is shown in D). E, fluoroscopic location of multi-electrode catheter (left top and bottom) and electroanatomical map (centre) in a patient with ischaemic cardiomyopathy. On the electroanatomical map, the purple areas represent unscarred tissue with normal voltage while the dense grey represents dense scar. All other colours represent border zones (tissue with bipolar voltage \times 0.5 mV but \times 1.5 mV). Graph on the right shows change in activation recovery interval (Δ ARI) from baseline in normal (NL) and cardiomyopathic (CM) hearts in response to isoproterenol and nitroprusside. The Δ ARI is greatest in the CM–NL and scar regions of the cardiomyopathic heart. Border zones within each patient are the least responsive to isoproterenol. In response to nitroprusside, the scar and the CM–NL zones are least responsive, or paradoxically increase ARI compared with the border zone regions.

Heart transplantation. There is no argument that the transplanted heart has been separated from all anatomical connections with higher centres of the nervous system. Using quantitative electron microscopic techniques applied to endomyocardial biopsies, Rowan & Billingham (1988) found very few nerves present and concluded that normal myocardial innervation is not likely to be restored in transplanted human hearts. The conventional wisdom that severed nerves have a limited capacity to heal supported the notion that reinnervation of the transplanted heart was unlikely to occur. However, this was challenged by reports of chest pain in patients with transplant-related coronary artery disease. This was strong, perhaps irrefutable, evidence that some patients were experiencing reinnervation of the transplanted heart. Other direct lines of evidence came from the description of vasovagal reflexes in heart transplant recipients (Fitzpatrick et al. 1993). It subsequently became established that both sympathetic and parasympathetic reinnervation occurred in some patients (Kim et al. 2004), and was associated with improved peak oxygen uptake, heart rate and contractile function during exercise (Schwaiblmair et al. 1999; Bengel et al. 2001). An autopsy study of 29 patients who died after heart transplantation (Kim et al. 2004) found a decrease in total number of nerves (\$100-positive) over time, but an increase in growth associated protein 43 (GAP43) positive and tyrosine hydroxylase-positive nerves over time. These findings provided the first histopathological evidence of nerve sprouting (GAP43-positive) and specifically sympathetic nerve sprouting in the transplanted heart (Fig. 5A). More importantly, the neural remodelling that occurs in the transplanted human heart has a beneficial effect on cardiac performance (Schwaiblmair et al. 1999; Bengel et al. 2001). Unfortunately, sudden cardiac death is not uncommon in heart transplant patients (Vaseghi et al. 2009). The role of denervation/reinnervation in these deaths remains to be determined.

Ischaemic heart disease. Remodelling of the heart after myocardial infarction (MI) plays an important role in patient outcome. The pioneering work of Drs Jan and Marc Pfeffer established that pharmacological prevention of negative left ventricular remodelling after MI improves survival (Pfeffer & Pfeffer, 1988). In experimental models, much has been learned about remodelling of cardiac and extra-cardiac nerves after coronary occlusion (Vracko et al. 1991; Cao et al. 2000a,b; Ajijola et al. 2015). Studies of electrophysiological changes within and outside of the infarcted zone have been thoroughly reviewed (Vaseghi & Shivkumar, 2008). There are relatively few morphological studies of neural changes within the human heart after MI. Vracko et al. (1991) described the pattern of nerve fibres in human hearts with myocardial scars using S-100 immunohistochemical staining. This method detects Schwann

cells but not axons. Nerves devoid of Schwann cells are not decorated, and whether the nerves are sympathetic or parasympathetic cannot be determined. Vracko et al. observed increased nerve fibres at the edges of the scar tissue, aligned and oriented in the long axis of myofibres and adjacent to the scars. The investigators postulated that nerve fibres within the infarcted region were destroyed, and that the fibres observed at the edges of the scar tissue were the result of fibre regeneration. Trophic factors, such as nerve growth factor (NGF) and GAP43 have been shown in experimental animals to play a role in the nerve sprouting (Zhou et al. 2004). Unfortunately, in experimental systems of ischaemic heart disease, reinnervation is heterogeneous and hyperadrenergic, conditions known to be arrhythmogenic. Infusion of NGF into the stellate ganglion may cause increased cardiac sympathetic nerve sprouting and can facilitate sudden death in dogs with MI (Cao et al. 2000a) (Fig. 5B). Perhaps this is why β -adrenergic blockade reduces the risk of sudden death after MI.

In ischaemic heart disease, neural remodelling is not limited to myocardial structures. Indeed, in experimental systems, neurons and ganglia on the surface of the heart and remote from the heart demonstrate degenerative changes. In an elegant morphological study (Armour et al. 1997), epicardial ganglia from patients undergoing coronary artery bypass surgery were removed and studied by electron microscopy. Three types of ultrastructural abnormalities were identified: (1) lamellated inclusions, (2) membrane-bound whorls and parallel arrays of lightly stained membrane with fine granular material, and (3) concentric layers of lightly stained membranes with a darker granular core. These findings help to explain the recognized dysfunction of cardiac ganglionated plexi associated with experimental models of ischaemic heart disease (Huang et al. 1993a).

While much has been learned about morphological changes in the heart after ischaemic injury, knowledge is incomplete, and the process is complex. There are many factors that promote the degeneration and regeneration of neural structures after injury. It should not be surprising that neural stem cells participate in this process. In rat hearts studied 1 week after coronary occlusion, Drapeau et al. (2004) observed nestin-expressing neural stem cells in the myocardial scars. The findings of that study suggest recruitment of stem cells into the infarcted region, contributing to nerve fibre growth in the region of the infarct.

Atrial fibrillation. AF is the most common cardiac arrhythmia, and its complications are associated with significant morbidity and mortality. Morphological changes in the atria include dilatation, fibrosis and degenerative changes in atrial myocytes. A landmark study (Haissaguerre *et al.* 1998) established the importance

of the pulmonary veins in the generation of AF. The pulmonary veins not only contain cardiac-type myocytes, but are also highly innervated (Tan *et al.* 2006). There is growing evidence that remodelling in the left atrium and pulmonary veins contributes to the generation of AF. Much less is known about neural remodelling of the heart and its relationship to AF.

ANS activation has been shown to induce atrial tachyarrhythmias, including AF. The atria are the most richly innervated of the cardiac chambers. There are both intraand extra-cardiac autonomic nerves and ganglia that have the potential to remodel under a variety of conditions (Tan et al. 2006; Arora et al. 2007). In fact, labelled sympathetic nerve terminals in dogs with pacing-induced AF showed increased heterogeneous atrial sympathetic innervation (Jayachandran et al. 2000). In a similar model, Chang et al. (2001) confirmed the increase in sympathetic nerve density. Interestingly, AF and atrial nerve sprouting and sympathetic hyperinnervation also occur in experimental models of MI (Miyauchi et al. 2001). Nguyen et al. (2009) confirmed these findings in human biopsy specimens of patients who were having coronary artery bypass surgery and suffered from AF (Fig. 5C). Currently of great interest is the finding that remodelling of extra-cardiac nerves also occurs in patients with MI, AF and other conditions (Nguyen et al. 2011; Ajijola et al. 2015). Stimulation or inhibition of selective extra-cardiac neural structures may be a useful therapeutic option of cardiac arrhythmias and other abnormalities.

However, the story is even more complex. Cajal-like cells (Gherghiceanu *et al.* 2008), node-like cells (Masani, 1986), Purkinje-like cells (Chou *et al.* 2005), P cells and transitional cells (Perez-Lugones *et al.* 2003), and melanocyte-like cells (Levin *et al.* 2009) have been identified in human pulmonary veins (Fig. 5*D*). These cells have the potential to contribute to atrial arrhythmogenesis, but a direct contribution of such cells to human AF has not been proven.

Aerobic exercise, an important modulator of cardiac autonomic function, has a variety of beneficial effects on cardiovascular health. However, association between level and intensity of exercise and AF risk is known to exist (Andersen *et al.* 2013). Athletes, particularly as they get older, may be at increased risk for AF (Wilhelm, 2014). Whether this relationship is related directly or indirectly to autonomic function and/or neural remodelling is unknown.

Heart failure. There is ample evidence that neurohormonal activation plays a role in clinical heart failure (CHF). CHF is associated with increased levels of circulating catecholamines and decreased cardiac catecholamine levels. In 1993, in an experimental model of CHF, Himura *et al.* used glyoxylic acid (sucrose–potassium phosphate–glyoxylic acid)-induced histofluorescence and tyrosine hydroxylase immunohistochemical staining to show loss of noradrenergic nerve terminals in the failing ventricles (Himura et al. 1993). Cha et al. (2008) induced CHF in dogs by rapid ventricular pacing for 5 weeks. All paced dogs had atrial fibrosis, depressed left ventricular function and chamber dilatation. Dogs with CHF had marked heterogeneity of nerve density as measured by GAP43 immuno-staining with an overall decrease in nerve density. Of interest, dogs with the most nerves had a greater incidence of sudden death, while those with the fewest nerves had the lowest cardiac output. The investigators treated some dogs with omapatrilat, a vasopeptidase inhibitor that inhibits the renin-angiotensin system as well. The omapatrilat administration prevented the functional decline and negative myocardial and neural remodelling in CHF animals. While elevated sympathetic activity appears to be detrimental to the heart, parasympathetic stimulation appears to be cardioprotective, providing the basis for investigation of vagal stimulation for the treatment of cardiac disease (Li et al. 2004).

Myocardial infarction, in addition to causing formation of scars, leads to partial denervation of the fibres that innervate the myocardium at sites of injury. Given that the majority of the neuronal cell bodies whose axons have been damaged lie within the intrinsic cardiac ganglia and remain intact (Janes et al. 1986; Huang et al. 1993b; Yuan et al. 1994; Beaumont et al. 2013; Rajendran et al. 2016), re-innervation occurs after myocardial infarction, with evidence of nerve sprouts observed at the border zones of infarct (Cao et al. 2000a,b). These regions of denervation with nerve sprouts create heterogeneity in innervation, which is associated SCD risk. However, the functionality of these nerve sprouts remains unknown. In 1983, Barber and colleagues reported that infarcted in addition to viable regions distal to the infarct in a canine model no longer showed evidence of effective refractory period shortening in response to stellate stimulation, implying that many regions of the infarcted heart remained denervated. Furthermore, these regions had an exaggerated response to noradrenaline infusion, called denervation supersensitivity, suggesting that significant heterogeneity in response to circulating catecholamines might also be present in infarcted hearts. Whether these phenomena, however, existed in humans with MI and ischaemic cardiomyopathy (ICM) remained unknown. Vaseghi and colleagues studied the response to sympathetic activation in patients with ICM and ventricular tachycardia and compared them to control patients with normal hearts. They assessed the response of scar, border zone and viable regions in infarcted hearts to normal hearts after infusing isoproterenol, a direct β -1 and β -2 receptor agonist, and nitroprusside, which cause reflex activation of sympathetic nerve fibres in response to hypotension. They measured activation recovery interval (ARI), a surrogate of action

potential duration, and found that in infarcted hearts, ARI did not shorten in response to nitroprusside, even in viable peri-infarct regions, suggesting significant denervation, while all regions in control hearts showed evidence of ARI shortening (Fig. 5*E*). Furthermore, in response to isoproterenol, all regions in infarcted hearts showed a greater response (greater decrease in ARI) as compared to control hearts, confirming that denervation supersensitivity exists in patients with ICM and ventricular arrhythmias (Vaseghi *et al.* 2012).

Extra-cardiac neuronal remodelling. Following ischaemic and non-ischaemic cardiomyopathy (ICM and non-ischaemic cardiomyopathy (NICM), respectively), alterations in the structure, neurochemical phenotype, and function of neurons within the cardiac neural hierarchy but extrinsic to the heart occur (Fukuda *et al.* 2015). These sites include the stellate ganglion (SG), brainstem and higher brain centres.

Intra-thoracic ganglia. Studies of the intrathoracic ganglia, such as the stellate ganglia, have demonstrated enlargement, increased dendritic sprouting and neurochemical changes in SG neurons in animal models of ICM and NICM (Kanazawa *et al.* 2010; Han *et al.* 2011; Ajijola *et al.* 2015). From a mechanistic standpoint, studies in animal models in part link deregulated nitric oxide signalling (Li *et al.* 2007, 2013), and its subsequent impact on calcium handling in SG neurons (Wang *et al.* 2007; Lu *et al.* 2015), with enhanced sympathetic outflow seen in the hypertension, with implications for other causes and types of cardiovascular disease.

In humans, evidence of histological changes in ganglia associated with cardiac disease (sudden death in this case), was first reported by James et al. (1979). They reported on two women with sudden cardiac death, and post-mortem findings of inflammation within cardiac ganglia (i.e. epicardial and intra-myocardial ganglia). The term 'ganglionitis' was used to describe this phenomenon, and its associated malignant cardiac rhythm. A decade later, Pfeiffer et al. (1989) described the first case of a young patient with a structurally normal heart, but with severe inflammation of the SG, linking viral-mediated neuronal dysfunction in SGs to malignant arrhythmogenesis in this patient. Recently, in patients with congenital long QT syndrome (LQTS) and catecholaminergic polymorphic ventricular tachycardia (CPVT), Rizzo et al. (2014) observed evidence of T cell-mediated inflammation. In ganglia from patients with LQTS and CPVT, when compared to SG from control subjects with accidental death, CD3⁺ and CD8⁺ cells were present in greater quantities. Further, there was evidence of inflammation albeit low grade in all LQTS/CPVT patients studied.

Histological evidence of remodelling in stellate ganglia of patients with cardiopulmonary disease was first

presented by Docimo *et al.* (2008). They observed an increase in the number of nerve cell bodies within the left SG (LSG) of cadaveric subjects with cardiopulmonary disease, compared to other causes of death. In a follow-up study (Wood *et al.* 2010), the authors correlated fibrosis within the myocardium of cadaveric subjects (a marker of ICM), with the number of nerve cell bodies in the LSG, the implication being that ICM is associated with increased SG neuronal count. While differentiated stellate ganglion neurons are recognized to be post-mitotic, and not expected to divide, the mechanism underlying this increase remains elusive.

Consistent with findings in pre-clinical studies demonstrating neuronal enlargement and dendritic sprouting in SGs of animals with myocardial infarction, stellate ganglion neurons from humans with arrhythmias, and both ICM and NICM exhibit remodelling (Ajijola *et al.* 2012*b*). Mean neuronal size was greater in SGs from ICM and NICM patients compared to controls, with NICM demonstrating the largest neurons in the population. In this study the numbers of neurons did not differ across the groups, in contrast to Wood *et al.* (2010) and Docimo *et al.* (2008). Rather, the difference in neuronal size appeared to be a population shift from smaller to larger neurons. Specific findings in addition to neuronal size include *no* difference in the degree of intra-ganglionic fibrosis, synaptic density, or nerve sprouting amongst the groups.

Higher centres. Significant components of aberrant cardiovascular symptoms can arise from changes in higher central nervous system control over medullary sympathetic and parasympathetic output sites. Among the most common syndromes, neural injury that accompanies obstructive sleep apnoea (OSA) and heart failure (HF) have been well documented, and the resulting functional consequences from such damage are substantial. The impaired outcomes in both conditions typically include enhanced and unvarying sympathetic discharge, often a reduction in parasympathetic output, with resulting influences on cardiovascular characteristics, including hypertension, and phase-shifted or unresponsive blood pressure changes to autonomic, ventilatory, or motor challenges.

The sites of injury in OSA and HF preferentially target autonomic regulatory areas, as revealed by a number of structural magnetic resonance imaging (MRI) techniques, including voxel-based-morphometry (Macey et al. 2002; Woo et al. 2003), T2-relaxometry (Woo et al. 2009), and diffusion tensor imaging procedures (Kumar et al. 2011, 2012, 2014; Woo et al. 2015). Damage is often lateralized (Woo et al. 2003); since autonomic regulatory areas are unequally distributed in the left and right sides of the brain, the asymmetric nature of damage contributes to the inordinate sympathetic effects. Thus, in both OSA and HF patients, forebrain areas, such as the right insular cortex,

which plays a significant role in baroreflex modulation, are especially damaged, as is the right ventral medial prefrontal cortex, an essential component for blood pressure mediation, and the hippocampus, a major player in blood pressure control, along with mood and cognition (Woo et al. 2009; Kumar et al. 2011, 2012, 2014; Woo et al. 2015). Although injury is preferentially lateralized, often the damage is bilateral, with asymmetric severity. Other forebrain areas are similarly targeted, including portions of the cingulate cortex and cingulum bundle, and especially the anterior cingulate (Macey et al. 2008; Kumar et al. 2012), the entire extent of the hypothalamus, including the anterior hypothalamus and basal forebrain areas (Kumar et al. 2011). Midbrain and medullary sites are not spared; the raphe system, especially more-caudal raphe nuclei, and the right ventral lateral medullary nuclei are affected (Kumar et al. 2011). The cerebellar cortex and deep autonomic nuclei, the fastigial nuclei, which serve to dampen extremes of hyper- and hypotension (Lutherer et al. 1983, 1989), are extensively damaged in both OSA and HF subjects (Macey et al. 2002, 2008; Woo et al. 2003, 2009, 2015; Kumar et al. 2011, 2012, 2014). An overview of near-midline structural injury in HF is shown in Fig. 6A. Figure 6B shows a subset of axonal injury, derived from diffusion tensor imaging procedures in HF subjects.

Close examination of particular autonomic sites, especially the insular cortices, shows a topographical organization within cortical areas that is altered in OSA. The consequences of such injury are found in phase-delayed and muted heart rate responses to a range of autonomic challenges (cold-pressor, hand grip, Valsalva manoeuvre) (Macey *et al.* 2013, 2014). Both phase shifts

and amplitude changes are shown in Fig. 6*C*; heart rate responses between OSA and control subjects are shown in Fig. 6*D*.

The autonomic effects extend beyond responses to transient manipulation. The sustained increase in sympathetic nerve discharge in OSA (Fatouleh et al. 2014; Lundblad et al. 2014), and the poorly controlled hypertension in both OSA and HF patients are well described. In addition, AF is common in OSA (Holmqvist et al. 2015), and poorly regulated glucose control, possibly arising from impaired sympathetic regulation of glucagon, as well as other hormonal regulatory deficits in the syndrome, is exceptionally frequent (Rajan & Greenberg, 2015). The mechanisms underlying the remodelling of brain structures, which affect cardiovascular and other autonomic action, remain unclear. Heart failure patients show a high incidence of sleep disordered breathing, and may be exposed to similar epochs of intermittent hypoxia as OSA patients; intermittent hypoxia is especially injurious to both cerebellar Purkinje cells (Pae et al. 2005) and cardiac autonomic ganglia in neonatal animals (Soukhova-O'Hare et al. 2006; Ai et al. 2009). However, many other processes accompany sleep disordered breathing, including significant perfusion changes, alterations in CO₂, and large, transient changes in blood pressure with interrupted breathing. A critical aspect is maintaining the integrity of the blood-brain barrier function, which prevents entry of neurotoxic substances into the brain; however, that integrity is breached in OSA (Palomares et al. 2015) and HF subjects.

The evidence suggests that the injuries introduced by OSA and HF modify cardiovascular action, as




Figure 6. Remodelling in higher brain centres induced by chronic heart failure

A, mean diffusivity, a measure of tissue injury, in a midline and near-midline sagittal view of the brain in 16 HF patients; injury appears in basal forebrain (a, b), hypothalamic (c), cerebellar (d, e), posterior and mid cingulate (f, g), and dorsal medullary regions. B, axonal injury, indicated by axial diffusivity measures, showing image in axons from the hypothalamus (c), through the medial forebrain bundle (d), to the pons, and from the midline raphe to the cerebellum (a). Injury also appears in the corpus callosum (b). C, averaged functional MRI signals during the course of three Valsalva manoeuvres from 16 HF patients (red) vs. 33 healthy controls (blue); * indicates significance at P < 0.05. HF signals are unable to decline to the challenge, and are phase delayed. D, heart rate responses in 16 HF subjects (red) vs. 33 healthy age- and sex-comparable controls (blue) to three successive Valsalva challenges. The HF subjects show muted and phase-shifted heart rate changes to the challenges.

demonstrated by flawed neural responses to transient challenges, distorted cardiovascular responses to those challenges, and altered baseline cardiovascular activity. However, the injury is not all-or-none; the data suggest that the injury is progressive, leaving the possibility that early intervention can halt or reverse the damage. Both OSA and HF are accompanied by profuse sweating and other aspects of high fluid loss, leaving the possibility of high nutrient loss that can be corrected. Interventions to support the blood–brain barrier and acute damage may provide a mechanism to prevent such injury.

Ouestions and controversies

- What are the mechanisms underlying neuronal damage in OSA?
- Can these changes be ameliorated by neuromodulation?
- Is the reversal of pathological neuronal remodelling a return to baseline?
- Why do most OSA subjects remain hypertensive even after breathing treatment?
- Does brain tissue injury recover after heart transplant in HF subjects, or after interventions for breathing in OSA?

Autonomic testing

A number of indices are available to assess autonomic function in humans. Key measurement techniques include heart rate, heart rate variability (HRV), plasma catecholamine spillover and microneurographic recordings of peripheral sympathetic activity. Each of these techniques has strengths and limitations, and the interpretive, logistical and technical challenges of using them vary by several orders of magnitude. Importantly these techniques can be coupled with specific manoeuvres that provide a homeostatic challenge and can be used to diagnose specific autonomic disorders in patients and the response to therapy (Low, 2003). In this context, the utility of several commonly employed tests will be highlighted next.

Sympathoexcitatory stressors. Several forms of acute sympathoexcitatory stress can be used to test autonomic responses in humans. Most commonly these include mental stress of various forms, the cold pressor test and isometric handgrip exercise (Liu *et al.* 2009). After a period of baseline measurements several minutes of stress are imposed and the responses of interest are measured. These responses might be as simple as blood pressure and heart rate or include the more complex measures mentioned above. Importantly, in normotensive subjects the pressor

response to acute sympathoexcitatory stress is highly predictive of the future risk of hypertension (Matthews et al. 2004). Additionally, isometric handgrip exercise and several minutes of post-exercise ischaemia can be used to physiologically dissect the mechanisms contributing to the rise in blood pressure associated with this intervention. The initial rise in pressure is driven largely by central command signals from the motor cortex followed by a major contribution from chemo-sensitive afferents in the contracting muscles driving a robust increase in MSNA (Mark et al. 1985).

Similar approaches have been used to study the rise in pressure and peripheral vasodilatation in the human forearm during mental stress. In humans mental stress does not cause a clear and consistent increase in sympathetic vasodilator outflow to the forearm but does evoke marked NO-dependent forearm vasodilatation (Dietz et al. 1994; Carter & Ray, 2009). This outcome contrasts with many animal species where well-described sympathetic dilator nerves exist in skeletal muscle (Joyner & Dietz, 2003). It is also important for any investigator using mental stress interventions to appreciate that there are complex interactions between the type and intensity of the stressor (puzzles, Stroop colour-word test, mental arithmetic, etc.) and the subject's perception of stress (Callister et al. 1992; Joyner & Casey, 2015). Some individuals might respond robustly to one form of stress but not another (Roddie, 1977).

Baroreflex testing. The ability of the autonomic and cardiovascular systems to respond to acute perturbations in blood pressure is largely regulated by stretch-sensitive receptors in the carotid sinus and aortic arch, the so-called arterial baroreflexes (Donald & Shepherd, 1980). These receptors respond to differing amounts of stretch caused by changes in arterial blood pressure and when stimulated by higher pressure they generally evoke reflex responses that serve to lower blood pressure including a reduction in heart rate and reduced sympathetic outflow, with directionally opposite responses seen when blood pressure falls. There are also stretch-sensitive receptors in the heart, great veins and lungs that respond to a number of stimuli related to the 'fullness' of the central circulation (Hainsworth, 2014).

By evaluating the physiological responses to beat-to-beat changes in blood pressure it is possible to use analytical techniques from engineering to calculate the threshold, gain, set-point and saturation characteristics of various baroreflex responses. This can be done using observational techniques and looking for 'spontaneous' sequences of rising or falling blood pressure that are associated with directionally opposite changes in heart rate (Diaz & Taylor, 2006). One general problem with this approach is that during most activities of life when

blood pressure is rising or falling, heart rate is also rising or falling in parallel with the changes in pressure. This means that only a limited number of heart rate and blood pressure sequences can be used in this type of analysis. Oscillations in arterial blood pressure occurring typically at 0.1 Hz in humans, known as Mayer waves, have been linked to cardiac autonomic tone (Julien, 2006). These oscillations are slower than respiratory frequency, and exhibit the strongest coherence to MSNA. Mayer waves have also been reported in other species (0.3 Hz in rabbits, 0.4 Hz in rats and 0.1 Hz in cats). Although the origin of Mayer waves remains debated, it is known that surgical denervation of aortic and carotid sinus baroreceptors eliminates these oscillations, suggesting that the baroreflex causes or at least modulates Mayer waves (Julien, 2006). During states of sympathetic activation, Mayer waves are enhanced, and have been investigated as a marker of sympathoexcitation (Cohen & Taylor, 2002). The clinical usefulness of Mayer waves remain to be determined. While the oscillations show correlation to sympathetic nerve activity in normal subjects, conditions such as heart failure, characterized by chronic sympathetic activation exhibit diminished oscillations. Mayer wave oscillations have been identified preceding vasovagal syncope (Lepicovska et al. 1992; ten Harkel et al. 1993); however, whether they cause or reflect the underlying haemodynamic and autonomic disturbance also remains

Another approach to baroreflex testing is to use vasoactive drugs to raise and lower blood pressure and then measure the heart rate or MSNA responses to the changes in pressure (Smyth et al. 1969; Rudas et al. 1999). The advantage of this technique is that a wide range of pressures can be studied and a comprehensive set of indices of baroreflex behaviour can be generated. However, the stimulus is non-specific with contributions from essentially all of the barosensitive areas noted above. Another issue is that while drug infusions evoke reflex changes in both heart rate and sympathetic activity, these responses are not correlated (Dutoit et al. 2010). This means that caution must be used in making broad categorical statements about baroreflex function based on a given test or single physiological variable. When MSNA is the outcome variable of interest, it is possible to link the probability of a sympathetic burst with a given level of blood pressure and generate a relationship that provides insight into variables like gain typically obtained via drug infusions (Kienbaum et al. 2001; Wehrwein et al. 2010). Selective pressure or suction can also be applied to the neck to temporarily stimulate the carotid baroreceptors in relative isolation (Fadel et al. 2003).

It is also possible to evoke baroreflex responses by applying subatmospheric (e.g. negative) pressure to the lower body (LBNP) and cause graded venous pooling and simulate blood loss in humans (Cooke *et al.* 2004). For many years it was felt that low levels of LBNP primarily unloaded the cardiopulmonary receptors and could be used to study their responses. The idea was that 10–20 mmHg of LBNP did not cause measurable changes in arterial pressure and thus provided a 'selective' stimulus to the cardiopulmonary receptors. However, imaging studies demonstrated that even low levels of LBNP caused deformation of the aortic arch and raised questions about the selectivity of low levels of LBNP (Taylor *et al.* 1995).

Finally, tilt table testing and the Valsalva manoeuvre are two other techniques frequently used to study the behaviour of the ANS (Deng et al. 1998; Low, 2008). These are especially valuable in the testing of patients for diagnostic purposes, but in general are not as selective as the other laboratory-based techniques noted above. Elevated sympathetic tone and diminished parasympathetic tone are proarrhythmic and the evaluation of the ANS provides information beyond the use of the left ventricular ejection fraction (LVEF). These tests include HRV (Feldman et al. 2013; Guyenet et al. 2013), baroreflex sensitivity (BRS) (Billman et al. 1984; La Rovere et al. 1988; La Rovere et al. 1998; La Rovere et al. 1998), heart rate turbulence (HRT) (Schmidt et al. 1999; Ghuran et al. 2002), heart rate deceleration capacity (Bauer et al. 2006) and T wave alternans (TWA) (Ikeda et al. 2006).

Although a variety of tests exist to probe ANS function, they require standardization for accuracy and reproducibility in each lab (Malik, 1996). Some tests are time consuming (e.g. baroreflex testing), unpleasant to the patient (e.g. microneurographic recordings and cold pressor test), and are performed with extensive variability at many institutions. Despite these challenges, reliable and reproducible testing of cardiac autonomic function may have a significant impact in early detection and management of cardiovascular diseases.

Risk stratification

Alterations in autonomic function are often detected in patients with cardiac, and even non-cardiac, medical disease. These alterations are often associated with negative prognostic implications. The most commonly used measures to evaluate cardiac autonomic function rely on quantitating autonomic input at the level of the sinus node, i.e. some evaluation of heart rate or heart rate change. The simplest parameter, heart rate, reflects the net parasympathetic and sympathetic inputs, which in rest conditions in normal subjects is heavily weighted to be predominantly parasympathetic. HRV examines beat-to-beat changes in heart rate, and can be quantified using time domain techniques,

frequency domain techniques, or nonlinear analyses. Other evaluations focus on assessing the response of the ANS to small perturbations, such as changes in blood pressure (baroreceptor sensitivity), premature ventricular complexes (heart rate turbulence), and orthostatic hypotension. As demonstrated in the Swedish Malmo Preventative Study, a prospective study of over 33,000 men (Fedorowski et al. 2010), orthostatic hypotension was present in 6.2%, and was associated with hypertension and diabetes. The presence of orthostatic hypotension was associated with increased mortality and coronary event risk, independent of traditional risk factors. Finally, parameters of heart rate acceleration or deceleration with exercise or other conditions also provide a measure of autonomic responsiveness. Virtually all of these techniques provide prognostic information (La Rovere et al. 1998; Jouven et al. 2005; Exner et al. 2007; Lahiri et al. 2008; Bauer et al. 2009). While the roles for sympathetic activation as a trigger for life-threatening ventricular arrhythmias (Albert et al. 2000, 2003; Lampert et al. 2002), and for parasympathetic activation's protective effect (Vanoli et al. 1991; Ng, 2014) are well established, none of the heart rate-based parameters provides strong enough risk prediction to be useful clinically, particularly for sudden arrhythmic cardiac death. Randomized clinical trials of implantable defibrillator therapy in the early post-myocardial infarction period that have incorporated heart rate and HRV among the inclusion criteria have been negative (Hohnloser et al. 2004; Steinbeck et al. 2009). While it has been suggested that these parameters are not very predictive of sudden arrhythmic death that is reversible by implantable defibrillator therapy, these trials also included a low left ventricular ejection fraction criterion. Just as low left ventricular ejection fraction may be predictive in other settings, so may these parameters. Yet, in a recent meta-analysis, none of the heart rate-based parameters (HRV, baroreceptor sensitivity, heart rate turbulence) were significant predictors of arrhythmic events in patients with NICM (Goldberger et al. 2014).

It is therefore important that alterations in autonomic function are likely to track the severity of the underlying cardiac disease and may serve as a strong or stronger predictor of overall mortality rather than arrhythmic mortality. The trigger for ventricular tachyarrhythmias is often associated with sympathetic activation (Lampert et al. 2002), but many of the parameters studied focus on rest conditions or only mild perturbations of the ANS, raising the possibility that these evaluations may not be sensitive enough. In addition, how alterations in autonomic function at the level of the sinus node translate to effects on ventricular electrophysiology, the presumed source for its effect on ventricular arrhythmogenesis, has not been clearly demonstrated.

Ouestions and controversies

- Why are clinical autonomic indices limited in prediction of sudden death?
- Have autonomic variables largely failed in clinical use due to one-time measurement, rather than dynamic measurements over time?
- How do we develop robust 'reflex' measures of autonomic activity that can be applied easily to a large population?
- What barriers limit translation of population studies of autonomic indices to risk prediction in individuals?
- Is post-exercise vagal tone a useful marker of sudden risk in humans?

In summary, there are several important questions to consider when evaluating the prognostic role of ANS control of the heart in patients with cardiac disease. It is possible that different parameters may be better suited to evaluate overall mortality versus arrhythmic mortality. In particular, evaluation of autonomic control in stress *versus* rest conditions could be an important factor delineating risk. Specifically, assessment of reflex measures of autonomic function provide more information than tonic measures, which are affected by many variables (Wellens *et al.* 2014). In addition, further work focusing on the effects of alterations in autonomic function on ventricular electrophysiology may provide better insight into the risk for developing ventricular tachyarrhythmias.

Clinical management: specific interventions for heart failure, arrhythmias and other cardiorespiratory conditions


Cardiac sympathetic denervation.

Channelopathies. Despite its rapidly growing use in the management of life-threatening arrhythmias, cardiac sympathetic denervation (CSD) is not a novel therapy. Following a series of pioneering interventions which started 100 years ago as reported in detail elsewhere (Schwartz, 2014), the resurgence of interest for this approach was associated with initial reports in 1961 by Estes and Izlar (Estes & Izlar, 1961) and in 1968 by Zipes and colleagues (Zipes et al. 1968) for recurrent ventricular tachycardia and subsequently for the congenital long QT syndrome (LQTS) (Moss & McDonald, 1971; Schwartz et al. 1975). It was later extended to another channelopathy, catecholaminergic polymorphic ventricular tachycardia (CPVT) (Wilde et al. 2008; De Ferrari et al. 2015).

To better understand the role of CSD, its anatomical and physiological basis is important. CSD involves removal of the lower half of the stellate ganglion as well as

T2–T4 thoracic ganglia, and is typically performed via video-assisted thoracoscopic surgery (VATS; Figure 7*A* and *B*; Ajijola *et al.* 2012*a*; Vaseghi *et al.* 2014). Thus, the upper half of the stellate ganglia, as well as the middle cervical ganglia, remain intact with this procedure. This also means that despite CSD, some post-ganglionic efferent sympathetic innervation to the heart is preserved. However, since the middle cervical ganglia contain afferent fibres that transit through the stellate ganglia before reaching the spinal cord, afferent neurotransmission through the middle cervical ganglia, in addition to the lower half of the stellate ganglia, is interrupted. Therefore, CSD affects both efferent and afferent sympathetic neurotransmission.

A strong rationale exists for this therapy based on a large series of experimental studies (Schwartz, 2014). The major mechanism underlying the antiarrhythmic efficacy of left CSD (LCSD) is the reduction of noradrenaline release at neural terminals in the ventricles, a factor that increases heterogeneity of repolarization and facilitates the occurrence of ventricular fibrillation (Han & Moe, 1964). The largest series reported so far includes 147 patients and was published in 2004 (Schwartz *et al.* 2004). The number of cardiac sympathectomy procedures (mostly LCSD) being performed has increased steadily, and can be safely done in children and adults (Collura *et al.* 2009). The reported series focused on patients at unusually high risk as demonstrated by the fact that



Relationship of stellate and middle cervical ganglia are shown. Both ganglia contain interneurons as well as efferent and afferent neurons that both synapse and traverse through the ganglia. Only the stellate ganglia are anatomically connected to the spinal cord. *B*, the left sympathetic chain including the left stellate ganglion as well as the T2–T4 ganglia are shown intra-operatively behind the parietal pleura via video-assisted thoracoscopic surgery. *C*, implantable cardioverter defibrillator (ICD) shock free survival is better after bilateral than left cardiac sympathetic denervation (CSD) in patients with refractory ventricular tachycardia (VT) and structural heart disease. *D*, after CSD, in patient with structural heart disease, the burden of ICD shocks significantly improves at approximately 1 year of follow-up. Aff, afferent; DH, dorsal horn; Eff, efferent; IML, interomediolateral cell column; MCG, middle cervical ganglia.

99% of patients were symptomatic, and among them 48% had survived a cardiac arrest, their QT interval was extremely prolonged (mean QTc 543 \pm 63 ms), and 75% continued to have major cardiac events despite full-dose beta-blockade. During an 8 year follow-up, 46% became completely asymptomatic, aborted cardiac arrest occurred in 16% and there was 7% mortality. The mean yearly rate of cardiac events dropped impressively by 91% (P < 0.001), and the patients with more than five cardiac events decreased from 55% to 8% (P < 0.001). Among patients with electrical storms and multiple implantable cardioverter defibrillator (ICD) discharges, the median number of ICD shocks after LCSD decreased by 95% (P = 0.002) from 25 to 0. The conclusion was that LCSD produced a major but incomplete abolition of cardiac events in the high-risk LQTS patients; it was and still is recommended that the primary indication for LCSD is for patients who continue to have syncope despite beta-blocker therapy and for patients with multiple ICD shocks (Schwartz & Ackerman, 2013). In this case LCSD would improve quality of life using the ICD as a safety net. Indeed, the impact of LCSD on quality of life is a very important consideration that often physicians overlook (Antiel et al. 2015; Schwartz, 2015). CPVT patients, especially young ones, are not always fully protected by beta-blockers and, as a consequence, usually end up receiving an ICD. Unfortunately for them, the risk of electrical storm is devastatingly high. They are exquisitely sensitive to catecholamines and as the first ICD shock through the pain and fear will lead to catecholamine release, this often initiates a vicious cycle with multiple shocks because of recurrent ventricular tachycardia (VT)/ventricular fibrillation (VF). In 2008 Wilde et al. reported for the first time the efficacy of LCSD in CPVT patients (Wilde et al. 2008). There were only three patients but the follow-up was very long, exceeding 10 years, and showed a complete suppression of the cardiac events, which in all patients were multiple cardiac arrests. As the most recent guidelines (Priori et al. 2013) had considered LCSD only as a Class IIB recommendation, it was felt necessary to provide stronger evidence. Collura et al. subsequently reported the benefit of LCSD in two patients with CPVT at mean follow up of 16.6 \pm 9.5 months (Collura et al. 2009) and a recent worldwide dataset has been reported (De Ferrari et al. 2015). This study involved 63 CPVT patients who underwent LCSD and had a mean follow-up of 4 years. There were nine asymptomatic patients, who remained asymptomatic, and 54 patients with major cardiac events. This was a high risk group, as indicated by the fact that 33% had survived ≥ 1 cardiac arrest, that 43% had ≥ 1 appropriate ICD shock, that 26% had ≥ 1 electrical storms. Among these 54 symptomatic patients, 38 continued to have cardiac events despite optimal medical treatment. Within this group the percentage of patients with cardiac events dropped from 100% to 32% (P < 0.001) and among the 29 patients with an ICD implanted before LCSD the rate of shocks dropped by 93% (from 3.6 to 0.6 shocks/person/year, P < 0.001). In the internal control analysis in which each patient served as his or her own control, there was a 92% reduction in major cardiac events. The important observation was also made that the few instances in which denervation was incomplete (i.e. ablating less than from the lower half of the left stellate ganglion down to the fourth thoracic ganglion) recurrent arrhythmias were noted; this exemplifies the critical notion of the 'adequate dose' of denervation. In 2015, a large Canadian study (Roston et al. 2015) on CPVT patients provided additional data on the efficacy of LCSD. In 18 patients who underwent LCSD, the success rate was 89%, almost identical to the worldwide report (Roston et al. 2015).

These data force a reassessment of the current clinical approach to the management of CPVT. Beta-blockers should remain the first-line therapy, as they are effective in two-thirds of the patients. The combination of beta-blockers and flecainide appears promising but more long-term data are needed. The present data, showing that 32% of the patients suffer serious ICD-related adverse events within 7 years, should lead to a revision of the guidelines. Indeed, for CPVT patients with syncope despite beta-blockers, it appears that left CSD prior to consideration of an ICD implant is warranted (De Ferrari et al. 2015).

Structural heart disease. Multiple accounts of the benefits of CSD have emerged in patients with refractory VT and structural heart disease. (Zipes et al. 1968; Nitter-Hauge & Storstein, 1973; Lloyd et al. 1974; Schoonmaker et al. 1975) However, with the introduction of medications targeting the sympathetic nervous system (including beta-blockers and angiotensin converting enzyme (ACE) inhibitors) that significantly improved the risk of SCD (Kober et al. 1995; Hjalmarson, 1997), CSD took a secondary role. It gradually became clear that despite the best available anti-arrhythmic medications and catheter ablation techniques, approximately 30-50% of patients continue to experience recurrent arrhythmias and ICD shocks (Kuck et al. 2010; Mathuria et al. 2012). In this setting, CSD was once again revived as the therapy to treat recurrent ventricular arrhythmias and ICD shocks.

Given the success of LCSD in hereditary arrhythmias (LQT and CPVT), LCSD was initially tried in a case series of patients with refractory VT/VF storm and structural heart disease (Bourke *et al.* 2010). However, approximately 44% of patients did not benefit from the left sided procedure at follow-up. Of note, in this series, thoracic epidural anaesthesia showed a greater rate of partial or complete response, with approximately 75% of patients having a greater than 80% reduction in their arrhythmias

burden, pointing to the possibility of bilateral CSD, rather than left alone, may provide better results.

To better assess the effects of bilateral CSD, as with any treatment modality with significant translational and retrospective data, randomized clinical trials are critical. To this end, the PRophylactic Cervicothoracic Sympathectomy for PrEVENTion of Ventricular TAchyarrhythmias (PREVENT) has been planned, and is a multi-centre randomized clinical trial designed to assess the effect of CSD in patients with cardiomyopathy who continue to experience ICD shocks despite catheter ablation and medical therapies. The trial, currently registered at clinical trials.gov, is undergoing a feasibility and surgical standardization phase, involving centres both inside and outside the United States, and will provide much needed answers in quantifying the benefit of CSD in the treatment of ventricular tachyarrhythmias.

Questions and controversies

Mechanism of benefit of CSD

- What is the contribution of cardiac de-afferentation to the success of this procedure?
- To what extent are cardiac chronotropy and inotropy preserved after bilateral CSD?
- Is removal of T3 and T4 paravertebral ganglia needed for procedural efficacy?

Role of CSD in other cardiovascular diseases

- Heart failure
- Refractory vasospastic angina

Thoracic epidural anaesthesia and stellate ganglion blockade. Anaesthetic interventional techniques such as thoracic epidural anaesthesia and stellate ganglion block allow modulation of sympathetic tone to the heart. In patients with sympathetically triggered arrhythmias, these neural modulation approaches have provided therapies for reducing the arrhythmia burden.

Selective neuraxial modulation of sympathetic signalling to the heart using a high thoracic epidural anaesthesia (TEA) technique is an effective antiarrhythmic approach in animals and humans (Kamibayashi *et al.* 1995; Oka *et al.* 2001; Mahajan *et al.* 2005; Bourke *et al.* 2010). In patients with refractory ventricular tachycardia (VT) unresponsive to medical therapy and cardiac ablation, Shivkumar and colleagues demonstrated that TEA was highly effective in reducing the VT burden in 75% of the patients (CI 51% to 91%, P = 0.05) (Bourke *et al.* 2010). Similarly, Oka and colleagues showed that in patients undergoing lung surgery, the incidence of postoperative supra-ventricular tachyarrhythmias was significantly lower in the group receiving TEA when

compared to controls (1/23 vs. 7/25, respectively, P = 0.05) (Oka *et al.* 2001).

TEA is performed by percutaneously administering local anaesthetics in the thoracic epidural space – a limited sympathectomy of the T1-T5 spinal segments is created by pharmacological block at the level of the spinal cord and spinal roots (Mahajan et al. 2005; Bourke et al. 2010). Since the spread of the drug is not restricted in the epidural space, TEA can block both afferent and efferent sympathetic signalling of the right and left spinal roots, significantly reducing the spinal cord influences. At the level of the heart, the antiarrhythmic effects of TEA are seen due to changes in myocardial excitability, including lengthening of repolarization and prolongation of refractory periods (Meissner et al. 2001). In patients with ischaemic heart disease, regional sympathetic block from TEA not only reduces ischaemic pain but also preserves coronary perfusion with the effect most pronounced in stenotic vessels (Olausson et al. 1997; Nygard et al. 2005). These results demonstrate the protective effects of TEA in mitigating ventricular arrhythmias mediated by increased sympathetic tone. While the effectiveness of TEA is evident, its therapeutic use is limited by its short-term use in patients.

LSG block is a percutaneous approach to block the peripheral sympathetic ganglia using local anaesthetics. The technique reduces sympathetically driven cardiac excitability in animal and human studies (Nademanee et al. 2000; Loyalka et al. 2011; Patel et al. 2011). Nademanee and colleagues reported that cardiac sympathetic blockade, including LSG block, is superior to the Advanced Cardiovascular Life Support recommended antiarrhythmic therapy in treating patients with electrical storm (Nademanee et al. 2000). Long-term survival in patients who were treated with sympathetic blockade was also better, compared to those treated with antiarrhythmics. Reduction in adrenergic tone secondary to blockade of both afferent and efferent sympathetic pathways is most likely responsible for the efficacy of the LSG block. The antiarrhythmic effects of LSG block have also been reported in patients with myocardial infarction (Garcia-Moran et al. 2013) Although the therapeutic effects of a single shot LSG block are short lived, Hayase and colleagues have described administering bi-weekly stellate ganglia blocks for 10 consecutive months for effectively suppressing ventricular arrhythmias and ICD shocks in a patient unresponsive to medical therapy and ablation (Hayase et al. 2013). In clinical practice, LSG block can induce transient reduction or abolition of VTs during electrical storm and should be considered a temporary measure (analogous to TEA). However it should be noted that the lack of response to LSG block has no impact on predicting the outcome of planned surgical CSD.

LSG block has some limitations. Unlike surgical sympathetic denervation, which selectively removes only

the lower third of the stellate ganglia along with the upper thoracic sympathetic chain, percutaneous LSG block with local anaesthetics is non-specific, and typically abolishes conduction to the heart as well as to the head, neck and diaphragm. This also limits the concurrent application of left and right stellate ganglion blocks, unless the patients are being mechanically ventilated.

Ouestions and controversies

Where in ACLS algorithms should TEA and SG blockade be placed in resuscitation and treatment of electrical storm?

Vagal stimulation - heart failure. Left-sided vagus nerve stimulation (VNS) has been extensively used clinically for the treatment of drug-refractory epilepsy (Ben-Menachem, 2001; Schachter, 2002). It was tested for the first time in patients with heart failure (HF) in a small single-centre feasibility study (Schwartz et al. 2008) that was followed by a multicentre international extension increasing the number of patients studied to a total of 32 (De Ferrari et al. 2011; De Ferrari & Dusi, 2015). Patients in New York Heart Association (NYHA) class II-III on optimal medical treatment were eligible in the study if they had LVEF $\leq 35\%$; presence of sinus rhythm with 24 h Holter heart rate of 60–110 beats min⁻¹ and were capable to perform a 6 min walk test. The primary end-point of the study was the occurrence of all system and procedure-related adverse events (AEs). Secondary end-points were the changes between baseline and the 6-month follow-up visit in the following: NYHA class, quality of life using Minnesota Living with Heart Failure Questionnaire (MLwHFQ), six-minute walk test, LV ejection fraction, LV end-diastolic and end-systolic volumes. Patients were implanted with an electrode on the right cervical vagus connected to a CardioFit stimulator. The CardioFit system (BioControl Medical Ltd, Yehud, Israel) is an implantable vagal neurostimulator system, designed to sense heart rate (via a standard intracardiac electrode) and deliver stimulation at a variable delay (70-325 ms) from the R wave. The stimulation lead is an asymmetric bipolar multi-contact cuff electrode specifically designed for cathodic induction of action potentials, while simultaneously applying asymmetrical anodal blocks that are expected to lead to preferential, but not exclusive, activation of vagal efferent fibres. Electrical stimulation of the right vagal nerve was started 2–4 weeks after implant with a single pulse synchronized per cardiac cycle (duty cycle was $21 \pm 5\%$). The stimulation current intensity was slowly increased to a maximum of 5.5 mA. The up-titration was generally limited by patient's discomfort or pain, and on average reached an intensity of 4.1 \pm 1.2 mA. Patients were followed at 3 and 6 months thereafter with optional 1 year follow-up.

With the exception of two device-related AEs (post-operative pulmonary oedema, need of surgical revision), the other 24 serious AEs, including three deaths, were adjudicated to be unrelated to the investigational device. Expected non-serious device-related AEs (cough, dysphonia, stimulation-related pain) occurred early but were reduced and disappeared after stimulation intensity adjustment. There were significant improvements in NYHA class, quality of life, 6-min walk test, LV ejection fraction (from $22 \pm 7\%$ to 29 ± 8) and LV systolic volumes at 6 months. These improvements were maintained (and even magnified) at 1 year (De Ferrari *et al.* 2011). The favourable outcomes regarding feasibility, safety, tolerability and preliminary efficacy of this proof-of-concept study led to additional clinical trials.

ANTHEM-HF (Autonomic Regulation Therapy for the improvement of Left Ventricular Function and Heart Failure Symptoms) was an open-label phase II multicentre study performed in India that assessed the feasibility, safety and tolerability of autonomic regulation therapy (ART), and compared right vs. left vagal nerve stimulation in 60 NYHA II–III patients with LVEF $\leq 40\%$ (Premchand et al. 2014). The stimulation device and electrode were similar to those used by Cyberonics (Houston TX, USA), and approved for the treatment of epilepsy and previously assessed in preclinical HF studies (Zhang et al. 2009); the system does not include an intracardiac electrode. Stimulation parameters were systematically adjusted over a 10-week titration period to a pulse width of 250 μ s and a pulse frequency of 10 Hz in all patients. The mean output current at the end of titration was 2.0 ± 0.6 mA, and this stimulation was maintained throughout the study. After 6 months of ART, the study confirmed the feasibility and safety of the procedure. At the end of 6 months, significant improvements were seen in LVEF, LV end systolic diameter, HRV and Minnesota living with HF questionnaire and the six-minute walk distance. NYHA class improved in 77% of patients. Overall, a modest trend was also observed toward a greater efficacy for the right-sided stimulation. An extension of the ANTHEM-HF patient follow-up for an additional 6 months (total of 12 months following titration) showed that the benefits of ART seen at 6 months persisted at 1 year with no further improvement in any of the objective outcome measures (Premchand et al. 2016).

NECTAR-HF (Neural Cardiac Therapy for Heart Failure Study) was a randomized sham-controlled trial multicentre European phase II study in which all patients (NYHA II–III, LVEF < 35%), were implanted with a right-sided vagal stimulator without intracardiac electrode, and randomized 2:1 to an active group that had the device switched 'ON' and an inactive group had it switched 'OFF' for the first 6 months (De Ferrari *et al.* 2014; Zannad *et al.* 2015b). The device, produced by Boston Scientific (Minneapolis, MN, USA) had showed favourable efficacy results in preclinical studies (Hamann

et al. 2013). The frequency of stimulation was 20 Hz and the current intensity only reached an average of 1.3 ± 0.8 mA, limited by side effects. Of the 96 patients enrolled, 87 had paired data at 6 months for analysis. No change in any hard endpoint was observed between the two groups, although there were favourable changes in NYHA (62% of patients in the active group vs. 43% in the inactive group improved NYHA class, P = 0.032) and in quality of life in the active group. However, the efficacy of blinding was incomplete among patients assigned to the active stimulation group. Recently, the NECTAR-HF investigators reported the 12 and 18 months follow-up data after therapy was switched 'ON' in the sham-control group at 6 months and remained 'ON' in the active group through out. Although the pre-specified safety endpoint was met and the adverse event and survival rates were within the expected range compared to historical controls, the LV structure and function improved equally in both groups with no difference between the groups (Zannad et al. 2015a).

How do we explain the differences in the results of VNS seen in CardioFit and ANTHEM-HF compared to those in NECTAR-HF? All the three trials studied very similar NYHA class II–III patients with depressed ejection fraction. Whereas CardioFit and ANTHEM-HF were small open-label studies, NECTAR-HF was a somewhat larger 96-patient randomized, sham-controlled study. The primary safety outcomes were very similar in all the three studies. None of the trials raised any safety concerns. However, whereas CardioFit and ANTHEM-HF showed improvements in cardiac function and HF symptoms, no improvement in LV function was seen in NECTAR-HF. The only major differences in the design of the three trials were in the stimulation parameters used. CardioFit used a stimulation frequency of 1-2 Hz and achieved a stimulation current of approximately 4 mA. ANTHEM-HF used 10 Hz, the natural frequency of the vagus nerve, and achieved a current output of 2 mA, while NECTAR-HF used a much higher frequency of 20 Hz and could only reach 1.2 mA. It is therefore likely that the low current amplitude VNS used in NECTAR-HF and the presence of a control group, albeit with incomplete blinding, may be responsible for the unfavourable study results of NECTAR-HF (De Ferrari, 2014). However, these explanations may be questioned because of the recent announcement of the premature termination of the INOVATE-HF trial for futility. This is a large international multi-centre randomized controlled trial phase III pivotal trial (Hauptman et al. 2012), based on the CardioFit pilot study and designed to assess safety and efficacy of vagus nerve stimulation in patients with heart failure with reduced ejection fraction, NYHA class III, LVEF < 40%, left ventricular end-diastolic diameter 50 and 80 mm, in sinus rhythm, receiving optimal medical therapy. The study randomized 725 patients from 80 sites in a 3:2 ratio to receive active treatment or standard optimal medical therapy. The primary endpoint of the study is the composite of all-cause mortality or hospitalization for HF. Although no data are available yet, it is possible that the inclusion of many NYHA class III patients with an implanted cardiac resynchronization therapy (CRT) device (and thus CRT non-responders) may have contributed to the neutral result of the study.

In conclusion, several pre-clinical studies have demonstrated the efficacy of VNS in a variety of heart failure models. Three small clinical studies have demonstrated the feasibility and safety of VNS in heart failure. However, the clinical efficacy results are mixed as noted above. VNS remains a promising therapy for heart failure. However, efficacy may be dependent on appropriate VNS parameters and perhaps patient selection.

Vagal stimulation - atrial fibrillation. Based on the observation that mildly enhanced efferent vagal activity, induced by hypertension, was capable of suppressing PV firing even in the face of little or no heart rate slowing (Tai et al. 2000), the Oklahoma group studied the anti-arrhythmic effects of low-level vagus nerve stimulation (LL-VNS) in various canine AF models (Li et al. 2009; Sha et al. 2011; Sheng et al. 2011). LL-VNS was delivered at stimulation voltages below the threshold slowing the sinus rate or AV conduction. Cervical LL-VNS prolonged the atrial and PV effective refractory period, shortened AF duration, and suppressed neural activity of the major atrial ganglionated plexi (GP), as well as right stellate ganglion function (Sha et al. 2011). Importantly, stimulating either the right or bilateral vagus nerve produced similar results. The mechanism by which low-level VNS suppresses AF has been implicated by Shen et al. demonstrating that the antiarrhythmic effects of LL-VNS may reside in its anti-adrenergic effects, as evidenced by suppression of the neural activity of the left stellate ganglion through upregulation of the small conductance Ca²⁺-activated potassium channels (Shen et al. 2011, 2013). Release of another neurotransmitter (vasostatin-1) by low-level VNS was also implicated as a potential mechanism responsible for its antiarrhythmic effects (Stavrakis et al. 2012).

Transcutaneous stimulation of the auricular branch of the vagus nerve (tragus stimulation). A major drawback of cervical VNS is its invasiveness, requiring surgical implantation of a pulse generator and a cuff electrode around the vagus nerve. Common side effects include voice changes, discomfort at implant site and infection (Spuck *et al.* 2010). An alternative to cervical VNS is to stimulate the auricular branch of the vagus nerve (ABVN) located at the external ear. Two locations of the external

ear are richly innervated by the ABVN: concha and tragus (the anterior protuberance of the ear) (Peuker & Filler, 2002). Immersing the central cut end of the ABVN in horseradish peroxidase to trace the projection of ABVN revealed that main nerve terminal labelling was observed in the brain stem nuclei such as nucleus tractus solitarii (NTS) and spinal trigeminal nucleus (Nomura & Mizuno, 1984). It is known that neural output from the NTS projects to the cardiovascular centre, vagal motor nucleus in the brain stem, hypothalamus and cerebral cortex, serving as a 'coordination centre' between afferent neural inputs from peripheral organs and the higher centres of the brain (Boon et al. 2001; Ben-Menachem, 2002). The central projection of the ABVN was illustrated by a recent functional MRI study in which widespread excitation of ipsilateral NTS, bilateral spinal trigeminal nucleus, dorsal raphe and locus coeruleus, as well as suppression of the activity of the hippocampus and hypothalamus was elicited by electrical stimulation of the ABVN at 25 Hz (Frangos et al. 2015).

Electrical stimulation of the tragus, but not earlobe, helix or antihelix, in human subjects elicited vagus somatosensory evoked potentials evidenced by electroencephalograpy recordings (Fallgatter et al. 2003; Polak et al. 2009). Clancy et al. applied VNS to the tragus (30 Hz, at sensory threshold) in healthy volunteers and documented that tragus VNS significantly decreased the low-frequency/high-frequency ratio of HRV, as well as the frequency and incidence of the MSNA, indicating a shift of cardiac autonomic balance toward parasympathetic predominance (Clancy et al. 2014). Based on the antiarrhythmic effects of the low-level cervical VNS, the Oklahoma group explored the idea of low-level tragus stimulation (LLTS) at the stimulation level, which did not slow the sinus rate or atrioventricular (AV) conduction. The antiarrhythmic effects were comparable with low-level cervical VNS. In a canine AF model of rapid atrial pacing, LLTS suppressed AF and reversed acute atrial remodelling by lengthening the effective refractory period (ERP) and reducing ERP dispersion, likely by inhibiting the neural activity of the GP (Yu et al. 2013). Importantly, antiarrhythmic effects were maintained by stimulation strengths as low as 80% below the threshold that slows the sinus rate or AV conduction. It should be noted that based on recent studies, current delivered at 80% of bradycardia threshold is still 50% above that required to activate descending efferent projections (Ardell et al. 2015; Yamakawa et al. 2015). These data point to the critical impact of low-level vagus nerve stimulation to impact multiple levels of the cardiac nervous system to impart cardioprotection (Kember, 2014).

Stavrakis *et al.* reported the first-in-man experience of the antiarrhythmic effects of LLTS on patients referred for catheter ablation of drug-refractory paroxysmal AF (Stavrakis *et al.* 2015). The discomfort threshold (tingling

sensation at the stimulation site) was measured before anaesthesia and the cardiac threshold (the stimulation voltage slowing the sinus rate or AV conduction) was determined after anaesthesia was started. With only 1 h of LLTS (20 Hz, 50% below the cardiac threshold), the right and left atrial ERP was prolonged, AF duration was abbreviated and systemic pro-inflammatory markers such as tumour necrotic factor- α (TNF- α) and C-reactive protein were suppressed. Notably, the discomfort threshold was significantly higher than the stimulation voltage of LLTS, indicating that LLTS may be used as a non-sentient, outpatient, non-invasive therapy to treat paroxysmal AF.

Spinal cord stimulation. Heart failure is associated with abnormal neuro-hormonal activation leading to therapy with beta-blockers and ACE inhibitors. Spinal cord stimulation (SCS) has been employed for many years to treat patients with intractable neuropathic pain syndromes, pain from extremity ischaemia and non-revascularizable angina. The beneficial mechanisms of SCS may relate in part to modulation of the (alpha) sympathetic tone and/or vagal stimulation. Other mechanisms may be, and probably are, operative.

The purpose of the DEFEAT-HF trial was to test the hypothesis that spinal cord stimulation would improve heart failure metrics including heart size and muscle wall thickness and functional capacity. The trial was a randomized (3:2), parallel, single blind controlled study in patients with (1) NYHA class III HF, (2) LVEF < 35%, (3) QRS duration < 120 ms and LVEDD > 55 mm. The primary objective of the DEFEAT-HF study was to evaluate the reduction in left-ventricular end systolic volume index (LVESVi) after 6 months of SCS therapy in the treatment arm compared to the control arm (Zipes *et al.* 2016).

Forty-two patients were randomized to SCS ON and 24 to SCS OFF. Among randomized patients, the mean age was 61 years; 79% were male; mean LVEF was 27% and mean QRS duration was 105 ms. The change in LVESVi over 6 months was not significantly different between randomization arms (SCS OFF: -2.2 [95% CI: -9.1, 4.6] versus SCS ON: 2.1 [95% CI: -2.7, 6.9]; P = 0.30). Analyses of secondary endpoints including peak V_{O_2} , N-terminal of the prohormone brain natriuretic peptide, freedom from first hospitalization with HF or death at 6 months, change in Minnesota Living with HF quality of life, change in New York Heart class, and change in six-minute hall-walk all showed no differences between groups. For this study, SCS was delivered with a duty cycle of 12 h on and 12 h off. While safety concerns were addressed by this study, the SCS protocol likely was not optimized for cardioprotection. The successful application of SCS is critically dependent on: (1) the location of the electrode interface, (2) the stimulation protocol employed, and (3) the substrate that is being targeted (neural and cardiac). Future studies should consider these benchmarks as well as evolve appropriate biomarkers to assess efficacy.

Questions and controversies

- What are the optimal stimulation parameters for the vagus nerve in humans?
- What are the specific stimulation parameters that preferentially activate efferent vagal fibres over afferent fibres?

Ganglionated plexi and vein of Marshall ablation. The intrinsic (ICN) and extrinsic cardiac innervations play a critical role regulating physiological modulations on the cardiac myocardium. In the atria, such modulation can turn maladaptive and lead to AF. The role of vagal activation in the pathogenesis of certain forms of AF has been long recognized (Garrey, 1924). Experimentally, vagal stimulation - or the administration of acetylcholine - can lead to inducibility of sustained AF (Burn et al. 1955; Allessie et al. 1977, 1984, 1985), and clinically, some forms of paroxysmal AF may be related to an elevated parasympathetic tone (Coumel et al. 1978). Mechanistically, vagal stimulation leads to spatially heterogeneous shortening of atrial refractoriness (Zipes et al. 1974; Liu & Nattel, 1997), and cholinergic agonists shorten atrial propagation wavelength (Smeets et al. 1986), both promoting the maintenance of AF.

In the anatomical–physiological hierarchy of autonomic control of the heart, the ICN – organized in macroscopically recognizable ganglionated plexi (GP) and ligament of Marshall (LOM) – are not only necessary relay stations from central influences, but are able to activate independently. Whether in response to central control, or as a result of local activation, the ICN are the final implementers of the autonomic – and profibrillatory – influences on the atrial myocardium. As such, they are attractive therapeutic targets in patients with AF. In canines, catheter ablation of certain components of the intrinsic cardiac nerves (third fat pad in the right pulmonary artery) leads to elimination of AF inducibility after vagal stimulation (Chiou *et al.* 1997; Schauerte *et al.* 2000).

Translation of these facts to the clinic has not been straightforward. With the advent of pulmonary vein (PV) isolation (PVI) as the established technique for catheter ablation of AF (Haissaguerre *et al.* 1998), its claimed mechanistic underpinning – elimination of PV ectopy as AF triggers – gained widespread acceptance. The ablation sites required for PVI crudely match the location of ICN, and soon after PVI became accepted, its autonomic effects became apparent, either as bradycardic events during ablation (Tsai *et al.* 1999), or as alterations of HRV long term (Hsieh *et al.* 1999). The PVI procedure includes

variable extents of ICN ablation, which raises the question of which procedural component is responsible for the therapeutic effect. Indeed, PVI and ICN ablation may be synergistic. Pappone *et al.* reported that when PVI was associated with bradycardic events, subsequent abolition of these reflexes by radiofrequency – an index of ICN denervation – led to superior rhythm control outcomes (Pappone *et al.* 2004). Interestingly, in an acute model of vagal stimulation AF, ablation of ICN was superior to PVI, supporting the hypothesis that the ICN, and not the PVI triggers, were mechanistically more relevant (Lemola *et al.* 2008).

Clinical techniques to target ICN have lacked specificity, in part because most clinical procedures have included variable extents of PVI, and in part because the tools for ICN ablation - most commonly radiofrequency necessarily destroy adjacent myocardium. Surgical series have reported a successful AF ablation using a combined approach of PVI plus GP and LOM ablation, with successful elimination of AF reported as 65% (Han et al. 2009), 81% (Edgerton et al. 2010), or up to 86% (Krul et al. 2011). In order to discern the relative merits of GP ablation vs. PVI, Katritsis et al. randomized patients with paroxysmal AF to PVI alone or in combination with additional radiofrequency ablation at sites where parasympathetic reflexes are elicited or at the expected anatomical locations of individual GPs. Success rates at 1 year were 60.6% in the PVI group vs. 85.3% in the PVI+GP ablation group (Katritsis et al. 2011). Results were confirmed in a larger trial that compared PVI alone, GP ablation alone, and PVI+GP ablation in patients with paroxysmal AF. Success rates were 56%, 48% and 74%, respectively, suggesting a synergistic effect (Katritsis et al. 2013). A similar comparison in patients with persistent AF showed that PVI combined with GP ablation was superior to PVI combined with linear lesions (68% success vs. 52%, respectively) (Steinberg et al. 2013).

Targeting the LOM as part of the GP has been traditionally performed surgically. A percutaneous approach using ethanol infusion in the vein of Marshall (VOM) showed that regional parasympathetic denervation followed (Valderrabano et al. 2009). The procedure was validated in humans: high-frequency stimulation in the VOM could trigger AF and AV nodal conduction slowing, demonstrating a GP-to-GP communication between VOM-GP and the right inferior GP (which controls the AV node). Such a response was eliminated after VOM ethanol infusion (Baez-Escudero et al. 2014). The role of ethanol ablation of the VOM in rhythm control outcomes is being studied in a randomized trial.

Finally, it should be noted that GP destruction is merely a crude approach. The GP contain multiple neuronal populations and various neurotransmitters that could be pharmacologically targeted for anti-fibrillatory purposes. The use of botulinum toxin to interfere with cholinergic neurotransmission is a first approach in this direction (Oh *et al.* 2011; Pokushalov *et al.* 2015).

Renal denervation. Initial inferences into neuro-regulation of renal function were made as far back as 1859, when transection of the greater splanchnic nerve resulted in greater ipsilateral diuresis. The opposite effect was seen with electrical stimulation of the peripheral end of the transected nerve, suggesting that renal sympathetic denervation promoted diuresis, while sympathoexcitation had an antiduretic effect (DiBona & Esler, 2010). This concept was replicated using renal blood flow, where denervation increased blood flow, and stimulation decreased blood flow (Starling, 1908). Utilizing renal sympathetic nerve activity (RSNA) measurements, increasing RSNA via electrical stimulation resulted in increased renin secretion, sodium reabsorption and renal vasoconstriction (DiBona & Kopp, 1997).

Adrenergic innervation of the kidneys, occurring via nerves running along the renal artery to the renal microvasculature and juxtaglomerular apparatus (Barajas, 1964; Barajas & Muller, 1973) therefore represents a target for interruption, to achieve increased blood flow, lower renin secretion rate, vasoconstriction and sodium reabsorption. It is hypothesized that in hypertension, these mechanisms are perturbed. Renal sympathetic outflow is now recognized to be elevated in patients with essential hypertension (DiBona & Esler, 2010; Esler, 2014), particularly the drug-resistant population. It is also known that afferent nerve fibres, utilizing a variety of neurotransmitters, convey mechanical and chemical information to the neuraxis (Liu & Barajas, 1993). Both excitatory and inhibitory reflexes involving RSNA, diuresis, and natriuresis have been described, mediated by afferent nerves (Kopp et al. 1985; Stella & Zanchetti, 1991; Ye et al. 2002). Removal of both kidneys, the source of afferent signals to the CNS (Converse et al. 1992), and dorsal rhizotomy (Campese & Kogosov, 1995) (removal of T9–L1 dorsal roots), resulting in preferential afferent denervation, were shown to decrease hypertension. In humans with end-stage renal failure on haemodialysis and with hypertension, in whom MSNA was elevated, nephrectomy resulted in significant reduction of blood pressure (Converse et al. 1992). In a hypertensive model in rats, dorsal rhizotomy abrogated hypertension (Campese & Kogosov, 1995).

These findings and others provided strong rationale for renal denervation (of both afferent and efferent nerves) in humans to reduce hypertension. Percutaneous renal denervation by catheter ablation in the renal arteries has emerged as a promising therapeutic approach in patients with treatment resistant hypertension. This therapy applies radiofrequency or ultrasonic waves from

an endoluminal catheter, through the walls of both renal arteries to ablate nerves running along the vessel wall. Two initial clinical trials (Simplicity HTN-1 and -2) (Krum et al. 2009; Esler et al. 2010) demonstrated promising results, leading to a larger third trial, Simplicity HTN-3 (Bhatt et al. 2014). This third trial, however, failed to show the same promising results as the two prior trials.

As a result, renal denervation is at a watershed, with its future uncertain as a treatment for hypertension. The broad body of published trials does indicate technical efficacy is essential, now that the Symplicity HTN-3 trial (Bhatt et al. 2014) is seen to be flawed (Esler, 2014; Kandzari et al. 2015). In the words of its co-chief investigator George Bakris, 'it is highly likely that RDN as it was performed in the HTN-3 was technically inconsistent at best, but technically inadequate at worst' (Nathan & Bakris, 2014). The current clinical and research practice of largely restricting renal denervation to the treatment of severe, drug-resistant hypertension flowed from the first Symplicity trial, in which ethical considerations dictated restriction of the procedure to this patient group, to optimize the benefit/risk balance. However, in the new round of trials, inclusion criteria are changing, based more on concepts of hypertension pathogenesis, to include milder, untreated essential hypertension, where a neural pathophysiology is prominent (Esler et al. 1990) and to exclude isolated systolic hypertension, where biophysical changes in the aorta are more important than sympathetic nervous system activation.

Procedural issues arising from renal denervation will provide key information for future trials. As an example, a larger number of energy applications per artery, in recent trials, has been reported to produce greater blood pressure lowering (Esler, 2014; Kandzari et al. 2015), an effect prominent in the otherwise negative Symplicity HTN-3 trial (Kandzari et al. 2015). Why should this be so? In a study in pigs, Tzafriri et al. (2015) have demonstrated that previously unexpected denervation technical failures can arise when administered intraluminal radiofrequency (RF) energy does not reach its target - the periarterial sympathetic nerves - owing to energy conductivity and temperature gradients being distorted by regional tissue microanatomical variations. Fibrous muscle sheaths and lymph nodes draw the electric field, increasing the lateral and circumferential extent of the ablation geometry, whereas veins act as an energy sink, limiting ablation depth (Tzafriri et al. 2015). The outflow of energy from the electrode toward the targeted nerves is not a predictable, symmetrical cone. Assessment of denervation efficacy with renal noradrenaline spillover measurements in clinical studies reflects this, where it has been documented that achieved denervation differs markedly between individual patients, sometimes being less than 25% (Esler, 2014). Given this unpredictability, concerning the number of RF energy applications, it appears that 'the more the merrier'.

Of perhaps even greater relevance is improved understanding of the anatomy of the renal nerves, generated from studies on pigs (Mahfoud & Luscher, 2015), dogs (Henegar et al. 2015), and humans (Sakakura et al. 2014). The renal sympathetic nerves are more distant from the renal artery lumen proximally, and converge on the distal renal artery and the renal artery divisions (Sakakura et al. 2014; Henegar et al. 2015; Mahfoud & Luscher, 2015), where they are more accessible to administered radiofrequency energy. The essence of this anatomical knowledge has been known for more than 60 years (Oldham, 1950) but typically was not acted on in renal denervation trials, where RF energy was inexplicably preferentially directed into the proximal renal artery (Esler, 2014; Kandzari et al. 2015). The evidence from experimental studies on this point is that RF energy delivery into the proximal part of the renal arteries, near the ostia, produces suboptimal denervation, whereas energy delivery into the distal renal arteries and renal artery divisions produces near-complete denervation, uniform between animals (Esler, 2014; Mahfoud & Luscher, 2015). Another important limitation of renal denervation in the long term is the potential for renal reinnervation (Booth et al. 2015). This not only has the potential to limit efficacy, but could be pathological if it occurs inappropriately. The 'smart' renal denervation trial of the future will be transformed by new procedural knowledge.


Questions and controversies

- Are there adverse consequences to ganglionated plexi ablation for atrial fibrillation or other conditions including vaso-vagal syncope?
- How does ligament of Marshall ablation impact intrinsic cardiac nervous system function?
- Are the effects of ganglionated plexi and ligament of Marshall ablation durable?
- Is there a future for catheter-based renal denervation through the renal arteries?
- How should future clinical trials be designed to avoid technical failures?
- What are optimal inclusion criteria for patients being enrolled into renal denervation trials? Should documentation of sympathetically driven hypertension be required?
- Do renal denervation trials reflect a regression to the mean?

Sleep apnoea interventions. As discussed earlier, patients with sleep apnoea demonstrate significant autonomic pathology as a consequence of apneic events during sleep

(Fig. 8A) (Somers et al. 1995). Indeed, blood pressure during the night often shows a non-dipper pattern with likely consequences for increased long-term cardiovascular risk. In some patients, acute hypoxaemia may also manifest as bradyarrhythmias and even prolonged systole due to the reflex vagal response to hypoxaemia and apnoea mentioned earlier (Fig. 3) (Daly et al. 1979; Thalmann et al. 2001). The acute consequences of nocturnal hypoxaemia, apnoea, vasoconstriction and blood pressure surges may also include nocturnal angina (Mooe et al. 2000), nocturnal myocardial infarction (Kuniyoshi et al. 2008) and nocturnal sudden death (Konecny et al. 2010). High sympathetic drive at night appears to carry over into the daytime so that patients with OSA have very high levels of resting sympathetic drive, despite the absence of any other co-morbidity and even though they are breathing normally (Fig. 8B) (Somers et al. 1995). Heightened tonic sympathetic activation in OSA may contribute to the increased risk for daytime hypertension. It is likely that tonic chemoreflex activation, even in normoxia, contributes to the increased daytime sympathetic activity since exposure to 100% oxygen elicits not only an attenuation of sympathetic outflow but also of heart rate and blood pressure (Narkiewicz et al. 1998). This speaks again to the importance of respiratory variables in modulating cardiac function. Indeed patients with borderline hypertension have enhanced chemoreflex responses to hypoxaemia, and tonic chemoreflex drive may be implicated in their high levels of sympathetic activation and elevated blood pressure (Trzebski et al. 1982; Somers et al. 1988).

Central sleep apnoea (CSA), in contrast to OSA, occurs because of an attenuation of central respiratory drive. These patients often have hypocapnia, and oxygen desaturation during apnoea is less severe than that seen in OSA. CSA is usually seen in patients with heart failure where it can also manifest as the crescendo-decrescendo breathing pattern known as Cheyne-Stokes respiration (CSR). Heart failure patients with CSA-CSR have been shown consistently to have a poorer prognosis, with increased mortality (Somers et al. 2008). A recent randomized trial, SERVE HF, investigated whether treatment of central sleep apnoea in stable heart failure patients with systolic dysfunction (LVEF \leq 45%), would be accompanied by reduced cardiovascular morbidity and mortality. Surprisingly this study showed that patients randomized to adaptive servo ventilation (ASV) (positive pressure administration using an algorithm that effectively treats CSA) were in fact associated with increased mortality in the CSA treated group (Cowie et al. 2015). Importantly, increased mortality was especially evident in patients with CSR as opposed to CSA suggesting first that these two breathing patterns may be different, and second that CSR may conceivably be protective in patients with systolic heart failure. Preliminary evidence suggests that increased






Figure 8. Autonomic dysfunction in obstructive sleep apnoea

A, recordings of the electrocardiogram (ECG), SNA, RESP and BP in a sleep apnoeic patient undergoing continuous positive airway pressure (CPAP) therapy during REM sleep. The arrow indicates a reduction of CPAP from 8 to 6 mmHg, allowing the development of obstructive apnoea, with consequent increased SNA and BP. Reproduced with permission from Somers *et al.* (1995). *B,* recordings of sympathetic nerve activity (SNA) during wakefulness in patients with obstructive sleep apnoea (OSA) and matched controls showing high levels of SNA in patients with sleep apnoea. Reproduced with permission from Somers *et al.* (1995).

cardiovascular deaths in the ASV treated group may have been because of an increased likelihood of sudden death, speaking to the possibility of a protective effect of CSR (Naughton, 2012), perhaps due to an anti-arrhythmic effect of the crescendo–decrescendo breathing pattern.

Questions and controversies

Central sleep apnoea (CSA)—Cheyne—Stokes respiration (CSR) in systolic heart failure

 Whether CSA is a cause or marker of adverse outcomes is not clear. While CSR is often seen in patients with systolic heart failure, and is indicative of a poorer prognosis, is the rhythmic breathing pattern of CSR in fact protective in heart failure, perhaps due to an anti-arrhythmic effect?

OSA and adverse cardiovascular events

 While OSA is associated with increased sympathetic activity, hypertension, atrial fibrillation and increased risk of myocardial infarction and sudden death, particularly at night, does treatment of OSA reduce risk of cardiac and vascular events?

Meditation, holistic interventions and aerobic exercise.

Holistic interventions are treatments that address the psychological, social and spiritual needs of an individual in addition to their physical needs. Holistic interventions integrate the context in which an individual's illness arises by carefully attending to the roles played by their values, beliefs, culture and community. They are routinely sought and received outside of traditional healthcare settings, in the form of complementary health approaches. In fact, out-of-pocket expenditure on complementary health approaches outweighs the amount spent on inpatient hospitalizations in the United States (Eisenberg et al. 1993; Barnes et al. 2004), and more than 30% of adults and 12% of children currently utilize them (Black et al. 2015; Clarke et al. 2015), often in parallel with allopathic healthcare. Holistic interventions are also being increasingly included within allopathic healthcare services, particularly those delivered by comprehensive care centres such as for cancer (Bultz et al. 2014) or terminal illness (Bercovitz et al. 2011). The 10 most commonly utilized complementary health approaches are natural products (dietary supplements other than vitamins and minerals) (17.7%), deep breathing (10.9%), yoga, tai chi or qigong (10.1%), chiropractic or osteopathic manipulation (8.4%), meditation (8.0%), massage (6.9%), special diets (3.0%), homeopathy (2.2%), progressive relaxation (2.1%) and guided imagery (1.7%) (Clarke et al. 2015). Other utilized approaches include acupuncture, and movement therapies such as pilates and rolfing. Based on these use patterns, it is possible to parse complementary health approaches into two general groups: mind and body practices and natural products practices.

Stress, whether physical or mental, has a major impact on the development of cardiovascular pathology (Nemeroff & Goldschmidt-Clermont, 2012; Thurston et al. 2013; Golbidi et al. 2015). Early clinical indicators of the link between stress and cardiac arrhythmias identified increased ventricular irritability, arrhythmogenicity and sudden cardiac death during acute mental stressors (Lown et al. 1976; DeSilva & Lown, 1978; Lown & DeSilva, 1978; see also Ziegelstein, 2007). Further, strong emotions, including anger are recognized to powerfully modulate ventricular electrophysiology, and risk of arrhythmogenesis (Gray et al. 2007; Taggart et al. 2011). It is not a rarity for sudden death to be precipitated by a strong emotional outburst. Depression is also associated with increased mortality following myocardial infarction (Frasure-Smith et al. 1993, 1995). This relationship appears to be specific to negative affect as opposed to anger or low social support (Frasure-Smith & Lesperance, 2003). Population scale studies have also confirmed causal relationships between strong emotional experiences and acute cardiac events. For example, examinations of emergency department records have identified increased rates of acute coronary syndrome during natural disasters such as earthquakes (Leor et al. 1996), but also during major sporting events such as World Cup soccer, when arousal elevations are presumably positively as well as negatively valenced (Wilbert-Lampen et al. 2008, 2010; Borges et al. 2013). While such occurrences have been sometimes characterized as 'precipitating events' resulting in earlier unmasking of chronic pathology (Schwenk, 2008), emotional stress can also trigger ventricular dysfunction in healthy individuals, through a mechanism ascribed to exaggerated sympathetic stimulation (so called 'takotsubo cardiomyopathy') (Wittstein et al. 2005; Templin *et al.* 2015).

It is important to note that several considerations have been raised about this nascent field. These include the observations that many studies have utilized smaller sample sizes or non-randomized interventions, or failed to include adequate comparator conditions to control for non-specific effects associated with intervention delivery (so called 'attentional controls') (Ospina et al. 2007; Goyal et al. 2014). While the risk of bias is an ever-present concern in any field, and the utilization of appropriate comparators is essential, the non-pharmacological nature of mind-body interventions necessitates the judicious application of principles and methods from the field of behavioural randomized controlled trials. In pragmatic terms, this can sometimes mean that selection of an attention control in clinical settings may not appropriately determine whether a novel intervention shows clinical efficacy at treating illness. In such cases a more suitable comparator condition might be one in which an intervention's efficacy is compared with usual care, or an enhanced usual care strategy (Freedland *et al.* 2011; Freedland, 2013). On the other hand, an attention control may be more appropriate in cases where the specificity of an efficacious clinical intervention is being investigated.

Investigations of mind and body practices on autonomic and myocardial function are still in the early stages of development. Certain practices seem to induce quiescent autonomic states (Telles et al. 1995; Chaya et al. 2006) while others induce dynamic changes (Peng et al. 2004; Cysarz & Bussing, 2005), presumably through modulations in vagal tone via respiratory sinus arrhythmia. Compassion training may reduce sympathetic nervous system reactivity and improve parasympathetic reactivity during acute mental stress in women (Arch et al. 2014). Mindfulness training may enhance resiliency in marines preparing for deployment, by speeding the return to autonomic baseline and dampening viscerosensory reactivity in the central nervous system (Johnson et al. 2014). Extending this notion further, an intriguing case report described suggested a novel effect of meditation on the ability to down-regulate elevated adrenergic arousal due to isoproterenol (Dimsdale & Mills, 2002). However, a follow up study failed to replicate this finding at the group or individual level (Khalsa et al. 2015b).

Recent evidence suggests that mind and body interventions can be beneficial in the treatment of cardiac arrhythmias. A 3-month face-to-face training in yoga and meditation in patients with paroxysmal AF was associated with reductions in episodes of symptomatic AF, symptomatic non-AF episodes, and asymptomatic AF episodes, as well as reductions in heart rate and blood pressure, and improved quality of life indicators (Lakkireddy et al. 2013). In a separate study by the same group, neurocardiogenic syncope patients reported reduced syncopal and presyncopal episodes after receiving electronic training in yoga via instructional videos (Gunda et al. 2015). A small pilot study showed that mindfulness-based stress reduction effectively reduced anxiety in adolescents with ICDs or pacemakers (Freedenberg et al. 2015). While some of these studies lacked comparator groups, and do not identify specific mechanisms underlying the observed effects, they demonstrate the promise and potential efficacy of mind and body interventions on cardiac and psychological outcomes in arrhythmia patients.

Aerobic exercise has long been recognized to improve autonomic balance, by both central and peripheral mechanisms (Billman *et al.* 2015; Haack & Zucker, 2015). In humans with heart failure, ischaemic cardiomyopathy, diabetes and other diseases where autonomic imbalance is known to occur, aerobic exercise is known to improve cardiac autonomic indices (Gordon *et al.* 1997; Laterza *et al.* 2007; Spierer *et al.* 2007; Kleiber *et al.* 2008; Mousa

et al. 2008; El Mhandi et al. 2011; Rodrigues et al. 2012, 2014; Zheng et al. 2012; Kaikkonen et al. 2014; Groehs et al. 2015; Masson et al. 2015; Sa et al. 2016). As exercise has been covered in an accompanying White Paper, we will not focus on it further.

Ouestions and controversies

Mind and body practices such as yoga and meditation have well-established roles in attenuating autonomic stress, but their role in improving cardiovascular health is less established.

- Do mind and body interventions or their active components modulate information flow (i.e. dysfunctional crosstalk) between interacting feedback loops from the cardiovascular periphery and autonomic and central nervous systems?
- Which mind and body interventions have the greatest reliability and efficacy at improving psychosocial functional impairments in cardiovascular disorders?
 What is the role of context, e.g. individual vs. group delivery? In person vs. electronic delivery? Traditional (unaltered) delivery vs. active components?

Conclusions

Clinical neurocardiology links the substrate of the heart and the neurohumoral control systems that regulate it, and provides clinically applicable avenues for neuroscience-based therapies. These include (1) identifying patients at high risk for future adverse outcome, (2) helping monitor disease progression and monitoring effect of therapies, and (3) providing novel targets for therapeutics. Neuromodulation strategies show real promise of sustaining cardiac function while maintaining electrical stability. This field requires special efforts in increasing scientific investments on a large scale. In this vein, initiatives such as the 'Stimulating Peripheral Activity for Relieving Conditions' (SPARC) under the guidance of the National Institutes of Health (NIH), will pave the way for improving the efficacy and adverse effects of clinical neuro-modulation.

References

Adám G (1998). Visceral Perception: Understanding Internal Cognition. Plenum Press, New York.

Ai J, Wurster RD, Harden SW & Cheng ZJ (2009). Vagal afferent innervation and remodeling in the aortic arch of young-adult fischer 344 rats following chronic intermittent hypoxia. *Neuroscience* **164**, 658–666.

Ajijola OA, Lellouche N, Bourke T, Tung R, Ahn S, Mahajan A & Shivkumar K (2012*a*). Bilateral cardiac sympathetic denervation for the management of electrical storm. *J Am Coll Cardiol* **59**, 91–92.

- Ajijola OA, Wisco JJ, Lambert HW, Mahajan A, Stark E, Fishbein MC & Shivkumar K (2012*b*). Extracardiac neural remodeling in humans with cardiomyopathy. *Circ Arrhythm Electrophysiol* 5, 1010–1116.
- Ajijola OA, Yagishita D, Reddy NK, Yamakawa K, Vaseghi M, Downs AM, Hoover DB, Ardell JL & Shivkumar K (2015). Remodeling of stellate ganglion neurons after spatially targeted myocardial infarction: Neuropeptide and morphologic changes. *Heart Rhythm* 12, 1027–1035.
- Albert CM, Chae CU, Grodstein F, Rose LM, Rexrode KM, Ruskin JN, Stampfer MJ & Manson JE (2003). Prospective study of sudden cardiac death among women in the United States. *Circulation* **107**, 2096–2101.
- Albert CM, Mittleman MA, Chae CU, Lee IM, Hennekens CH & Manson JE (2000). Triggering of sudden death from cardiac causes by vigorous exertion. *N Engl J Med* **343**, 1355–1361.
- Allessie MA, Bonke FI & Schopman FJ (1977). Circus movement in rabbit atrial muscle as a mechanism of tachycardia. III. The "leading circle" concept: a new model of circus movement in cardiac tissue without the involvement of an anatomical obstacle. *Circ Res* 41, 9–18.
- Allessie MA, Lammers WJ, Bonke IM & Hollen J (1984). Intra-atrial reentry as a mechanism for atrial flutter induced by acetylcholine and rapid pacing in the dog. *Circulation* **70**, 123–135.
- Allessie MA, Lammers WJEP, Bonke FIM & Hollen J (1985). Experimental evaluation of Moe's multiple wavelet hypothesis of atrial fibrillation. In *Cardiac Electrophysiology and Arrhythmias*, ed. Zipes DP & Jalife J, pp. 265–275. Grune and Straton, Inc., Orlando.
- Andersen K, Farahmand B, Ahlbom A, Held C, Ljunghall S, Michaelsson K & Sundstrom J (2013). Risk of arrhythmias in 52 755 long-distance cross-country skiers: a cohort study. *Eur Heart J* **34**, 3624–3631.
- Antiel RM, Bos JM, Joyce DD, Owen HJ, Roskos PL, Moir C & Ackerman MJ (2015). Quality of life after videoscopic left cardiac sympathetic denervation in patients with potentially life-threatening cardiac channelopathies/cardiomyopathies. *Heart Rhythm* **13**, 62–69.
- Arch JJ, Brown KW, Dean DJ, Landy LN, Brown KD & Laudenslager ML (2014). Self-compassion training modulates alpha-amylase, heart rate variability, and subjective responses to social evaluative threat in women. *Psychoneuroendocrinology* **42**, 49–58.
- Ardell JL, Butler CK, Smith FM, Hopkins DA & Armour JA (1991). Activity of in vivo atrial and ventricular neurons in chronically decentralized canine hearts. *Am J Physiol* **260**, H713–H721.
- Ardell JL, Rajendran PS, Nier HA, KenKnight BH & Armour JA (2015). Central-peripheral neural network interactions evoked by vagus nerve stimulation: functional consequences on control of cardiac function. *Am J Physiol Heart Circ Physiol* **309**, H1740–H1752.
- Armour JA (1985). Activity of in situ middle cervical ganglion neurons in dogs, using extracellular recording techniques. *Can J Physiol Pharmacol* **63**, 704–716.
- Armour JA (2004). Cardiac neuronal hierarchy in health and disease. *Am J Physiol Regul Integr Comp Physiol* **287**, R262–R271.

- Armour JA (2008). Potential clinical relevance of the 'little brain' on the mammalian heart. *Exp Physiol* **93**, 165–176.
- Armour JA & Ardell JL (2004). *Basic and Clinical Neurocardiology*. Oxford University Press, New York.
- Armour JA, Huang MH, Pelleg A & Sylven C (1994). Responsiveness of in situ canine nodose ganglion afferent neurones to epicardial mechanical or chemical stimuli. *Cardiovasc Res* **28**, 1218–1225.
- Armour JA, Murphy DA, Yuan BX, Macdonald S & Hopkins DA (1997). Gross and microscopic anatomy of the human intrinsic cardiac nervous system. *Anat Rec* **247**, 289–298.
- Arora R, Ng J, Ulphani J, Mylonas I, Subacius H, Shade G, Gordon D, Morris A, He X, Lu Y, Belin R, Goldberger JJ & Kadish AH (2007). Unique autonomic profile of the pulmonary veins and posterior left atrium. *J Am Coll Cardiol* **49**, 1340–1348.
- Arora RC, Ardell JL & Armour JA (2000). Cardiac denervation and cardiac function. Curr Interv Cardiol Rep 2, 188–195.
- AVID Investigators (1997). A comparison of antiarrhythmic-drug therapy with implantable defibrillators in patients resuscitated from near-fatal ventricular arrhythmias. The Antiarrhythmics versus Implantable Defibrillators (AVID) Investigators. *N Engl J Med* **337**, 1576–1583.
- Baez-Escudero JL, Keida T, Dave AS, Okishige K & Valderrabano M (2014). Ethanol infusion in the vein of Marshall leads to parasympathetic denervation of the human left atrium: implications for atrial fibrillation. *J Am Coll Cardiol* **63**, 1892–1901.
- Barajas L (1964). The innervation of the juxtaglomerular apparatus. An electron microscopic study of the innervation of the glomerular arterioles. *Lab Invest* **13**, 916–929.
- Barajas L & Muller J (1973). The innervation of the juxtaglomerular apparatus and surrounding tubules: a quantitative analysis by serial section electron microscopy. *J Ultrastruct Res* **43**, 107–132.
- Barnes PM, Powell-Griner E, McFann K & Nahin RL (2004). Complementary and alternative medicine use among adults: United States, 2002. *Adv Data* **343**, 1–19.
- Barrett LF & Simmons WK (2015). Interoceptive predictions in the brain. *Nat Rev Neurosci* **16**, 419–429.
- Barsky AJ, Ahern DK, Brener J, Surman OS, Ring C & Dec GW (1998). Palpitations and cardiac awareness after heart transplantation. *Psychosom Med* **60**, 557–562.
- Barsky AJ, Cleary PD, Brener J & Ruskin JN (1993). The perception of cardiac activity in medical outpatients. *Cardiology* **83**, 304–315.
- Bauer A, Barthel P, Schneider R, Ulm K, Muller A, Joeinig A, Stich R, Kiviniemi A, Hnatkova K, Huikuri H, Schomig A, Malik M & Schmidt G (2009). Improved Stratification of Autonomic Regulation for risk prediction in post-infarction patients with preserved left ventricular function (ISAR-Risk). *Eur Heart J* 30, 576–583.
- Bauer A, Kantelhardt JW, Barthel P, Schneider R, Makikallio T, Ulm K, Hnatkova K, Schomig A, Huikuri H, Bunde A, Malik M & Schmidt G (2006). Deceleration capacity of heart rate as a predictor of mortality after myocardial infarction: cohort study. *Lancet* 367, 1674–1681.

- Beaumont E, Salavatian S, Southerland EM, Vinet A, Jacquemet V, Armour JA & Ardell JL (2013). Network interactions within the canine intrinsic cardiac nervous system: implications for reflex control of regional cardiac function. *J Physiol* **591**, 4515–4533.
- Beaumont E, Southerland EM, Hardwick JC, Wright GL, Ryan S, Li Y, KenKnight BH, Armour JA & Ardell JL (2015). Vagus nerve stimulation mitigates intrinsic cardiac neuronal and adverse myocyte remodeling postmyocardial infarction. *Am J Physiol Heart Circ Physiol* **309**, H1198–H1206.
- Ben-Menachem E (2001). Vagus nerve stimulation, side effects, and long-term safety. *J Clin Neurophysiol* **18**, 415–418.
- Ben-Menachem E (2002). Vagus-nerve stimulation for the treatment of epilepsy. *Lancet Neurol* **1**, 477–482.
- Bengel FM, Ueberfuhr P, Schiepel N, Nekolla SG, Reichart B & Schwaiger M (2001). Myocardial efficiency and sympathetic reinnervation after orthotopic heart transplantation: a noninvasive study with positron emission tomography. *Circulation* **103**, 1881–1886.
- Bercovitz A, Sengupta M, Jones A & Harris-Kojetin LD (2011). Complementary and alternative therapies in hospice: The national home and hospice care survey: United States, 2007. *Natl Health Stat Report* **33**, 1–20.
- Bhatt DL, Kandzari DE, O'Neill WW, D'Agostino R, Flack JM, Katzen BT, Leon MB, Liu M, Mauri L, Negoita M, Cohen SA, Oparil S, Rocha-Singh K, Townsend RR & Bakris GL (2014). A controlled trial of renal denervation for resistant hypertension. *N Engl J Med* **370**, 1393–1401.
- Billman GE, Cagnoli KL, Csepe T, Li N, Wright P, Mohler PJ & Fedorov VV (2015). Exercise training-induced bradycardia: evidence for enhanced parasympathetic regulation without changes in intrinsic sinoatrial node function. *J Appl Physiol* **118**, 1344–1355.
- Billman GE, Schwartz PJ & Stone HL (1984). The effects of daily exercise on susceptibility to sudden cardiac death. *Circulation* **69**, 1182–1189.
- Black LI, Clarke TC, Barnes PM, Stussman BJ & Nahin RL (2015). Use of complementary health approaches among children aged 4-17 years in the United States: National Health Interview Survey, 2007-2012. *Natl Health Stat Report* **78**, 1–19.
- Boon P, Vonck K, De Reuck J & Caemaert J (2001). Vagus nerve stimulation for refractory epilepsy. *Seizure* **10**, 456–458; quiz 458–460.
- Booth LC, Nishi EE, Yao ST, Ramchandra R, Lambert GW, Schlaich MP & May CN (2015). Reinnervation following catheter-based radio-frequency renal denervation. *Exp Physiol* **100**, 485–490.
- Borges DG, Monteiro RA, Schmidt A & Pazin-Filho A (2013). World soccer cup as a trigger of cardiovascular events. *Arq Bras Cardiol* **100**, 546–552.
- Bosnjak ZJ & Kampine JP (1989). Cardiac sympathetic afferent cell bodies are located in the peripheral nervous system of the cat. *Circ Res* **64**, 554–562.
- Bourke T, Vaseghi M, Michowitz Y, Sankhla V, Shah M, Swapna N, Boyle NG, Mahajan A, Narasimhan C, Lokhandwala Y & Shivkumar K (2010). Neuraxial modulation for refractory ventricular arrhythmias: value of thoracic epidural anesthesia and surgical left cardiac sympathetic denervation. *Circulation* 121, 2255–2262.

- Brener J & Kluvitse C (1988). Heartbeat detection: judgments of the simultaneity of external stimuli and heartbeats. *Psychophysiology* **25**, 554–561.
- Buckley U, Yamakawa K, Takamiya T, Andrew Armour J, Shivkumar K & Ardell JL (2016). Targeted stellate decentralization: Implications for sympathetic control of ventricular electrophysiology. *Heart Rhythm* **13**, 282–288.
- Bultz BD, Cummings GG, Grassi L, Travado L, Hoekstra-Weebers J & Watson M (2014). 2013 President's Plenary International Psycho-oncology Society: embracing the IPOS standards as a means of enhancing comprehensive cancer care. *Psychooncology* **23**, 1073–1078.
- Burn JH, Williams EM & Walker JM (1955). The effects of acetylcholine in the heart-lung preparation including the production of auricular fibrillation. *J Physiol* **128**, 277–293.
- Callister R, Suwarno NO & Seals DR (1992). Sympathetic activity is influenced by task difficulty and stress perception during mental challenge in humans. *J Physiol* **454**, 373–387.
- Cameron OG (2002). Visceral Sensory Neuroscience: Interoception. Oxford University Press, New York.
- Cameron OG & Minoshima S (2002). Regional brain activation due to pharmacologically induced adrenergic interoceptive stimulation in humans. *Psychosom Med* **64**, 851–861.
- Campese VM & Kogosov E (1995). Renal afferent denervation prevents hypertension in rats with chronic renal failure. *Hypertension* **25**, 878–882.
- Canales-Johnson A, Silva C, Huepe D, Rivera-Rei A, Noreika V, Garcia Mdel C, Silva W, Ciraolo C, Vaucheret E, Sedeno L, Couto B, Kargieman L, Baglivo F, Sigman M, Chennu S, Ibanez A, Rodriguez E & Bekinschtein TA (2015). Auditory feedback differentially modulates behavioral and neural markers of objective and subjective performance when tapping to your heartbeat. *Cereb Cortex* **25**, 4490–4503.
- Cao JM, Chen LS, KenKnight BH, Ohara T, Lee MH, Tsai J, Lai WW, Karagueuzian HS, Wolf PL, Fishbein MC & Chen PS (2000*a*). Nerve sprouting and sudden cardiac death. *Circ Res* **86**, 816–821.
- Cao JM, Fishbein MC, Han JB, Lai WW, Lai AC, Wu TJ, Czer L, Wolf PL, Denton TA, Shintaku IP, Chen PS & Chen LS (2000b). Relationship between regional cardiac hyperinnervation and ventricular arrhythmia. *Circulation* 101, 1960–1969.
- Cardinal R, Page P, Vermeulen M, Ardell JL & Armour JA (2009). Spatially divergent cardiac responses to nicotinic stimulation of ganglionated plexus neurons in the canine heart. *Auton Neurosci* **145**, 55–62.
- Carlson MD, Geha AS, Hsu J, Martin PJ, Levy MN, Jacobs G & Waldo AL (1992). Selective stimulation of parasympathetic nerve fibers to the human sinoatrial node. *Circulation* **85**, 1311–1317.
- Carter JR & Ray CA (2009). Sympathetic neural responses to mental stress: responders, nonresponders and sex differences. *Am J Physiol Heart Circ Physiol* **296**, H847–853.
- Cha YM, Oh J, Miyazaki C, Hayes DL, Rea RF, Shen WK, Asirvatham SJ, Kemp BJ, Hodge DO, Chen PS & Chareonthaitawee P (2008). Cardiac resynchronization therapy upregulates cardiac autonomic control. *J Cardiovasc Electrophysiol* **10**, 1045–1052.

- Chang CM, Wu TJ, Zhou SM, Doshi RN, Lee MH, Ohara T, Fishbein MC, Karagueuzian HS, Chen PS & Chen LS (2001). Nerve sprouting and sympathetic hyperinnervation in a canine model of atrial fibrillation produced by prolonged right atrial pacing. *Circulation* **103**, 22–25.
- Charney DS, Heninger GR & Jatlow PI (1985). Increased anxiogenic effects of caffeine in panic disorders. Arch General Psychiatry 42, 233–243.
- Chaya MS, Kurpad AV, Nagendra HR & Nagarathna R (2006). The effect of long term combined yoga practice on the basal metabolic rate of healthy adults. BMC Complement Altern Med 6, 28.
- Chiou CW, Eble JN & Zipes DP (1997). Efferent vagal innervation of the canine atria and sinus and atrioventricular nodes. The third fat pad. *Circulation* **95**, 2573–2584.
- Chou CC, Nihei M, Zhou S, Tan AY, Kawase A, Macias ES, Fishbein MC, Lin SF & Chen PS (2005). Intracellular calcium dynamics and anisotropic reentry in isolated canine pulmonary veins and left atrium. *Circulation* 111, 2889–2297.
- Chugh SS, Reinier K, Teodorescu C, Evanado A, Kehr E, Al Samara M, Mariani R, Gunson K & Jui J (2008).
 Epidemiology of sudden cardiac death: clinical and research implications. *Prog Cardiovasc Dis* 51, 213–228.
- Clancy JA, Mary DA, Witte KK, Greenwood JP, Deuchars SA & Deuchars J (2014). Non-invasive vagus nerve stimulation in healthy humans reduces sympathetic nerve activity. *Brain Stimul* 7, 871–877.
- Clarke TC, Black LI, Stussman BJ, Barnes PM & Nahin RL (2015). Trends in the use of complementary health approaches among adults: United States, 2002–2012. *Natl Health Stat Report* **79**, 1–16.
- Cohen MA & Taylor JA (2002). Short-term cardiovascular oscillations in man: measuring and modeling the physiologies. *J Physiol* **542**, 669–683.
- Collura CA, Johnson JN, Moir C & Ackerman MJ (2009). Left cardiac sympathetic denervation for the treatment of long QT syndrome and catecholaminergic polymorphic ventricular tachycardia using video-assisted thoracic surgery. *Heart Rhythm* **6**, 752–759.
- Converse RL Jr, Jacobsen TN, Toto RD, Jost CM, Cosentino F, Fouad-Tarazi F & Victor RG (1992). Sympathetic overactivity in patients with chronic renal failure. *N Engl J Med* **327**, 1912–1918.
- Cooke WH, Ryan KL & Convertino VA (2004). Lower body negative pressure as a model to study progression to acute hemorrhagic shock in humans. *J Appl Physiol* **96**, 1249–1261.
- Coumel P, Krikler D, Rosen MR, Wellens HJ & Zipes DP (1978). Newer antiarrhythmic drugs. *Pacing Clin Electrophysiol* 1, 521–528.
- Couto B, Salles A, Sedeno L, Peradejordi M, Barttfeld P, Canales-Johnson A, Dos Santos YV, Huepe D, Bekinschtein T, Sigman M, Favaloro R, Manes F & Ibanez A (2013). The man who feels two hearts: the different pathways of interoception. *Soc Cogn Affect Neurosci* **9**, 1253–1260.
- Cowie MR, Woehrle H, Wegscheider K, Angermann C, d'Ortho MP, Erdmann E, Levy P, Simonds AK, Somers VK, Zannad F & Teschler H (2015). Adaptive servo-ventilation for central sleep apnea in systolic heart failure. *N Engl J Med* **373**, 1095–1105.

- Critchley HD, Wiens S, Rotshtein P, Ohman A & Dolan RJ (2004). Neural systems supporting interoceptive awareness. *Nat Neurosci* 7, 189–195.
- Cysarz D & Bussing A (2005). Cardiorespiratory synchronization during Zen meditation. *Eur J Appl Physiol* **95**, 88–95.
- Daly MD, Angell-James JE & Elsner R (1979). Role of carotid-body chemoreceptors and their reflex interactions in bradycardia and cardiac arrest. *Lancet* 1, 764–767.
- Davidson RJ, Horowitz ME, Schwartz GE & Goodman DM (1981). Lateral differences in the latency between finger tapping and the heart beat. *Psychophysiology* **18**, 36–41.
- De Ferrari GM (2014). Vagal stimulation in heart failure. J Cardiovasc Transl Res 7, 310–320.
- De Ferrari GM, Crijns HJ, Borggrefe M, Milasinovic G, Smid J, Zabel M, Gavazzi A, Sanzo A, Dennert R, Kuschyk J, Raspopovic S, Klein H, Swedberg K & Schwartz PJ (2011). Chronic vagus nerve stimulation: a new and promising therapeutic approach for chronic heart failure. *Eur Heart J* **32**, 847–855.
- De Ferrari GM & Dusi V (2015). [Vagus nerve stimulation for the treatment of heart failure]. *G Ital Cardiol* **16**, 147–154.
- De Ferrari GM, Dusi V, Spazzolini C, Bos JM, Abrams DJ, Berul CI, Crotti L, Davis AM, Eldar M, Kharlap M, Khoury A, Krahn AD, Leenhardt A, Moir CR, Odero A, Olde Nordkamp L, Paul T, Roses INF, Shkolnikova M, Till J, Wilde AA, Ackerman MJ & Schwartz PJ (2015). Clinical management of catecholaminergic polymorphic ventricular tachycardia: the role of left cardiac sympathetic denervation. *Circulation* 131, 2185–2193.
- De Ferrari GM, Tuinenburg AE, Ruble S, Brugada J, Klein H, Butter C, Wright DJ, Schubert B, Solomon S, Meyer S, Stein K, Ramuzat A & Zannad F (2014). Rationale and study design of the NEuroCardiac TherApy foR Heart Failure Study: NECTAR-HF. *Eur J Heart Fail* 16, 692–699.
- Denq JC, O'Brien PC & Low PA (1998). Normative data on phases of the Valsalva maneuver. *J Clin Neurophysiol* **15**, 535–540.
- DeSilva RA & Lown B (1978). Ventricular premature beats, stress, and sudden death. *Psychosomatics* **19**, 649–653, 656–661.
- Diaz T & Taylor JA (2006). Probing the arterial baroreflex: is there a 'spontaneous' baroreflex? *Clin Auton Res* **16**, 256–261.
- DiBona GF & Esler M (2010). Translational medicine: the antihypertensive effect of renal denervation. *Am J Physiol Regul Integr Comp Physiol* **298**, R245–R253.
- DiBona GF & Kopp UC (1997). Neural control of renal function. *Physiol Rev* 77, 75–197.
- Dietz NM, Rivera JM, Eggener SE, Fix RT, Warner DO & Joyner MJ (1994). Nitric oxide contributes to the rise in forearm blood flow during mental stress in humans. *J Physiol* **480**, 361–368.
- Dillon DJ, Gorman JM, Liebowitz MR, Fyer AJ & Klein DF (1987). Measurement of lactate-induced panic and anxiety. *Psychiatry Res* **20**, 97–105.
- Dimsdale JE & Mills PJ (2002). An unanticipated effect of meditation on cardiovascular pharmacology and physiology. *Am J Cardiol* **90**, 908–909.

- Docimo S, Piccolo C, Van Arsdale D & Elkowitz DE (2008). Pathology-dependent histological changes of the left stellate Ganglia: a cadaveric study. *Clin Med Pathol* **1**, 105–113.
- Domschke K, Stevens S, Pfleiderer B & Gerlach AL (2010). Interoceptive sensitivity in anxiety and anxiety disorders: an overview and integration of neurobiological findings. *Clin Psychol Rev* **30**, 1–11.
- Donald DE & Shepherd JT (1980). Autonomic regulation of the peripheral circulation. *Annu Rev Physiol* **42**, 429–439.
- Drapeau J, El Helou V, Clement R, Bel-Hadj S, Gosselin H, Trudeau LE, Villeneuve L & Calderone A (2004). Nestin-expressing neural stem cells identified in the scar following myocardial infarction. *J Cell Physiol* **204**, 51–62.
- Dutoit AP, Hart EC, Charkoudian N, Wallin BG, Curry TB & Joyner MJ (2010). Cardiac baroreflex sensitivity is not correlated to sympathetic baroreflex sensitivity within healthy, young humans. *Hypertension* **56**, 1118–1123.
- Dworkin BR (2007). Interoception. In *Handbook of Psychophysiology*, 3rd edn, ed. Cacciopo JT, Tassinary LG, Bernston GG, pp. 482–506. Cambridge University Press.
- Dworkin BR, Dworkin S & Tang X (2000). Carotid and aortic baroreflexes of the rat: I. Open-loop steady-state properties and blood pressure variability. *Am J Physiol Regul Integr Comp Physiol* **279**, R1910–R1921.
- Edgerton JR, Brinkman WT, Weaver T, Prince SL, Culica D, Herbert MA & Mack MJ (2010). Pulmonary vein isolation and autonomic denervation for the management of paroxysmal atrial fibrillation by a minimally invasive surgical approach. *J Thorac Cardiovasc Surg* **140**, 823–828.
- Ehlers A & Breuer P (1996). How good are patients with panic disorder at perceiving their heartbeats? *Biol Psychol* **42**, 165–182
- Ehlers A, Mayou RA, Sprigings DC & Birkhead J (2000). Psychological and perceptual factors associated with arrhythmias and benign palpitations. *Psychosom Med* **62**, 693–702.
- Eisenberg DM, Kessler RC, Foster C, Norlock FE, Calkins DR & Delbanco TL (1993). Unconventional medicine in the United States. Prevalence, costs, and patterns of use. *N Engl J Med* **328**, 246–252.
- El Mhandi L, Pichot V, Calmels P, Gautheron V, Roche F & Feasson L (2011). Exercise training improves autonomic profiles in patients with Charcot-Marie-Tooth disease. *Muscle Nerve* **44**, 732–736.
- Esler M (2014). Illusions of truths in the Symplicity HTN-3 trial: generic design strengths but neuroscience failings. *J Am Soc Hypertens* **8**, 593–598.
- Esler M, Lambert G & Jennings G (1990). Increased regional sympathetic nervous activity in human hypertension: causes and consequences. *J Hypertens* Suppl. **8**, S53–S57.
- Esler MD, Krum H, Sobotka PA, Schlaich MP, Schmieder RE & Bohm M (2010). Renal sympathetic denervation in patients with treatment-resistant hypertension (The Symplicity HTN-2 Trial): a randomised controlled trial. *Lancet* **376**, 1903–1909
- Estes EH Jr & Izlar HL Jr (1961). Recurrent ventricular tachycardia. A case successfully treated by bilateral cardiac sympathectomy. *Am J Med* **31**, 493–497.

- Exner DV, Kavanagh KM, Slawnych MP, Mitchell LB, Ramadan D, Aggarwal SG, Noullett C, Van Schaik A, Mitchell RT, Shibata MA, Gulamhussein S, McMeekin J, Tymchak W, Schnell G, Gillis AM, Sheldon RS, Fick GH & Duff HJ (2007). Noninvasive risk assessment early after a myocardial infarction the REFINE study. *J Am Coll Cardiol* **50**, 2275–2284.
- Fadel PJ, Ogoh S, Keller DM & Raven PB (2003). Recent insights into carotid baroreflex function in humans using the variable pressure neck chamber. *Exp Physiol* 88, 671–680.
- Fallavollita JA, Heavey BM, Luisi AJ Jr, Michalek SM, Baldwa S, Mashtare TL Jr, Hutson AD, Dekemp RA, Haka MS, Sajjad M, Cimato TR, Curtis AB, Cain ME & Canty JM Jr (2014). Regional myocardial sympathetic denervation predicts the risk of sudden cardiac arrest in ischemic cardiomyopathy. *J Am Coll Cardiol* **63**, 141–149.
- Fallgatter AJ, Neuhauser B, Herrmann MJ, Ehlis AC, Wagener A, Scheuerpflug P, Reiners K & Riederer P (2003). Far field potentials from the brain stem after transcutaneous vagus nerve stimulation. *J Neural Transm* **110**, 1437–1443.
- Fatouleh RH, Hammam E, Lundblad LC, Macey PM, McKenzie DK, Henderson LA & Macefield VG (2014). Functional and structural changes in the brain associated with the increase in muscle sympathetic nerve activity in obstructive sleep apnoea. *Neuroimage Clin* **6**, 275–283.
- Fedorowski A, Stavenow L, Hedblad B, Berglund G, Nilsson PM & Melander O (2010). Orthostatic hypotension predicts all-cause mortality and coronary events in middle-aged individuals (The Malmo Preventive Project). *Eur Heart J* 31, 85–91
- Feldman JL, Del Negro CA & Gray PA (2013). Understanding the rhythm of breathing: so near, yet so far. *Annu Rev Physiol* **75**, 423–452.
- Fitzpatrick AP, Banner N, Cheng A, Yacoub M & Sutton R (1993). Vasovagal reactions may occur after orthotopic heart transplantation. *J Am Coll Cardiol* **21**, 1132–1137.
- Flaker GC, Belew K, Beckman K, Vidaillet H, Kron J, Safford R, Mickel M & Barrell P (2005). Asymptomatic atrial fibrillation: demographic features and prognostic information from the Atrial Fibrillation Follow-up Investigation of Rhythm Management (AFFIRM) study. *Am Heart J* 149, 657–663.
- Florea VG & Cohn JN (2014). The autonomic nervous system and heart failure. *Circ Res* **114**, 1815–1826.
- Frangos E, Ellrich J & Komisaruk BR (2015). Non-invasive access to the vagus nerve central projections via electrical stimulation of the external ear: fMRI evidence in humans. *Brain Stimul* **8**, 624–636.
- Frasure-Smith N & Lesperance F (2003). Depression and other psychological risks following myocardial infarction. *Arch Gen Psychiatry* **60**, 627–636.
- Frasure-Smith N, Lesperance F & Talajic M (1993). Depression following myocardial infarction. Impact on 6-month survival. *JAMA* **270**, 1819–1825.
- Frasure-Smith N, Lesperance F & Talajic M (1995). Depression and 18-month prognosis after myocardial infarction. *Circulation* **91**, 999–1005.

- Freedenberg VA, Thomas SA & Friedmann E (2015). A pilot study of a mindfulness based stress reduction program in adolescents with implantable cardioverter defibrillators or pacemakers. *Pediatr Cardiol* **36**, 786–795.
- Freedland KE (2013). Demanding attention: reconsidering the role of attention control groups in behavioral intervention research. *Psychosom Med* **75**, 100–102.
- Freedland KE, Mohr DC, Davidson KW & Schwartz JE (2011). Usual and unusual care: existing practice control groups in randomized controlled trials of behavioral interventions. *Psychosom Med* **73**, 323–335.
- Fukuda K, Kanazawa H, Aizawa Y, Ardell JL & Shivkumar K (2015). Cardiac innervation and sudden cardiac death. *Circ Res* 116, 2005–2019.
- Garcia-Moran E, Sandin-Fuentes MG, Alvarez Lopez JC, Duro-Aguado I, Uruena-Martinez N & Hernandez-Luis C (2013). Electrical storm secondary to acute myocardial infarction and heart failure treated with left stellate ganglion block. *Rev Esp Cardiol* **66**, 595–597.
- Garimella RS, Chung EH, Mounsey JP, Schwartz JD, Pursell I & Gehi AK (2015). Accuracy of patient perception of their prevailing rhythm: A comparative analysis of monitor data and questionnaire responses in patients with atrial fibrillation. *Heart Rhythm* **12**, 658–665.
- Garrey WE (1924). Auricular fibrillation. Physiol Rev 4, 215–250.
- Gherghiceanu M, Hinescu ME, Andrei F, Mandache E, Macarie CE, Faussone-Pellegrini MS & Popescu LM (2008). Interstitial Cajal-like cells (ICLC) in myocardial sleeves of human pulmonary veins. *J Cell Mol Med* 12, 1777–1781.
- Ghuran A, Reid F, La Rovere MT, Schmidt G, Bigger JT Jr, Camm AJ, Schwartz PJ & Malik M (2002). Heart rate turbulence-based predictors of fatal and nonfatal cardiac arrest (The Autonomic Tone and Reflexes After Myocardial Infarction substudy). *Am J Cardiol* **89**, 184–190.
- Golbidi S, Frisbee JC & Laher I (2015). Chronic stress impacts the cardiovascular system: animal models and clinical outcomes. *Am J Physiol Heart Circ Physiol* **308**, H1476–H1498.
- Goldberger JJ, Subacius H, Patel T, Cunnane R & Kadish AH (2014). Sudden cardiac death risk stratification in patients with nonischemic dilated cardiomyopathy. *J Am Coll Cardiol* **63**, 1879–1889.
- Gordon A, Tyni-Lenne R, Jansson E, Kaijser L, Theodorsson-Norheim E & Sylven C (1997). Improved ventilation and decreased sympathetic stress in chronic heart failure patients following local endurance training with leg muscles. *J Card Fail* 3, 3–12.
- Goyal M, Singh S, Sibinga EM, Gould NF, Rowland-Seymour A, Sharma R, Berger Z, Sleicher D, Maron DD, Shihab HM, Ranasinghe PD, Linn S, Saha S, Bass EB & Haythornthwaite JA (2014). Meditation programs for psychological stress and well-being: a systematic review and meta-analysis. *JAMA Intern Med* 174, 357–368.
- Gray AL, Johnson TA, Ardell JL & Massari VJ (2004). Parasympathetic control of the heart. II. A novel interganglionic intrinsic cardiac circuit mediates neural control of heart rate. *J Appl Physiol* **96**, 2273–2278.

- Gray MA, Taggart P, Sutton PM, Groves D, Holdright DR, Bradbury D, Brull D & Critchley HD (2007). A cortical potential reflecting cardiac function. *Proc Natl Acad Sci* **104**, 6818–6823.
- Groehs RV, Toschi-Dias E, Antunes-Correa LM, Trevizan PF, Rondon MU, Oliveira P, Alves MJ, Almeida DR, Middlekauff HR & Negrao CE (2015). Exercise training prevents the deterioration in the arterial baroreflex control of sympathetic nerve activity in chronic heart failure patients. *Am J Physiol Heart Circ Physiol* **308**, H1096–H1102.
- Gunda S, Kanmanthareddy A, Atkins D, Bommana S, Pimentel R, Drisko J, Dibiase L, Beheiry S, Hao S, Natale A & Lakkireddy D (2015). Role of yoga as an adjunctive therapy in patients with neurocardiogenic syncope: a pilot study. *J Interv Card Electrophysiol* **43**, 105–110.
- Gurguis GN, Vitton BJ & Uhde TW (1997). Behavioral, sympathetic and adrenocortical responses to yohimbine in panic disorder patients and normal controls. *Psychiatry Res* **71**, 27–39.
- Guyenet PG, Stornetta RL, Bochorishvili G, Depuy SD, Burke PG & Abbott SB (2013). C1 neurons: the body's EMTs. *Am J Physiol Regul Integr Comp Physiol* **305**, R187–R204.
- Haack KK & Zucker IH (2015). Central mechanisms for exercise training-induced reduction in sympatho-excitation in chronic heart failure. *Auton Neurosci* 188, 44–50.
- Hainsworth R (2014). Cardiovascular control from cardiac and pulmonary vascular receptors. *Exp Physiol* **99**, 312–319.
- Haissaguerre M, Jais P, Shah DC, Takahashi A, Hocini M, Quiniou G, Garrigue S, Le Mouroux A, Le Metayer P & Clementy J (1998). Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. *N Engl J Med* **339**, 659–666.
- Hamann JJ, Ruble SB, Stolen C, Wang M, Gupta RC, Rastogi S & Sabbah HN (2013). Vagus nerve stimulation improves left ventricular function in a canine model of chronic heart failure. *Eur J Heart Fail* **15**, 1319–1326.
- Han FT, Kasirajan V, Kowalski M, Kiser R, Wolfe L, Kalahasty G, Shepard RK, Wood MA & Ellenbogen KA (2009). Results of a minimally invasive surgical pulmonary vein isolation and ganglionic plexi ablation for atrial fibrillation: single-center experience with 12-month follow-up. *Circ Arrhythm Electrophysiol* 2, 370–377.
- Han J & Moe GK (1964). Nonuniform recovery of excitability in ventricular muscle. *Circ Res* **14**, 44–60.
- Han S, Kobayashi K, Joung B, Piccirillo G, Maruyama M & Vinters H (2011). Electroanatomical remodeling of the left stellate ganglion after myocardial infarction. *J Am Coll Cardiol* **59**, 954–961.
- Harper C (2009). The neuropathology of alcohol-related brain damage. *Alcohol Alcohol* **44**, 136–140.
- Harper RM, Macey PM, Henderson LA, Woo MA, Macey KE, Frysinger RC, Alger JR, Nguyen KP & Yan-Go FL (2003). fMRI responses to cold pressor challenges in control and obstructive sleep apnea subjects. *J Appl Physiol* 94, 1583–1595.
- Hauptman PJ, Schwartz PJ, Gold MR, Borggrefe M, Van Veldhuisen DJ, Starling RC & Mann DL (2012). Rationale and study design of the increase of vagal tone in heart failure study: INOVATE-HF. *Am Heart J* 163, 954–962.e1.

- Hayase J, Patel J, Narayan SM & Krummen DE (2013). Percutaneous stellate ganglion block suppressing VT and VF in a patient refractory to VT ablation. *J Cardiovasc Electrophysiol* **24**, 926–928.
- Heistad DD, Abboud FM, Mark AL & Schmid PG (1974). Interaction of baroreceptor and chemoreceptor reflexes. Modulation of the chemoreceptor reflex by changes in baroreceptor activity. *J Clin Invest* **53**, 1226–1236.
- Henderson LA, Woo MA, Macey PM, Macey KE, Frysinger RC, Alger JR, Yan-Go F & Harper RM (2003). Neural responses during Valsalva maneuvers in obstructive sleep apnea syndrome. *J Appl Physiol* **94**, 1063–1074.
- Henegar JR, Zhang Y, Hata C, Narciso I, Hall ME & Hall JE (2015). Catheter-based radiofrequency renal denervation: location effects on renal norepinephrine. *Am J Hypertens* 28, 909–914.
- Henning RJ, Khalil IR & Levy MN (1990). Vagal stimulation attenuates sympathetic enhancement of left ventricular function. *Am J Physiol Heart Circ Physiol* **258**, H1470–H1475.
- Herring N, Cranley J, Lokale MN, Li D, Shanks J, Alston EN, Girard BM, Carter E, Parsons RL, Habecker BA & Paterson DJ (2012). The cardiac sympathetic co-transmitter galanin reduces acetylcholine release and vagal bradycardia: implications for neural control of cardiac excitability. *J Mol Cell Cardiol* **52**, 667–676.
- Herring N, Lokale MN, Danson EJ, Heaton DA & Paterson DJ (2008). Neuropeptide Y reduces acetylcholine release and vagal bradycardia via a Y2 receptor-mediated, protein kinase C-dependent pathway. *J Mol Cell Cardiol* **44**, 477–485.
- Himura Y, Felten SY, Kashiki M, Lewandowski TJ, Delehanty JM & Liang CS (1993). Cardiac noradrenergic nerve terminal abnormalities in dogs with experimental congestive heart failure. *Circulation* **88**, 1299–1309.
- Hindricks G, Piorkowski C, Tanner H, Kobza R, Gerds-Li JH, Carbucicchio C & Kottkamp H (2005). Perception of atrial fibrillation before and after radiofrequency catheter ablation: relevance of asymptomatic arrhythmia recurrence. *Circulation* **112**, 307–313.
- Hjalmarson A (1997). Effects of beta blockade on sudden cardiac death during acute myocardial infarction and the postinfarction period. *Am J Cardiol* **80**, 35J–39J.
- Hohnloser SH, Kuck KH, Dorian P, Roberts RS, Hampton JR, Hatala R, Fain E, Gent M & Connolly SJ (2004). Prophylactic use of an implantable cardioverter-defibrillator after acute myocardial infarction. *N Engl J Med* **351**, 2481–2488.
- Holmqvist F, Guan N, Zhu Z, Kowey PR, Allen LA, Fonarow GC, Hylek EM, Mahaffey KW, Freeman JV, Chang P, Holmes DN, Peterson ED, Piccini JP & Gersh BJ (2015). Impact of obstructive sleep apnea and continuous positive airway pressure therapy on outcomes in patients with atrial fibrillation—Results from the Outcomes Registry for Better Informed Treatment of Atrial Fibrillation (ORBIT-AF). *Am Heart J* 169, 647–654.e2.
- Hoover DB, Isaacs ER, Jacques F, Hoard JL, Page P & Armour JA (2009). Localization of multiple neurotransmitters in surgically derived specimens of human atrial ganglia. *Neuroscience* **164**, 1170–1179.

- Hopkins DA & Armour JA (1984). Localization of sympathetic postganglionic and parasympathetic preganglionic neurons which innervate different regions of the dog heart. *J Comp Neurol* **229**, 186–198.
- Hopkins DA & Armour JA (1989). Ganglionic distribution of afferent neurons innervating the canine heart and cardiopulmonary nerves. *J Auton Nerv Syst* **26**, 213–222.
- Hopkins DA, Bieger D, deVente J & Steinbusch WM (1996). Vagal efferent projections: viscerotopy, neurochemistry and effects of vagotomy. *Prog Brain Res* **107**, 79–96.
- Hsieh MH, Chiou CW, Wen ZC, Wu CH, Tai CT, Tsai CF, Ding YA, Chang MS & Chen SA (1999). Alterations of heart rate variability after radiofrequency catheter ablation of focal atrial fibrillation originating from pulmonary veins. *Circulation* **100**, 2237–2243.
- Huang MH, Ardell JL, Hanna BD, Wolf SG & Armour JA (1993*a*). Effects of transient coronary artery occlusion on canine intrinsic cardiac neuronal activity. *Integr Physiol Behav Sci* **28**, 5–21.
- Huang MH, Sylven C, Pelleg A, Smith FM & Armour JA (1993b). Modulation of in situ canine intrinsic cardiac neuronal activity by locally applied adenosine, ATP, or analogues. *Am J Physiol Regul Integr Comp Physiol* **265**, R914–R922.
- Ikeda T, Yoshino H, Sugi K, Tanno K, Shimizu H, Watanabe J, Kasamaki Y, Yoshida A & Kato T (2006). Predictive value of microvolt T-wave alternans for sudden cardiac death in patients with preserved cardiac function after acute myocardial infarction: results of a collaborative cohort study. J Am Coll Cardiol 48, 2268–2274.
- James TN, Zipes DP, Finegan RE, Eisele JW & Carter JE (1979). Cardiac ganglionitis associated with sudden unexpected death. Ann Intern Med 91, 727–730.
- Janes RD, Brandys JC, Hopkins DA, Johnstone DE, Murphy DA & Armour JA (1986). Anatomy of human extrinsic cardiac nerves and ganglia. Am J Cardiol 57, 299–309.
- Janig W (1996). Neurobiology of visceral afferent neurons: neuroanatomy, functions, organ regulations and sensations. *Biol Psychol* **42**, 29–51.
- Janig W (2006). The Integrative Action of the Autonomic Nervous System: Neurobiology of Homeostasis. Cambridge University Press, Cambridge.
- Jayachandran JV, Sih HJ, Winkle W, Zipes DP, Hutchins GD & Olgin JE (2000). Atrial fibrillation produced by prolonged rapid atrial pacing is associated with heterogeneous changes in atrial sympathetic innervation. *Circulation* **101**, 1185–1191.
- Johnson DC, Thom NJ, Stanley EA, Haase L, Simmons AN, Shih PA, Thompson WK, Potterat EG, Minor TR & Paulus MP (2014). Modifying resilience mechanisms in at-risk individuals: a controlled study of mindfulness training in Marines preparing for deployment. Am J Psychiatry 171, 844–853.
- Jones GE (1994). Perception of visceral sensations: a review of recent findings, methodologies, and future directions. In *Advances in Psychophysiology: A research annual*, ed. Jennings JR and Ackles PK, vol. 5, pp. 155–192. Jessica Kingsley Publishers, London.

- Jones GE & Hollandsworth JG (1981). Heart rate discrimination before and after exercise-induced augmented cardiac activity. *Psychophysiology* **18**, 252–257.
- Jones GE, Jones KR, Cunningham RA & Caldwell JA (1985). Cardiac awareness in infarct patients and normals. *Psychophysiology* **22**, 480–487.
- Jouven X, Empana JP, Schwartz PJ, Desnos M, Courbon D & Ducimetiere P (2005). Heart-rate profile during exercise as a predictor of sudden death. *N Engl J Med* **352**, 1951–1958.
- Joyner MJ & Casey DP (2015). Regulation of increased blood flow (hyperemia) to muscles during exercise: a hierarchy of competing physiological needs. *Physiol Rev* **95**, 549–601.
- Joyner MJ & Dietz NM (2003). Sympathetic vasodilation in human muscle. *Acta Physiol Scand* 177, 329–336.
- Julien C (2006). The enigma of Mayer waves: facts and models. *Cardiovasc Res* **70**, 12–21.
- Kaikkonen KM, Korpelainen RI, Tulppo MP, Kaikkonen HS, Vanhala ML, Kallio MA, Keinanen-Kiukaanniemi SM & Korpelainen JT (2014). Physical activity and aerobic fitness are positively associated with heart rate variability in obese adults. *J Phys Act Health* 11, 1614–1621.
- Kamibayashi T, Hayashi Y, Mammoto T, Yamatodani A, Taenaka N & Yoshiya I (1995). Thoracic epidural anesthesia attenuates halothane-induced myocardial sensitization to dysrhythmogenic effect of epinephrine in dogs. *J Phys Act Health* **82**, 129–134.
- Kammerling JJ, Green FJ, Watanabe AM, Inoue H, Barber MJ, Henry DP & Zipes DP (1987). Denervation supersensitivity of refractoriness in noninfarcted areas apical to transmural myocardial infarction. *Circulation* **76**, 383–393.
- Kanazawa H, Ieda M, Kimura K, Arai T, Kawaguchi-Manabe H, Matsuhashi T, Endo J, Sano M, Kawakami T, Kimura T, Monkawa T, Hayashi M, Iwanami A, Okano H, Okada Y, Ishibashi-Ueda H, Ogawa S & Fukuda K (2010). Heart failure causes cholinergic transdifferentiation of cardiac sympathetic nerves via gp130-signaling cytokines in rodents. *J Clin Invest* 120, 408–421.
- Kandzari DE, Bhatt DL, Brar S, Devireddy CM, Esler M, Fahy M, Flack JM, Katzen BT, Lea J, Lee DP, Leon MB, Ma A, Massaro J, Mauri L, Oparil S, O'Neill WW, Patel MR, Rocha-Singh K, Sobotka PA, Svetkey L, Townsend RR & Bakris GL (2015). Predictors of blood pressure response in the SYMPLICITY HTN-3 trial. *Eur Heart J* 36, 219–227.
- Katkin ES, Blascovich J & Goldband S (1981). Empirical assessment of visceral self-perception: individual and sex differences in the acquisition of heartbeat discrimination. *J Pers Soc Psychol* **40**, 1095–1101.
- Katkin ES, Morell MA, Goldband S, Bernstein GL & Wise JA (1982). Individual differences in heartbeat discrimination. *Psychophysiology* **19**, 160–166.
- Katkin ES, Wiens S & Ohman A (2001). Nonconscious fear conditioning, visceral perception, and the development of gut feelings. *Psychol Sci* **12**, 366–370.
- Katritsis DG, Giazitzoglou E, Zografos T, Pokushalov E, Po SS & Camm AJ (2011). Rapid pulmonary vein isolation combined with autonomic ganglia modification: a randomized study. *Heart Rhythm* **8**, 672–678.

- Katritsis DG, Pokushalov E, Romanov A, Giazitzoglou E, Siontis GC, Po SS, Camm AJ & Ioannidis JP (2013). Autonomic denervation added to pulmonary vein isolation for paroxysmal atrial fibrillation: a randomized clinical trial. *J Am Coll Cardiol* **62**, 2318–2325.
- Kember G, Ardell JL, Armour JA & Zamir M (2014). Vagal nerve stimulation therapy: What is being stimulated? *PLoS One* **9**, e114498.
- Kember G, Armour JA & Zamir M (2013*a*). Dynamic neural networking as a basis for plasticity in the control of heart rate. *I Theo Biol* **317**, 39–46.
- Kember G, Armour JA & Zamir M (2013*b*). Neural control hierarchy of the heart has not evolved to deal with myocardial ischemia. *Physiol Genomics* **45**, 638–644.
- Kember GC, Zamir M & Armour JA (2004). "Smart" baroreception along the aortic arch, with reference to essential hypertension. *Phys Rev E Stat Nonlin Soft Matter Phys* **70**, 051914.
- Kemp AH, Quintana DS, Gray MA, Felmingham KL, Brown K & Gatt JM (2010). Impact of depression and antidepressant treatment on heart rate variability: a review and meta-analysis. *Biol Psychiatry* **67**, 1067–1074.
- Kern M, Aertsen A, Schulze-Bonhage A & Ball T (2013). Heart cycle-related effects on event-related potentials, spectral power changes, and connectivity patterns in the human ECoG. *Neuroimage* 81, 178–190.
- Khalsa S, Rudrauf D & Tranel D (2009*a*). Interoceptive awareness declines with age. *Psychophysiology* **46**, 1130–1136.
- Khalsa SS, Craske MG, Li W, Vangala S, Strober M & Feusner JD (2015*a*). Altered interoceptive awareness in anorexia nervosa: Effects of meal anticipation, consumption and bodily arousal. *Int J Eat Disord* **48**, 889–897.
- Khalsa SS, Rudrauf D, Damasio AR, Davidson RJ, Lutz A & Tranel D (2008). Interoceptive awareness in experienced meditators. *Psychophysiology* **45**, 671–677.
- Khalsa SS, Rudrauf D, Davidson RJ & Tranel D (2015*b*). The effect of meditation on regulation of internal body states. *Front Psychol* **6**, 924.
- Khalsa SS, Rudrauf D, Sandesara C, Olshansky B & Tranel D (2009*a*). Bolus isoproterenol infusions provide a reliable method for assessing interoceptive awareness. *Int J Psychophysiol* **72**, 34–45.
- Khalsa SS, Rudrauf D, Feinstein JS & Tranel D (2009*b*). The pathways of interoceptive awareness. *Nat Neurosci* **12**, 1494–1496.
- Kienbaum P, Karlssonn T, Sverrisdottir YB, Elam M & Wallin BG (2001). Two sites for modulation of human sympathetic activity by arterial baroreceptors? *J Physiol* **531**, 861–869.
- Kim DT, Luthringer DJ, Lai AC, Suh G, Czer L, Chen LS, Chen PS & Fishbein MC (2004). Sympathetic nerve sprouting after orthotopic heart transplantation. *J Heart Lung Transplant* **23**, 1349–1358.
- Kleckner IR, Wormwood JB, Simmons WK, Barrett LF & Quigley KS (2015). Methodological recommendations for a heartbeat detection-based measure of interoceptive sensitivity. *Psychophysiology* **52**, 1432–1440.

- Kleiber AC, Zheng H, Schultz HD, Peuler JD & Patel KP (2008). Exercise training normalizes enhanced glutamate-mediated sympathetic activation from the PVN in heart failure. *Am J Physiol Regul Integr Comp Physiol* **294**, R1863–R1872.
- Kober L, Torp-Pedersen C, Carlsen JE, Bagger H, Eliasen P, Lyngborg K, Videbaek J, Cole DS, Auclert L & Pauly NC (1995). A clinical trial of the angiotensin-converting-enzyme inhibitor trandolapril in patients with left ventricular dysfunction after myocardial infarction. Trandolapril Cardiac Evaluation (TRACE) Study Group. N Engl J Med 333, 1670–1676.
- Konecny T, Kuniyoshi FH, Orban M, Pressman GS, Kara T,
 Gami A, Caples SM, Lopez-Jimenez F & Somers VK (2010).
 Under-diagnosis of sleep apnea in patients after acute myocardial infarction. *J Am Coll Cardiol* 56, 742–743.
- Kopp UC, Smith LA & DiBona GF (1985). Renorenal reflexes: neural components of ipsilateral and contralateral renal responses. *Am J Physiol Renal Physiol* **249**, F507–F517.
- Koroboki E, Zakopoulos N, Manios E, Rotas V, Papadimitriou G & Papageorgiou C (2010). Interoceptive awareness in essential hypertension. *Int J Psychophysiol* **78**, 158–162.
- Kostreva DR, Armour JA & Bosnjak ZJ (1985). Metabolic mapping of a cardiac reflex mediated by sympathetic ganglia in dogs. *Am J Physiol Regul Integr Comp Physiol* **249**, R317–R322.
- Krul SP, Driessen AH, van Boven WJ, Linnenbank AC, Geuzebroek GS, Jackman WM, Wilde AA, de Bakker JM & de Groot JR (2011). Thoracoscopic video-assisted pulmonary vein antrum isolation, ganglionated plexus ablation, and periprocedural confirmation of ablation lesions: first results of a hybrid surgical-electrophysiological approach for atrial fibrillation. *Circ Arrhythm Electrophysiol* 4, 262–270.
- Krum H, Schlaich M, Whitbourn R, Sobotka PA, Sadowski J, Bartus K, Kapelak B, Walton A, Sievert H, Thambar S, Abraham WT & Esler M (2009). Catheter-based renal sympathetic denervation for resistant hypertension: a multicentre safety and proof-of-principle cohort study. *Lancet* 373, 1275–1281.
- Kuck KH, Schaumann A, Eckardt L, Willems S, Ventura R, Delacretaz E, Pitschner HF, Kautzner J, Schumacher B, Hansen PS; VTACH study group (2010). Catheter ablation of stable ventricular tachycardia before defibrillator implantation in patients with coronary heart disease (VTACH): a multicentre randomised controlled trial. *Lancet* **375**, 31–40.
- Kumar R, Birrer BV, Macey PM, Woo MA, Gupta RK, Yan-Go FL & Harper RM (2008). Reduced mammillary body volume in patients with obstructive sleep apnea. *Neurosci Lett* **438**, 330–334.
- Kumar R, Chavez AS, Macey PM, Woo MA, Yan-Go FL & Harper RM (2012). Altered global and regional brain mean diffusivity in patients with obstructive sleep apnea. *J Neurosci Res* **90**, 2043–2052.
- Kumar R, Pham TT, Macey PM, Woo MA, Yan-Go FL & Harper RM (2014). Abnormal myelin and axonal integrity in recently diagnosed patients with obstructive sleep apnea. *Sleep* **37**, 723–732.
- Kumar R, Woo MA, Birrer BV, Macey PM, Fonarow GC, Hamilton MA & Harper RM (2009). Mammillary bodies and fornix fibers are injured in heart failure. *Neurobiol Dis* **33**, 236–242.

- Kumar R, Woo MA, Macey PM, Fonarow GC, Hamilton MA & Harper RM (2011). Brain axonal and myelin evaluation in heart failure. *J Neurol Sci* 307, 106–113.
- Kuniyoshi FH, Garcia-Touchard A, Gami AS, Romero-Corral A, van der Walt C, Pusalavidyasagar S, Kara T, Caples SM, Pressman GS, Vasquez EC, Lopez-Jimenez F & Somers VK (2008). Day-night variation of acute myocardial infarction in obstructive sleep apnea. *J Am Coll Cardiol* **52**, 343–346.
- Kuntz A & Morehouse A (1930). Thoracic sympathetic cardiac nerves in man: Their relation to cervical sympathetic ganglionectomy. *Arch Surg* 20, 607–613.
- La Rovere MT, Bigger JT Jr, Marcus FI, Mortara A & Schwartz PJ (1998). Baroreflex sensitivity and heart-rate variability in prediction of total cardiac mortality after myocardial infarction. ATRAMI (Autonomic Tone and Reflexes After Myocardial Infarction) Investigators. *Lancet* 351, 478–484.
- La Rovere MT, Specchia G, Mortara A & Schwartz PJ (1988). Baroreflex sensitivity, clinical correlates, and cardiovascular mortality among patients with a first myocardial infarction. A prospective study. *Circulation* **78**, 816–824.
- Lahiri MK, Kannankeril PJ & Goldberger JJ (2008). Assessment of autonomic function in cardiovascular disease: physiological basis and prognostic implications. *J Am Coll Cardiol* **51**, 1725–1733.
- Lakkireddy D, Atkins D, Pillarisetti J, Ryschon K, Bommana S, Drisko J, Vanga S & Dawn B (2013). Effect of yoga on arrhythmia burden, anxiety, depression, and quality of life in paroxysmal atrial fibrillation. *J Am Coll Cardiol* **61**, 1177–1182.
- Lampert R, Joska T, Burg MM, Batsford WP, McPherson CA & Jain D (2002). Emotional and physical precipitants of ventricular arrhythmia. *Circulation* **106**, 1800–1805.
- Lane RD, McRae K, Reiman EM, Chen K, Ahern GL & Thayer JF (2009). Neural correlates of heart rate variability during emotion. *Neuroimage* 44, 213–222.
- Laterza MC, de Matos LD, Trombetta IC, Braga AM, Roveda F, Alves MJ, Krieger EM, Negrao CE & Rondon MU (2007). Exercise training restores baroreflex sensitivity in never-treated hypertensive patients. *Hypertension* **49**, 1298–1306.
- Lemola K, Chartier D, Yeh YH, Dubuc M, Cartier R, Armour A, Ting M, Sakabe M, Shiroshita-Takeshita A, Comtois P & Nattel S (2008). Pulmonary vein region ablation in experimental vagal atrial fibrillation: role of pulmonary veins versus autonomic ganglia. *Circulation* 117, 470–477.
- Leor J, Poole WK & Kloner RA (1996). Sudden cardiac death triggered by an earthquake. *N Engl J Med* **334**, 413–419.
- Lepicovska V, Novak P & Nadeau R (1992). Time-frequency dynamics in neurally mediated syncope. *Clin Auton Res* **2**, 317–326.
- Levin MD, Lu MM, Petrenko NB, Hawkins BJ, Gupta TH, Lang D, Buckley PT, Jochems J, Liu F, Spurney CF, Yuan LJ, Jacobson JT, Brown CB, Huang L, Beermann F, Margulies KB, Madesh M, Eberwine JH, Epstein JA & Patel VV (2009). Melanocyte-like cells in the heart and pulmonary veins contribute to atrial arrhythmia triggers. *J Clin Invest* 119, 3420–3436.
- Levy MN & Zieske H (1969). Autonomic control of cardiac pacemaker activity and atrioventricular transmission. *J Appl Physiol* **27**, 465–470.

- Li D, Nikiforova N, Lu CJ, Wannop K, McMenamin M, Lee CW, Buckler KJ & Paterson DJ (2013). Targeted neuronal nitric oxide synthase transgene delivery into stellate neurons reverses impaired intracellular calcium transients in prehypertensive rats. *Hypertension* **61**, 202–207.
- Li D, Wang L, Lee CW, Dawson TA & Paterson DJ (2007). Noradrenergic cell specific gene transfer with neuronal nitric oxide synthase reduces cardiac sympathetic neurotransmission in hypertensive rats. *Hypertension* **50**, 69–74.
- Li M, Zheng C, Sato T, Kawada T, Sugimachi M & Sunagawa K (2004). Vagal nerve stimulation markedly improves long-term survival after chronic heart failure in rats. *Circulation* **109**, 120–124.
- Li S, Scherlag BJ, Yu L, Sheng X, Zhang Y, Ali R, Dong Y, Ghias M & Po SS (2009). Low-level vagosympathetic stimulation: a paradox and potential new modality for the treatment of focal atrial fibrillation. *Circ Arrhythm Electrophysiol* **2**, 645–651
- Liu L & Barajas L (1993). The rat renal nerves during development. *Anat Embryol* **188**, 345–361.
- Liu L & Nattel S (1997). Differing sympathetic and vagal effects on atrial fibrillation in dogs: role of refractoriness heterogeneity. Am J Physiol Heart Circ Physiol 273, H805–H816.
- Liu Z, Hesse C, Curry TB, Pike TL, Issa A, Bernal M,
 Charkoudian N, Joyner MJ & Eisenach JH (2009).
 Ambulatory arterial stiffness index is not correlated with the pressor response to laboratory stressors in normotensive humans. *J Hypertens* 27, 763–768.
- Lloyd R, Okada R, Stagg J, Anderson R, Hattler B & Marcus F (1974). The treatment of recurrent ventricular tachycardia with bilateral cervico-thoracic sympathetic-ganglionectomy. A report of two cases. *Circulation* **50**, 382–388.
- Low PA (2003). Testing the autonomic nervous system. *Semin Neurol* **23**, 407–421.
- Low PA (2008). Prevalence of orthostatic hypotension. *Clin Auton Res* **18** Suppl 1, 8–13.
- Lown B & DeSilva RA (1978). Roles of psychologic stress and autonomic nervous system changes in provocation of ventricular premature complexes. *Am J Cardiol* **41**, 979–985.
- Lown B, Temte JV, Reich P, Gaughan C, Regestein Q & Hal H (1976). Basis for recurring ventricular fibrillation in the absence of coronary heart disease and its management. *N Engl J Med* **294**, 623–629.
- Loyalka P, Hariharan R, Gholkar G, Gregoric ID, Tamerisa R, Nathan S & Kar B (2011). Left stellate ganglion block for continuous ventricular arrhythmias during percutaneous left ventricular assist device support. *Texas Heart Inst J* 38, 409–411.
- Lu CJ, Hao G, Nikiforova N, Larsen HE, Liu K, Crabtree MJ, Li D, Herring N & Paterson DJ (2015). CAPON modulates neuronal calcium handling and cardiac sympathetic neurotransmission during dysautonomia in hypertension. *Hypertension* **65**, 1288–1297.
- Ludwick-Rosenthal R & Neufeld RW (1985). Heart beat interoception: a study of individual differences. *Int J Psychophysiol* **3**, 57–65.

- Lundblad LC, Fatouleh RH, Hammam E, McKenzie DK, Macefield VG & Henderson LA (2014). Brainstem changes associated with increased muscle sympathetic drive in obstructive sleep apnoea. *Neuroimage* **103**, 258–266.
- Lutherer LO, Lutherer BC, Dormer KJ, Janssen HF & Barnes CD (1983). Bilateral lesions of the fastigial nucleus prevent the recovery of blood pressure following hypotension induced by hemorrhage or administration of endotoxin. *Brain Res* **269**, 251–257.
- Lutherer LO, Williams JL & Everse SJ (1989). Neurons of the rostral fastigial nucleus are responsive to cardiovascular and respiratory challenges. *J Auton Nerv Syst* 27, 101–111.
- Macey PM, Henderson LA, Macey KE, Alger JR, Frysinger RC, Woo MA, Harper RK, Yan-Go FL & Harper RM (2002). Brain morphology associated with obstructive sleep apnea. *Am J Respir Crit Care Med* **166**, 1382–1387.
- Macey PM, Kumar R, Ogren JA, Woo MA & Harper RM (2014). Global brain blood-oxygen level responses to autonomic challenges in obstructive sleep apnea. *PLoS One* **9**, e105261.
- Macey PM, Kumar R, Woo MA, Valladares EM, Yan-Go FL & Harper RM (2008). Brain structural changes in obstructive sleep apnea. *Sleep* **31**, 967–977.
- Macey PM, Kumar R, Woo MA, Yan-Go FL & Harper RM (2013). Heart rate responses to autonomic challenges in obstructive sleep apnea. *PLoS One* **8**, e76631.
- Mahajan A, Moore J, Cesario DA & Shivkumar K (2005). Use of thoracic epidural anesthesia for management of electrical storm: a case report. *Heart Rhythm* **2**, 1359–1362.
- Mahfoud F & Luscher TF (2015). Renal denervation: symply trapped by complexity? *Eur Heart J* **36**, 199–202.
- Malik M (1996). Heart rate variability: standards of measurement, physiological interpretation and clinical use. Task Force of the European Society of Cardiology and the North American Society of Pacing and Electrophysiology. *Circulation* **93**, 1043–1065.
- Malliani A, Recordati G & Schwartz PJ (1973). Nervous activity of afferent cardiac sympathetic fibres with atrial and ventricular endings. *J Physiol* **229**, 457–469.
- Mark AL, Victor RG, Nerhed C & Wallin BG (1985). Microneurographic studies of the mechanisms of sympathetic nerve responses to static exercise in humans. *Circ Res* 57, 461–469.
- Masani F (1986). Node-like cells in the myocardial layer of the pulmonary vein of rats: an ultrastructural study. *J Anat* **145**, 133–142
- Masson GS, Nair AR, Silva Soares PP, Michelini LC & Francis J (2015). Aerobic training normalizes autonomic dysfunction, HMGB1 content, microglia activation and inflammation in hypothalamic paraventricular nucleus of SHR. *Am J Physiol Heart Circ Physiol* **309**, H1115–H1122.
- Mathuria N, Tung R & Shivkumar K (2012). Advances in ablation of ventricular tachycardia in nonischemic cardiomyopathy. *Curr Cardiol Rep* **14**, 577–583.
- Matthews KA, Katholi CR, McCreath H, Whooley MA, Williams DR, Zhu S & Markovitz JH (2004). Blood pressure reactivity to psychological stress predicts hypertension in the CARDIA study. *Circulation* **110**, 74–78.

- Meissner A, Eckardt L, Kirchhof P, Weber T, Rolf N, Breithardt G, Van Aken H & Haverkamp W (2001). Effects of thoracic epidural anesthesia with and without autonomic nervous system blockade on cardiac monophasic action potentials and effective refractoriness in awake dogs. *Anesthesiology* **95**, 132–138.
- Mitchell GAG (1956). *Cardiovascular Innervation*. E. & S. Livingstone, Edinburgh.
- Miyauchi Y, Zhou S, Miyauchi M, Omichi C, Okuyama Y, Hamabe A, Hayashi H, Mandel WJ, Fishbein MC, Chen LS, Chen PS & Karagueuzian HS (2001). Induction of atrial sympathetic nerve sprouting and increased vulnerability to atrial fibrillation by chronic left ventricular myocardial infarction. *Circulation* **104**, II–17.
- Mizeres NJ (1963). The cardiac plexus in man. *Am J Anat* **112**, 141–151.
- Montgomery WA, Jones GE & Hollandsworth JG Jr (1984). The effects of physical fitness and exercise on cardiac awareness. *Biol Psychol* **18**, 11–22.
- Mooe T, Franklin KA, Wiklund U, Rabben T & Holmstrom K (2000). Sleep-disordered breathing and myocardial ischemia in patients with coronary artery disease. *Chest* 117, 1597–1602.
- Moss AJ & McDonald J (1971). Unilateral cervicothoracic sympathetic ganglionectomy for the treatment of long QT interval syndrome. *N Engl J Med* **285**, 903–904.
- Mousa TM, Liu D, Cornish KG & Zucker IH (2008). Exercise training enhances baroreflex sensitivity by an angiotensin II-dependent mechanism in chronic heart failure. *J Appl Physiol* **104**, 616–624.
- Murphy DA, O'Blenes S, Hanna BD & Armour JA (1994). Functional capacity of nicotine-sensitive canine intrinsic cardiac neurons to modify the heart. *Am J Physiol Regul Integr Comp Physiol* **266**, R1127–R1135.
- Murphy DA, Thompson GW, Ardell JL, McCraty R, Stevenson RS, Sangalang VE, Cardinal R, Wilkinson M, Craig S, Smith FM, Kingma JG & Armour JA (2000). The heart reinnervates after transplantation. *Ann Thoracic Surg* **69**, 1769–1781.
- Nademanee K, Taylor R, Bailey WE, Rieders DE & Kosar EM (2000). Treating electrical storm: sympathetic blockade versus advanced cardiac life support-guided therapy. *Circulation* **102**, 742–747.
- Narkiewicz K, van de Borne PJ, Montano N, Dyken ME, Phillips BG & Somers VK (1998). Contribution of tonic chemoreflex activation to sympathetic activity and blood pressure in patients with obstructive sleep apnea. *Circulation* **97**, 943–945.
- Nathan S & Bakris GL (2014). The future of renal denervation in resistant hypertension. *Curr Hypertens Rep* **16**, 494.
- Naughton MT (2012). Cheyne-Stokes respiration: friend or foe? *Thorax* **67**, 357–360.
- Nemeroff CB & Goldschmidt-Clermont PJ (2012). Heartache and heartbreak—the link between depression and cardiovascular disease. *Nat Rev Cardiol* **9**, 526–539.
- Ng GA (2014). Vagal modulation of cardiac ventricular arrhythmia. *Exp Physiol* **99**, 295–299.
- Nguyen BL, Fishbein MC, Chen LS, Chen PS & Masroor S (2009). Histopathological substrate for chronic atrial fibrillation in humans. *Heart Rhythm* **6**, 454–460.

- Nguyen BL, Li H, Fishbein MC, Lin SF, Gaudio C, Chen PS & Chen LS (2011). Acute myocardial infarction induces bilateral stellate ganglia neural remodeling in rabbits. *Cardiovasc Pathol* **21**, 143–148.
- Nitter-Hauge S & Storstein O (1973). Surgical treatment of recurrent ventricular tachycardia. *Br Heart J* **35**, 1132–1135.
- Nomura S & Mizuno N (1984). Central distribution of primary afferent fibers in the Arnold's nerve (the auricular branch of the vagus nerve): a transganglionic HRP study in the cat. *Brain Res* **292**, 199–205.
- Norris JE, Foreman RD & Wurster RK (1974). Responses of the canine heart to stimulation of the first five ventral thoracic roots. *Am J Physiol* **227**, 9–12.
- Norris JE, Lippincott D & Wurster RD (1977). Responses of canine endocardium to stimulation of the upper thoracic roots. *Am J Physiol Heart Circ Physiol* **233**, H655–H659.
- Nygard E, Kofoed KF, Freiberg J, Holm S, Aldershvile J, Eliasen K & Kelbaek H (2005). Effects of high thoracic epidural analgesia on myocardial blood flow in patients with ischemic heart disease. *Circulation* **111**, 2165–2170.
- O'Brien WH, Reid GJ & Jones KR (1998). Differences in heartbeat awareness among males with higher and lower levels of systolic blood pressure. *Int J Psychophysiol* **29**, 53–63.
- Ogren JA, Macey PM, Kumar R, Fonarow GC, Hamilton MA, Harper RM & Woo MA (2012). Impaired cerebellar and limbic responses to the valsalva maneuver in heart failure. *Cerebellum* 11, 931–938.
- Oh S, Choi EK, Zhang Y & Mazgalev TN (2011). Botulinum toxin injection in epicardial autonomic ganglia temporarily suppresses vagally mediated atrial fibrillation. *Circ Arrhythm Electrophysiol* **4**, 560–565.
- Oka T, Ozawa Y & Ohkubo Y (2001). Thoracic epidural bupivacaine attenuates supraventricular tachyarrhythmias after pulmonary resection. *Anesth Analg* **93**, 253–259.
- Olausson K, Magnusdottir H, Lurje L, Wennerblom B, Emanuelsson H & Ricksten SE (1997). Anti-ischemic and anti-anginal effects of thoracic epidural anesthesia versus those of conventional medical therapy in the treatment of severe refractory unstable angina pectoris. *Circulation* **96**, 2178–2182.
- Oldham JB (1950). Denervation of the kidney. *Ann R Coll Surg Engl* 7, 222–245.
- Olex S, Newberg A & Figueredo VM (2013). Meditation: should a cardiologist care? *Int J Cardiol* **168**, 1805–1810.
- O'Regan RG & Majcherczyk S (1982). Role of peripheral chemoreceptors and central chemosensitivity in the regulation of respiration and circulation. *J Exp Biol* **100**, 23–40.
- Ospina MB, Bond K, Karkhaneh M, Tjosvold L, Vandermeer B, Liang Y, Bialy L, Hooton N, Buscemi N, Dryden DM & Klassen TP (2007). Meditation practices for health: state of the research. *Evid Rep Technol Assess*, 1–263.
- Pae EK, Chien P & Harper RM (2005). Intermittent hypoxia damages cerebellar cortex and deep nuclei. *Neurosci Lett* **375**, 123–128.
- Pagani M, Montano N, Porta A, Malliani A, Abboud FM, Birkett C & Somers VK (1997). Relationship between spectral components of cardiovascular variabilities and direct measures of muscle sympathetic nerve activity in humans. *Circulation* **95**, 1441–1448.

- Palomares JA, Tummala S, Wang DJ, Park B, Woo MA, Kang DW, Lawrence KS, Harper RM & Kumar R (2015). Water exchange across the blood-brain barrier in obstructive sleep apnea: an MRI diffusion-weighted pseudo-continuous arterial spin labeling study. *J Neuroimaging* 25, 900–905.
- Pappone C, Santinelli V, Manguso F, Vicedomini G, Gugliotta F, Augello G, Mazzone P, Tortoriello V, Landoni G,
 Zangrillo A, Lang C, Tomita T, Mesas C, Mastella E & Alfieri O (2004). Pulmonary vein denervation enhances long-term benefit after circumferential ablation for paroxysmal atrial fibrillation. *Circulation* 109, 327–334.
- Patel RA, Priore DL, Szeto WY & Slevin KA (2011). Left stellate ganglion blockade for the management of drug-resistant electrical storm. *Pain Med* **12**, 1196–1198.
- Paulus MP & Stein MB (2010). Interoception in anxiety and depression. *Brain Struct Funct* **214**, 451–463.
- Pauza DH, Rysevaite-Kyguoliene K, Vismantaite J, Brack KE, Inokaitis H, Pauza AG, Rimasauskaite-Petraitiene V, Pauzaite JI & Pauziene N (2014). A combined acetylcholinesterase and immunohistochemical method for precise anatomical analysis of intrinsic cardiac neural structures. *Ann Anat* 196, 430–440.
- Peng CK, Henry IC, Mietus JE, Hausdorff JM, Khalsa G, Benson H & Goldberger AL (2004). Heart rate dynamics during three forms of meditation. *Int J Cardiol* **95**, 19–27.
- Perez-Lugones A, McMahon JT, Ratliff NB, Saliba WI, Schweikert RA, Marrouche NF, Saad EB, Navia JL, McCarthy PM, Tchou P, Gillinov AM & Natale A (2003). Evidence of specialized conduction cells in human pulmonary veins of patients with atrial fibrillation. *J Cardiovasc Electrophysiol* 14, 803–809.
- Peuker ET & Filler TJ (2002). The nerve supply of the human auricle. *Clin Anat* **15**, 35–37.
- Pfeffer JM & Pfeffer MA (1988). Angiotensin converting enzyme inhibition and ventricular remodeling in heart failure. *Am J Med* **84**, 37–44.
- Pfeiffer D, Fiehring H, Henkel HG, Rostock KJ & Rathgen K (1989). Long QT syndrome associated with inflammatory degeneration of the stellate ganglia. *Clin Cardiol* **12**, 222–224.
- Pohl R, Yeragani VK, Balon R, Rainey JM, Lycaki H, Ortiz A, Berchou R & Weinberg P (1988). Isoproterenol-induced panic attacks. *Biol Psychiatry* **24**, 891–902.
- Pokushalov E, Kozlov B, Romanov A, Strelnikov A, Bayramova S, Sergeevichev D, Bogachev-Prokophiev A, Zheleznev S, Shipulin V, Lomivorotov VV, Karaskov A, Po SS & Steinberg JS (2015). Long-term suppression of atrial fibrillation by botulinum toxin injection into epicardial fat pads in patients undergoing cardiac surgery: one-year follow-up of a randomized pilot study. *Circ Arrhythm Electrophysiol* **8**, 1334–1341.
- Polak T, Markulin F, Ehlis AC, Langer JB, Ringel TM & Fallgatter AJ (2009). Far field potentials from brain stem after transcutaneous vagus nerve stimulation: optimization of stimulation and recording parameters. *J Neural Transm* (*Vienna*) **116**, 1237–1242.

- Premchand RK, Sharma K, Mittal S, Monteiro R, Dixit S, Libbus I, DiCarlo LA, Ardell JL, Rector TS, Amurthur B, KenKnight BH & Anand IS (2014). Autonomic regulation therapy via left or right cervical vagus nerve stimulation in patients with chronic heart failure: results of the ANTHEM-HF trial. *J Cardiac Fail* **20**, 808–816.
- Premchand RK, Sharma K, Mittal S, Monteiro R, Dixit S, Libbus I, DiCarlo LA, Ardell JL, Rector TS, Amurthur B, KenKnight BH & Anand IS (2016). Extended follow-up of patients with heart failure receiving autonomic regulation therapy in the ANTHEM-HF study. *J Cardiac Fail* DOI: 10.1016/j.cardfail.2015.11.002.
- Priori SG, Wilde AA, Horie M, Cho Y, Behr ER, Berul C, Blom N, Brugada J, Chiang CE, Huikuri H, Kannankeril P, Krahn A, Leenhardt A, Moss A, Schwartz PJ, Shimizu W, Tomaselli G & Tracy C (2013). HRS/EHRA/APHRS expert consensus statement on the diagnosis and management of patients with inherited primary arrhythmia syndromes: document endorsed by HRS, EHRA, and APHRS in May 2013 and by ACCF, AHA, PACES, and AEPC in June 2013. *Heart Rhythm* 10, 1932–1963.
- Rajan P & Greenberg H (2015). Obstructive sleep apnea as a risk factor for type 2 diabetes mellitus. *Nat Sci Sleep* **7**, 113–125.
- Rajendran PS, Nakamura K, Ajijola OA, Vaseghi M, Armour JA, Ardell JL & Shivkumar K (2016). Myocardial infarction induces structural and functional remodelling of the intrinsic cardiac nervous system. *J Physiol* **594**, 321–341.
- Randall WC, Armour JA, Geis WP & Lippincott DB (1972). Regional cardiac distribution of the sympathetic nerves. *Fed Proc* **31**, 1199–1208.
- Rassovsky Y & Kushner MG (2003). Carbon dioxide in the study of panic disorder: issues of definition, methodology, and outcome. *J Anxiety Disord* 17, 1–32.
- Re RN (2004). Mechanisms of disease: local renin-angiotensin-aldosterone systems and the pathogenesis and treatment of cardiovascular disease. *Nat Clin Pract Cardiovasc Med* 1, 42–47.
- Ring C & Brener J (1992). The temporal locations of heartbeat sensations. *Psychophysiology* **29**, 535–545.
- Ring C, Brener J, Knapp K & Mailloux J (2015). Effects of heartbeat feedback on beliefs about heart rate and heartbeat counting: a cautionary tale about interoceptive awareness. *Biol Psychol* **104**, 193–198.
- Rizzo S, Basso C, Troost D, Aronica E, Frigo AC, Driessen AH, Thiene G, Wilde AA & van der Wal AC (2014). T-cell-mediated inflammatory activity in the stellate ganglia of patients with ion-channel disease and severe ventricular arrhythmias. *Circ Arrhythm Electrophysiol* 7, 224–229.
- Roddie IC (1977). Human responses to emotional stress. *Ir J Med Sci* **146**, 395–417.
- Rodrigues B, Jorge L, Mostarda CT, Rosa KT, Medeiros A, Malfitano C, de Souza AL Jr, Viegas KA, Lacchini S, Curi R, Brum PC, De Angelis K & Irigoyen MC (2012). Aerobic exercise training delays cardiac dysfunction and improves autonomic control of circulation in diabetic rats undergoing myocardial infarction. *J Card Fail* **18**, 734–744.

- Rodrigues F, Feriani DJ, Barboza CA, Abssamra ME, Rocha LY, Carrozi NM, Mostarda C, Figueroa D, Souza GI, De Angelis K, Irigoyen MC & Rodrigues B (2014). Cardioprotection afforded by exercise training prior to myocardial infarction is associated with autonomic function improvement. *BMC Cardiovasc Disord* 14, 84.
- Roston TM, Vinocur JM, Maginot KR, Mohammed S, Salerno JC, Etheridge SP, Cohen M, Hamilton RM, Pflaumer A, Kanter RJ, Potts JE, LaPage MJ, Collins KK, Gebauer RA, Temple JD, Batra AS, Erickson C, Miszczak-Knecht M, Kubus P, Bar-Cohen Y, Kantoch M, Thomas VC, Hessling G, Anderson C, Young ML, Cabrera Ortega M, Lau YR, Johnsrude CL, Fournier A, Kannankeril PJ & Sanatani S (2015). Catecholaminergic polymorphic ventricular tachycardia in children: analysis of therapeutic strategies and outcomes from an international multicenter registry. *Circ Arrhythm Electrophysiol* 8, 633–642.
- Rouse CH, Jones GE & Jones KR (1988). The effect of body composition and gender on cardiac awareness. *Psychophysiology* **25**, 400–407.
- Rowan RA & Billingham ME (1988). Myocardial innervation in long-term heart transplant survivors: a quantitative ultrastructural survey. *J Heart Transplant* 7, 448–452.
- Rudas L, Crossman AA, Morillo CA, Halliwill JR, Tahvanainen KU, Kuusela TA & Eckberg DL (1999). Human sympathetic and vagal baroreflex responses to sequential nitroprusside and phenylephrine. *Am J Physiol Heart Circ Physiol* **276**, H1691–H1698.
- Sa JC, Costa EC, da Silva E, Tamburus NY, Porta A, Medeiros LF, Lemos TM, Soares EM & Azevedo GD (2016). Aerobic exercise improves cardiac autonomic modulation in women with polycystic ovary syndrome. *Int J Cardiol* **202**, 356–361.
- Saccomanno G (1943). The components of the upper thoracic sympathetic nerves. *J Comp Neurol* **79**, 355–378.
- Sakakura K, Ladich E, Cheng Q, Otsuka F, Yahagi K, Fowler DR, Kolodgie FD, Virmani R & Joner M (2014). Anatomic assessment of sympathetic peri-arterial renal nerves in man. *J Am Coll Cardiol* **64**, 635–643.
- Schachter SC (2002). Vagus nerve stimulation therapy summary: five years after FDA approval. *Neurology* **59**, S15–S20.
- Schandry R (1981). Heart beat perception and emotional experience. *Psychophysiology* **18**, 483–488.
- Schandry R, Bestler M & Montoya P (1993). On the relation between cardiodynamics and heartbeat perception. *Psychophysiology* **30**, 467–474.
- Schauerte P, Scherlag BJ, Pitha J, Scherlag MA, Reynolds D, Lazzara R & Jackman WM (2000). Catheter ablation of cardiac autonomic nerves for prevention of vagal atrial fibrillation. *Circulation* **102**, 2774–2780.
- Schmidt G, Malik M, Barthel P, Schneider R, Ulm K, Rolnitzky L, Camm AJ, Bigger JT Jr & Schomig A (1999). Heart-rate turbulence after ventricular premature beats as a predictor of mortality after acute myocardial infarction. *Lancet* **353**, 1390–1396.
- Schneider TR, Ring C & Katkin ES (1998). A test of the validity of the method of constant stimuli as an index of heartbeat detection. *Psychophysiology* **35**, 86–89.

- Schoonmaker FW, Carey T & Grow JB Sr (1975). Treatment of tachyarrhythmias and bradyarrhythmias by cardiac sympathectomy and permanent ventricular pacing. *Ann Thoracic Surg* **19**, 80–87.
- Schroeder S, Gerlach AL, Achenbach S & Martin A (2015). The relevance of accuracy of heartbeat perception in noncardiac and cardiac chest pain. *Int J Behav Med* **22**, 258–267.
- Schunck T, Erb G, Mathis A, Gilles C, Namer IJ, Hode Y, Demaziere A, Luthringer R & Macher JP (2006). Functional magnetic resonance imaging characterization of CCK-4-induced panic attack and subsequent anticipatory anxiety. *Neuroimage* **31**, 1197–1208.
- Schwaiblmair M, von Scheidt W, Uberfuhr P, Ziegler S, Schwaiger M, Reichart B & Vogelmeier C (1999). Functional significance of cardiac reinnervation in heart transplant recipients. *J Heart Lung Transplant* 18, 838–845.
- Schwartz PJ (2014). Cardiac sympathetic denervation to prevent life-threatening arrhythmias. Nat Rev Cardiol 11, 346–353.
- Schwartz PJ (2015). When the risk is sudden death, does quality of life matter? *Heart Rhythm* **13**, 70–71.
- Schwartz PJ & Ackerman MJ (2013). The long QT syndrome: a transatlantic clinical approach to diagnosis and therapy. *Eur Heart J* **34**, 3109–3116.
- Schwartz PJ, De Ferrari GM, Sanzo A, Landolina M, Rordorf R, Raineri C, Campana C, Revera M, Ajmone-Marsan N, Tavazzi L & Odero A (2008). Long term vagal stimulation in patients with advanced heart failure: first experience in man. *Eur J Heart Fail* **10**, 884–891.
- Schwartz PJ, Periti M & Malliani A (1975). The long Q-T syndrome. *Am Heart J* **89**, 378–390.
- Schwartz PJ, Priori SG, Cerrone M, Spazzolini C, Odero A, Napolitano C, Bloise R, De Ferrari GM, Klersy C, Moss AJ, Zareba W, Robinson JL, Hall WJ, Brink PA, Toivonen L, Epstein AE, Li C & Hu D (2004). Left cardiac sympathetic denervation in the management of high-risk patients affected by the long-QT syndrome. *Circulation* **109**, 1826–1833.
- Schwenk TL (2008). Cardiovascular events during World Cup soccer. *N Engl J Med* **358**, 2408.
- Seth AK & Critchley HD (2013). Extending predictive processing to the body: Emotion as interoceptive inference. *Behav Brain Sci* **36**, 227–228.
- Sha Y, Scherlag BJ, Yu L, Sheng X, Jackman WM, Lazzara R & Po SS (2011). Low-level right vagal stimulation: anticholinergic and antiadrenergic effects. *J Cardiovasc Electrophysiol* **22**, 1147–1153.
- Shen MJ, Hao-Che C, Park HW, George Akingba A, Chang PC, Zheng Z, Lin SF, Shen C, Chen LS, Chen Z, Fishbein MC, Chiamvimonvat N & Chen PS (2013). Low-level vagus nerve stimulation upregulates small conductance calcium-activated potassium channels in the stellate ganglion. *Heart Rhythm* **10**, 910–915.
- Shen MJ, Shinohara T, Park HW, Frick K, Ice DS, Choi EK, Han S, Maruyama M, Sharma R, Shen C, Fishbein MC, Chen LS, Lopshire JC, Zipes DP, Lin SF & Chen PS (2011). Continuous low-level vagus nerve stimulation reduces stellate ganglion nerve activity and paroxysmal atrial tachyarrhythmias in ambulatory canines. *Circulation* 123, 2204–2212.

- Shen MJ & Zipes DP (2014). Role of the autonomic nervous system in modulating cardiac arrhythmias. *Circ Res* **114**, 1004–1021.
- Sheng X, Scherlag BJ, Yu L, Li S, Ali R, Zhang Y, Fu G, Nakagawa H, Jackman WM, Lazzara R & Po SS (2011). Prevention and reversal of atrial fibrillation inducibility and autonomic remodeling by low-level vagosympathetic nerve stimulation. *J Am Coll Cardiol* 57, 563–571.
- Sherrington C (1961). The Integrative Action of the Nervous System. Yale University Press, New Haven.
- Sica AL, Gootman PM, Gootman N & Armour JA (1994). Neuronal activity of the stellate ganglia in neonatal swine. *J Auton Nerv Syst* 48, 273–277.
- Smeets JL, Allessie MA, Lammers WJ, Bonke FI & Hollen J (1986). The wavelength of the cardiac impulse and reentrant arrhythmias in isolated rabbit atrium. The role of heart rate, autonomic transmitters, temperature, and potassium. *Circ Res* **58**, 96–108.
- Smyth HS, Sleight P & Pickering GW (1969). Reflex regulation of arterial pressure during sleep in man. A quantitative method of assessing baroreflex sensitivity. *Circ Res* **24**, 109–121.
- Somers VK, Dyken ME, Clary MP & Abboud FM (1995). Sympathetic neural mechanisms in obstructive sleep apnea. *J Clin Invest* **96**, 1897–1904.
- Somers VK, Dyken ME, Mark AL & Abboud FM (1992). Parasympathetic hyperresponsiveness and bradyarrhythmias during apnoea in hypertension. *Clin Auton Res* **2**, 171–176.
- Somers VK, Mark AL & Abboud FM (1988). Potentiation of sympathetic nerve responses to hypoxia in borderline hypertensive subjects. *Hypertension* 11, 608–612.
- Somers VK, Mark AL & Abboud FM (1991). Interaction of baroreceptor and chemoreceptor reflex control of sympathetic nerve activity in normal humans. *J Clin Invest* **87**, 1953–1957.
- Somers VK, Mark AL, Zavala DC & Abboud FM (1989). Contrasting effects of hypoxia and hypercapnia on ventilation and sympathetic activity in humans. *J Appl Physiol* **67**, 2101–2106.
- Somers VK, White DP, Amin R, Abraham WT, Costa F, Culebras A, Daniels S, Floras JS, Hunt CE, Olson LJ, Pickering TG, Russell R, Woo M & Young T (2008). Sleep apnea and cardiovascular disease: an American Heart Association/American College Of Cardiology Foundation Scientific Statement from the American Heart Association Council for High Blood Pressure Research Professional Education Committee, Council on Clinical Cardiology, Stroke Council, and Council On Cardiovascular Nursing. In collaboration with the National Heart, Lung, and Blood Institute National Center on Sleep Disorders Research (National Institutes of Health). *Circulation* 118, 1080–1111.
- Soukhova-O'Hare GK, Cheng ZJ, Roberts AM & Gozal D (2006). Postnatal intermittent hypoxia alters baroreflex function in adult rats. *Am J Physiol Heart Circ Physiol* **290**, H1157–1164.
- Spierer DK, DeMeersman RE, Kleinfeld J, McPherson E, Fullilove RE, Alba A & Zion AS (2007). Exercise training improves cardiovascular and autonomic profiles in HIV. Clin Auton Res 17, 341–348.

- Spuck S, Tronnier V, Orosz I, Schonweiler R, Sepehrnia A, Nowak G & Sperner J (2010). Operative and technical complications of vagus nerve stimulator implantation. *Neurosurgery* **67**, 489–494.
- Starling EH (1908). The chemical control of the body. *JAMA* L, 835–840.
- Stavrakis S, Humphrey MB, Scherlag BJ, Hu Y, Jackman WM, Nakagawa H, Lockwood D, Lazzara R & Po SS (2015). Low-level transcutaneous electrical vagus nerve stimulation suppresses atrial fibrillation. *J Am Coll Cardiol* **65**, 867–875.
- Stavrakis S, Scherlag BJ, Fan Y, Liu Y, Liu Q, Mao J, Cai H, Lazzara R & Po SS (2012). Antiarrhythmic effects of vasostatin-1 in a canine model of atrial fibrillation. *J Cardiovasc Electrophysiol* **23**, 771–777.
- Steinbeck G, Andresen D, Seidl K, Brachmann J, Hoffmann E, Wojciechowski D, Kornacewicz-Jach Z, Sredniawa B, Lupkovics G, Hofgartner F, Lubinski A, Rosenqvist M, Habets A, Wegscheider K & Senges J (2009). Defibrillator implantation early after myocardial infarction. N Engl J Med 361, 1427–1436.
- Steinberg JS, Pokushalov E & Mittal S (2013). Renal denervation for arrhythmias: hope or hype? *Curr Cardiol Rep* **15**, 392.
- Stella A & Zanchetti A (1991). Functional role of renal afferents. *Physiol Rev* **71**, 659–682.
- Taggart P, Boyett MR, Logantha S & Lambiase PD (2011). Anger, emotion, and arrhythmias: from brain to heart. *Front Physiol* **2**, 67.
- Tai CT, Chiou CW, Wen ZC, Hsieh MH, Tsai CF, Lin WS, Chen CC, Lin YK, Yu WC, Ding YA, Chang MS & Chen SA (2000). Effect of phenylephrine on focal atrial fibrillation originating in the pulmonary veins and superior vena cava. *J Am Coll Cardiol* **36**, 788–793.
- Tan AY, Li H, Wachsmann-Hogiu S, Chen LS, Chen PS & Fishbein MC (2006). Autonomic innervation and segmental muscular disconnections at the human pulmonary veinatrial junction: implications for catheter ablation of atrial-pulmonary vein junction. *J Am Coll Cardiol* **48**, 132–143.
- Taylor JA, Halliwill JR, Brown TE, Hayano J & Eckberg DL (1995). 'Non-hypotensive' hypovolaemia reduces ascending aortic dimensions in humans. *J Physiol* **483**, 289–298.
- Telles S, Nagarathna R & Nagendra HR (1995). Autonomic changes during "OM" meditation. *Indian J Physiol Pharmacol* **39**, 418–420.
- Templin C, Ghadri JR, Diekmann J, Napp LC, Bataiosu DR, Jaguszewski M, Cammann VL, Sarcon A, Geyer V, Neumann CA, Seifert B, Hellermann J, Schwyzer M, Eisenhardt K, Jenewein J, Franke J, Katus HA, Burgdorf C, Schunkert H, Moeller C, Thiele H, Bauersachs J, Tschope C, Schultheiss HP, Laney CA, Rajan L, Michels G, Pfister R, Ukena C, Bohm M, Erbel R, Cuneo A, Kuck KH, Jacobshagen C, Hasenfuss G, Karakas M, Koenig W, Rottbauer W, Said SM, Braun-Dullaeus RC, Cuculi F, Banning A, Fischer TA, Vasankari T, Airaksinen KE, Fijalkowski M, Rynkiewicz A, Pawlak M, Opolski G, Dworakowski R, MacCarthy P, Kaiser C, Osswald S, Galiuto L, Crea F, Dichtl W, Franz WM, Empen K, Felix SB, Delmas C, Lairez O, Erne P, Bax JJ, Ford I, Ruschitzka F, Prasad A & Luscher TF (2015). Clinical features and outcomes of takotsubo (stress) cardiomyopathy. N Engl J Med 373, 929-938.

- ten Harkel AD, van Lieshout JJ, Karemaker JM & Wieling W (1993). Differences in circulatory control in normal subjects who faint and who do not faint during orthostatic stress. *Clin Auton Res* **3**, 117–124.
- Thalmann M, Trampitsch E, Haberfellner N, Eisendle E, Kraschl R & Kobinia G (2001). Resuscitation in near drowning with extracorporeal membrane oxygenation. *The Ann Thorac Surg* **72**, 607–608.
- Thayer JF & Lane RD (2000). A model of neurovisceral integration in emotion regulation and dysregulation. *J Affect Disord* **61**, 201–216.
- Thayer JF & Lane RD (2009). Claude Bernard and the heart-brain connection: further elaboration of a model of neurovisceral integration. *Neurosci Biobehav Rev* **33**, 81–88.
- Thoren PN (1977). Characteristics of left ventricular receptors with nonmedullated vagal afferents in cats. *Circ Res* **40**, 415–421.
- Thurston RC, Rewak M & Kubzansky LD (2013). An anxious heart: anxiety and the onset of cardiovascular diseases. *Prog Cardiovasc Dis* **55**, 524–537.
- Tomney PA, Hopkins DA & Armour JA (1985). Axonal branching of canine sympathetic postganglionic cardiopulmonary neurons. A retrograde fluorescent labeling study. *Brain Res Bull* 14, 443–452.
- Trzebski A, Tafil M, Zoltowski M & Przybylski J (1982). Increased sensitivity of the arterial chemoreceptor drive in young men with mild hypertension. *Cardiovasc Res* **16**, 163–172.
- Tsai CF, Chen SA, Tai CT, Chiou CW, Prakash VS, Yu WC, Hsieh MH, Ding YA & Chang MS (1999). Bezold-Jarisch-like reflex during radiofrequency ablation of the pulmonary vein tissues in patients with paroxysmal focal atrial fibrillation. *J Cardiovasc Electrophysiol* **10**, 27–35.
- Tzafriri AR, Keating JH, Markham PM, Spognardi AM, Stanley JR, Wong G, Zani BG, Highsmith D, O'Fallon P, Fuimaono K, Mahfoud F & Edelman ER (2015). Arterial microanatomy determines the success of energy-based renal denervation in controlling hypertension. *Sci Transl Med* 7, 285ra265.
- Valderrabano M, Chen HR, Sidhu J, Rao L, Ling Y & Khoury DS (2009). Retrograde ethanol infusion in the vein of Marshall: regional left atrial ablation, vagal denervation and feasibility in humans. *Circ Arrhythm Electrophysiol* **2**, 50–56.
- Vance WH & Bowker RC (1983). Spinal origins of cardiac afferents from the region of the left anterior descending artery. *Brain Res* **258**, 96–100.
- Van der Does AJW, Antony MM, Ehlers A & Barsky AJ (2000). Heartbeat perception in panic disorder: a reanalysis. *Behav Res Ther* **38**, 47–62.
- Vanoli E, De Ferrari GM, Stramba-Badiale M, Hull SS Jr, Foreman RD & Schwartz PJ (1991). Vagal stimulation and prevention of sudden death in conscious dogs with a healed myocardial infarction. *Circ Res* **68**, 1471–1481.
- Vaseghi M, Gima J, Kanaan C, Ajijola OA, Marmureanu A, Mahajan A & Shivkumar K (2014). Cardiac sympathetic denervation in patients with refractory ventricular arrhythmias or electrical storm: intermediate and long-term follow-up. *Heart Rhythm* 11, 360–366.
- Vaseghi M, Lellouche N, Ritter H, Fonarow GC, Patel JK, Moriguchi J, Fishbein MC, Kobashigawa JA & Shivkumar K (2009). Mode and mechanisms of death after orthotopic heart transplantation. *Heart Rhythm* **6**, 503–509.

- Vaseghi M, Lux RL, Mahajan A & Shivkumar K (2012). Sympathetic stimulation increases dispersion of repolarization in humans with myocardial infarction. Am J Physiol Heart Circ Physiol 302, H1838–H1846.
- Vaseghi M & Shivkumar K (2008). The role of the autonomic nervous system in sudden cardiac death. *Prog Cardiovasc Dis* **50**, 404–419.
- Vracko R, Thorning D & Frederickson RG (1991). Nerve fibers in human myocardial scars. *Hum Pathol* **22**, 138–146.
- Wang L, Henrich M, Buckler KJ, McMenamin M, Mee CJ, Sattelle DB & Paterson DJ (2007). Neuronal nitric oxide synthase gene transfer decreases [Ca²⁺]_i in cardiac sympathetic neurons. *J Mol Cell Cardiol* **43**, 717–725.
- Wehrwein EA, Joyner MJ, Hart EC, Wallin BG, Karlsson T & Charkoudian N (2010). Blood pressure regulation in humans: calculation of an "error signal" in control of sympathetic nerve activity. *Hypertension* **55**, 264–269.
- Wellens HJ, Schwartz PJ, Lindemans FW, Buxton AE, Goldberger JJ, Hohnloser SH, Huikuri HV, Kaab S, La Rovere MT, Malik M, Myerburg RJ, Simoons ML, Swedberg K, Tijssen J, Voors AA & Wilde AA (2014). Risk stratification for sudden cardiac death: current status and challenges for the future. *Eur Heart J* 35, 1642–1651.
- Whitehead WE, Drescher VM & Heiman P (1977). Relation of heart rate control to heartbeat perception. *Biofeedback Self Regul* **2**, 371–392.
- Wilbert-Lampen U, Leistner D, Greven S, Pohl T, Sper S, Volker C, Guthlin D, Plasse A, Knez A, Kuchenhoff H & Steinbeck G (2008). Cardiovascular events during World Cup soccer. *N Engl J Med* **358**, 475–483.
- Wilbert-Lampen U, Nickel T, Leistner D, Guthlin D, Matis T, Volker C, Sper S, Kuchenhoff H, Kaab S & Steinbeck G (2010). Modified serum profiles of inflammatory and vasoconstrictive factors in patients with emotional stress-induced acute coronary syndrome during World Cup Soccer 2006. *J Am Coll Cardiol* 55, 637–642.
- Wilde AA, Bhuiyan ZA, Crotti L, Facchini M, De Ferrari GM, Paul T, Ferrandi C, Koolbergen DR, Odero A & Schwartz PJ (2008). Left cardiac sympathetic denervation for catecholaminergic polymorphic ventricular tachycardia. *N Engl J Med* **358**, 2024–2029.
- Wilhelm M (2014). Atrial fibrillation in endurance athletes. *Eur J Prev Cardiol* **21**, 1040–1048.
- Windmann S, Schonecke OW, Frohlig G & Maldener G (1999). Dissociating beliefs about heart rates and actual heart rates in patients with cardiac pacemakers. *Psychophysiology* **36**, 339–342.
- Wittstein IS, Thiemann DR, Lima JA, Baughman KL, Schulman SP, Gerstenblith G, Wu KC, Rade JJ, Bivalacqua TJ & Champion HC (2005). Neurohumoral features of myocardial stunning due to sudden emotional stress. *N Engl J Med* **352**, 539–548.
- Woo MA, Kumar R, Macey PM, Fonarow GC & Harper RM (2009). Brain injury in autonomic, emotional, and cognitive regulatory areas in patients with heart failure. *J Card Fail* 15, 214–223.
- Woo MA, Macey PM, Fonarow GC, Hamilton MA & Harper RM (2003). Regional brain gray matter loss in heart failure. *J Appl Physiol* **95**, 677–684.

- Woo MA, Macey PM, Keens PT, Kumar R, Fonarow GC, Hamilton MA & Harper RM (2005). Functional abnormalities in brain areas that mediate autonomic nervous system control in advanced heart failure. *J Card Fail* 11, 437–446.
- Woo MA, Macey PM, Keens PT, Kumar R, Fonarow GC, Hamilton MA & Harper RM (2007). Aberrant central nervous system responses to the Valsalva maneuver in heart failure. *Congest Heart Fail* **13**, 29–35.
- Woo MA, Palomares JA, Macey PM, Fonarow GC, Harper RM & Kumar R (2015). Global and regional brain mean diffusivity changes in patients with heart failure. *J Neurosci Res* **93**, 678–685.
- Wood A, Docimo S & Elkowitz DE (2010). Cardiovascular disease and its association with histological changes of the left stellate ganglion. *Clin Med Insights Pathol* **3**, 19–24.
- Yamakawa K, Rajendran PS, Takamiya T, Yagishita D, So EL, Mahajan A, Shivkumar K & Vaseghi M (2015). Vagal nerve stimulation activates vagal afferent fibers that reduce cardiac efferent parasympathetic effects. *Am J Physiol Heart Circ Physiol* **309**, H1579–1590.
- Yancy CW, Jessup M, Bozkurt B, Butler J, Casey DE Jr,
 Drazner MH, Fonarow GC, Geraci SA, Horwich T, Januzzi
 JL, Johnson MR, Kasper EK, Levy WC, Masoudi FA,
 McBride PE, McMurray JJ, Mitchell JE, Peterson PN, Riegel
 B, Sam F, Stevenson LW, Tang WH, Tsai EJ & Wilkoff BL
 (2013). 2013 ACCF/AHA guideline for the management of
 heart failure: a report of the American College of Cardiology
 Foundation/American Heart Association Task Force on
 practice guidelines. *Circulation* 128, e240–e327.
- Ye S, Zhong H, Yanamadala V & Campese VM (2002). Renal injury caused by intrarenal injection of phenol increases afferent and efferent renal sympathetic nerve activity. *Am J Hypertens* **15**, 717–724.
- Yu L, Scherlag BJ, Li S, Fan Y, Dyer J, Male S, Varma V, Sha Y, Stavrakis S & Po SS (2013). Low-level transcutaneous electrical stimulation of the auricular branch of the vagus nerve: a noninvasive approach to treat the initial phase of atrial fibrillation. *Heart Rhythm* 10, 428–435.
- Yuan BX, Ardell JL, Hopkins DA, Losier AM & Armour JA (1994). Gross and microscopic anatomy of the canine intrinsic cardiac nervous system. *Anat Rec* **239**, 75–87.
- Zamir M, Kimmerly DS & Shoemaker JK (2012). Cardiac mechanoreceptor function implicated during premature ventricular contraction. *Auton Neurosci* **167**, 50–55.
- Zannad F, De Ferrari GM, Tuinenburg AE, Wright D, Brugada J, Butter C, Klein H, Neuzil P, Botman C, Castel MA, D'Onofrio A, De Borst GJ, Solomon S, Stein KM, Meyer S, Schubert B, Stalsberg K, Wold N, Ruble S & Stolen C (2015a). Long term safety and efficacy results of the NEural Cardiac TherApy foR Heart Failure (NECTAR-HF) trial. *J Cardiac Fail* 21, 936.
- Zannad F, De Ferrari GM, Tuinenburg AE, Wright D, Brugada J, Butter C, Klein H, Stolen C, Meyer S, Stein KM, Ramuzat A, Schubert B, Daum D, Neuzil P, Botman C, Castel MA, D'Onofrio A, Solomon SD, Wold N & Ruble SB (2015b). Chronic vagal stimulation for the treatment of low ejection fraction heart failure: results of the NEural Cardiac TherApy foR Heart Failure (NECTAR-HF) randomized controlled trial. *Eur Heart J* 36, 425–433.

- Zhang Y, Popovic ZB, Bibevski S, Fakhry I, Sica DA, Van Wagoner DR & Mazgalev TN (2009). Chronic vagus nerve stimulation improves autonomic control and attenuates systemic inflammation and heart failure progression in a canine high-rate pacing model. *Circ Heart Fail* **2**, 692–699.
- Zheng H, Sharma NM, Liu X & Patel KP (2012). Exercise training normalizes enhanced sympathetic activation from the paraventricular nucleus in chronic heart failure: role of angiotensin II. *Am J Physiol Regul Integr Comp Physiol* **303**, R387–R394.
- Zhou S, Chen LS, Miyauchi Y, Miyauchi M, Kar S, Kangavari S, Fishbein MC, Sharifi B & Chen PS (2004). Mechanisms of cardiac nerve sprouting after myocardial infarction in dogs. *Circ Res* **95**, 76–83.
- Ziegelstein RC (2007). Acute emotional stress and cardiac arrhythmias. *JAMA* **298**, 324–329.
- Zipes DP, Festoff B, Schaal SF, Cox C, Sealy WC & Wallace AG (1968). Treatment of ventricular arrhythmia by permanent atrial pacemaker and cardiac sympathectomy. *Ann Intern Med* **68**, 591–597.
- Zipes DP, Mihalick MJ & Robbins GT (1974). Effects of selective vagal and stellate ganglion stimulation of atrial refractoriness. *Cardiovasc Res* **8**, 647–655.
- Zipes DP, Neuzil P, Theres H, Caraway D, Mann DL, Mannheimer C, Van Buren P, Linde C, Linderoth B, Kueffer F, Sarazin SA & DeJongste MJ (2016). Determining the feasibility of spinal cord neuromodulation for the treatment of chronic systolic heart failure: The DEFEAT-HF study. *JACC Heart Fail* 4, 129–136.
- Zoellner LA & Craske MG (1999). Interoceptive accuracy and panic. *Behav Res Ther* **37**, 1141–1158.

Additional information

Competing interests

None declared.

Author contributions

All authors have approved the final version of the manuscript and agree to be accountable for all aspects of the work. All persons designated as authors qualify for authorship, and all those who qualify for authorship are listed.

Acknowledgements

This work was supported by National Institutes of Health grants (HL084261 to K.S.; HL125730 to O.A.A.; HL083947 to M.J.J., HL71140 and HL78931 to P.-S.C.; NS090407, HL113251, NR015038 and NR013625 to R.H. and R.K.; HL65176, HL114024 and HL114676 to V.K.S.; and HL132356 to M.V.); Strategic Research Initiative of the Indiana University Health/Indiana University School of Medicine (P.-S.C.); National Health and Medical Research Council (1081143, 1036352 and 1070386 to M.E.).