

Circulation Research Compendium on Sudden Cardiac Death

The Spectrum of Epidemiology Underlying Sudden Cardiac Death

Sudden Cardiac Death Risk Stratification

Genetics of Sudden Cardiac Death

Mechanisms of Sudden Cardiac Death: Oxidants and Metabolism

Role of Sodium and Calcium Dysregulation in Tachyarrhythmias in Sudden Cardiac Death

Ion Channel Macromolecular Complexes in Cardiomyocytes: Roles in Sudden Cardiac Death

Finding the Rhythm of Sudden Cardiac Death: New Opportunities Using Induced Pluripotent Stem Cell–Derived Cardiomyocytes

Cardiac Innervation and Sudden Cardiac Death

Clinical Management and Prevention of Sudden Cardiac Death

Cardiac Arrest: Resuscitation and Reperfusion

Gordon Tomaselli, Editor

Role of Sodium and Calcium Dysregulation in Tachyarrhythmias in Sudden Cardiac Death

Stefan Wagner, Lars S. Maier, Donald M. Bers

Abstract: Despite improvements in the therapy of underlying heart disease, sudden cardiac death is a major cause of death worldwide. Disturbed Na and Ca handling is known to be a major predisposing factor for life-threatening tachyarrhythmias. In cardiomyocytes, many ion channels and transporters, including voltage-gated Na and Ca channels, cardiac ryanodine receptors, Na/Ca-exchanger, and SR Ca-ATPase are involved in this regulation. We have learned a lot about the pathophysiological relevance of disturbed ion channel function from monogenetic disorders. Changes in the gating of a single ion channel and the activity of an ion pump suffice to dramatically increase the propensity for arrhythmias even in structurally normal hearts. Nevertheless, patients with heart failure with acquired dysfunction in many ion channels and transporters exhibit profound dysregulation of Na and Ca handling and Ca/calmodulin-dependent protein kinase and are especially prone to arrhythmias. A deeper understanding of the underlying arrhythmic principles is mandatory if we are to improve their outcome. This review addresses basic tachyarrhythmic mechanisms, the underlying ionic mechanisms and the consequences for ion homeostasis, and the situation in complex diseases like heart failure. (*Circ Res.* 2015;116:1956-1970. DOI: 10.1161/CIRCRESAHA.116.304678.)

Key Words: alternans ■ arrhythmias ■ calcium ■ calcium/calmodulin–dependent kinase II ■ sodium

Sudden cardiac death (SCD) refers to a sudden unexpected death from cardiac causes and is a major cause of death worldwide.¹ In patients with structural heart disease, arrhythmias are the main cause of death.² Despite improvements in primary and secondary prevention with substantial decline

in mortality of coronary heart disease in recent years,^{1,3} SCD rates have declined to a much lesser extent.⁴ Implantable cardioverter-defibrillators can reduce cardiovascular mortality in high risk patients, but prevention of SCD is particularly challenging because the majority of cases occur in individuals

Original received February 14, 2015; revision received April 17, 2015; accepted May 7, 2015. In April 2015, the average time from submission to first decision for all original research papers submitted to *Circulation Research* was 13.84 days.

From the Department of Internal Medicine II, University Hospital Regensburg, Regensburg, Germany (S.W., L.S.M.); and Department of Pharmacology, University of California, Davis, CA (D.M.B.).

Correspondence to Lars S. Maier, FAHA, FESC, Head of the Department for Internal Medicine II, University Hospital Regensburg, Franz-Josef-Strauß-Allee 11, 93053 Regensburg, Germany. E-mail lars.maier@ukr.de

© 2015 American Heart Association, Inc.

Circulation Research is available at <http://circres.ahajournals.org>

DOI: 10.1161/CIRCRESAHA.116.304678

Nonstandard Abbreviations and Acronyms

AP	action potential
APD	action potential duration
CaMKII	Ca/Calmodulin-dependent protein kinase II
CPVT	catecholaminergic polymorphic ventricular tachycardia
CV	conduction velocity
DAD	delayed afterdepolarization
EAD	early afterdepolarization
HF	heart failure
I_{Na}	Na current
I_{NCX}	Na/Ca exchange current
MERLIN-TIMI-36 trial	Metabolic Efficiency With Ranolazine for Less Ischemia in Non ST-Elevation Acute Coronary Syndrome Thrombolysis in Myocardial Infarction 36 trial
PLN	phospholamban
ROS	reactive oxygen species
RyR2	cardiac ryanodine receptor 2
SCD	sudden cardiac death
SERCA2a	SR Ca ATPase 2a
SR	sarcoplasmic reticulum
TWA	T wave alternans
VF	ventricular fibrillation
V_m	membrane potential
VT	ventricular tachycardia

without a prior diagnosis of cardiac disease or other clear risk factors for SCD. Thus, understanding the mechanisms of SCD is crucial for both identification of patients at risk and also for developing novel medical or interventional treatment.

SCD is descriptive of a clinical event, but is not a well-defined disease with one specific underlying mechanistic pathophysiology. In contrast to contractile failure, SCD is a consequence of the immediate and complete inability of the heart to maintain the circulation. The most likely SCD causes are lethal cardiac arrhythmias like ventricular tachycardia (VT) or ventricular fibrillation (VF), although bradyarrhythmias also play an important role.⁵ The imprecise definition may also include pulseless electric activity, for example, as a consequence of large pulmonary embolism.

The underlying mechanisms of life-threatening ventricular tachyarrhythmias are manifold. They can occur as a consequence of isolated channelopathies in a structurally normal heart. However, more frequently, they are secondary to substantial dysregulation of intracellular Ca and Na handling that accompanies contractile dysfunction in heart failure (HF). There is also a growing body of evidence that Ca/calmodulin-dependent kinase II (CaMKII) is involved. CaMKII regulates many Na and Ca handling proteins (and is also regulated by Ca), thereby influencing both contractile function and arrhythmogenesis. This review will discuss the mechanisms linking Ca and Na handling to arrhythmogenesis with special focus on CaMKII.

Excitation–Contraction Coupling

To understand arrhythmogenesis, basic knowledge about the fundamental principles of excitation–contraction coupling is




Figure 1. Schematic diagram of cellular Ca fluxes during excitation–contraction coupling. Ca entry via L-type Ca channels triggers Ca-induced Ca release from the sarcoplasmic reticulum (SR), which results in myofilament activation. For relaxation, cytosolic Ca is transported into the SR via SR Ca-ATPase (SERCA2a) and into the extracellular space via sarcolemmal Na/Ca exchanger. Reprinted from Bers⁶ with permission of the publisher. Copyright ©2000, Wolters Kluwer Health.

required (Figure 1). Cardiac excitation is based on sarcolemmal ion fluxes that are tightly coupled to cytosolic Ca and contraction.⁷ Excitation is initiated by opening of voltage-gated Na channels and Na current influx (I_{Na}), leading to the rapid action potential (AP) upstroke (phase 0).^{7,8} Phase 0 is limited by fast I_{Na} inactivation and the activation of transient outward K channels ($K_v4.2$, $K_v4.3$, and $K_v1.4$). These K channels generate transient outward K current (I_{to}), resulting in an early and partial repolarization, that is, the AP notch (phase 1), which is followed by a plateau phase (phase 2). During the latter, membrane conductance falls dramatically as inward and transient outward currents inactivate. Because of the time- and voltage-dependent gating properties of delayed and inward rectifier K channels, K current fluxes are small and only capable of balancing the relatively small remaining inward Ca current (I_{Ca}) via L-type Ca channels ($Ca_v1.2$). This is why the repolarization rate at plateau voltages is slow. Moreover, because of high membrane resistance, even small currents, such as Na/Ca exchange current (I_{NCX}), Na/K-ATPase current (I_{NaK}), or I_{Ca} reactivation, can destabilize the plateau membrane potential, resulting in arrhythmias. I_{Ca} (which activates more slowly than I_{Na}) also progressively decays during the plateau. In ventricular cells, phase 3 repolarization ensues as K currents slowly activate (I_{Ks} , carried by $K_v7.1$, $KCNQ1$) or undergo voltage-dependent recovery from inactivation (I_{Kr} , carried by hERG, $KCNH2$) and I_{K1} (carried by $Kir2.x$ channels) progressively contributes as the membrane repolarizes. I_{K1} also stabilizes diastolic V_m near -80 mV during phase 4. In pacemaker cells, the relative paucity of stabilizing I_{K1} ,⁹ together with the prominence of the hyperpolarization-activated nonspecific cation current I_f (HCN channels) and Ca release driven I_{NCX} , are mainly responsible for diastolic depolarization, leading to AP initiation and pacemaker function.⁷

The cardiomyocyte membrane (sarcolemma) has transverse invaginations called T-tubules (≈ 200 nm diameter) that reach deep into the myocytes interior. T-tubules make close contact with enlarged junctional sarcoplasmic reticulum (SR)

cisternae, with only a 14 nm wide dyadic cleft that separates the SR and T-tubular membranes. L-type Ca channels are highly concentrated in the T-tubules at these junctional clefts and are in close proximity to cardiac ryanodine receptors (RyR2), the SR Ca release channel (Ca release microdomain). This L-type Ca channels-RyR2 proximity at the dyadic cleft means that Ca ions that enter via I_{Ca} cause local cleft [Ca] to be high, sufficient to trigger the opening of RyR2 clusters responsible for SR Ca release during excitation-contraction coupling. This Ca-induced Ca release amplifies the sarcolemmal Ca influx, and depending on species and heart rate, the SR Ca release accounts for 70% to 92% of total Ca increase during systole. Total Ca is heavily buffered ($\approx 100:1$), but free cytosolic [Ca] typically increases from 100 to 600 nmol/L, which leads to myofilament activation and contraction (systole). The $[Ca]_i$ rise is transient because I_{Ca} inactivation and RyR2 closure limit cytosolic Ca entry, and at the same time, Ca removal pathways are activated. For Ca removal, the 2 major pathways are SR Ca-ATPase (SERCA2a) and sarcolemmal Na/Ca exchange, although a tiny fraction of Ca can be extruded by the sarcolemmal Ca-ATPase and mitochondrial Ca uniporter. During steady state, the proportion of Ca transported into SR and out of the cell correspond exactly to the amount of Ca released from the SR and sarcolemmal Ca influx, respectively.

CaMKII

CaMKII is a serine/threonine kinase that can be activated by Ca/calmodulin binding to its regulatory domain. This activation results in the phosphorylation of target proteins, including RyR2, $Na_v1.5$, L-type Ca channels, and phospholamban (the

negative regulator of SERCA2a).^{10,11} Therefore, CaMKII is crucially involved in the regulation of excitation-contraction coupling.

CaMKII assembles into a dodecameric holoenzyme consisting of 2 stacked hexamers. One of the target proteins of CaMKII is the regulatory subunit of an adjacent CaMKII monomer. This auto-phosphorylation results in augmented activation that persists even after dissociation from Ca/CaM and provides an integrated response from the short-term fluctuations in intracellular Ca. It was also shown that reactive oxygen species (ROS)-dependent oxidation of CaMKII at 2 methionine residues at the regulatory domain (M281/282) results in autonomous activation similar to autophosphorylation.¹² Thus, CaMKII can link increased ROS to their downstream effectors. Increased expression and activity of CaMKII^{13,14} has been linked to contractile dysfunction and arrhythmias in HF and on conditions of increased ROS production.^{15–21}

Mechanisms of Electric Instability

Ventricular tachycardia or fibrillation are the result of cell membrane hyperexcitability, disturbed repolarization, or defective conduction of the electric wavefront across the myocardium.²² The presence of an electrically inactive scar tissue that often develops after myocardial infarction forces the wavefront to propagate around this line of block. The wavefront then enters the area behind the scar. If that wavefront reaches the back of the obstacle from multiple directions simultaneously (or near the same time), then the waves will annihilate (Figure 2A). However, if unidirectional block (often coupled with decremental conduction) exists around one side, the wavefront can




Figure 2. Proarrhythmogenic mechanisms. **A, Reentry.** **Upper panel,** Normal conduction around an obstacle. The latter exhibits electric properties that differ dramatically from the rest of the myocardium (slowed conduction and repolarization). The wavefront propagates around this line of block and reaches the back from multiple directions simultaneously, resulting in wave annihilation. If, however, unidirectional block (often coupled with decremental conduction) exists around one side (**lower panel**), the wavefront can reenter and re-excite tissue in front of the unidirectional block. This can lead to stable reentry. **B, Triggered activity.** Increased depolarizing currents that result in prolongation of action potential duration (APD) can favor L-type Ca channel reactivation. Increased Ca entry via I_{Ca} can also lead to sarcoplasmic reticulum (SR) Ca overload and spontaneous SR Ca release during AP plateau phase. This may generate transient inward (I_{Ti}) current by Na/Ca exchange. Both Ca channel reactivation and I_{Ti} may depolarize the membrane during AP plateau phase, which can result in an early afterdepolarization (EAD). In addition, SR Ca overload may also result in diastolic SR Ca release. The consequent I_{Ti} may lead to depolarization from the resting membrane potential, which can result in a delayed afterdepolarization (DAD). Partially derived from Bers.²³

reenter and re-excite tissue in front of the unidirectional block (Figure 2A). This can lead to stable macroscopic re-entry. Reentry can also occur around other functional obstacles (eg, depolarized tissue) and can take the form of a rotor.^{22,24} Stable rotors lead to monomorphic VT. If the waves that propagate outward from a rotor develop wavebreaks, additional rotors are generated, which may lead to polymorphic VT and VF.²⁴ This development is much more likely under conditions of altered excitation, repolarization, or conduction because of disturbed ion channel function (Figure 3). A typical cause for this electric remodeling is structural remodeling associated with cardiac disease secondary to advanced age, increased oxidant stress, hypertension, diabetes mellitus, tissue injury, or inflammation.²² However, the disturbed function of ion channels that underlies these changes in excitation, repolarization, or conduction can also occur in structurally normal myocardium. Abnormal impulse conduction with consequent reentrant excitation can also be a consequence of Ca alternans (Figure 3). This is an example of how Ca dysregulation can cause reentry, which is the most common mechanism of electric instability.

Beside reentry, triggered activity is another important mechanism of electric instability. The latter is the result of an imbalance of ion currents within a single cardiomyocyte. Such imbalances can be classified as either early or delayed depending on their occurrence during the AP. Early afterdepolarizations (EADs) occur during the plateau phase of a cardiac AP, whereas delayed afterdepolarizations (DADs) disturb the diastolic membrane potential after AP repolarization is complete (Figure 2B). Both EADs and DADs can reach the threshold inducing premature ventricular contractions, which can degenerate into monomorphic or polymorphic VTs.

Early Afterdepolarization

EADs occur during the AP and are a consequence of increased inward currents or reduced repolarization reserve (reduced outward K currents).⁷ The plateau phase of the AP is especially vulnerable because repolarizing and depolarizing currents are small and nearly balanced. During this phase, small alterations in the amplitude of even one ionic current can result in

the generation of an EADs. We have learned much about the fundamental mechanisms of triggered activity by studying familial disorders with prolonged repolarization. For example, congenital long QT syndrome 3 is characterized by profound AP prolongation because of increased persistent or late Na current (late I_{Na}),^{16,25} EADs are a major trigger for tachyarrhythmias, but can also occur at low heart rates (where AP duration [APD] is intrinsically long). The long APD allows for reactivation of I_{Ca} at plateau potentials. The resulting inward current creates an EAD as a depolarization in the otherwise monotonically decaying plateau voltage.^{26,27} Among the long QT syndromes, the Timothy syndrome is a rare form, caused by a mutation in $Ca_v1.2$ that prevents normal I_{Ca} inactivation. This disease is characterized by AP prolongation, increased QT intervals, life-threatening arrhythmias, and multisystem defects. Similar to other causes of AP prolongation, I_{Ca} reactivation is a major cause of EAD formation. Interestingly, pathological AP prolongation is a prominent feature of a proarrhythmic electric remodeling in HF,^{28,29} but can also occur in response to drugs that inhibit K channels (especially I_{Kr}). This highlights the importance of this fundamental arrhythmic mechanism.

Beside these well-studied EADs arising during long AP plateaus via I_{Ca} reactivation, there is now compelling evidence that some EADs are initiated by SR Ca rerelease during the AP that causes inward I_{NCX} . These EADs are more common during repolarization and are sometimes called phase 3 EADs. The inward I_{NCX} may only cause a weak depolarization, but that can be amplified by reactivation of I_{Ca} ,^{30,31} depending on the voltage. That is, in atrial (and rodent ventricular) APs with I_{NCX} -prolonged late and plateau at voltages below -40 mV (where I_{Ca} reactivation does not occur), EADs may be driven by nonequilibrium reactivation of I_{Na} , which is facilitated by increased SR Ca release.³²

Delayed Afterdepolarization

DADs are a consequence of cytosolic and SR Ca overload.²³ Both increased cytosolic and SR luminal Ca increase the diastolic open probability of cardiac RyR2.⁷ Ca sparks are elementary SR Ca release events that occur if a cluster of RyR2




Figure 3. Overview of proarrhythmic consequences of ion channel dysfunction.

Congenital or acquired dysfunction of ion channels can result in (1) slowed conduction, (2) repolarization, and (3) disturbed intracellular Na and Ca homeostasis. Slowed conduction velocity (CV; with or without CV alternans) or repolarization (with or without repolarization alternans) causes spatiotemporal differences in CV and action potential duration (APD) within the myocardium. These regional differences can lead to unidirectional block with reentry and rotor formation. The consequent monomorphic ventricular tachycardia (VT) may result in additional rotors by heart rate–dependent alternans causing polymorphic VT and ventricular fibrillation (VF). Disturbed Na and Ca homeostasis, on the other hand, may lead to Ca alternans, which is linked to repolarization alternans. Moreover, prolongation in APD duration causes early afterdepolarization (EADs) and increased Ca leak results in delayed afterdepolarizations (DADs), both of which underlie ectopic activity. The latter can also lead to monomorphic or polymorphic VT and VF.

open. During systolic Ca-induced Ca release, Ca sparks are synchronized within the cell and summate to form the Ca transient. Spontaneous Ca sparks are mainly responsible for diastolic SR Ca release and diastolic SR Ca leak via RyR2.^{33,34} Because RyR2 is in relatively close proximity to the NCX, this localized $[Ca]_i$ elevation drives inward I_{NCX} . Note that the stoichiometry of Na/Ca exchange ($3Na^+:1Ca^{2+}$) meaning that Ca extrusion is accompanied by a net inward movement of one positive charge via I_{NCX} which is almost entirely responsible for what used to be called transient inward current I_{ti} .²³ If this I_{NCX} is large enough, it can cause an appreciable DAD, which can trigger an arrhythmic AP. Individual Ca sparks normally do not produce enough I_{NCX} to produce a measurable DAD because they are isolated unsynchronized events. However, at higher SR Ca content, the Ca sparks are larger in amplitude and nearby RyR2 clusters are more sensitive to activation, resulting in a cell-wide Ca wave with sufficient I_{NCX} to cause DADs and triggered APs. Sympathetic stimulation can drive up SR Ca content and increase the propensity for Ca waves and triggered beats.

In the intact heart, the I_{NCX} produced by a Ca wave in a single myocyte is insufficient to trigger an appreciable DAD or AP because all of the neighboring cells can effectively clamp that single cell at the diastolic voltage. However, when these Ca waves and consequent I_{NCX} are synchronized regionally among many cells (by the prior AP and similar RyR2 recovery kinetics), that region can initiate triggered beats as premature ventricular contractions for the whole heart.^{35,36} The heart is relatively protected from these triggered arrhythmias initiated via EADs and DADs, but under pathological conditions caused by either genetic ion channel/transporter mutations or acquired diseases like HF, both the cellular propensity for EADs and DADs and their ability to cause whole heart premature ventricular contractions can be greatly increased.^{35–38}

Cardiac Alternans

Cardiac alternans is a known risk factor for cardiac arrhythmias and SCD.^{39–41} The transition from monomorphic VT to polymorphic VT or VF requires breaks in the wave propagating from a stable rotor. During VT, cardiomyocyte Ca cycles at high rates. The time for intracellular Ca removal is drastically shortened and can elevate diastolic $[Ca]_i$ and limit SR Ca load.⁴² This can also encroach upon the recovery time of the I_{Ca} -induced SR Ca release.^{43–46} Normally, $[Ca]_i$ is tightly controlled, but Ca dysregulation can predispose to afterdepolarizations and cardiac alternans.^{47–50} Cardiac alternans are alternating beats with large/small amplitude Ca transient and long/short APD (Figure 4A). Most common in large mammals are electromechanically (or V_m -Ca) concordant alternans in which the large Ca transient and contraction is associated with the longer APD. However, V_m -Ca discordant alternans can also occur. Alternans can also be spatially concordant, where all regions of the ventricle are in phase with each other, or spatially discordant where different regions are out of phase with each other. As we will see the Ca and V_m changes are usually functionally linked, but both can occur independently.⁴⁷

Repolarization Alternans

Alternation in APD is called repolarization alternans. Because this type of alternans involves membrane potential, it can be clinically observed as T-wave alternans (TWA).⁵¹ TWA has been shown to be associated with cardiac arrhythmias and SCD.^{52–54} TWA is not restricted to a specific underlying disease but can be observed in HF, Brugada syndrome, and long QT syndrome.⁵⁵

TWA can result from changes in sarcolemmal ion current recovery (typically of I_{Na} or I_{Ca}) manifested by a steep slope of cellular APD restitution.⁵⁶ This type of TWA typically occurs at high heart rates with reduced diastolic intervals.^{51,57,58} The dependence of APD and conduction velocity (CV) on the preceding diastolic interval are called APD restitution and CV restitution, respectively. Shorter diastolic intervals result in shorter APD and slower CV.⁴⁸ Both APD and CV restitution critically depend on the speed of recovery from inactivation of sarcolemmal ion channels. Nevertheless, other mechanisms, for instance SR Ca release, can also contribute to APD restitution changes. Intracellular Ca released from the SR inactivates I_{Ca} . A large SR Ca release could, therefore, result in a more pronounced Ca-dependent inactivation of I_{Ca} , which would influence APD restitution.

CV restitution, on the other hand, mainly depends on Na channel recovery.⁵⁹ This is because CV depends on AP upstroke velocity, and the latter is determined by the magnitude of I_{Na} . Na channel recovery is usually fast; thus, CV restitution occurs only at high heart rates, as would be the influence of Na channel recovery on APD restitution. However, under conditions of slowed Na channel recovery, the impact on CV restitution (and also APD restitution) may already occur at much lower heart rates. This is important because slowed Na channel recovery has been shown to be a feature of many cardiac diseases like HF^{15,60} or ischemia^{59,61,62} (where CaMKII may be involved)¹⁵ and Brugada syndrome.⁶³

As mentioned above, APD restitution depends on sarcolemmal ion channel recovery. Among them, L-type Ca channels are very important. I_{Ca} is the major inward current during the AP plateau.⁷ As stated earlier, AP plateau is the most vulnerable phase of the AP. This explains why inhibition of I_{Ca} has been shown to reduce the slope of the APD restitution curve. In addition, the recovery of Ca channels is slower compared with, for instance, Na channels. Therefore, Ca channels influence APD restitution already at lower heart rates. Interestingly, because I_{Ca} also determines SR Ca load, and vice versa, APD alternans and Ca alternans are functionally linked.

Repolarization alternans can be either spatially concordant or discordant.^{51,64} It was shown that Na channel recovery is critically involved in spatially discordant alternans.^{64,65} Because spatially discordant alternans is accompanied by changes in CV, both T wave and QRS are affected, resulting in T-wave and QRS alternans (Figure 6). Repolarization alternans can be proarrhythmic by 2 mechanisms. First, spatially discordant alternans results in spatial dispersion of repolarization providing a substrate for reentry.⁵¹ Second, at high heart rates, alternans can promote conduction block, which may trigger rotor formation. Alternans-dependent arrhythmias are, therefore, most likely generated under conditions of increased




Figure 4. Characteristics of cardiac alternans. **A**, Simultaneous recordings of action potential (AP) and intra-sarcoplasmic reticulum (SR) free $[Ca]$ in isolated Langendorff-perfused rabbit hearts by optical mapping. At shorter pacing intervals (190 ms), alternans of the AP duration as well as SR Ca release alternans was observed. **B**, SR Ca release alternans can occur at heart rates where action potential duration (APD) alternans is not yet detectable. **C**, Ca release alternans can be amplified by alternating changes in SR Ca load, that is, SR Ca load alternans. Reprinted from Picht et al⁴³ with permission of the publisher. Copyright ©2006, Wolters Kluwer Health.

heart rate like during exercise or occur secondary to a stable rotor-dependent ventricular tachycardia. In fact, alternans may mechanistically explain why a stable ventricular tachycardia may degenerate into a multiple wavelet-driven polymorphic ventricular tachycardia or VF.^{66,45}

Importantly, if APD is substantially prolonged, which is a typical feature of HF^{28,29} or long QT syndrome,⁴⁶ ion channel recovery is already impaired at lower heart rates. Even without AP prolongation, if Na channel recovery is impaired like in HF,^{15,60} ischemia,^{59,61,62} and Brugada syndrome,⁶³ alternans-dependent arrhythmias can occur at much lower heart rates.

Ca Alternans

From the surface electric measurements of an ECG, it is impossible to state whether APD alternans is accompanied by Ca alternans or whether one leads to the other. However, extensive work over the past years has shown that Ca and APD alternans usually coexist, are mechanistically linked, and that




Figure 5. Proarrhythmogenic mechanisms of enhanced late I_{Na} . In systole (**upper panel**), enhanced late I_{Na} leads to action potential (AP) prolongation (1). The longer AP plateau phase increases the likelihood of I_{Ca} reactivation (2), which may lead to early afterdepolarizations (EAD, 3). **Lower panel**, The increased amount of Na influx also results in increased intracellular Na (1), which impairs Ca elimination (2) by the Na/Ca exchanger (either less forward or even increased reverse mode activity). In diastole, the increased intracellular Ca facilitates SR Ca leak (3), which could lead to transient inward current (I_{Ti}) by the Na/Ca exchanger (4). The latter can result in delayed afterdepolarizations (DAD, 5).




Figure 6. Ca/calmodulin-dependent kinase II (CaMKII)-dependent mechanisms of triggered activity. CaMKII is activated by pathophysiologically relevant stimuli, that is, increased reactive oxygen species (ROS), increased intracellular Ca, hyperglycemia. CaMKII-dependent phosphorylation of L-type Ca channels (1) may increase I_{Ca} window current predisposing to early afterdepolarizations (EADs). Increased CaMKII-dependent RyR2 phosphorylation (2) results in increased diastolic Ca leak. CaMKII-dependent phospholamban (PLN) phosphorylation (3) maintains SR Ca content, which may also stimulate diastolic RyR2 openings from the luminal side. Diastolic SR Ca release triggers transient inward current (I_{Ti} , 4) and delayed afterdepolarizations (DADs). Increased CaMKII-dependent phosphorylation of $Na_v1.5$ (5) leads to enhanced late I_{Na} , which predisposes for both EADs and DADs (Figure 5).

Ca-driven alternans seems to be more clinically relevant in the setting of heart disease.⁵⁶

Alternans start to occur as pacing rate is increased, and Ca alternans occur even when myocytes are voltage clamped with identical V_m waveforms, consistent with Ca-driven alternans.^{47,48,67}

In the absence of voltage clamp, those Ca alternans also cause APD alternans, and there are logical reasons why a larger versus smaller Ca transient would alter APD. The 2 most prominent $[Ca]_i$ -dependent currents are I_{Ca} and I_{NCX} . A large Ca transient would (a) strengthen Ca-dependent inactivation of I_{Ca} , which would shorten APD, and (b) drive increased inward I_{NCX} , which would prolong APD. In mammals with prominent AP plateaus, concordant V_m -Ca alternans usually predominate, suggesting that the I_{NCX} effects are more powerful contributors to alternans. However, some conditions could shift this balance to favor a predominant role for I_{Ca} inactivation and discordant V_m -Ca alternans. As stimulation rate increases, alternans start at a certain threshold and the alternans ratio (large:small Ca transient amplitude or APD) increases progressively. This can be seen as alternans in the amount of SR Ca release in direct measurements of intra-SR free $[Ca]$ ($[Ca]_{SR}$), in both isolated myocytes and whole heart (Figure 4A).^{43,44}

Several factors contribute to alternans, and they seem to follow a sequence. The first step seems to be an encroachment on RyR2 recovery so that SR Ca release decreases, despite unaltered APD, I_{Ca} , and $[Ca]_{SR}$.^{43,44,68}

This is because RyR2 restitution is much slower than normal I_{Ca} restitution. The partial failure of SR Ca release at this first small beat allows improved RyR2 recovery at the next large beat, but then the cycle repeats. SR Ca release alternans can appear at heart rates where the consequent APD alternans are not yet detectable (Figure 4B). As Ca release alternans grow, they are amplified by alternating changes in SR Ca load, that is, SR Ca load alternans (Figure 4C). The small release limits Ca-dependent inactivation of I_{Ca} , thereby increasing Ca influx and load for the next (large) beat. The small Ca transient also drives less Ca extrusion via NCX, which limits Ca loss at the small beat. Then at the large beat, there is greater Ca-dependent I_{Ca} inactivation and greater Ca efflux via NCX, which reduces cell and SR Ca load, setting the scene for stable alternans, ≤ 6 to 7 Hz (150 ms cycle length) in rabbit hearts (Figure 4C). The steep nonlinear dependence of SR Ca release on diastolic SR Ca load^{42,69} enhances the impact of the SR load alternans on Ca release alternans.⁴⁹ At even higher stimulation rates, shorter diastolic intervals or more depolarized diastolic voltage, one can encroach on I_{Ca} (or even I_{Na}) restitution, which may also exacerbate a sort of Ca-driven alternans (smaller I_{Ca} causes smaller Ca transient and APD). But here the lines become blurred with the electric, restitution-driven alternans involving APD and CV restitution, as discussed above.

Several factors can influence the pacing rate at which alternans is observed. Inhibition of glycolysis, mitochondrial energetic limitations, decreased SERCA function, and redox modification of RyR2 can all favor alternans.^{70–73} All of these factors may be relevant in pathophysiological conditions like HF, thus enhancing the propensity for alternans and arrhythmogenic sequelae in these conditions.

Similar to repolarization alternans, Ca alternans can be spatially in phase (concordant) or out of phase, that is, discordant, in different regions of a single myocytes^{74–76} or hearts, and spatially discordant alternans have been associated with lethal VTs and VF in patients.^{51,77}

Arrhythmias in HF and Ischemia/Reperfusion

Despite improvements in HF therapy and ischemia/reperfusion associated myocardial damage, it has proven difficult to develop antiarrhythmic treatments and prevent SCD. Conventional ion channel blockers that are used as antiarrhythmic drugs are known to induce arrhythmias in structural heart disease.

There is an increasing body of evidence that HF and ischemic heart disease are accompanied by alterations in intracellular Na and Ca handling. Both conditions are characterized by an increased generation of ROS,^{78–80} which may contribute to disturbed Na and Ca handling.⁸¹ Changes in intracellular Na and Ca handling are associated with electric instability, and many of these alterations are linked to both contractile dysfunction and arrhythmias.

Late Na Current

As stated earlier, APD is increased in HF.²⁹ Beside reduction of K current expression, increased magnitude of late I_{Na} has been shown to contribute to AP prolongation in HF⁸² and after myocardial infarction.⁸³ Late I_{Na} is generated by dysfunctional inactivation of cardiac voltage-gated Na channels $Na_v1.5$.⁸⁴ The detailed molecular mechanism is not fully understood.⁸⁵

Although late I_{Na} has a small amplitude compared with peak I_{Na} , it persists for hundreds of milliseconds during the cardiac AP,⁸⁶ providing a source for increasing $[Na]_i$.^{15,16} Increased $[Na]_i$ is a well-known feature of HF, ischemia/reperfusion, or other conditions of increased ROS production and can contribute to contractile dysfunction and arrhythmias.^{16,19,25,87–91}

The late I_{Na} -dependent prolongation of the AP plateau renders the membrane potential vulnerable for EADs (Figure 5).¹⁶ A similar arrhythmic mechanism can be found in familial long QT syndrome 3. Mutations in the gene encoding for $Na_v1.5$ (*SCN5A*) have been shown to increase late I_{Na} , leading to AP prolongation and EADs.^{92,93} Transmural differences in late I_{Na} might also increase dispersion of repolarization,⁹⁴ which underlies the development of torsade de pointes tachyarrhythmias. In addition, late I_{Na} is involved in cardiac alternans: ischemia-induced spatially discordant repolarization alternans has been shown to be prevented by inhibition of late I_{Na} .⁹⁵

Beside AP prolongation, late I_{Na} -dependent Na overload leads to intracellular Ca accumulation either by reduced Ca exit because of limitations to Ca extrusion by NCX or by additional Ca entry via reverse mode NCX.¹⁶ Intracellular Ca accumulation is associated with increased diastolic SR Ca leak and DADs (Figure 5). Moreover, it contributes to diastolic dysfunction in HF and under conditions of increased ROS production,^{16,90} and it may even lead to cellular hypercontracture.⁹⁶

AP-clamp experiments revealed that late I_{Na} -dependent Na entry also substantially contributes to arrhythmias in long QT syndrome 3 syndrome.⁹⁷ Consistent with this evidence, blocking late I_{Na} and Ca entry via NCX have been shown to inhibit afterdepolarizations.^{25,98}

What are the mechanisms that lead to increased late I_{Na} ? ROS have been shown to increase $[Na]_i$, prolong APD,

and induce EADs, and ROS-enhanced late I_{Na} may be involved.^{16,25,91} Ahern and colleagues showed that nitrosylation of $Na_v1.5$ increased late I_{Na} under physiological and pathophysiologic conditions.⁹⁹ Over recent years, however, a body of evidence suggests that CaMKII associates with and phosphorylates $Na_v1.5$, leading to increased late Na current, and some ROS and nitric oxide effects may be mediated via CaMKII activation (Figure 6).^{15,16,60,100–104}

Because late I_{Na} is strongly linked to cardiac arrhythmogenesis, strategies aimed at inhibiting late I_{Na} may have strong antiarrhythmic potential and may possibly be used to prevent SCD. The clinically approved antianginal drug ranolazine was found to inhibit late I_{Na} .^{105,106} Within the therapeutic range, which varies between 2 and 8 $\mu\text{mol/L}$, ranolazine also inhibits I_{Ca} by 30%, which could limit Ca entry,¹⁰⁶ and I_{K_r} , which can prolong APD. Depending of the actual ionic conditions, the effects of ranolazine on APD may vary. Inhibition of late I_{Na} by ranolazine has been shown, for instance, to reduce ROS-dependent AP prolongation, as well as intracellular Na and Ca accumulation, diastolic dysfunction, and arrhythmias.^{16,25,89,107,108} Ranolazine has been demonstrated to inhibit afterdepolarizations, to reduce transmural dispersion of repolarization, and to inhibit spatially discordant repolarization alternans in various models of enhanced late I_{Na} .^{95,105,106,109–111} It also prevented pacing-induced reentry and multifocal VF.¹¹² In experimental models of systolic HF, ranolazine treatment also improved systolic function,¹¹³ possibly by reducing late I_{Na} -dependent diastolic SR Ca leak.¹⁹

The clinical safety and efficacy of ranolazine are well investigated. A large clinical outcome trial investigating the efficacy of ranolazine to reduce cardiovascular death, myocardial infarction, or recurrent ischemia in >6000 patients with acute coronary syndrome (Metabolic Efficiency With Ranolazine for Less Ischemia in Non ST-Elevation Acute Coronary Syndrome Thrombolysis in Myocardial Infarction 36 trial [MERLIN-TIMI-36]) was not able to show an improvement in outcome.¹¹⁴ Nevertheless, subgroup analysis revealed that in placebo-treated patients, prolonged QTc was a significant independent predictor for SCD, whereas this was not the case in patients treated with ranolazine,¹¹⁵ suggesting a potential beneficial effect in the prevention of SCD. Investigation of the 7-day Holter monitoring data acquired in the MERLIN-TIMI-36 trial revealed a reduction in the incidence of ventricular tachycardia in the ranolazine treated patients.¹¹⁶ Ranolazine also reduced VT burden and implantable cardioverter-defibrillator shocks in a small observational study of patients with refractory VT and previous implantable cardioverter-defibrillator shocks.¹¹⁷ In a small study of patients with long QT syndrome 3, ranolazine shortened the QTc interval in a concentration-dependent manner.¹¹⁸ Moreover, in patients with atrial fibrillation, a dose-finding randomized controlled trial showed a decreased rate of in overall atrial fibrillation recurrence.¹¹⁹

CaMKII

CaMKII is a central regulator of many Na and Ca (and K) channels and transporters. It is well established that transgenic overexpression of CaMKII results in the development of HF and arrhythmias^{15,120} and that CaMKII inhibition prevented afterdepolarizations and arrhythmias on β -adrenergic stimulation.¹⁸

Increased CaMKII autophosphorylation was observed in rabbit chronic atrioventricular block models of left ventricular dysfunction, acquired long QT syndrome, and incessant ventricular tachycardia.^{121,122} In addition, CaMKII inhibition prevents the development of structural heart disease or arrhythmias on myocardial infarction,¹²³ increased ROS formation,¹⁶ pressure overload,¹²⁴ or pacing-induced incessant VT.¹²² Calcineurin overexpressing mice with increased CaMKII activity show contractile dysfunction and arrhythmias.¹²⁵ Also here, CaMKII inhibition improved contractile function and suppressed arrhythmias.¹²⁵

Oxidized CaMKII has been shown to be involved in ROS-induced enhancement of late I_{Na} (Figure 6), leading to intracellular Na and Ca accumulation, contractile dysfunction, and triggered activity.^{16,126} Increased oxidative activation of CaMKII because of enhanced NADPH oxidase 2 may also be important for angiotensin II-dependent arrhythmogenesis.^{127,128} Moreover, oxidized CaMKII plays a role in diabetes mellitus-dependent increased mortality after myocardial infarction possibly by sinus node dysfunction.¹²⁹ A novel CaMKII activation pathway by O-linked glycosylation at serine 279 was demonstrated during acute hyperglycemia. This activation confers similar autonomous activity as autophosphorylation or oxidation.¹³⁰ O-linked glycosylation on diabetic hyperglycemia was shown to result in increased diastolic SR Ca leak and be proarrhythmogenic.¹³⁰ CaMKII can be also nitrosylated at another site in the regulatory domain to cause autonomous activation.¹³¹ Nitrosylation-dependent CaMKII activation has been implicated in increased SR Ca leak in cardiac myocytes.^{132,133} Neurohumoral activation that occurs in HF has been shown to activate CaMKII. Both β -adrenergic stimulation and angiotensin II exposure can activate CaMKII either by increased cytosolic Ca or by increased ROS.^{12,123,127,134–137}

The mechanisms of CaMKII-dependent arrhythmias are complex because CaMKII regulates a variety of ion channels and transporters (Figure 6). CaMKII was first identified to increase peak L-type Ca current and slow I_{Ca} inactivation,^{138–140} which may predispose to EADs.^{141–143}

CaMKII is also known to associate with and phosphorylate cardiac $Na_v1.5$.^{15,16,60,100–104} Beside increased late I_{Na} that may contribute to AP prolongation, increased EAD and DAD formation, and repolarization alternans, CaMKII-dependent phosphorylation also enhanced I_{Na} intermediate inactivation and slowed recovery from inactivation.^{15,60,103,104} Both effects reduce the steady-state availability of I_{Na} , which could lead to repolarization alternans at high heart rates. Reduced Na channel availability can also increase transmural dispersion of repolarization and slow intraventricular conduction (Figure 7).¹⁵ This peculiar phenotype of CaMKII-dependent gain in Na channel function (enhanced late I_{Na}) and loss of Na channel function (reduced I_{Na} availability) may sound incongruous. However, similar changes in I_{Na} gating have been observed in a *SCN5A* mutation (1795InsD) associated with features of both Brugada and long QT syndrome.^{144,145} Thus, increased CaMKII activity in HF may lead to an acquired form of combined long QT and Brugada syndrome.

There is an overwhelming body of evidence that CaMKII regulates RyR2 and SERCA2a. Although RyR2 is directly




Figure 7. Ca/calmodulin-dependent kinase II (CaMKII)-dependent regulation of I_{Na} gating and alternans. **A**, CaMKII-dependent phosphorylation of $Na_v1.5$ has been shown to enhance I_{Na} intermediate inactivation (reproduced from Wagner et al¹⁴ with permission; whole-cell patch clamp in rabbit ventricular myocytes). As a result, the number of available Na channels is reduced especially at shorter diastolic intervals. **B**, Consequences of enhanced I_{Na} intermediate inactivation: CV and repolarization alternans. Reduced Na channel availability results in slowed intramural conduction and slowed AP upstroke velocity (1) evident as broader QRS complex on surface ECG (1). In addition, K channel expression is larger in epicardium. Therefore, repolarization is faster in epicardium, especially if Na current is reduced (2). This leads to increased transmural dispersion of repolarization evident as larger and wider T wave on surface ECG (2). Interestingly, Na channels in intermediate inactive state cannot be activated (and become refractory) during the excitation. Thus, these channels are available for the consecutive excitation. Consequently, conduction velocity and AP upstroke velocity will be larger, AP duration longer for the consecutive excitation: the typical pattern of CV and repolarization alternans.

phosphorylated at serine 2814,¹⁴⁶ the activity of SERCA2a is indirectly influenced by phosphorylation of phospholamban (PLN) at threonine 17. The latter results in relief of PLN-dependent SERCA2a inhibition and activation of SERCA2a.¹⁴⁷ In HF, increased diastolic Ca leak through RyR2 occurs in the face of a reduced SR Ca reuptake because of a smaller SERCA2a/PLN expression ratio. This results in a reduced SR Ca content, which is an important determinant for the impaired Ca transient amplitude of failing cardiomyocytes.^{7,148,149} In HF, CaMKII-dependent diastolic SR Ca leak is a major cause for DADs by activating forward mode NCX.^{150,151}

Zhang et al¹⁵² tried to rescue cardiac function in $CaMKII\delta_c$ transgenic with reduced SR Ca load by PLN ablation. Although this improved SR Ca load and Ca transients, this occurred at the expense of increased CaMKII-dependent SR Ca leak, mitochondrial Ca loading, and myocyte death. Thus, despite improved myocyte function, cardiac function, HF progression, and mortality were worsened by the PLN ablation.

More insights into the role of CaMKII-dependent SR Ca leak for contractile dysfunction and arrhythmias in HF comes from animal models with increased afterload or myocardial infarction. It was shown that afterload-induced HF development and arrhythmias were inhibited in knock-in mice harboring only a mutant RyR2 (S2814A) that cannot be phosphorylated by CaMKII at that site.^{153,154} This knock-in, however, did not prevent HF postmyocardial infarction.¹⁵³ In contrast, recent

data show similar CaMKII-dependent SR Ca leak in isolated human ventricular cardiomyocytes from ischemic and dilated cardiomyopathy.¹⁵⁵

Also, KI mice with a CaMKII-phosphomimetic mutation (S2814D) exhibited increased SR Ca leak and developed sustained ventricular tachycardia and SCD on β -adrenergic stimulation, programmed electric stimulation, and increased afterload.¹⁵⁴ This sort of CaMKII-dependent SR Ca leak has also been observed on β -adrenergic stimulation^{156–158} and by activation of NADPH oxidase resulting in DADs.¹²⁷ Although much attention has been on ventricular muscle, it was also shown that CaMKII-dependent SR Ca leak is crucially involved in atrial arrhythmogenesis.^{159–162} For ROS-induced SR Ca leak, however, a direct oxidation of the RyR2 has also been proposed.^{16,163–165} Thus, CaMKII inhibition may be a novel strategy to prevent SCD.

Catecholaminergic Polymorphic Ventricular Tachycardia

Increased SR Ca leak is causally involved in catecholaminergic polymorphic ventricular tachycardia (CPVT), a rare familial condition characterized by ventricular arrhythmias that are associated with exercise or catecholaminergic stimulation in a structurally normal heart.¹⁶⁶ CPVT occurs at young ages. For example, the majority of 101 patients with CPVT had symptoms before the age of 21.¹⁶⁷ Importantly, CPVT

accounts for 15% of all sudden unexplained deaths in young people. CPVT is linked to mutations in *RyR2* and the intra-SR Ca binding and *RyR2*-associated proteins calsequestrin (*CASQ2*) and triadin (*TRND*).^{168–172} For a substantial fraction of patients, however, the disease-causing gene has remained elusive to date. As a result of the irregular SR Ca release,¹⁷³ electrogenic NCX is activated causing DADs. Clinically, patients exhibit monomorphic ventricular premature beats and more severe bidirectional ventricular tachycardia, which can degenerate into polymorphic VT and VF. Interestingly, pharmacological CaMKII inhibition has recently been shown to inhibit stress-induced arrhythmias and triggered activity in a mouse model of CPVT (*RyR2* R4496C^{+/−}).¹⁷⁴ The opposite, transgenic CaMKII overexpression in *RyR2* R4496C^{+/−} mice resulted in increase SR Ca leak, DADs, arrhythmias on β -adrenergic stimulation, and increased mortality possibly because of SCD.¹⁷⁵ Because disturbed CaMKII-dependent regulation of SR Ca release is a frequent feature of HF, exercise-induced arrhythmias in HF may be related to the same underlying mechanism. For more information about this important genetic arrhythmogenic disease, the reader is referred to these more comprehensive reviews.^{176–178}

Epigenetic Modifications Related to SCD

Epigenetic regulation of gene expression is increasingly recognized as important contributor arrhythmias.¹⁷⁹ This emerging area includes microRNAs, DNA methylation, and histone modifications (eg, acetylation/deacetylation). Apropos to the Ca/CaMKII focus here, it is known that CaMKII can importantly influence the nuclear export of class II histone deacetylases (eg, HDAC4 and HDAC5) and that it can modulate transcription of key proteins involved in hypertrophic signaling and arrhythmias.^{180–182} CaMKII can also directly phosphorylate histone H3 and contribute to hypertrophic changes in gene expression.^{183,184} Thus, CaMKII seems to be integrally involved in epigenetic mechanisms of cardiac hypertrophy, and some of the consequent changes in protein expression (of ion channels and Ca regulatory proteins) may also contribute to enhanced propensity for arrhythmias.

Conclusions

Cardiac myocyte Na and Ca fluxes and concentrations are tightly controlled, but also change dynamically as part of normal physiological modulation of cardiac electric, contractile, and energetic state. There is tight coupling between Na, Ca, electric, mechanical, and energetic properties in the heart, only some of which we touched upon here. Many arrhythmias are a result of dysregulation of this complex system. To understand how deranged Na and Ca homeostasis comes about in pathophysiological states and how it contributes to electric and contractile dysfunction, such as HF, arrhythmias and SCD, is a challenge. But this understanding is essential for progress in novel therapeutic approaches to these major clinical problems.

Sources of Funding

S. Wagner and L.S. Maier are funded by Deutsche Forschungsgemeinschaft (DFG) through an International Research Training Group GRK 1816. L.S. Maier is funded by the Fondation

Leducq Transatlantic Network on “Redox and Nitrosative Regulation of Cardiac Remodeling”. L.S. Maier, and S. Wagner are funded by the DZHK (Deutsches Zentrum für Herz-Kreislauf-Forschung-German Centre for Cardiovascular Research). D.M. Bers is funded by National Institute of Health grants (R37-HL30077, P01-HL080101, and R01-HL081562).

Disclosures

L.S. Maier receives research grants from Gilead and is involved in clinical trials with Gilead and Menarini. L.S. Maier received speaker's honoraria from Berlin-Chemie. The other authors report no conflicts.

References

- Go AS, Mozaffarian D, Roger VL, et al; American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2014 update: a report from the American Heart Association. *Circulation*. 2014;129:e28–e292. doi: 10.1161/01.cir.0000441139.02102.80.
- Lehnart SE, Ackerman MJ, Benson DW Jr, et al. Inherited arrhythmias: a National Heart, Lung, and Blood Institute and Office of Rare Diseases workshop consensus report about the diagnosis, phenotyping, molecular mechanisms, and therapeutic approaches for primary cardiomyopathies of gene mutations affecting ion channel function. *Circulation*. 2007;116:2325–2345. doi: 10.1161/CIRCULATIONAHA.107.711689.
- Ford ES, Ajani UA, Croft JB, Critchley JA, Labarthe DR, Kottke TE, Giles WH, Capewell S. Explaining the decrease in U.S. deaths from coronary disease, 1980–2000. *N Engl J Med*. 2007;356:2388–2398. doi: 10.1056/NEJMs053935.
- Dudas K, Lappas G, Stewart S, Rosengren A. Trends in out-of-hospital deaths due to coronary heart disease in Sweden (1991 to 2006). *Circulation*. 2011;123:46–52. doi: 10.1161/CIRCULATIONAHA.110.964999.
- Huikuri HV, Castellanos A, Myerburg RJ. Sudden death due to cardiac arrhythmias. *N Engl J Med*. 2001;345:1473–1482. doi: 10.1056/NEJMra000650.
- Bers DM. Calcium fluxes involved in control of cardiac myocyte contraction. *Circ Res*. 2000;87:275–281.
- Bers DM. Cardiac excitation-contraction coupling. *Nature*. 2002;415:198–205. doi: 10.1038/415198a.
- Grandi E, Pasqualini FS, Bers DM. A novel computational model of the human ventricular action potential and Ca transient. *J Mol Cell Cardiol*. 2010;48:112–121. doi: 10.1016/j.jmcc.2009.09.019.
- Cho HS, Takano M, Noma A. The electrophysiological properties of spontaneously beating pacemaker cells isolated from mouse sinoatrial node. *J Physiol*. 2003;550:169–180. doi: 10.1113/jphysiol.2003.040501.
- Swaminathan PD, Purohit A, Hund TJ, Anderson ME. Calmodulin-dependent protein kinase II: linking heart failure and arrhythmias. *Circ Res*. 2012;110:1661–1677. doi: 10.1161/CIRCRESAHA.111.243956.
- Anderson ME, Brown JH, Bers DM. CaMKII in myocardial hypertrophy and heart failure. *J Mol Cell Cardiol*. 2011;51:468–473. doi: 10.1016/j.jmcc.2011.01.012.
- Erickson JR, Joiner ML, Guan X, et al. A dynamic pathway for calcium-independent activation of CaMKII by methionine oxidation. *Cell*. 2008;133:462–474. doi: 10.1016/j.cell.2008.02.048.
- Kirchhefer U, Schmitz W, Scholz H, Neumann J. Activity of cAMP-dependent protein kinase and Ca²⁺/calmodulin-dependent protein kinase in failing and nonfailing human hearts. *Cardiovasc Res*. 1999;42:254–261.
- Hoch B, Meyer R, Hetzer R, Krause EG, Karczewski P, Max Delbrück Center for Molecular Medicine B-BG. Identification and expression of δ -isoforms of the multifunctional Ca²⁺/calmodulin-dependent protein kinase in failing and nonfailing human myocardium. *Circ Res*. 1999;84(6):713–721.
- Wagner S, Dybkova N, Rasenack EC, Jacobshagen C, Fabritz L, Kirchhof P, Maier SK, Zhang T, Hasenfuss G, Brown JH, Bers DM, Maier LS. Ca²⁺/calmodulin-dependent protein kinase II regulates cardiac Na⁺ channels. *J Clin Invest*. 2006;116:3127–3138. doi: 10.1172/JCI26620.
- Wagner S, Ruff HM, Weber SL, Bellmann S, Sowa T, Schulte T, Anderson ME, Grandi E, Bers DM, Backs J, Belardinelli L, Maier LS. Reactive oxygen species-activated Ca/calmodulin kinase II δ is required for late I_{Na} augmentation leading to cellular Na and Ca overload. *Circ Res*. 2011;108:555–565. doi: 10.1161/CIRCRESAHA.110.221911.
- Toischer K, Rokita AG, Unsöld B, et al. Differential cardiac remodeling in preload versus afterload. *Circulation*. 2010;122:993–1003. doi: 10.1161/CIRCULATIONAHA.110.943431.

18. Sag CM, Wadsack DP, Khabbazzadeh S, Abesser M, Greffe C, Neumann K, Opiela MK, Backs J, Olson EN, Brown JH, Neef S, Maier SK, Maier LS. Calcium/calmodulin-dependent protein kinase II contributes to cardiac arrhythmogenesis in heart failure. *Circ Heart Fail*. 2009;2:664–675. doi: 10.1161/CIRCHEARTFAILURE.109.865279.
19. Sag CM, Mallwitz A, Wagner S, Hartmann N, Schotola H, Fischer TH, Ungeheuer N, Herting J, Shah AM, Maier LS, Sossalla S, Unsöld B. Enhanced late INa induces proarrhythmic SR Ca leak in a CaMKII-dependent manner. *J Mol Cell Cardiol*. 2014;76:94–105. doi: 10.1016/j.yjmcc.2014.08.016.
20. Sag CM, Köhler AC, Anderson ME, Backs J, Maier LS. CaMKII-dependent SR Ca leak contributes to doxorubicin-induced impaired Ca handling in isolated cardiac myocytes. *J Mol Cell Cardiol*. 2011;51:749–759. doi: 10.1016/j.yjmcc.2011.07.016.
21. Sag CM, Wolff HA, Neumann K, et al. Ionizing radiation regulates cardiac Ca handling via increased ROS and activated CaMKII. *Basic Res Cardiol*. 2013;108:385. doi: 10.1007/s00395-013-0385-6.
22. Pandit SV, Jalife J. Rotors and the dynamics of cardiac fibrillation. *Circ Res*. 2013;112:849–862. doi: 10.1161/CIRCRESAHA.111.300158.
23. Bers DM. *Excitation-Contraction Coupling and Cardiac Contractile Force*. Dordrecht, The Netherlands: Kluwer Academic Publishers; 2001.
24. Weiss JN, Qu Z, Chen PS, Lin SF, Karagueuzian HS, Hayashi H, Garfinkel A, Karma A. The dynamics of cardiac fibrillation. *Circulation*. 2005;112:1232–1240. doi: 10.1161/CIRCULATIONAHA.104.529545.
25. Song Y, Shryock JC, Wagner S, Maier LS, Belardinelli L. Blocking late sodium current reduces hydrogen peroxide-induced arrhythmogenic activity and contractile dysfunction. *J Pharmacol Exp Ther*. 2006;318:214–222. doi: 10.1124/jpet.106.101832.
26. January CT, Riddle JM. Early afterdepolarizations: mechanism of induction and block. A role for L-type Ca²⁺ current. *Circ Res*. 1989;64:977–990.
27. Antoons G, Volders PG, Stankovicova T, Bito V, Stengl M, Vos MA, Sipido KR. Window Ca²⁺ current and its modulation by Ca²⁺ release in hypertrophied cardiac myocytes from dogs with chronic atrioventricular block. *J Physiol*. 2007;579:147–160. doi: 10.1113/jphysiol.2006.124222.
28. Käb S, Nuss HB, Chiamvimonvat N, O'Rourke B, Pak PH, Kass DA, Marban E, Tomaselli GF. Ionic mechanism of action potential prolongation in ventricular myocytes from dogs with pacing-induced heart failure. *Circ Res*. 1996;78:262–273.
29. Beuckelmann DJ, Näbauer M, Erdmann E. Alterations of K⁺ currents in isolated human ventricular myocytes from patients with terminal heart failure. *Circ Res*. 1993;73:379–385.
30. Burashnikov A, Antzelevitch C. Acceleration-induced action potential prolongation and early afterdepolarizations. *J Cardiovasc Electrophysiol*. 1998;9:934–948.
31. Volders GP, Vos MA, Szabo B, Sipido KR, de Groot SH, Gorgels AP, Wellens HJ, Lazzara R. Progress in the understanding of cardiac early afterdepolarizations and torsades de pointes: time to revise current concepts. *Cardiovasc Res*. 2000;46:376–392.
32. Edwards AG, Grandi E, Hake JE, Patel S, Li P, Miyamoto S, Omens JH, Heller Brown J, Bers DM, McCulloch AD. Nonequilibrium reactivation of Na⁺ current drives early afterdepolarizations in mouse ventricle. *Circ Arrhythm Electrophysiol*. 2014;7:1205–1213. doi: 10.1161/CIRCEP.113.001666.
33. Bassani RA, Bers DM. Rate of diastolic Ca release from the sarcoplasmic reticulum of intact rabbit and rat ventricular myocytes. *Biophys J*. 1995;68:2015–2022. doi: 10.1016/S0006-3495(95)80378-4.
34. Bers DM. Cardiac sarcoplasmic reticulum calcium leak: basis and roles in cardiac dysfunction. *Annu Rev Physiol*. 2014;76:107–127. doi: 10.1146/annurev-physiol-020911-153308.
35. Xie Y, Sato D, Garfinkel A, Qu Z, Weiss JN. So little source, so much sink: requirements for afterdepolarizations to propagate in tissue. *Biophys J*. 2010;99:1408–1415. doi: 10.1016/j.bpj.2010.06.042.
36. Myles RC, Wang L, Kang C, Bers DM, Ripplinger CM. Local β -adrenergic stimulation overcomes source-sink mismatch to generate focal arrhythmia. *Circ Res*. 2012;110:1454–1464. doi: 10.1161/CIRCRESAHA.111.262345.
37. Pogwizd SM, Schlotthauer K, Li L, Yuan W, Bers DM. Arrhythmogenesis and contractile dysfunction in heart failure: Roles of sodium-calcium exchange, inward rectifier potassium current, and residual beta-adrenergic responsiveness. *Circ Res*. 2001;88:1159–1167.
38. Myles RC, Wang L, Bers DM, Ripplinger CM. Decreased inward rectifying K⁺ current and increased ryanodine receptor sensitivity synergistically contribute to sustained focal arrhythmia in the intact rabbit heart. *J Physiol*. 2015;593:1479–1493. doi: 10.1113/jphysiol.2014.279638.
39. Walker ML, Rosenbaum DS. Repolarization alternans: implications for the mechanism and prevention of sudden cardiac death. *Cardiovasc Res*. 2003;57:599–614.
40. Comtois P, Nattel S. Atrial repolarization alternans as a path to atrial fibrillation. *J Cardiovasc Electrophysiol*. 2012;23:1013–1015. doi: 10.1111/j.1540-8167.2012.02391.x.
41. Verrier RL, Klingenstein T, Malik M, El-Sherif N, Exner DV, Hohnloser SH, Ikeda T, Martínez JP, Narayan SM, Nieminen T, Rosenbaum DS. Microvolt T-wave alternans testing has a role in arrhythmia risk stratification. *J Am Coll Cardiol*. 2012;59:1572–1573. doi: 10.1016/j.jacc.2012.03.008.
42. Bassani JW, Yuan W, Bers DM. Fractional SR Ca release is regulated by trigger Ca and SR Ca content in cardiac myocytes. *Am J Physiol*. 1995;268:C1313–C1319.
43. Picht E, DeSantiago J, Blatter LA, Bers DM. Cardiac alternans do not rely on diastolic sarcoplasmic reticulum calcium content fluctuations. *Circ Res*. 2006;99:740–748. doi: 10.1161/01.RES.0000244002.88813.91.
44. Wang L, Myles RC, De Jesus NM, Ohlendorf AK, Bers DM, Ripplinger CM. Optical mapping of sarcoplasmic reticulum Ca²⁺ in the intact heart: ryanodine receptor refractoriness during alternans and fibrillation. *Circ Res*. 2014;114:1410–1421. doi: 10.1161/CIRCRESAHA.114.302505.
45. Garfinkel A, Kim YH, Voroshilovsky O, Qu Z, Kil JR, Lee MH, Karagueuzian HS, Weiss JN, Chen PS. Preventing ventricular fibrillation by flattening cardiac restitution. *Proc Natl Acad Sci U S A*. 2000;97:6061–6066. doi: 10.1073/pnas.090492697.
46. Wang Q, Chen Q, Li H, Towbin JA. Molecular genetics of long QT syndrome from genes to patients. *Curr Opin Cardiol*. 1997;12:310–320.
47. Chudin E, Goldhaber J, Garfinkel A, Weiss J, Kogan B. Intracellular Ca(2+) dynamics and the stability of ventricular tachycardia. *Biophys J*. 1999;77:2930–2941. doi: 10.1016/S0006-3495(99)77126-2.
48. Goldhaber JI, Xie LH, Duong T, Motter C, Khuu K, Weiss JN. Action potential duration restitution and alternans in rabbit ventricular myocytes: the key role of intracellular calcium cycling. *Circ Res*. 2005;96:459–466. doi: 10.1161/01.RES.0000156891.66893.83.
49. Díaz ME, O'Neill SC, Eisner DA. Sarcoplasmic reticulum calcium content fluctuation is the key to cardiac alternans. *Circ Res*. 2004;94:650–656. doi: 10.1161/01.RES.0000119923.64774.72.
50. Alvarez-Lacalle E, Cantalapiedra IR, Peñaranda A, Cinca J, Hove-Madsen L, Echebarria B. Dependency of calcium alternans on ryanodine receptor refractoriness. *PLoS One*. 2013;8:e55042. doi: 10.1371/journal.pone.0055042.
51. Pastore JM, Girouard SD, Laurita KR, Akar FG, Rosenbaum DS. Mechanism linking T-wave alternans to the genesis of cardiac fibrillation. *Circulation*. 1999;99:1385–1394.
52. Rosenbaum DS, Jackson LE, Smith JM, Garan H, Ruskin JN, Cohen RJ. Electrical alternans and vulnerability to ventricular arrhythmias. *N Engl J Med*. 1994;330:235–241. doi: 10.1056/NEJM199401273300402.
53. Armoundas AA, Tomaselli GF, Esperer HD. Pathophysiological basis and clinical application of T-wave alternans. *J Am Coll Cardiol*. 2002;40:207–217.
54. Narayan SM. T-wave alternans and the susceptibility to ventricular arrhythmias. *J Am Coll Cardiol*. 2006;47:269–281. doi: 10.1016/j.jacc.2005.08.066.
55. Qu Z, Xie Y, Garfinkel A, Weiss JN. T-wave alternans and arrhythmogenesis in cardiac diseases. *Front Physiol*. 2010;1:154. doi: 10.3389/fphys.2010.00154.
56. Pruvot EJ, Katta RP, Rosenbaum DS, Laurita KR. Role of calcium cycling versus restitution in the mechanism of repolarization alternans. *Circ Res*. 2004;94:1083–1090. doi: 10.1161/01.RES.0000125629.72053.95.
57. Mironov S, Jalife J, Tolkacheva EG. Role of conduction velocity restitution and short-term memory in the development of action potential duration alternans in isolated rabbit hearts. *Circulation*. 2008;118:17–25. doi: 10.1161/CIRCULATIONAHA.107.737254.
58. Christini DJ, Riccio ML, Culianu CA, Fox JJ, Karma A, Gilmour RF Jr. Control of electrical alternans in canine cardiac purkinje fibers. *Phys Rev Lett*. 2006;96:104101.
59. Qu Z, Karagueuzian HS, Garfinkel A, Weiss JN. Effects of Na(+) channel and cell coupling abnormalities on vulnerability to reentry: a simulation study. *Am J Physiol Heart Circ Physiol*. 2004;286:H1310–H1321. doi: 10.1152/ajpheart.00561.2003.
60. Koval OM, Snyder JS, Wolf RM, et al. Ca²⁺/calmodulin-dependent protein kinase II-based regulation of voltage-gated Na⁺ channel in cardiac disease. *Circulation*. 2012;126:2084–2094. doi: 10.1161/CIRCULATIONAHA.112.105320.

61. Pu J, Boyden PA. Alterations of Na⁺ currents in myocytes from epicardial border zone of the infarcted heart. A possible ionic mechanism for reduced excitability and postrepolarization refractoriness. *Circ Res*. 1997;81:110–119.
62. Joyner RW, Ramza BM, Osaka T, Tan RC. Cellular mechanisms of delayed recovery of excitability in ventricular tissue. *Am J Physiol*. 1991;260:H225–H233.
63. Wang DW, Makita N, Kitabatake A, Balser JR, George AL Jr. Enhanced Na⁺ channel intermediate inactivation in Brugada syndrome. *Circ Res*. 2000;87:E37–E43.
64. Hayashi H, Shiferaw Y, Sato D, Nihei M, Lin SF, Chen PS, Garfinkel A, Weiss JN, Qu Z. Dynamic origin of spatially discordant alternans in cardiac tissue. *Biophys J*. 2007;92:448–460. doi: 10.1529/biophysj.106.091009.
65. Cao JM, Qu Z, Kim YH, Wu TJ, Garfinkel A, Weiss JN, Karagueuzian HS, Chen PS. Spatiotemporal heterogeneity in the induction of ventricular fibrillation by rapid pacing: importance of cardiac restitution properties. *Circ Res*. 1999;84:1318–1331.
66. Wu TJ, Lin SF, Baher A, Qu Z, Garfinkel A, Weiss JN, Ting CT, Chen PS. Mother rotors and the mechanisms of D600-induced type 2 ventricular fibrillation. *Circulation*. 2004;110:2110–2118. doi: 10.1161/01.CIR.0000143834.51102.91.
67. Qu Z, Nivala M, Weiss JN. Calcium alternans in cardiac myocytes: order from disorder. *J Mol Cell Cardiol*. 2013;58:100–109. doi: 10.1016/j.yjmcc.2012.10.007.
68. Kornyevev D, Petrosky AD, Zepeda B, Ferreiro M, Knollmann B, Escobar AL. Calsequestrin 2 deletion shortens the refractoriness of Ca²⁺ release and reduces rate-dependent Ca²⁺-alternans in intact mouse hearts. *J Mol Cell Cardiol*. 2012;52:21–31. doi: 10.1016/j.yjmcc.2011.09.020.
69. Shannon TR, Ginsburg KS, Bers DM. Potentiation of fractional sarcoplasmic reticulum calcium release by total and free intra-sarcoplasmic reticulum calcium concentration. *Biophys J*. 2000;78:334–343. doi: 10.1016/S0006-3495(00)76596-9.
70. Hüser J, Wang YG, Sheehan KA, Cifuentes F, Lipsius SL, Blatter LA. Functional coupling between glycolysis and excitation-contraction coupling underlies alternans in cat heart cells. *J Physiol*. 2000;524 Pt 3:795–806.
71. Florea SM, Blatter LA. The role of mitochondria for the regulation of cardiac alternans. *Front Physiol*. 2010;1:141. doi: 10.3389/fphys.2010.00141.
72. Cutler MJ, Wan X, Laurita KR, Hajjar RJ, Rosenbaum DS. Targeted SERCA2a gene expression identifies molecular mechanism and therapeutic target for arrhythmogenic cardiac alternans. *Circ Arrhythm Electrophysiol*. 2009;2:686–694. doi: 10.1161/CIRCEP.109.863118.
73. Belevych AE, Terentyev D, Viatchenko-Karpinski S, Terentyeva R, Sridhar A, Nishijima Y, Wilson LD, Cardounel AJ, Laurita KR, Carnes CA, Billman GE, Gyorke S. Redox modification of ryanodine receptors underlies calcium alternans in a canine model of sudden cardiac death. *Cardiovasc Res*. 2009;84:387–395. doi: 10.1093/cvr/cvp246.
74. Aistrup GL, Kelly JE, Kapur S, Kowalczyk M, Sysman-Wolpin I, Kadish AH, Wasserstrom JA. Pacing-induced heterogeneities in intracellular Ca²⁺ signaling, cardiac alternans, and ventricular arrhythmias in intact rat heart. *Circ Res*. 2006;99:e65–e73. doi: 10.1161/01.RES.0000244087.36230.bf.
75. Aistrup GL, Shiferaw Y, Kapur S, Kadish AH, Wasserstrom JA. Mechanisms underlying the formation and dynamics of subcellular calcium alternans in the intact rat heart. *Circ Res*. 2009;104:639–649. doi: 10.1161/CIRCRESAHA.108.181909.
76. Tian Q, Kaestner L, Lipp P. Noise-free visualization of microscopic calcium signaling by pixel-wise fitting. *Circ Res*. 2012;111:17–27. doi: 10.1161/CIRCRESAHA.112.266403.
77. Gehi AK, Stein RH, Metz LD, Gomes JA. Microvolt T-wave alternans for the risk stratification of ventricular tachyarrhythmic events: a meta-analysis. *J Am Coll Cardiol*. 2005;46:75–82. doi: 10.1016/j.jacc.2005.03.059.
78. Tsutsui H, Ide T, Hayashidani S, Suematsu N, Utsumi H, Nakamura R, Egashira K, Takeshita A. Greater susceptibility of failing cardiac myocytes to oxygen free radical-mediated injury. *Cardiovasc Res*. 2001;49:103–109.
79. Kinugawa S, Tsutsui H, Hayashidani S, Ide T, Suematsu N, Satoh S, Utsumi H, Takeshita A. Treatment with dimethylthiourea prevents left ventricular remodeling and failure after experimental myocardial infarction in mice: role of oxidative stress. *Circ Res*. 2000;87:392–398.
80. Ide T, Tsutsui H, Kinugawa S, Suematsu N, Hayashidani S, Ichikawa K, Utsumi H, Machida Y, Egashira K, Takeshita A. Direct evidence for increased hydroxyl radicals originating from superoxide in the failing myocardium. *Circ Res*. 2000;86:152–157.
81. Mallat Z, Philip I, Lebreton M, Chatel D, Maclouf J, Tedgui A. Elevated levels of 8-iso-prostaglandin F₂α in pericardial fluid of patients with heart failure: a potential role for in vivo oxidant stress in ventricular dilatation and progression to heart failure. *Circulation*. 1998;97:1536–1539.
82. Maltsev VA, Sabbah HN, Higgins RS, Silverman N, Lesch M, Undrovinas AI. Novel, ultraslow inactivating sodium current in human ventricular cardiomyocytes. *Circulation*. 1998;98:2545–2552.
83. Huang B, El-Sherif T, Gidh-Jain M, Qin D, El-Sherif N. Alterations of sodium channel kinetics and gene expression in the postinfarction remodeled myocardium. *J Cardiovasc Electrophysiol*. 2001;12:218–225.
84. Maltsev VA, Undrovinas AI. A multi-modal composition of the late Na⁺ current in human ventricular cardiomyocytes. *Cardiovasc Res*. 2006;69:116–127. doi: 10.1016/j.cardiores.2005.08.015.
85. Moreno JD, Clancy CE. Pathophysiology of the cardiac late Na current and its potential as a drug target. *J Mol Cell Cardiol*. 2012;52:608–619. doi: 10.1016/j.yjmcc.2011.12.003.
86. Saint DA, Ju YK, Gage PW. A persistent sodium current in rat ventricular myocytes. *J Physiol*. 1992;453:219–231.
87. Pieske B, Maier LS, Piacentino V III, Weisser J, Hasenfuss G, Houser S. Rate dependence of [Na⁺], and contractility in nonfailing and failing human myocardium. *Circulation*. 2002;106:447–453.
88. Despa S, Islam MA, Weber CR, Pogwizd SM, Bers DM. Intracellular Na⁺ concentration is elevated in heart failure but Na/K pump function is unchanged. *Circulation*. 2002;105:2543–2548.
89. Sossalla S, Wagner S, Rasenack EC, Ruff H, Weber SL, Schöndube FA, Tirilomis T, Tenderich G, Hasenfuss G, Belardinelli L, Maier LS. Ranolazine improves diastolic dysfunction in isolated myocardium from failing human hearts—role of late sodium current and intracellular ion accumulation. *J Mol Cell Cardiol*. 2008;45:32–43. doi: 10.1016/j.yjmcc.2008.03.006.
90. Sossalla S, Maurer U, Schotola H, Hartmann N, Didié M, Zimmermann WH, Jacobshagen C, Wagner S, Maier LS. Diastolic dysfunction and arrhythmias caused by overexpression of CaMKIIδ can be reversed by inhibition of late Na⁺ current. *Basic Res Cardiol*. 2011;106:263–272. doi: 10.1007/s00395-010-0136-x.
91. Ward CA, Giles WR. Ionic mechanism of the effects of hydrogen peroxide in rat ventricular myocytes. *J Physiol*. 1997;500 (Pt 3):631–642.
92. Dumaine R, Wang Q, Keating MT, Hartmann HA, Schwartz PJ, Brown AM, Kirsch GE. Multiple mechanisms of Na⁺ channel-linked long-QT syndrome. *Circ Res*. 1996;78:916–924.
93. Viswanathan PC, Balser JR. Inherited sodium channelopathies: a continuum of channel dysfunction. *Trends Cardiovasc Med*. 2004;14:28–35.
94. Antzelevitch C, Belardinelli L. The role of sodium channel current in modulating transmural dispersion of repolarization and arrhythmogenesis. *J Cardiovasc Electrophysiol*. 2006;17(suppl 1):S79–S85. doi: 10.1111/j.1540-8167.2006.00388.x.
95. Bonatti R, Silva AF, Batatinha JA, Sobrado LF, Machado AD, Varone BB, Nearing BD, Belardinelli L, Verrier RL. Selective late sodium current blockade with GS-458967 markedly reduces ischemia-induced atrial and ventricular repolarization alternans and ECG heterogeneity. *Heart Rhythm*. 2014;11:1827–1835. doi: 10.1016/j.hrthm.2014.06.017.
96. Wagner S, Seidler T, Picht E, Maier LS, Kazanski V, Teucher N, Schillinger W, Pieske B, Isenberg G, Hasenfuss G, Kögler H. Na⁺-Ca²⁺ exchanger overexpression predisposes to reactive oxygen species-induced injury. *Cardiovasc Res*. 2003;60:404–412.
97. Fredj S, Lindegar N, Sampson KJ, Carmeliet P, Kass RS. Altered Na⁺ channels promote pause-induced spontaneous diastolic activity in long QT syndrome type 3 myocytes. *Circ Res*. 2006;99:1225–1232. doi: 10.1161/01.RES.0000251305.25604.b0.
98. Nagy N, Kormos A, Kohajda Z, Szebeni Á, Szepesi J, Pollesello P, Levijoki J, Acsai K, Virág L, Nánási PP, Papp JG, Varró A, Tóth A. Selective Na⁺/Ca²⁺ exchanger inhibition prevents Ca²⁺ overload-induced triggered arrhythmias. *Br J Pharmacol*. 2014;171:5665–5681. doi: 10.1111/bph.12867.
99. Ahern GP, Hsu SF, Klyachko VA, Jackson MB. Induction of persistent sodium current by exogenous and endogenous nitric oxide. *J Biol Chem*. 2000;275:28810–28815. doi: 10.1074/jbc.M003090200.
100. Maltsev VA, Reznikov V, Undrovinas NA, Sabbah HN, Undrovinas A. Modulation of late sodium current by Ca²⁺, calmodulin, and CaMKII in normal and failing dog cardiomyocytes: similarities and differences. *Am J Physiol Heart Circ Physiol*. 2008;294:H1597–H1608. doi: 10.1152/ajpheart.00484.2007.
101. Aiba T, Hesketh GG, Liu T, Carlisle R, Villa-Abrille MC, O'Rourke B, Akar FG, Tomaselli GF. Na⁺ channel regulation by Ca²⁺/calmodulin and Ca²⁺/calmodulin-dependent protein kinase II in guinea-pig ventricular myocytes. *Cardiovasc Res*. 2010;85:454–463. doi: 10.1093/cvr/cvp324.
102. Yao L, Fan P, Jiang Z, Viatchenko-Karpinski S, Wu Y, Kornyevev D, Hirakawa R, Budas GR, Rajamani S, Shryock JC, Belardinelli

- L. Nav1.5-dependent persistent Na⁺ influx activates CaMKII in rat ventricular myocytes and N1325S mice. *Am J Physiol Cell Physiol*. 2011;301:C577–C586. doi: 10.1152/ajpcell.00125.2011.
103. Ashpole NM, Herren AW, Ginsburg KS, Brogan JD, Johnson DE, Cummins TR, Bers DM, Hudmon A. Ca²⁺/calmodulin-dependent protein kinase II (CaMKII) regulates cardiac sodium channel NaV1.5 gating by multiple phosphorylation sites. *J Biol Chem*. 2012;287:19856–19869. doi: 10.1074/jbc.M111.322537.
 104. Hund TJ, Koval OM, Li J, Wright PJ, Qian L, Snyder JS, Gudmundsson H, Kline CF, Davidson NP, Cardona N, Rasband MN, Anderson ME, Mohler PJ. A β (IV)-spectrin/CaMKII signaling complex is essential for membrane excitability in mice. *J Clin Invest*. 2010;120:3508–3519. doi: 10.1172/JCI43621.
 105. Undrovinas AI, Belardinelli L, Undrovinas NA, Sabbah HN. Ranolazine improves abnormal repolarization and contraction in left ventricular myocytes of dogs with heart failure by inhibiting late sodium current. *J Cardiovasc Electrophysiol*. 2006;17(suppl 1):S169–S177. doi: 10.1111/j.1540-8167.2006.00401.x.
 106. Antzelevitch C, Belardinelli L, Zygmunt AC, Burashnikov A, Di Diego JM, Fish JM, Cordeiro JM, Thomas G. Electrophysiological effects of ranolazine, a novel antianginal agent with antiarrhythmic properties. *Circulation*. 2004;110:904–910. doi: 10.1161/01.CIR.0000139333.83620.5D.
 107. Sossalla S, Maurer U, Schotola H, Hartmann N, Didié M, Zimmermann WH, Jacobshagen C, Wagner S, Maier LS. Diastolic dysfunction and arrhythmias caused by overexpression of CaMKII δ can be reversed by inhibition of late Na⁺ current. *Basic Res Cardiol*. 2011;106:263–272. doi: 10.1007/s00395-010-0136-x.
 108. Hwang H, Arcidi JM Jr, Hale SL, Simkhovich BZ, Belardinelli L, Dhalla AK, Shryock JC, Kloner RA. Ranolazine as a cardioplegia additive improves recovery of diastolic function in isolated rat hearts. *Circulation*. 2009;120:S16–S21. doi: 10.1161/CIRCULATIONAHA.108.844167.
 109. Song Y, Shryock JC, Wu L, Belardinelli L. Antagonism by ranolazine of the pro-arrhythmic effects of increasing late I_{Na} in guinea pig ventricular myocytes. *J Cardiovasc Pharmacol*. 2004;44:192–199.
 110. Wu L, Shryock JC, Song Y, Li Y, Antzelevitch C, Belardinelli L. Antiarrhythmic effects of ranolazine in a guinea pig in vitro model of long-QT syndrome. *J Pharmacol Exp Ther*. 2004;310:599–605. doi: 10.1124/jpet.104.066100.
 111. Dhalla AK, Wang WQ, Dow J, Shryock JC, Belardinelli L, Bhandari A, Kloner RA. Ranolazine, an antianginal agent, markedly reduces ventricular arrhythmias induced by ischemia and ischemia-reperfusion. *Am J Physiol Heart Circ Physiol*. 2009;297:H1923–H1929. doi: 10.1152/ajpheart.00173.2009.
 112. Morita N, Lee JH, Xie Y, Sovari A, Qu Z, Weiss JN, Karagueuzian HS. Suppression of re-entrant and multifocal ventricular fibrillation by the late sodium current blocker ranolazine. *J Am Coll Cardiol*. 2011;57:366–375. doi: 10.1016/j.jacc.2010.07.045.
 113. Rastogi S, Sharov VG, Mishra S, Gupta RC, Blackburn B, Belardinelli L, Stanley WC, Sabbah HN. Ranolazine combined with enalapril or metoprolol prevents progressive LV dysfunction and remodeling in dogs with moderate heart failure. *Am J Physiol Heart Circ Physiol*. 2008;295:H2149–H2155. doi: 10.1152/ajpheart.00728.2008.
 114. Morrow DA, Scirica BM, Karwowska-Prokopczuk E, Murphy SA, Budaj A, Varshavsky S, Wolff AA, Skene A, McCabe CH, Braunwald E; MERLIN-TIMI 36 Trial Investigators. Effects of ranolazine on recurrent cardiovascular events in patients with non-ST-elevation acute coronary syndromes: the MERLIN-TIMI 36 randomized trial. *JAMA*. 2007;297:1775–1783. doi: 10.1001/jama.297.16.1775.
 115. Karwowska-Prokopczuk E, Wang W, Cheng ML, Zeng D, Schwartz PJ, Belardinelli L. The risk of sudden cardiac death in patients with non-ST elevation acute coronary syndrome and prolonged QTc interval: effect of ranolazine. *Europace*. 2013;15:429–436. doi: 10.1093/europace/eus400.
 116. Scirica BM, Morrow DA, Hod H, Murphy SA, Belardinelli L, Hedgepeth CM, Molhoek P, Verheugt FW, Gersh BJ, McCabe CH, Braunwald E. Effect of ranolazine, an antianginal agent with novel electrophysiological properties, on the incidence of arrhythmias in patients with non ST-segment elevation acute coronary syndrome: results from the Metabolic Efficiency With Ranolazine for Less Ischemia in Non ST-Elevation Acute Coronary Syndrome Thrombolysis in Myocardial Infarction 36 (MERLIN-TIMI 36) randomized controlled trial. *Circulation*. 2007;116:1647–1652. doi: 10.1161/CIRCULATIONAHA.107.724880.
 117. Bunch TJ, Mahapatra S, Murdock D, Molden J, Weiss JP, May HT, Bair TL, Mader KM, Crandall BG, Day JD, Osborn JS, Muhlestein JB, Lappe DL, Anderson JL. Ranolazine reduces ventricular tachycardia burden and ICD shocks in patients with drug-refractory ICD shocks. *Pacing Clin Electrophysiol*. 2011;34:1600–1606. doi: 10.1111/j.1540-8159.2011.03208.x.
 118. Moss AJ, Zareba W, Schwarz KQ, Rosero S, McNitt S, Robinson JL. Ranolazine shortens repolarization in patients with sustained inward sodium current due to type-3 long-QT syndrome. *J Cardiovasc Electrophysiol*. 2008;19:1289–1293. doi: 10.1111/j.1540-8167.2008.01246.x.
 119. De Ferrari GM, Maier LS, Mont L, et al.; RAFFAELLO Investigators (see Online Supplementary Appendix for List of Participating Centers and Investigators). Ranolazine in the treatment of atrial fibrillation: Results of the dose-ranging RAFFAELLO (Ranolazine in Atrial Fibrillation Following An Electrical Cardioversion) study. *Heart Rhythm*. 2015;12:872–878. doi: 10.1016/j.hrthm.2015.01.021.
 120. Maier LS, Zhang T, Chen L, DeSantiago J, Brown JH, Bers DM. Transgenic CaMKII δ overexpression uniquely alters cardiac myocyte Ca²⁺ handling: reduced SR Ca²⁺ load and activated SR Ca²⁺ release. *Circ Res*. 2003;92:904–911. doi: 10.1161/01.RES.0000069685.20258.F1.
 121. Qi X, Yeh YH, Chartier D, Xiao L, Tsuji Y, Brundel BJ, Kodama I, Nattel S. The calcium/calmodulin/kinase system and arrhythmogenic afterdepolarizations in bradycardia-related acquired long-QT syndrome. *Circ Arrhythm Electrophysiol*. 2009;2:295–304. doi: 10.1161/CIRCEP.108.815654.
 122. Tsuji Y, Hojo M, Voigt N, El-Armouche A, Inden Y, Murohara T, Dobrev D, Nattel S, Kodama I, Kamiya K. Ca(2+)-related signaling and protein phosphorylation abnormalities play central roles in a new experimental model of electrical storm. *Circulation*. 2011;123:2192–2203. doi: 10.1161/CIRCULATIONAHA.110.016683.
 123. Zhang R, Khoo MS, Wu Y, et al. Calmodulin kinase II inhibition protects against structural heart disease. *Nat Med*. 2005;11:409–417. doi: 10.1038/nm1215.
 124. Ling H, Zhang T, Pereira L, Means CK, Cheng H, Gu Y, Dalton ND, Peterson KL, Chen J, Bers D, Brown JH, Heller Brown J. Requirement for Ca²⁺/calmodulin-dependent kinase II in the transition from pressure overload-induced cardiac hypertrophy to heart failure in mice. *J Clin Invest*. 2009;119:1230–1240. doi: 10.1172/JCI38022.
 125. Khoo MS, Li J, Singh MV, et al. Death, cardiac dysfunction, and arrhythmias are increased by calmodulin kinase II in calcineurin cardiomyopathy. *Circulation*. 2006;114:1352–1359. doi: 10.1161/CIRCULATIONAHA.106.644583.
 126. Xie LH, Chen F, Karagueuzian HS, Weiss JN. Oxidative-stress-induced afterdepolarizations and calmodulin kinase II signaling. *Circ Res*. 2009;104:79–86. doi: 10.1161/CIRCRESAHA.108.183475.
 127. Wagner S, Dantz C, Flebbe H, Azizian A, Sag CM, Engels S, Möllencamp J, Dybkova N, Islam T, Shah AM, Maier LS. NADPH oxidase 2 mediates angiotensin II-dependent cellular arrhythmias via PKA and CaMKII. *J Mol Cell Cardiol*. 2014;75:206–215. doi: 10.1016/j.yjmcc.2014.07.011.
 128. Zhao Z, Fefelova N, Shanmugam M, Bishara P, Babu GJ, Xie LH. Angiotensin II induces afterdepolarizations via reactive oxygen species and calmodulin kinase II signaling. *J Mol Cell Cardiol*. 2011;50:128–136. doi: 10.1016/j.yjmcc.2010.11.001.
 129. Luo M, Guan X, Luczak ED, et al. Diabetes increases mortality after myocardial infarction by oxidizing CaMKII. *J Clin Invest*. 2013;123:1262–1274. doi: 10.1172/JCI65268.
 130. Erickson JR, Pereira L, Wang L, Han G, Ferguson A, Dao K, Copeland RJ, Despa F, Hart GW, Ripplinger CM, Bers DM. Diabetic hyperglycaemia activates CaMKII and arrhythmias by O-linked glycosylation. *Nature*. 2013;502:372–376. doi: 10.1038/nature12537.
 131. Coultrap SJ, Bayer KU. Nitric oxide induces Ca²⁺-independent activity of the Ca²⁺/calmodulin-dependent protein kinase II (CaMKII). *J Biol Chem*. 2014;289:19458–19465. doi: 10.1074/jbc.M114.558254.
 132. Gutierrez DA, Fernandez-Tenorio M, Ogradnik J, Niggli E. NO-dependent CaMKII activation during β -adrenergic stimulation of cardiac muscle. *Cardiovasc Res*. 2013;100:392–401. doi: 10.1093/cvr/cvt201.
 133. Jian Z, Han H, Zhang T, et al. Mechanochemotransduction during cardiomyocyte contraction is mediated by localized nitric oxide signaling. *Sci Signal*. 2014;7:ra27. doi: 10.1126/scisignal.2005046.
 134. Dybkova N, Wagner S, Backs J, Hund TJ, Mohler PJ, Sowa T, Nikolaev VO, Maier LS. Tubulin polymerization disrupts cardiac β -adrenergic regulation of late I_{Na}. *Cardiovasc Res*. 2014;103:168–177. doi: 10.1093/cvr/cvu120.
 135. Mangmool S, Shukla AK, Rockman HA. beta-Arrestin-dependent activation of Ca(2+)/calmodulin kinase II after beta(1)-adrenergic receptor stimulation. *J Cell Biol*. 2010;189:573–587. doi: 10.1083/jcb.200911047.

136. Pereira L, Cheng H, Lao DH, Na L, van Oort RJ, Brown JH, Wehrens XH, Chen J, Bers DM. Epac2 mediates cardiac β 1-adrenergic-dependent sarcoplasmic reticulum Ca^{2+} leak and arrhythmia. *Circulation*. 2013;127:913–922. doi: 10.1161/CIRCULATIONAHA.12.148619.
137. Erickson JR, Patel R, Ferguson A, Bossuyt J, Bers DM. Fluorescence resonance energy transfer-based sensor Camui provides new insight into mechanisms of calcium/calmodulin-dependent protein kinase II activation in intact cardiomyocytes. *Circ Res*. 2011;109:729–738. doi: 10.1161/CIRCRESAHA.111.247148.
138. Anderson ME, Braun AP, Schulman H, Premack BA. Multifunctional Ca^{2+} /calmodulin-dependent protein kinase mediates Ca^{2+} -induced enhancement of the L-type Ca^{2+} current in rabbit ventricular myocytes. *Circ Res*. 1994;75:854–861.
139. Xiao RP, Cheng H, Lederer WJ, Suzuki T, Lakatta EG. Dual regulation of Ca^{2+} /calmodulin-dependent kinase II activity by membrane voltage and by calcium influx. *Proc Natl Acad Sci U S A*. 1994;91:9659–9663.
140. Yuan W, Bers DM. Ca^{2+} -dependent facilitation of cardiac Ca^{2+} current is due to Ca^{2+} -calmodulin-dependent protein kinase. *Am J Physiol*. 1994;267:H982–H993.
141. Anderson ME, Braun AP, Wu Y, Lu T, Wu Y, Schulman H, Sung RJ. KN-93, an inhibitor of multifunctional Ca^{2+} /calmodulin-dependent protein kinase, decreases early afterdepolarizations in rabbit heart. *J Pharmacol Exp Ther*. 1998;287:996–1006.
142. Wu Y, MacMillan LB, McNeill RB, Colbran RJ, Anderson ME. Ca^{2+} kinase augments cardiac L-type Ca^{2+} current: a cellular mechanism for long Q-T arrhythmias. *Am J Physiol*. 1999;276:H2168–H2178.
143. Koval OM, Guan X, Wu Y, Joiner ML, Gao Z, Chen B, Grumbach IM, Luczak ED, Colbran RJ, Song LS, Hund TJ, Mohler PJ, Anderson ME. $\text{CaV}1.2$ beta-subunit coordinates CaMKII-triggered cardiomyocyte death and afterdepolarizations. *Proc Natl Acad Sci U S A*. 2010;107:4996–5000. doi: 10.1073/pnas.0913760107.
144. Bezzina C, Veldkamp MW, van Den Berg MP, Postma AV, Rook MB, Viersma JW, van Langen IM, Tan-Sindhunata G, Bink-Boelkens MT, van Der Hout AH, Mannens MM, Wilde AA. A single Na^{+} channel mutation causing both long-QT and Brugada syndromes. *Circ Res*. 1999;85:1206–1213.
145. Veldkamp MW, Viswanathan PC, Bezzina C, Baartscheer A, Wilde AA, Balser JR. Two distinct congenital arrhythmias evoked by a multidysfunctional Na^{+} channel. *Circ Res*. 2000;86:E91–E97.
146. Wehrens XH, Lehnart SE, Reiken SR, Marks AR. Ca^{2+} /calmodulin-dependent protein kinase II phosphorylation regulates the cardiac ryanodine receptor. *Circ Res*. 2004;94:e61–e70. doi: 10.1161/01.RES.0000125626.33738.E2.
147. Simmerman HK, Collins JH, Theibert JL, Wegener AD, Jones LR. Sequence analysis of phospholamban. Identification of phosphorylation sites and two major structural domains. *J Biol Chem*. 1986;261:13333–13341.
148. Bers DM, Eisner DA, Valdivia HH. Sarcoplasmic reticulum Ca^{2+} and heart failure: roles of diastolic leak and Ca^{2+} transport. *Circ Res*. 2003;93:487–490. doi: 10.1161/01.RES.0000091871.54907.6B.
149. Lindner M, Erdmann E, Beuckelmann DJ. Calcium content of the sarcoplasmic reticulum in isolated ventricular myocytes from patients with terminal heart failure. *J Mol Cell Cardiol*. 1998;30:743–749. doi: 10.1006/jmcc.1997.0626.
150. Ai X, Curran JW, Shannon TR, Bers DM, Pogwizd SM. Ca^{2+} /calmodulin-dependent protein kinase modulates cardiac ryanodine receptor phosphorylation and sarcoplasmic reticulum Ca^{2+} leak in heart failure. *Circ Res*. 2005;97:1314–1322. doi: 10.1161/01.RES.0000194329.41863.89.
151. Curran J, Brown KH, Santiago DJ, Pogwizd S, Bers DM, Shannon TR. Spontaneous Ca^{2+} waves in ventricular myocytes from failing hearts depend on Ca^{2+} -calmodulin-dependent protein kinase II. *J Mol Cell Cardiol*. 2010;49:25–32. doi: 10.1016/j.jmcc.2010.03.013.
152. Zhang T, Guo T, Mishra S, Dalton ND, Kranias EG, Peterson KL, Bers DM, Brown JH. Phospholamban ablation rescues sarcoplasmic reticulum Ca^{2+} handling but exacerbates cardiac dysfunction in CaMKII δ transgenic mice. *Circ Res*. 2010;106:354–362. doi: 10.1161/CIRCRESAHA.109.207423.
153. Respress JL, van Oort RJ, Li N, et al. Role of RyR2 phosphorylation at S2814 during heart failure progression. *Circ Res*. 2012;110:1474–1483. doi: 10.1161/CIRCRESAHA.112.268094.
154. van Oort RJ, McCauley MD, Dixit SS, Pereira L, Yang Y, Respress JL, Wang Q, De Almeida AC, Skapura DG, Anderson ME, Bers DM, Wehrens XH. Ryanodine receptor phosphorylation by calcium/calmodulin-dependent protein kinase II promotes life-threatening ventricular arrhythmias in mice with heart failure. *Circulation*. 2010;122:2669–2679. doi: 10.1161/CIRCULATIONAHA.110.982298.
155. Fischer TH, Eiringhaus J, Dybkova N, Förster A, Herting J, Kleinwächter A, Ljubojevic S, Schmitto JD, Streckfuß-Bömeke K, Renner A, Gummert J, Hasenfuss G, Maier LS, Sossalla S. Ca^{2+} /calmodulin-dependent protein kinase II equally induces sarcoplasmic reticulum Ca^{2+} leak in human ischaemic and dilated cardiomyopathy. *Eur J Heart Fail*. 2014;16:1292–1300. doi: 10.1002/ehf.163.
156. Oestreich EA, Malik S, Goonasekera SA, Blaxall BC, Kelley GG, Dirksen RT, Smrcka AV. Epac and phospholipase Cepsilon regulate Ca^{2+} release in the heart by activation of protein kinase Cepsilon and calcium-calmodulin kinase II. *J Biol Chem*. 2009;284:1514–1522. doi: 10.1074/jbc.M806994200.
157. Curran J, Hinton MJ, Ríos E, Bers DM, Shannon TR. Beta-adrenergic enhancement of sarcoplasmic reticulum calcium leak in cardiac myocytes is mediated by calcium/calmodulin-dependent protein kinase. *Circ Res*. 2007;100:391–398. doi: 10.1161/01.RES.0000258172.74570.e6.
158. Pereira L, Métrich M, Fernández-Velasco M, Lucas A, Leroy J, Perrier R, Morel E, Fischmeister R, Richard S, Bénitah JP, Lezoualc'h F, Gómez AM. The cAMP binding protein Epac modulates Ca^{2+} sparks by a Ca^{2+} /calmodulin kinase signalling pathway in rat cardiac myocytes. *J Physiol*. 2007;583:685–694. doi: 10.1113/jphysiol.2007.133066.
159. Neef S, Dybkova N, Sossalla S, Ort KR, Fluschnik N, Neumann K, Seipelt R, Schöndube FA, Hasenfuss G, Maier LS. CaMKII-dependent diastolic SR Ca^{2+} leak and elevated diastolic Ca^{2+} levels in right atrial myocardium of patients with atrial fibrillation. *Circ Res*. 2010;106:1134–1144. doi: 10.1161/CIRCRESAHA.109.203836.
160. Purohit A, Rokita AG, Guan X, et al. Oxidized Ca^{2+} /calmodulin-dependent protein kinase II triggers atrial fibrillation. *Circulation*. 2013;128:1748–1757. doi: 10.1161/CIRCULATIONAHA.113.003313.
161. Voigt N, Li N, Wang Q, Wang W, Trafford AW, Abu-Taha I, Sun Q, Wieland T, Ravens U, Nattel S, Wehrens XH, Dobrev D. Enhanced sarcoplasmic reticulum Ca^{2+} leak and increased Na^{+} - Ca^{2+} exchanger function underlie delayed afterdepolarizations in patients with chronic atrial fibrillation. *Circulation*. 2012;125:2059–2070. doi: 10.1161/CIRCULATIONAHA.111.067306.
162. Li N, Wang T, Wang W, Cutler MJ, Wang Q, Voigt N, Rosenbaum DS, Dobrev D, Wehrens XH. Inhibition of CaMKII phosphorylation of RyR2 prevents induction of atrial fibrillation in FKBP12.6 knockout mice. *Circ Res*. 2012;110:465–470. doi: 10.1161/CIRCRESAHA.111.253229.
163. Xu L, Eu JP, Meissner G, Stamler JS. Activation of the cardiac calcium release channel (ryanodine receptor) by poly-S-nitrosylation. *Science*. 1998;279:234–237.
164. Prosser BL, Ward CW, Lederer WJ. X-ROS signaling: rapid mechano-chemo transduction in heart. *Science*. 2011;333:1440–1445. doi: 10.1126/science.1202768.
165. Bovo E, Lipsius SL, Zima AV. Reactive oxygen species contribute to the development of arrhythmogenic Ca^{2+} waves during β -adrenergic receptor stimulation in rabbit cardiomyocytes. *J Physiol*. 2012;590:3291–3304. doi: 10.1113/jphysiol.2012.230748.
166. Leenhardt A, Lucet V, Denjoy I, Grau F, Ngoc DD, Coumel P. Catecholaminergic polymorphic ventricular tachycardia in children. A 7-year follow-up of 21 patients. *Circulation*. 1995;91:1512–1519.
167. Hayashi M, Denjoy I, Extramiana F, Maltret A, Buisson NR, Lupoglazoff JM, Klug D, Hayashi M, Takatsuki S, Villain E, Kamblock J, Messali A, Guicheney P, Lunardi J, Leenhardt A. Incidence and risk factors of arrhythmic events in catecholaminergic polymorphic ventricular tachycardia. *Circulation*. 2009;119:2426–2434. doi: 10.1161/CIRCULATIONAHA.108.829267.
168. Sy RW, Gollob MH, Klein GJ, Yee R, Skanes AC, Gula LJ, Leong-Sit P, Gow RM, Green MS, Birmie DH, Krahn AD. Arrhythmia characterization and long-term outcomes in catecholaminergic polymorphic ventricular tachycardia. *Heart Rhythm*. 2011;8:864–871. doi: 10.1016/j.hrthm.2011.01.048.
169. Priori SG, Napolitano C, Tiso N, Memmi M, Vignati G, Bloise R, Sorrentino V, Danieli GA. Mutations in the cardiac ryanodine receptor gene (hRyR2) underlie catecholaminergic polymorphic ventricular tachycardia. *Circulation*. 2001;103:196–200.
170. Dalal D, Nasir K, Bomma C, Prakasa K, Tandri H, Piccini J, Roguin A, Tichnell C, James C, Russell SD, Judge DP, Abraham T, Spevak PJ, Blumek DA, Calkins H. Arrhythmogenic right ventricular dysplasia: a United States experience. *Circulation*. 2005;112:3823–3832. doi: 10.1161/CIRCULATIONAHA.105.542266.
171. Lahat H, Pras E, Olender T, Avidan N, Ben-Asher E, Man O, Levy-Nissenbaum E, Khoury A, Lorber A, Goldman B, Lancet D, Eldar M. A missense mutation in a highly conserved region of CASQ2 is associated with autosomal recessive catecholamine-induced polymorphic ventricular tachycardia in Bedouin families from Israel. *Am J Hum Genet*. 2001;69:1378–1384. doi: 10.1086/324565.

172. Roux-Buisson N, Cacheux M, Fourest-Lieuvin A, et al. Absence of triadin, a protein of the calcium release complex, is responsible for cardiac arrhythmia with sudden death in human. *Hum Mol Genet.* 2012;21:2759–2767. doi: 10.1093/hmg/dds104.
173. Sedej S, Heinzel FR, Walther S, Dybkova N, Wakula P, Groborz J, Gronau P, Maier LS, Vos MA, Lai FA, Napolitano C, Priori SG, Kockskämper J, Pieske B. Na⁺-dependent SR Ca²⁺ overload induces arrhythmogenic events in mouse cardiomyocytes with a human CPVT mutation. *Cardiovasc Res.* 2010;87:50–59. doi: 10.1093/cvr/cvq007.
174. Liu N, Ruan Y, Denegri M, Bachetti T, Li Y, Colombi B, Napolitano C, Coetzee WA, Priori SG. Calmodulin kinase II inhibition prevents arrhythmias in RyR2(R4496C+/-) mice with catecholaminergic polymorphic ventricular tachycardia. *J Mol Cell Cardiol.* 2011;50:214–222. doi: 10.1016/j.yjmcc.2010.10.001.
175. Dybkova N, Sedej S, Napolitano C, Neef S, Rokita AG, Hünlich M, Brown JH, Kockskämper J, Priori SG, Pieske B, Maier LS. Overexpression of CaMKIIδ in RyR2R4496C^{+/-} knock-in mice leads to altered intracellular Ca²⁺ handling and increased mortality. *J Am Coll Cardiol.* 2011;57:469–479. doi: 10.1016/j.jacc.2010.08.639.
176. Venetucci L, Denegri M, Napolitano C, Priori SG. Inherited calcium channelopathies in the pathophysiology of arrhythmias. *Nat Rev Cardiol.* 2012;9:561–575. doi: 10.1038/nrcardio.2012.93.
177. van der Werf C, Wilde AA. Catecholaminergic polymorphic ventricular tachycardia: from bench to bedside. *Heart.* 2013;99:497–504. doi: 10.1136/heartjnl-2012-302033.
178. Leenhardt A, Denjoy I, Guicheney P. Catecholaminergic polymorphic ventricular tachycardia. *Circ Arrhythm Electrophysiol.* 2012;5:1044–1052. doi: 10.1161/CIRCEP.111.962027.
179. Duygu B, Poels EM, da Costa Martins PA. Genetics and epigenetics of arrhythmia and heart failure. *Front Genet.* 2013;4:219. doi: 10.3389/fgene.2013.00219.
180. Bers DM. Ca²⁺-calmodulin-dependent protein kinase II regulation of cardiac excitation-transcription coupling. *Heart Rhythm.* 2011;8:1101–1104. doi: 10.1016/j.hrthm.2011.01.030.
181. Backs J, Olson EN. Control of cardiac growth by histone acetylation/deacetylation. *Circ Res.* 2006;98:15–24. doi: 10.1161/01.RES.0000197782.21444.8f.
182. Wu X, Zhang T, Bossuyt J, Li X, McKinsey TA, Dedman JR, Olson EN, Chen J, Brown JH, Bers DM. Local InsP3-dependent perinuclear Ca²⁺ signaling in cardiac myocyte excitation-transcription coupling. *J Clin Invest.* 2006;116:675–682. doi: 10.1172/JCI27374.
183. Awad S, Al-Haffar KM, Marashly Q, Quijada P, Kunhi M, Al-Yacoub N, Wade FS, Mohammed SF, Al-Dayel F, Sutherland G, Assiri A, Sussman M, Bers D, Al-Habeeb W, Poizat C. Control of histone H3 phosphorylation by CaMKIIδ in response to haemodynamic cardiac stress. *J Pathol.* 2015;235:606–618. doi: 10.1002/path.4489.
184. Awad S, Kunhi M, Little GH, Bai Y, An W, Bers D, Kedes L, Poizat C. Nuclear CaMKII enhances histone H3 phosphorylation and remodels chromatin during cardiac hypertrophy. *Nucleic Acids Res.* 2013;41:7656–7672. doi: 10.1093/nar/gkt500.