



## CLINICAL REVIEW

## Circadian rhythms and cardiovascular health

Francesco Portaluppi<sup>a,b,\*</sup>, Ruana Tiseo<sup>a,b</sup>, Michael H. Smolensky<sup>c</sup>, Ramón C. Hermida<sup>d</sup>,  
Diana E. Ayala<sup>d</sup>, Fabio Fabbian<sup>b</sup>

<sup>a</sup> Hypertension Center, University Hospital “S. Anna” of Ferrara, Ferrara, Italy

<sup>b</sup> Section of “Clinica Medica”, Department of Clinical and Experimental Medicine, University of Ferrara, Ferrara, Italy

<sup>c</sup> Department of Biomedical Engineering, University of Texas at Austin, Austin, TX, USA

<sup>d</sup> Bioengineering & Chronobiology Laboratories, University of Vigo, Campus Universitario, Vigo, Spain

## ARTICLE INFO

## Article history:

Received 3 March 2011

Accepted 27 April 2011

Available online 8 June 2011

## Keywords:

Blood pressure

Hypertension

Myocardial ischemia

Acute myocardial infarction

Sudden cardiac death

Arrhythmias

Circadian rhythm

Chronobiology

Chronotherapy

## SUMMARY

The functional organization of the cardiovascular system shows clear circadian rhythmicity. These and other circadian rhythms at all levels of organization are orchestrated by a central biological clock, the suprachiasmatic nuclei of the hypothalamus. Preservation of the normal circadian time structure from the level of the cardiomyocyte to the organ system appears to be essential for cardiovascular health and cardiovascular disease prevention. Myocardial ischemia, acute myocardial infarct, and sudden cardiac death are much greater in incidence than expected in the morning. Moreover, supraventricular and ventricular cardiac arrhythmias of various types show specific day–night patterns, with atrial arrhythmias – premature beats, tachycardias, atrial fibrillation, and flutter – generally being of higher frequency during the day than night – and ventricular fibrillation and ventricular premature beats more common, respectively, in the morning and during the daytime activity than sleep span. Furthermore, different circadian patterns of blood pressure are found in arterial hypertension, in relation to different cardiovascular morbidity and mortality risk. Such temporal patterns result from circadian periodicity in pathophysiological mechanisms that give rise to predictable-in-time differences in susceptibility-resistance to cyclic environmental stressors that trigger these clinical events. Circadian rhythms also may affect the pharmacokinetics and pharmacodynamics of cardiovascular and other medications. Knowledge of 24-h patterns in the risk of cardiac arrhythmias and cardiovascular disease morbidity and mortality plus circadian rhythm-dependencies of underlying pathophysiologic mechanisms suggests the requirement for preventive and therapeutic interventions is not the same throughout the day and night, and should be tailored accordingly to improve outcomes.

© 2011 Elsevier Ltd. All rights reserved.

## Introduction

Biological processes and functions oscillate rhythmically in time. Circadian rhythms, which are of particular relevance to everyday life and clinical medicine, are controlled by an inherited master clock residing in the paired suprachiasmatic nuclei (SCN) of the hypothalamus.<sup>1</sup> Rhythmic activities the SCN clock genes *Per1*, *Per2*, *Per3*, *Bmal*, *Clock*, and *Cry* and their gene products comprise the central time-keeping mechanism. Transcription factors *CLOCK* and *BMAL1* drive the expression of *Per1*, *Per2*, *Cry1*, *Cry2*, plus a variety of clock-controlled genes, i.e., target genes that are not integral

clock components and that do not feedback on *CLOCK/BMAL1*, via E-box sequences in their promoters. *PER* and *CRY* proteins negatively feedback on the transcriptional activity of *CLOCK:BMAL1*, which results in a circadian rhythm in expression of the *CLOCK:BMAL1*-driven clock and various clock-controlled genes. Precision of the period and staging of circadian rhythms is achieved via cyclic environmental time cues. The ambient light–dark cycle is the most important cue under normal conditions, with the retinohypothalamic neural projection relaying information sensed by specialized non-cone and non-rod receptors of the retina to the SCN about the timing of light onsets and offsets during each 24-h period. The biological time-keeping system also includes the multitude of peripheral cell, tissue, and organ circadian clocks that are regulated and coordinated by the master SCN clock. The output of the central and peripheral circadian clocks is mediated by various clock-controlled genes, giving rise to the body's circadian time structure (CTS) that is appropriately staged by environmental

\* Corresponding author. Hypertension Center, University Hospital “S. Anna” of Ferrara, Corso Giovecca 203, I-44100 Ferrara, Italy. Tel.: +39 0532 236631; fax: +39 0532 236622.

E-mail address: [prf@unife.it](mailto:prf@unife.it) (F. Portaluppi).

**Abbreviations**

ABPM	ambulatory BP monitoring	MI	myocardial ischemia
ACEI	angiotensin-converting enzyme inhibitor	MSA	multiple system atrophy
AF	atrial fibrillation	nREM	non rapid eye movement
AMI	acute myocardial infarction	OSAS	obstructive sleep apnea syndrome
ANS	autonomic nervous system	PAF	pure autonomic failure
ARB	angiotensin II receptor blocker	PD	pharmacodynamics
AV	atrio-ventricular	PK	pharmacokinetics
BP	blood pressure	PSVT	paroxysmal supraventricular tachycardia
CAD	coronary artery disease	REM	rapid eye movement
CTS	circadian time structure	SBP	systolic blood pressure
CVD	cardiovascular disease	SCD	sudden cardiac death
DBP	diastolic blood pressure	SCN	suprachiasmatic nuclei
ECG	electrocardiogram	SN	sinus node
FFI	fatal familial insomnia	SNS	sympathetic nervous system
HR	heart rate	VF	ventricular fibrillation
HRV	heart rate variability	VPB	ventricular premature beat
		VT	ventricular tachycardia

time cues to support optimal human metabolic and performance efficiency and capability during diurnal activity and repair and rejuvenation during nighttime rest/sleep.

Circadian clocks have been identified in nearly all mammalian cells investigated, including cardiomyocytes, fibroblasts, vascular smooth muscle cells, and endothelial cells.<sup>2–6</sup> Although 24-h rhythms in heart rate (HR), blood pressure (BP), and cardiac output are classically attributed to rhythms in neuroendocrine constituents, rhythms at the cellular level also play an important role.<sup>7–9</sup> The evidence for this comes from a series of different studies. Genetic manipulation of circadian clock components, such as CLOCK and BMAL1,<sup>10</sup> variations within a tandem repeat of the human clock gene Per3,<sup>11</sup> and genetic ablation of the circadian clock within endothelial or vascular smooth muscle cells,<sup>12</sup> either alter, markedly attenuate, or abolish HR and/or BP circadian rhythms. Furthermore, studies on the cardiomyocyte-specific clock mutant mouse model, in which the cardiomyocyte circadian clock is temporally locked to the commencement of the inactive/sleep phase, show the cardiomyocyte clock differentially regulates cardiac metabolism and contractile function during the 24 h.<sup>13–16</sup> Furthermore, rodent models reveal it directly modulates myocardial ischemia/reperfusion tolerance in a circadian rhythm-dependent manner.<sup>14</sup> Recent findings also implicate the disruption (desynchronization) of circadian clocks in the pathogenesis of cardiovascular disease (CVD).<sup>8,17</sup> Circadian clocks are altered in numerous animal models of increased CVD risk, including aging, diet-induced obesity, diabetes mellitus, hypertension- and pressure overload-induced hypertrophy, simulated shift work, and ischemia/reperfusion.<sup>18–24</sup> Circadian desynchrony of the organism from its environment, either in humans through rotating shift-work schedules or in rodent through light/dark cycle or genetic alteration, augments CVD development.<sup>25–28</sup> Furthermore, rodent studies show chronic desynchrony of the circadian clock results in enhancement of cardiac mass, cardiomyocyte size, and expressed hypertrophic markers, thereby suggesting it influences responsiveness to pro-hypertrophic stimuli.<sup>29</sup>

Beyond the cellular level, circadian rhythms of cardiovascular physiology and function are very well established<sup>30</sup>; moreover, clear circadian rhythmicity is found in pathophysiological mechanisms that underlie 24-h patterning of morbid and mortal CVD events. Moreover, the administration time, relative to the staging of involved circadian rhythms, of various classes of medications used to manage CVD risk may impact, sometimes quite dramatically, their pharmacokinetics (PK) and pharmacodynamics (PD).<sup>31</sup> This

infers it is necessary to tailor preventive and therapeutic interventions to circadian rhythm determinants to optimize intended outcomes.<sup>32</sup>

Recognition of the importance of circadian rhythms in cardiovascular functions and their involvement in 24-h patterns of CVD conditions and events has lead to renewed scientific interest in chronobiology, i.e., the study of biological rhythms. Indeed, investigation of the temporal structure of the sources and mechanisms of cardiovascular rhythms is a necessary preliminary step in understanding their clinical implications, especially in regard to CVD risk and its control through therapeutic interventions. The main sources of the time-dependency of cardiovascular physiology and pathophysiology are cyclic variation in external stimuli and demands, especially physical and mental activity and stress, and endogenous circadian rhythms, even though it is often impossible to clearly separate the relative contribution of each. This article discusses the relevance of circadian rhythms to the prevention and management of CVD, focusing on the three major clinical entities of arterial hypertension, myocardial ischemia (MI), and cardiac arrhythmias. Published pathophysiologic, epidemiologic, and clinico-therapeutic evidence clearly establishes the importance of a chronobiologic approach, both to uncovering new insight into the maintenance of cardiovascular health and to improving the prognostic assessment and therapeutic management of CVD patients.

### Circadian rhythms in arterial hypertension

Systolic (SBP) and diastolic (DBP) BP exhibit distinct, although different, 24-h patterning among patients. In so-called normal dippers, the sleep-time BP mean is reduced by 10–20% relative to the daytime mean.<sup>33</sup> During the daytime, typically two daytime peaks are manifested, the first more prominent one shortly after morning awakening and the second late in the afternoon or early evening, with a small mid-afternoon nadir. In healthy young adults, the immediate morning SBP rise amounts to about 20–25 mmHg, but in the elderly, who have less compliant and elastic arteries, it can be as great as 40–60 mmHg. Bed-bound subjects, whether normotensive<sup>34</sup> or hypertensive,<sup>35</sup> and those with fixed HR<sup>36</sup> also exhibit significant BP decline during sleep, it being more pronounced in women than men.<sup>37</sup>

Non-dipping (<10% BP decrease in sleep-time mean relative to daytime mean) and even rising (sleep-time BP mean > daytime mean) 24-h patterns are common in secondary hypertension, which is caused by a co-existing medical condition. Super-dipping BP

patterns (>20% decrease in sleep-time BP mean relative to daytime mean) may occur in persons with autonomic nervous system (ANS) dysfunction, those treated too aggressively with hypertension medications, particularly when ingested in the evening, or in those with 'white coat' (i.e., pseudo) hypertension, abnormally high BP only in the presence of medical personnel in clinical settings.<sup>38</sup> Blunted or reversed BP sleep-time decline has been reported in orthostatic ANS failure,<sup>39</sup> Shy-Drager syndrome,<sup>40</sup> vascular dementia,<sup>41</sup> Alzheimer-type dementia,<sup>42</sup> cerebral atrophy,<sup>43</sup> cerebrovascular disease,<sup>44,45</sup> ischemic arterial disease after carotid endoarterectomy,<sup>46</sup> neurogenic hypertension,<sup>47</sup> fatal familial insomnia (FFI),<sup>48</sup> diabetes,<sup>49</sup> catecholamine-producing tumors,<sup>50,51</sup> Cushing's syndrome,<sup>52</sup> exogenous glucocorticoid administration,<sup>53</sup> mineralocorticoid excess syndromes,<sup>54</sup> Addison's disease,<sup>55</sup> pseudohypoparathyroidism,<sup>56</sup> obstructive sleep apnea syndrome (OSAS),<sup>57</sup> normotensive and hypertensive asthma,<sup>58</sup> chronic renal failure,<sup>59,60</sup> severe hypertension,<sup>61</sup> salt-sensitive essential hypertension,<sup>62</sup> gestational hypertension,<sup>63</sup> toxemia of pregnancy,<sup>64</sup> essential hypertension with left ventricular hypertrophy,<sup>65</sup> renal, liver, or cardiac transplant<sup>66–68</sup> related to immunosuppressive treatment, congestive heart failure,<sup>69,70</sup> and recombinant human erythropoietin therapy.<sup>71</sup> Moreover, a circadian profile characterized by daytime hypertension and sleep-time hypotension has been described in hemodynamic brain infarction associated with prolonged blood–brain barrier disturbance,<sup>72</sup> the range of variation between daytime activity and nighttime sleep being significantly increased from expected.

The circadian HR pattern closely parallels that of SBP and DBP in normal conditions and shows strong genetic dependency in terms of the daytime mean, amplitude of variation, and peak time during the 24 h.<sup>73</sup> There is no doubt the circadian HR rhythm is intrinsic and driven by 24-h variation in ANS activity. However, in all hypertensive conditions that result in loss or reversal of normal nocturnal BP reduction, nocturnal bradycardia is at least partially preserved, indicating the modulating influences of the two variables are only partially coincident. Further support of this notion comes from study of rats made hypertensive by transgenic implantation of the mouse salivary gland renin gene (TGR(mRen-2) 27)<sup>74</sup>; the HR and BP circadian rhythms persist but their phasing manifests in complete opposition.

### **Circadian changes in the pathophysiological mechanisms of arterial hypertension**

The 24-h BP variation results primarily from the cyclic exogenous alteration of physical and mental activity, stress, among others, coupled with the sleep–wake cycle. However, endogenous neurohumoral and other circadian rhythms also play a role,<sup>75</sup> although it is impossible to clearly separate the relative influence of the former from the latter. For sure, the effects of physical and mental activity account for the predominant proportion of the day–night variation,<sup>76</sup> as demonstrated by studies of shift workers who show a close linkage between physical activity and BP even on the first 24 h of night work.<sup>77,78</sup>

Endogenous mechanisms of the 24-h BP pattern also are clearly influenced by genetic factors.<sup>79,80</sup> BP dipping is a heritable trait which can be mapped by linkage analysis, with genetic control of arterial stiffness being the likely mechanism.<sup>79,80</sup> The endogenous basis for the 24-h BP variation is also supported by studies showing lesioning of the SCN of laboratory rodents abolishes BP and HR circadian rhythms without affecting sleep–wake and motor activity cycles.<sup>81,82</sup> Rodents are nocturnally active animals, and their BP circadian rhythm is characterized by elevated nighttime and decreased daytime BP values. Thus, the relationship of BP level to activity level is underlined in both human beings and laboratory rodents.

### *Sleep-related alterations of circadian BP rhythms*

Sleep is the most important and consistent source of circadian BP variation.<sup>83</sup> BP variation varies with sleep stage<sup>84,85</sup>; lowest BP levels occur during deepest sleep, i.e., stages 3 and 4, whereas relatively higher BP levels, although generally not as high as awake-state BP levels, occur during less deep sleep, i.e., stages 1 and 2 and REM sleep. Significant BP increase takes place in the morning in diurnally active persons. Deep (delta or slow-wave) sleep prevails during the first half of nighttime rest coincident with nocturnal BP decline. During the last half of nighttime sleep, REM-related and brief arousal-related episodes of BP and HR rises increase in frequency. This is the likely explanation of why intra-arterial data aligned to the time of waking show the morning BP rise commences prior to awakening; thus, its attribution solely to awakening seems highly implausible.<sup>86</sup> Wrist actigraphy, a tool commonly used in sleep studies, has helped identify two distinct patterns of morning BP rise: one defined by gradual rise commencing before awakening and common in young persons and the other defined by steep BP rise commencing after waking and common in elderly persons who show pronounced responses to mental stress.<sup>87</sup>

Important additional influences of sleep on BP may be exerted through variation in respiration. During sleep onset and light NREM sleep, breathing is regular and periodic with only sporadic central apneas, and BP and HR decline. Oscillations of 3–5 s and 20–30 s in breathing are synchronous with oscillations of cortical electroencephalographic activity, BP, HR, and oxygen saturation. During REM sleep, breathing and HR become irregular and central apneas or hypopneas of a few seconds occur sporadically, often in association with bursts of REM. Hypertensive BP spikes, as great as 30–40 mmHg from baseline, occur abruptly and continue for the duration of the REM episode. During sleep, alveolar ventilation decreases by 0.4–1.5 l/min, and pulmonary arterial pressure rises by 4–5 mmHg; nonetheless, these variables remain within the normal range. Changes in sensitivity and functioning of homeostatic mechanisms account for these oscillatory phenomena, which also occur with similar periodicity in other pathophysiologic states<sup>88,89</sup> or with sleep onset.<sup>90</sup> Respiratory instability associated with the transition from mixed neurogenic and metabolic control of breathing typical of wakefulness to the purely metabolic state typical of NREM sleep is the underlying cause of the periodic oscillations.<sup>91</sup>

The stage organization of sleep is also reflected in ANS function,<sup>92</sup> a well established driver of the circadian BP profile.<sup>93–95</sup> Sympathetic nervous system (SNS) activity during deep NREM sleep is lower than during wakefulness. However, it increases in REM sleep to the same, or greater, level seen in wakefulness. During NREM sleep, activity of the sympathoadrenal branch of the SNS decreases, probably due to entrainment of the sleep–wake cycle, whereas low activity of the noradrenergic branches seems to depend mainly on the assumption of horizontal body posture during sleep.<sup>96</sup> During REM sleep, concomitant reduction in coronary blood flow<sup>97</sup> and increased coronary spasm<sup>98</sup> can be demonstrated. Not surprisingly, therefore, REM sleep is associated with elevated MI risk among individuals with coronary artery disease (CAD) or vasospastic angina.<sup>98,99</sup> Thus, REM sleep constitutes a period of potential CVD risk due to dramatic phasic SNS, BP, and HR fluctuation.

Loss or reversal of the expected normal physiologic sleep-time BP decline may result from the large number of medical disorders that cause insomnia. Unfortunately, objective studies have not been performed in the majority of these conditions, even though polysomnographic data demonstrate sleep disturbances in CVD,<sup>100,101</sup> ANS failure,<sup>40,102–105</sup> rheumatoid arthritis possibly in relation to

corticosteroid therapy,<sup>106,107</sup> chronic renal failure,<sup>108</sup> hyperthyroidism,<sup>109</sup> Cushing's disease and exogenous glucocorticoid administration,<sup>110–113</sup> and FFI.<sup>114,115</sup> It is also of relevance that sleep disturbances may be induced by a variety of acute and chronic common medical conditions, e.g., allergic rhinitis (sleep-time exacerbation of nasal obstruction often accompanied by sleep-disordered breathing),<sup>116,117</sup> atopic dermatitis,<sup>118,119</sup> asthma (a nocturnal condition in 75% or so of sufferers),<sup>120,121</sup> osteoarthritis (arthritic condition characterized by inflammation and pain that builds during daytime activity and being most severe at night),<sup>122</sup> chronic pain syndrome (lowest pain threshold overnight),<sup>123</sup> peptic ulcer disease and gastric-esophageal reflux disorder (most severe late evening and very early morning hours after midnight),<sup>124</sup> congestive heart failure (dyspnea and increased work of breathing worse overnight),<sup>125</sup> nocturnal polyuria,<sup>126–128</sup> and voluntary chronic sleep loss/deprivation.<sup>129</sup> If chronic, these medical conditions may result in persistent sleep-onset insomnia and/or disturbed and discontinuous sleep, the consequence being increased risk of higher than normal overnight BP level and potential nocturnal hypertension with associated elevated hazard of end-organ injury to blood vessels of the brain, eye, kidney, and heart. It is usually assumed that anxiety and discomfort are the major causes of poor sleep, sometimes leading to generalized stress from sleep disruption.<sup>130–132</sup> Three conditions, namely sleep-disordered breathing, ANS failure, and FFI, for which the relationships between sleep and BP have been extensively, though not completely investigated, deserve detailed discussion for the relationships they imply between cardiovascular physiology rhythmicity and CVD disease prevention and treatment.

#### *Sleep-disordered breathing and circadian BP rhythms*

Sleep-disordered breathing is characterized by snoring and sleep apnea of various severity.<sup>133</sup> An abnormal breathing event is defined as snoring (clear inspiratory noise over the trachea), apnea (complete cessation of airflow  $\geq 10$  s), or hypopnea ( $\geq 50\%$  decrease in airflow  $> 10$  s). The snoring and apnea/hypopnea indices, i.e., average number of respective events/hour of sleep, serves to quantify sleep-disordered breathing. By convention, a snoring index  $\geq 10$  defines the presence of snoring, and an apnea/hypopnea index  $\geq 10$  defines the presence of sleep apnea. By means of these criteria, nonapneic snoring can be differentiated from OSAS. It is in OSAS that major alterations of the circadian BP profile are found, with obliteration of the sleep-related BP reduction and increase in BP variability.<sup>134</sup>

Cardiovascular changes that accompany OSAS events are now well recognized. HR, systemic BP, and pulmonary arterial pressure initially decline with the onset of apnea but then rise acutely with its termination. As a result, not only is BP variability increased, but also average BP level.<sup>134</sup> Cardiovascular reflex studies<sup>135</sup> reveal in subjects at rest significantly higher HR and plasma levels of norepinephrine, higher response of BP to head-up tilting, and by the cold face test significantly less respiratory arrhythmia, lower baroreflex sensitivity index, and lower Valsalva ratio associated with enhanced HR decrease. These findings in OSAS patients suggest sympathetic overactivity associated with blunting of reflexes dependent on baroreceptor or pulmonary afferents and normal or greater than normal cardiac vagal efferent activity.

Half or more of all OSAS patients are hypertensive, not only during nighttime sleep span but also during daytime activity.<sup>136,137</sup> Conversely, several studies have found an approximately 30% prevalence of OSAS among patients with primary hypertension.<sup>138,139</sup> Several candidate mechanisms have been advanced to explain how snoring and OSAS lead to sustained daytime elevation of BP and ultimately daytime hypertension. These include direct

effects of episodic hypoxemia and hypercapnia on chemoreceptors and SNS overactivity,<sup>140</sup> cardiovascular system modification (including fluid balance) in response to marked fluctuations in intrathoracic pressure during obstructed breathing,<sup>141</sup> generalized stress from sleep disruption (i.e., arousal effect),<sup>130,132</sup> genetic factors, and age. The relative contribution of each factor remains unknown, and the role of recurrent hypoxia is particularly controversial. Conceivably, concerted action of several factors, e.g., hypoxia, negative intrathoracic pressure, intermittent BP changes, sleep arousal, and metabolic or endocrine factors, possibly only in combination with a particular age and specific genetically-determined characteristics, could bring about hypertension.<sup>142</sup> Whatever the causal factors, the common mechanism shared by hypertension and OSAS is activation of the SNS,<sup>132,135</sup> which somehow becomes chronic or acts as an intermediary, changing hormonal balance or causing vascular remodeling.

Undiagnosed sleep-disordered breathing might play a role also in the genesis of BP pattern alteration of some, if not all, nondippers diagnosed as essential hypertensives. This is more likely for male patients, since in the adult population snoring and OSAS are of much higher prevalence among men than women.<sup>143</sup> In this regard, we previously documented by polysomnographic study of a representative population of male essential hypertensive nondippers the manifestation of blunted sleep-related BP decline and increased BP variability in 10 of 11 apneic snorers.<sup>144</sup> Hence, undiagnosed apneic snoring could play an important role in determining changes in the circadian BP profile toward non-dipping.

#### *Autonomic failure and circadian BP rhythms*

Altered BP level and 24-h patterning can result from ANS failure in the form of a "pure" autonomic failure (PAF) without other neurological signs, previously termed "idiopathic orthostatic hypotension",<sup>145</sup> or from neurodegenerative disease, i.e., multiple system atrophy (MSA). The term MSA, first used by Graham and Oppenheimer in 1969,<sup>146</sup> refers to a sporadic adult-onset neurodegenerative disease that had previously been described variously as striatonigral degeneration, olivopontocerebellar atrophy, or Shy-Drager syndrome according to the clinical predominance of parkinsonism, cerebellar signs, and ANS failure, respectively.<sup>147–149</sup> The clinical picture<sup>150</sup> of MSA patients with ANS failure is highly variable, at least at its onset; it may include parkinsonism poorly responsive to levodopa, cerebellar syndrome, progressive ANS failure (postural hypotension, defective sweating, constipation, and erectile dysfunction in men), or variable combination of these syndromes. The prognosis of MSA with ANS failure is usually determined by progression of neurologic features over the course of about 6 years. In contrast, the natural history of PAF is of a slow progression over some 10–15 years.

All ANS failure patients manifest severe symptomatic postural hypotension. Those with MSA and ANS failure exhibit normal norepinephrine plasma levels when resting in supine position and show no significant increase when subjected to orthostatic stress. PAF subjects differ by having very low plasma levels of norepinephrine when resting in supine position and also show no elevation when subjected to orthostatic stress.<sup>134</sup> Complete obliteration of the circadian BP pattern is found in PAF patients, while reversal of the normal circadian BP profile is found in MSA patients with ANS failure. However, MSA patients without ANS failure express a normal BP rhythm.<sup>134</sup> These observations are consistent with the notion that normal day–night BP variation depends on proper SNS functioning.

Sleep abnormalities, (e.g., sleep fragmentation with greater than normal sleep latency, altered muscle tone, and abnormal movements; decreases in REM and stage 3 and 4 sleep; REM sleep



behavior disorder) are common in MSA, irrespective of ANS failure. However, such sleep abnormalities are absent in PAF patients, suggesting sleep disruption is mainly the consequence of central nervous system lesions<sup>134</sup> and ANS failure can induce nocturnal BP dysregulation independently of sleep abnormalities.

#### *FFI and circadian BP rhythms*

FFI is a hereditary prion disease characterized by a mutation at codon 178 of the prion protein gene cosegregating with the methionine polymorphism at codon 129 of the mutated allele. FFI presents as disturbances of the wake–sleep cycle, dysautonomia, hypertension, and somatomotor manifestations of myoclonus, ataxia, dysarthria, and spasticity.<sup>151</sup> The cardinal feature of FFI, total and sustained disruption of the sleep–wake cycle, constitutes a unique, albeit dramatic, opportunity to observe how the chronic absence of sleep impacts human circadian rhythms. In FFI, dominant 24-h rhythmicity is detectable both in BP and HR.<sup>48</sup> However, as a combined result of its decreased amplitude and shifted phase, the nocturnal BP fall is lost early during disease evolution, even though nocturnal bradycardia is still preserved. A rhythmic BP component persists, although of lower amplitude and shifted phase, for months after the total cessation of the sleep–wake circadian rhythm. Only in its terminal stage is there complete obliteration of 24-h BP variation. In addition, no association with the pattern of motor activity or meals can be detected. These findings support the existence of an intrinsic component of the BP rhythm that is independent of the sleep–wake and rest–activity cycles.

Dysautonomia is an important feature in FFI; however, changes in pituitary–adrenal functioning also seem to contribute significantly to the pathogenesis of the hypertensive condition. In FFI, abnormal nocturnal peaks are detected in the circadian rhythm of plasma ACTH and cortisol concentration, commencing in its earliest stages, i.e., before development of hypertension and clinical and instrumental signs of severe dysautonomia.<sup>48</sup> It is of interest that nocturnal peaks of these hormones, similar to those found in FFI and representing reversal of the normal sleep-related inhibition of secretion, have been produced in healthy subjects by acute sleep deprivation.<sup>152–154</sup>

FFI changes occur not only in rhythms of the cardiovascular system and functions,<sup>48</sup> but also in neurohumoral rhythms,<sup>155,156</sup> indicating this prion disease disturbs and eventually obliterates the endogenous CTS. Although nothing can be drawn from our data regarding basic mechanisms underlying progressive obliteration of the CTS found in FFI, it is very tempting to hypothesize the prion protein plays a role in the origin and modulation of human circadian rhythms, analogous to the role of the per protein in *Drosophila*.<sup>157</sup> Recent demonstration that mice devoid of prion protein exhibit clear CTS alteration<sup>157</sup> further suggests it plays such a role. Hence, future research in this direction is warranted.

#### *Prognostic and therapeutic implications*

Physiological temporal balance with predominant circadian rhythmicity exists between the various neurohumoral factors integrating BP regulation.<sup>158–160</sup> Loss of this temporal balance may be a pathogenetic mechanism of hypertension and should be taken into account in its clinical management.<sup>60,70,161,162</sup> Moreover, it is bound to differentially affect the response of individuals to pharmacologic treatment and constitutes, in part, the basis for the chronotherapy of hypertension, i.e., delivery of medications in synchrony with biological rhythm determinants to enhance desired outcomes and/or minimize/avert adverse effects.<sup>163</sup> During the past two decades, specific features of the 24-h BP pattern, as determined

by around-the-clock ambulatory BP monitoring (ABPM), have been assessed as potential sources of injury to target tissues and triggers of cardiac and cerebrovascular events. Thus, a growing number of studies<sup>164–167</sup> have consistently shown association between blunted sleep-time BP decline and increased incidence of fatal and non-fatal CVD events. Independent prospective studies and recent meta-analyses have also found the asleep BP mean to better predict CVD risk than the awake or 24-h BP means.<sup>166,168–172</sup> Most important, a recent prospective randomized trial involving hypertensive subjects systematically evaluated by periodic (at least annually) ABPM for over 5 years has documented the progressive decrease in asleep BP and increase in sleep-time relative BP decline toward the normal dipping pattern, two new novel therapeutic targets best achieved with bedtime hypertension therapy, to be the most significant predictors of CVD event-free survival.<sup>173</sup>

Appreciable ingestion-time differences in the PK – absorption, distribution, metabolism, and elimination – and PD – independent of PK phenomena – of BP-lowering medications are well known.<sup>83,174</sup> The former result not from meal effects or posture, but circadian rhythms in gastric pH and emptying, gastrointestinal motility, biliary function, hepatic enzyme activity, blood flow to the duodenum and other organs of the gastrointestinal tract, and glomerular filtration rate, among others.<sup>175</sup> Hence, one might expect antihypertensive medications to be cleared more slowly overnight, thereby potentially prolonging their duration of action when ingested at bedtime than upon morning awakening.<sup>176</sup> Administration-time differences in the PD of BP medications, in the absence of differences in PK, are also known<sup>177</sup>; they most likely result from circadian rhythms in circulating drug-free fraction, rate-limiting steps of key biochemical and metabolic processes, receptor number and conformation, and/or second messenger and signaling pathways.<sup>178</sup> A number of published clinical trials, reviewed in-depth elsewhere,<sup>176,177,179</sup> document a variety of highly prescribed calcium channel blocker, angiotensin-converting enzyme inhibitor (ACEI), angiotensin II receptor blocker (ARB),  $\alpha$ -blocker,  $\beta$ -blocker, and diuretic medications when studied by around-the-clock ABPM methods display very significant, and sometimes dramatic, morning–evening dosing-time differences in BP-lowering effects. In particular, once-daily evening, in comparison to morning, scheduling of ARBs and ACEIs results in stronger therapeutic effect on asleep BP without loss of efficacy during the hours of diurnal activity. This administration-time effect is not dependent on the terminal half-life of the individual medications trialed, which ranges from 6 to 9 h for the ARB valsartan to ~40 h for the ACEI spirapril.<sup>180,181</sup>

The chronotherapeutic approach to hypertension is justified by the fact that BP is usually lowest at night as is sodium excretion. When sodium intake is excessive or its daytime excretion hampered, nocturnal BP is adjusted higher via the pressure/natriuresis mechanism to a level required for compensation overnight, the result being non-dipping 24-h BP patterning.<sup>182</sup> In diurnally active persons, the entire circadian BP pattern may be reset to a lower mean level and to a “more normal” day–night variation, simply by enhancing natriuresis nocturnally – the time-of-day when it can be most effective. A modification as simple and inexpensive as switching the ingestion time of one or more prescribed hypertension medications from morning to evening, may alone be sufficient to normalize nighttime BP, exerting an effect like that of sodium restriction.<sup>183</sup>

#### **Twenty-four-hour pattern in MI**

MI is the underlying pathogenetic mechanism of multiple clinical manifestations of CVD events: transient ischemic episodes, acute myocardial infarction (AMI), and sudden cardiac death (SCD).

MI may result from either restricted and insufficient oxygen supply or increased oxygen demand. A number of physiologic variables are crucial in determining mismatch between myocardial oxygen supply and demand, and predictable circadian changes are exhibited by all of them.<sup>32</sup> Such predictable-in-time differences in the susceptibility to MI, on the one hand, and in the pathogenetic mechanisms of MI, on the other hand, result in corresponding time-of-day differences in its overt expression and manifestations.

#### *Twenty-four-hour pattern of transient MI events*

Analysis of 24-h electrocardiogram (ECG) recordings of untreated diurnally active patients reveals episodes of transient ST-segment depression, including silent, i.e., non-symptomatic, MI are two-to-three-fold more frequent in the morning, between 06:00 h and noon, than at night.<sup>184,185</sup> A secondary vulnerability to transient ischemic episodes is sometimes observed later in the day, between 18:00 and 20:00 h.<sup>186</sup> The same circadian distribution is shown by symptomatic, i.e., chest-pain associated episodes of ST-segment elevations.<sup>187</sup> Moreover, greater ST-segment displacements are found when exercise is performed at 16:00 h than at 08:00 h,<sup>188</sup> indicating circadian rhythm-dependent difference in ischemic response of the myocardium to exercise. Finally, so-called variant angina, a form of MI that results from coronary artery vasospasm and expressed as ST-segment elevation, shows different circadian patterning, being most frequent during the second half of the nocturnal sleep span, between 02:00 and 04:00 h,<sup>189</sup> and more frequent during morning than afternoon exercise.<sup>190</sup>

#### *Twenty-four-hour pattern of AMI*

Published case series and population-based studies of temporal patterns in AMI have relied solely on the clock time of events. It is assumed such data are representative of events of at-risk individuals adhering to a common diurnal activity-nighttime sleep routine. However, since a significant proportion of the working population in Western countries is employed in night or rotating shift work, this assumption is invalid. Nonetheless, a great number of reports substantiate AMI onset is lowest during the first part of the night, increased during the second part of the night, and highest during the initial hours of diurnal activity, between 06:00 h and noon,<sup>191,192</sup> when the peak incidence rate of 40% (relative risk: 1.38) of the total episodes is observed.<sup>192</sup> The temporal relationship with awakening, rather than clock time, is verified by studies on shift workers with a reversed sleep–wake routine and also evening-types ('owls') with very late nocturnal bed and diurnal waking times.<sup>25,193</sup> A different pattern in AMI onset defined only by a minor evening peak or absence of 24-h variation has been reported in cohorts of patients with history of smoking, diabetes,<sup>194,195</sup> CVD, stroke,<sup>196</sup> or previous other non-Q-wave type AMI,<sup>195</sup> congestive heart failure, or non-Q-wave type AMI.<sup>194,195</sup> Unfortunately, most studies have failed to take into account possible effects of differences in the sleep–wake routine of cases (in the United States about 15–20% of the work force is likely to be engaged in night or rotating shift work) and also scheduled medications. Prominence of the morning peak in AMI onset, despite the potential effects of these and other deterministic variables, suggests the actual morning risk of AMI may be underestimated.

#### *Twenty-four-hour pattern of SCD*

SCD, an extreme manifestation of MI, exhibits marked 24-h variability in occurrence, with prominent morning peak between 06:00 h and noon (when the incidence rate of SCD is 29%; relative risk: 1.29), secondary afternoon peak, and nighttime trough.<sup>192,197</sup>

### **Circadian changes in the pathophysiological mechanisms of MI**

MI is thought to be triggered by several pathophysiological mechanisms, particularly the sudden morning increase of BP, HR, SNS activity, basal vascular tone, vasoconstrictive hormones, pro-thrombotic tendency, platelet aggregability, plasma viscosity, and hematocrit.<sup>198</sup> Each of these variables exhibits prominent circadian rhythmicity in phase with the reported 24-h pattern of the ischemic events.

#### *Role of sleep*

More than 80% of the episodes of ambulatory MI are associated with substantial increases in HR.<sup>199,200</sup> The relative tachycardia that accompanies morning arousal from sleep at night results in increased myocardial oxygen demand<sup>199</sup> and contributes to the excess morning frequency of MI, AMI, and SCD.<sup>200,201</sup> However, ischemic events occur also during sleep, especially in persons with severe CAD and vasospastic (variant) angina. As previously discussed, the strong circadian relationships between sleep, BP, and HR help explain why MI may be triggered overnight during REM sleep.<sup>98,99</sup> This risk is time-dependent, considering that episodes of REM sleep occur preferentially during the last half of the nocturnal rest span. Thus, the time-dependent early morning increase in MI risk appears to be determined, at least in part, by the circadian rhythm of REM sleep propensity through increased levels of "SNS activity", HR, and BP.

#### *Role of neurohumoral factors*

Other endogenous circadian rhythms are also involved in the elevated morning-time risk of MI. Plasma norepinephrine level<sup>202,203</sup> and plasma renin activity, which exhibit profound circadian variation and achieve peak levels in the morning,<sup>204</sup> induce coronary vasoconstriction.<sup>205</sup> A circadian rhythm in vascular basal tone is also demonstrated<sup>206</sup> in relation to increased morning  $\alpha$ -sympathetic vasoconstrictor activity. In addition, ischemic threshold is lower at this time,<sup>207</sup> suggesting increased ischemia-induced coronary vascular resistance. This finding is further supported by the similar temporal variation in post-ischemic forearm vascular resistance.<sup>207</sup> Thus, the low morning ischemic threshold presumably mirrors the corresponding-in-time elevated coronary vascular resistance. In addition, the circadian pattern of plasma cortisol secretion, characterized in diurnally active persons by a sharp morning rise,<sup>208</sup> contributes to reduced coronary blood flow by increasing sensitivity of epicardial vessels to vasoconstrictor stimuli.<sup>209</sup>

#### *Prognostic and therapeutic implications*

Only limited data are available on the prognostic implications of the temporal variation in the inducing mechanisms of MI. However, it is known the (biological) time of AMI onset seems to affect its clinical course and outcome.<sup>210</sup> On average, AMIs that commence in the morning between 06:00 h and noon in presumably day-active persons are of greater infarct size than those that commence at other times of the day or night, whereas those that occur between noon and 06:00 h carry a significantly lower risk of circulatory arrests from ventricular arrhythmias than ones that commence at other times.<sup>211</sup> Moreover, the circadian time of AMI affects the success rate of thrombolysis; it is much poorer in patients who manifest an AMI in the morning than in those who manifest it at other times during the 24 h.<sup>212</sup>

All types of medications may show administration time (i.e., circadian rhythm)-dependent differences in their PK and/or PD<sup>213</sup>; however, medications used to treat MI patients are also bound to be influenced by the circadian staging of rhythms in underlying physiopathogenic mechanisms. It is noteworthy that a number of anti-ischemic medications show clinically significant ingestion-time differences in their PK and PD. A detailed review of all the temporal issues of antianginal medications is beyond the scope of the present paper, but can be found elsewhere.<sup>214,215</sup>

The 24-h rhythmic organization of cardiovascular functions is such that the defense mechanisms against MI are incapable of providing the same degree of protection throughout the day and night. Instead, temporal gates of excessive susceptibility exist, particularly in the morning and to a lesser extent in the evening, to aggressive ischemia-inducing mechanisms through which overt clinical manifestations may be triggered. Peak levels of critical physiologic variables, such as BP, HR, rate-pressure product (SBP  $\times$  HR, surrogate measure of ventricular work and myocardial oxygen demand), SNS, and plasma concentrations of endogenous vasoconstricting substances, are aligned together at the same circadian time, in the morning in day-active persons, giving rise to heightened risk of ischemic events then. Thus, even relatively minor environmental stimuli that are usually harmless, like physical and mental stress, may precipitate AMI or SCD in the morning. Accordingly, the requirement for preventive and therapeutic interventions varies predictably during the 24 h. We feel that therapeutic strategies ought to be tailored to prevent the excessive risk of morning as well as late afternoon/evening AMI events. This implies the delivery of anti-ischemic medications, either alone or in combination, needs to be synchronized in time during the 24 h in proportion to predicted need, as documented by established rhythmic patterns in MI risk, and in a manner that will avert or minimize undesired side effects conventionally monitored in the clinic as well as those indicative of disruption of biological time-keeping systems. Thus far, only a few selected MI chronotherapies have been developed and approved by governmental agencies for marketing. The relative merit of these in comparison to conventional therapies in reducing ischemic events, especially in the morning when risk is greatest, and improving patient quality of life remains unresolved, since large prospective trials have yet to be conducted.<sup>214</sup>

### Circadian rhythms in arrhythmias

Circadian variations have been established also in the pattern of presentation of both supraventricular<sup>216,217</sup> and ventricular cardiac arrhythmias,<sup>218–220</sup> irrespective of the presence or absence of concomitant medications.<sup>221</sup> In clinical practice, many arrhythmic episodes are observed as a consequence of MI, a pathophysiologic event that exhibits profound 24-h patterning, as previously related. However, the distribution during the 24 h of malignant arrhythmias is similar in patients with and without evidence of ischemic heart disease,<sup>222</sup> suggesting the importance of other factors.

Interpretation of data reported in the medical literature on circadian patterns in cardiac arrhythmias is complicated. This is because the conduct of previous studies were confounded, at least to some extent, by key factors extraneous to intrinsic arrhythmogenic activity that were overlooked by investigators, such as consumption of alcohol, caffeine, and other sympathomimetic substances, time and dose of prescription and perhaps illicit drugs, and differences in sleep–wake routine between subjects of various cohort studies. Because of these and other confounders, the findings of some studies may have been biased resulting in inappropriate conclusions. Moreover, the failure to recognize the capability of concomitant diseases, e.g., diabetes, renal failure, and heart

failure, among others, to alter the circadian pattern of certain functional triggers of arrhythmias, like ANS activity and HR and BP, may also occasionally contribute to seemingly contradictory findings.

### Circadian rhythms of atrial arrhythmias

Circadian patterning in atrial arrhythmias has been little studied, probably because of their relatively benign nature. Atrial arrhythmias – premature beats, tachycardias, atrial fibrillation (AF), and flutter – appear to exhibit circadian patterning, being of higher frequency during the day than night and with the abnormal atrial foci under the same long-term ANS regulation as normal pacemaker tissue.<sup>223</sup> A notable exception is AF. Paroxysmal AF is classified into two types according to onset time and ANS tone. Vagotonic ones typically occur at night, and adrenergic ones during the day.<sup>224,225</sup> Coumel et al.<sup>224</sup> found patients with vagally mediated AF tend to be younger, always have idiopathic AF, experience occurrences at night, and do not have a tendency toward persistent AF. In contrast, the adrenergically mediated type is opposite to the vagal type in all respects. Persistent AF, as studied with implanted atrial defibrillators, shows circadian distribution of onsets, i.e., predominant during the day.<sup>225</sup>

Pritchett et al.<sup>226</sup> documented paroxysmal supraventricular tachycardia (PSVT) by telephone transmission of the ECG of 14 apparently diurnally active patients treated with the calcium channel blocker medications of verapamil or diltiazem. Measuring the “tachycardia-free periods” between commencement of therapy and first recurrence of tachycardia as well as between successive recurrences, they found PSVT episodes were uniformly distributed during the span of wakefulness, between 09:00 h and midnight, whereas no tachycardia was noted between midnight and 09:00 h. Moreover, the occurrence of PSVT was uniformly distributed throughout medication dosing intervals without predilection for tachycardia during the time of presumed plasma trough drug levels. However, different results were obtained for untreated diurnally active patients.<sup>216</sup> The PSVTs were atrio-ventricular (AV) nodal reentrant or AV reciprocating tachycardia types. Highest relative incidence of PSVT was at 16:00 h, when PSVT was five times more likely than at 04:00 h, the time of lowest relative incidence. If the relative incidence of attacks of PSVT is confirmed in future studies to be highest in the afternoon, one would expect targeting antiarrhythmic medications to this time of highest risk to improve therapeutic outcome.<sup>216</sup> In patients referred to an emergency department, Kupari et al.<sup>227</sup> observed two peaks in the frequency of PSVT onset, one in the morning between 06:00 h and noon and the other in the evening between 18:00 h and midnight. Neither the etiology of the arrhythmia nor preceding alcohol consumption appeared to modify this tendency, although administration of  $\beta$ -adrenergic blocking medication shifted the morning surge of arrhythmias to the nighttime, between midnight and 06:00 h.

Another study found symptomatic patients referred to a mobile coronary care unit<sup>228</sup> experienced greatest frequency of PSVT during the daytime and least frequency between midnight and 08:00 h. AF was also investigated in this study<sup>228</sup>; peak incidence occurred between midnight and 02:00 h, with secondary peaks in the morning between 08:00 and 09:00 h and afternoon between 14:00 and 16:00 h. Isolated AF, as well as AF associated with organic heart disease, exhibited the same 24-h pattern. Moreover, circadian fluctuation of the ventricular response to AF, when investigated by ambulatory Holter monitoring outside the hospital,<sup>229</sup> revealed prominent circadian patterning, with an early afternoon peak and nocturnal trough, likely due to increased vagal tone during sleep, which would be expected to increase AV node refractory period.

Several factors may contribute to the above-discussed differences between studies in the circadian pattern of PSVT. Different clinical settings (mobile vs. hospital coronary care unit, vs. emergency departments) and patients of different countries and cultures are bound to determine variable thresholds for seeking emergency care, which in turn becomes a selection bias in determining which arrhythmic episode will be recorded (if sufficiently symptomatic) and which will not (if relatively asymptomatic). Differences in treatment strategies, including class, dose, and schedule of medications, of treated cohorts and also the timing of sleep and wakefulness during the 24 h also are of major importance. Nonetheless, all the studies are consistent in reporting low incidence of PSVT during presumed nighttime sleep.

#### *Circadian rhythms of AV block*

A 24-h pattern, with morning preference, in the onset of symptomatic third-degree AV block was demonstrated in an emergency room setting,<sup>230</sup> similar to the one in SCD and cardiogenic cardiac arrest.<sup>231</sup> The etiology of this 24-h pattern is unknown; it may be an expression of intrinsic biological rhythmicity within cardiac tissue or its control system and/or the timing of environmental triggers culminating in coronary ischemia.

#### *Circadian rhythms of ventricular premature beats (VPBs)*

The frequency of VPBs, primarily based on studies conducted in patients affected by ischemic heart disease, appears to be decreased by as much as 50% during nighttime sleep relative to diurnal activity,<sup>232,233</sup> with the minimum number of events recorded between midnight and 02:00 h and greater number widely dispersed throughout the day, but without a clear peak time.

The morning increase in SCD as previously discussed might be related, at least in part, to increased frequency of ventricular arrhythmias, since highest prevalence (about one-third) of complex or frequent ventricular arrhythmias, as well as higher mean number of VPBs/hour, are observed between 06:00 h and noon.<sup>234</sup> In contrast with this conclusion are the findings of Goldstein et al.,<sup>235</sup> who did not find any relationship between the circadian rhythm of ventricular ectopic activity and cardiac mortality, prompting the conclusion the circadian rhythm in VPBs is not predictive of sudden death in patients with a high frequency of this type of arrhythmia.

The observed nocturnal decrease in VPBs was found to be reproducible, as established through replicate 24-h ECG monitoring studies.<sup>233,236</sup> Only two studies<sup>237,238</sup> reported no nocturnal decrease in VPBs; however, altered sleep patterns<sup>238</sup> or absence of circadian rhythmicity in ANS activity<sup>237</sup> may have contributed to these contrasting findings. Another study documented a nocturnal peak of VPBs in only one-half the patient sample, but these findings were probably affected by hypertrophic cardiomyopathy. Treatment for six-weeks with the  $\beta$ -blocker medication propranolol did not reduce the frequency of VPBs in recent AMI patients.<sup>239</sup> However, propranolol (60 or 80 mg three times a day (t.i.d.)) was protective against the morning surge in VPBs associated in time with awakening from nocturnal sleep, with persistence of the therapeutic effect throughout the day.<sup>239,240</sup>

#### *Circadian rhythms of ventricular tachyarrhythmias*

Ventricular tachycardia (VT) and ventricular fibrillation (VF) show circadian variability in their occurrence. For example, Lucente et al.<sup>241</sup> demonstrated a significant circadian rhythm of VT based on Holter ECG tracings of AMI patients studied before antiarrhythmic treatment. A late-morning peak prevailed among patients who

suffered an AMI in the distant past, while an afternoon peak in VT prevailed among patients who had an AMI recently. Unfortunately, patients with recent AMI were studied in the hospital whereas those with an earlier AMI were studied at home. Thus, difference in study conditions confounds the comparison of the respective patterns in VT between the two patient groups. In another study, Twidale et al.<sup>242</sup> studied the clock time of VT onset in 68 AMI patients. VT was not associated with AMI in 53 of them, as documented by 12-lead ECG. The remaining 15 patients, who did not have a previous medical history of sustained VT, had a medical history of syncope or presyncope of unknown etiology and experienced sustained monomorphic VT induced during electrophysiological study. In this patient subset, peak VT onset occurred between 10:00 h and noon.

Widespread use of implantable defibrillators set to automatically detect, record, and terminate episodes of VF facilitates study of their temporal patterns. Such derived data reveal a circadian pattern of VF, with primary morning peak between 07:00 and 11:00 h and much smaller secondary peak between 16:00 and 20:00 h.<sup>220,243</sup> This typical 24-h pattern characterized by major morning and secondary afternoon peaks was confirmed by Englund et al., who found it to be closely similar in both ischemic and non-ischemic heart disease patients, suggesting triggering mechanisms of VF may be similar and independent of underlying heart disease.<sup>222</sup>

Indirect confirmation of the daytime prevalence of ventricular tachyarrhythmias comes from a large cohort study demonstrating much greater proportion of cardiac arrests associated with VF or VT in public settings, assumingly the hours of daytime activity, than at home, assumingly the evening and nighttime.<sup>244</sup> Interestingly, cardiac arrest survival was much greater (odds ratio: 2.49) for those VF and VT events that occurred in public settings, i.e., daytime, than at home, i.e., evening and nighttime, findings consistent with ones of an experimental murine model demonstrating ischemic damage consequent to cardiac arrest is much worse when occurring during the rest than active phase of the 24 h.<sup>245</sup>

Fries et al. compared the circadian pattern of VF with that of the parameters derived from the spectral analysis of HR variability (HRV), a marker of ANS activity, in recipients of implantable cardioverter-defibrillators.<sup>246</sup> Twenty consecutive recipients were studied, comparing the circadian rhythms of HRV in patients with and without the typical morning peak of VF. The HRV marker of vagal activity displayed physiological circadian variability, i.e., highest values during nighttime sleep, only in patients without the morning peak of VF, and inverse circadian variability, i.e., lowest values during sleep, in patients showing the clear morning peak in tachyarrhythmias. As pronounced adrenergic hyperactivity in heart failure is the probable explanation for the paradoxically reversed HRV circadian pattern, the morning peak frequency of CVD events may be interpreted as a sign of cardiovascular overload caused by the natural change from sleep to activity upon awakening.

#### **Circadian changes in the pathophysiological mechanisms of arrhythmias**

Many functional, e.g., neurohumoral, electrolytic, hemodynamic, metabolic, etc., factors, aside from MI, may trigger and maintain arrhythmic episodes. Mechanisms by which these factors interact to determine an arrhythmogenic stimulus remain incompletely understood, constituting a major problem not only for the characterization of the determinants of the 24-h pattern of arrhythmias, but also for devising optimal conventional or chronotherapeutic and chronopreventive strategies. Arrhythmias are diagnosed by ECG, and it is of interest that many of ECG indices display significant circadian changes. In healthy subjects, P wave duration and its area, P–R interval, QRS duration, and QT interval



each shows circadian variation with the peak-to-trough difference equal to about 20% of the respective 24-h mean.<sup>247–250</sup> During the daytime, when SNS output is enhanced and HR increased, these indices are decreased, such that in diurnally active persons minimal values are expected between 10:00 h and 14:00 h.<sup>247–250</sup> During the night, following sympathetic withdrawal and parasympathetic dominance, these indices increase, attaining peak values between midnight and 06:00 h. These inherent changes during the 24 h, both in electrical activities and mechanical function, enable the heart to optimally meet workload demands during the daytime and favor or reflect repair and rejuvenation during overnight rest.

#### *Circadian rhythms in electrical properties of the myocardium*

Not only atrial, but also ventricular, rates exhibit circadian rhythmicity, with daytime peaks and nighttime troughs.<sup>251,252</sup> Circadian rhythmicity is also demonstrated in sinus node (SN) function, AV nodal and myocardial refractoriness, QT interval duration, and R and T wave voltage.<sup>247,253,254</sup> Temporal changes in SN function, with coefficients of variation over the 24 h as great as 10%, follow the ANS circadian rhythm.<sup>254</sup> SN recovery time, main index of SN function, is circadian rhythmic, with slowest recovery time between midnight and 07:00 h. The QT interval also exhibits similar variation, with longer QT interval during sleep than waking.<sup>247,255</sup> Moreover, the dispersion of the QT intervals (defined as the inter-lead QT variability in a 12-lead ECG), a simple method of evaluating repolarization heterogeneity of the ventricular myocardium, is greater during diurnal activity than nocturnal sleep,<sup>256</sup> both in CAD and non-CAD subjects.<sup>257</sup> In contrast, Manolis et al.<sup>258</sup> found a circadian rhythm of QT dispersion in heart failure patients, but with values higher during the night than the day; although, no such day–night difference was detected in subjects free of organic heart disease.

Ventricular refractoriness, i.e., duration of time required by ventricular fibers to recover excitability after depolarization, more directly reflects myocardial repolarization. Ventricular refractoriness is shorter during diurnal activity than nocturnal sleep.<sup>255,259</sup> In one study,<sup>259</sup> the ventricular refractory period was shortest around morning awakening, i.e., when SCD risk is highest.<sup>260</sup> Thus, circadian changes in the electrophysiological properties of the myocardium may play a very significant role in the complex pathogenesis both of arrhythmia initiation and SCD.

The circadian influence on refractoriness of normal cardiac tissues and accessory pathways exerts nocturnal protection against electrical induction of reciprocating tachycardia, but facilitates transient arrhythmia in the evening. In fact, there is 24-h variation in the frequency and pattern of ventricular arrhythmias, with significant reduction during sleep and absence during physical activity.<sup>261</sup> In patients with left-sided Kent bundles, i.e., bypass fibers between the atrium and ventricle thereby constituting an accessory pathway outside the normal conduction system and typical of Wolff–Parkinson–White syndrome, nocturnal protection against electrical induction of reciprocating tachycardia is associated with prolongation of atrial, AV nodal, ventricular, and Kent bundle refractoriness.<sup>262</sup> A greater number of VPBs are induced by hypoglycemia during daytime wakefulness than nighttime sleep.<sup>263</sup> HR<sup>264</sup> and left ventricular function<sup>265</sup> are major determinants of 24-h variation in the frequency of ventricular premature depolarization in subsets of patients; only those with premature beats of the HR-dependent type or those with left ventricular ejection fraction >30% show circadian rhythmicity. A circadian periodicity is demonstrated also in bradyarrhythmias<sup>266</sup> and ventricular response to AF.<sup>229</sup>

T-wave alternans, an integral component of congenital long QT syndrome, is a phenomenon of beat-to-beat variability in T-wave

amplitude, morphology, and sometimes polarity that can trigger life-threatening ventricular arrhythmias. Based on a single report, this phenomenon too shows circadian variation, with highest occurrence (216 out of 320 episodes) between 07:00 h and 13:00 h. in presumably diurnally active persons.<sup>267</sup>

The possible influence of the circadian BP rhythm on arrhythmias has been studied, but with unclear results. A strong correlation between the incidence of VPBs and BP and HR was found in patients with left ventricular hypertrophy, while in the absence of hypertrophy supraventricular premature beats also exhibited higher incidence during peaks of BP and HR.<sup>268</sup> In another study, an independent positive correlation between BP and VPBs was found in 8 of 12 study subjects, while the HR was a negative, although nonsignificant, factor for ectopic beats.<sup>269</sup>

Altogether, the findings of the various studies suggest a primary role for neural activity in modulating the electrical disturbances underlying ventricular ectopy, even though at least one study reported no significant circadian variation in any electrophysiologic measure of ventricular electrical instability.<sup>270</sup> A contribution of other hormonal factors, especially adrenocortical ones, cannot be excluded.

#### *Role of sleep on the electrical properties of the myocardium and arrhythmias*

Bradyarrhythmias, particularly sinus bradycardia and AV blocks, are common during sleep.<sup>271</sup> Indeed, sinus pauses, second degree AV block, and greater frequency of PVBs and VT have been reported in OSAS<sup>271–273</sup> and obstructive lung disease.<sup>274</sup> They appear to be associated with hypoxia and can be eliminated by oxygenation, as demonstrated in the rat,<sup>275</sup> being more resistant to systemic hypoxia and reoxygenation during the activity than rest span, which is the only time when systemic hypoxia is proarrhythmogenic.<sup>275</sup> During sleep, PVBs appear to significantly decline in number, whereas premature atrial contractions are sometimes found to be increased.<sup>232</sup> The sleep-time reduction in PVBs correlates more closely with nocturnal HR than arousal level decline. Changes that occur during sleep can be explained by changes in autonomic mediation, with the sympathetic limb of the ANS exerting greater effect than the vagal limb.<sup>276</sup> Sleep appears to be protective against arrhythmias, regardless of when it occurs, since HR and BP are decreased even during an afternoon nap or siesta.<sup>277</sup> As expected, loss of ANS circadian rhythmicity, e.g., as observed in AMI patients,<sup>237</sup> is reflected by absence of nocturnal decrease in VPBs, confirming again the protective role of sleep and its associated physiology against arrhythmias.

In AF patients, the frequency of ventricular response is significantly lower during the night than day, implying the AV node refractory period duration is longer and AV conduction faster nocturnally.<sup>278,279</sup> In diurnally active persons, lowest ventricular response generally occurs between 03:00 and 05:00 h, and greatest ventricular response generally occurs between 22:00 h and midnight, a pattern consistent with circadian activities of the ANS.

Significant nocturnal prolongation of SN rate and recovery time, QT interval, and effective refractory period of the atria, AV node, and right ventricle has been demonstrated electrophysiologically. AV nodal and myocardial refractoriness follow a circadian rhythm, the peak being between 23:00 h and 08:00 h.<sup>253</sup> Moreover, shorter refractory periods during the day than night<sup>280</sup> may promote initiation of reentrant circuits within the myocardium, facilitating AF, and VF in high-risk patients. This mechanism may also explain why  $\beta$ -blocker therapy interacts with the circadian variation of electrophysiological properties of the myocardium,<sup>259</sup> possibly leading to suppression of the morning peak of malignant arrhythmias<sup>281,282</sup> and protection against SCD.<sup>283</sup>

The pacing threshold for ventricular capture increases by 5% during the nighttime relative to daytime.<sup>284</sup> In nocturnally active laboratory rodents, the electrical stability of the heart, as reflected by the threshold for VF induction by programmed electrical stimulation, is greatest during the first half of the activity span, with the total peak-to-trough circadian variation in VF threshold amounting to 25%.<sup>285</sup> A circadian rhythm in the threshold of malignant ventricular arrhythmias may contribute to some extent to the unequal distribution of SCD during the 24 h. In this regard, Venditti et al.,<sup>286</sup> in a review of recordings from implantable cardioverter-defibrillators, found a morning peak in defibrillation threshold and corresponding morning peak in failed first shock frequency. These findings suggest cardiac electrical instability is increased in the morning, thus facilitating VF induction and impeding VF termination.<sup>285,286</sup> An opposite pattern, with a morning nadir, was observed in the threshold of atrial defibrillation,<sup>287</sup> suggesting different regulatory electrophysiological mechanisms in AF versus VF.

#### *Role of neurohumoral factors in cardiac arrhythmias*

Fluctuations in ANS activity constitute major triggers of cardiac arrhythmias.<sup>288</sup> In addition to circadian rhythms in ANS activity, as discussed in previous sections, both norepinephrine and epinephrine plasma concentrations peak in the morning, around the time when diurnal activity commences, and are lowest during nighttime sleep.<sup>289,290</sup> A strong relationship exists between the circadian changes in plasma dopamine and norepinephrine and epinephrine, thus indicating dopaminergic modulation of circadian variation in SNS activity.<sup>291</sup> Increased SNS activity accelerates HR, favors spontaneous cardiac tissue depolarization, shortens ventricular effective refractory period, and decreases VF threshold.<sup>271</sup> In contrast, increased parasympathetic activity slows HR, decreases AV nodal conduction, and, in the presence of baseline SNS activity, increases both ventricular refractory period and VF threshold.<sup>292</sup> Spontaneous episodes of VT are often preceded by changes in HRV,<sup>293</sup> a reliable index of ANS activity.<sup>294,295</sup>

Nocturnal HR decline is an important physiological phenomenon. In monkeys, prevention by atrial pacing of the normal nocturnal HR decline results in rapidly progressing heart failure due to sustained elevated left ventricular workload.<sup>296</sup> This suggests alteration of normal nocturnal bradycardia should be regarded as an undesirable adverse effect of pharmacotherapy. However, no clinical data are yet available on the prognostic and therapeutic implications of such attributes of the HR circadian rhythm.  $\beta$ -Blockade eliminates the 24-h rhythm in ventricular refractory period, even though there is no clear relationship between ventricular refractoriness and plasma catecholamine levels.<sup>259</sup> This infers the day–night oscillation in SNS activity (which does not directly correlate with plasma norepinephrine concentrations) may be responsible for these observations. Circadian variation is found also in the atrial and ventricular refractory periods, with longer refractory periods during nighttime sleep than daytime activity.<sup>280</sup> Moreover, a clear relationship exists between ANS tone and circadian variation of refractory period.<sup>280</sup> Atrial and ventricular refractory periods are shorter when sympathetic tone is high, i.e., during the day, and are substantially lengthened when parasympathetic tone is high, i.e., during the night. Thus, the higher the sympathetic or parasympathetic activity of the ANS, the shorter or longer, respectively, is the refractory period. This relationship is also expressed as a strong negative correlation between atrial and ventricular refractory periods and ratio of the spectral power of the low and high frequency band (LF/HF) of HRV used as a measure of sympathovagal balance. In patients without structural heart disease, ANS activity (determined by HRV spectral analysis)

significantly influences the time of AF onset. Both vagal and sympathetic activity increase immediately prior to AF onset in patients experiencing nocturnal episodes; however, vagal tone is more dominant, as shown by increase of the HF and LF components in the absence of significant variation in LF/HF ratio. Conversely, sympathetic tone increases immediately before AF in daytime-onset episodes.<sup>297,298</sup>

#### *Prognostic and therapeutic implications of circadian rhythmicity in cardiac arrhythmias*

Knowledge of circadian rhythms in cardiac arrhythmias and findings of tests used to determine their triggering thresholds are important, both for the optimal design of drug-delivery formulations and their accurate assessment. Evaluation of new dosage forms could be confounded by choice of an inappropriate biological time to conduct preclinical patient trials. Arrhythmogenesis appears to be suppressed during nighttime sleep, a phenomenon which is bound to influence results of studies done to assess efficacy of antiarrhythmic medications in relation to their administration time. Even defibrillator energy requirements of cardiac patients show circadian variation, supporting the potential merit of a chronotherapeutic approach to cardiac arrhythmias prevention. Since several types of common cardiac arrhythmias manifest predictable-in-time patterns, a chronotherapeutic approach to their prevention and treatment seems warranted.<sup>299</sup> Antiarrhythmic therapy tailored to be more intense during the hours of elevated risk signified by the peak frequency of arrhythmias seems promising. Yet, the advantage, i.e., increased drug efficacy and safety, of such a chronotherapeutic versus conventional (which aims at achieving constancy in drug concentration) approach, remains to be explored. We know of no studies that have been undertaken to assess the differential timing of antiarrhythmic drug administration as a means to improve therapeutic outcomes and increase patient tolerance. As a minimum, existing conventional formulations ought to be investigated for ingestion-time differences in antiarrhythmic efficacy. Up to now, relatively few conventional once-a-day anti-ischemic and anti-hypertensive medications – oral nitrates, calcium channel blockers, and  $\alpha$ -adrenoceptor and  $\beta$ -adrenoceptor antagonists – have been compared for their differential effect and safety when administered in the morning versus evening.<sup>214,300</sup> Some of these medications may be considered antiarrhythmic in as much as they can prevent arrhythmias induced by MI episodes and/or HR and BP surges. Further research is warranted with more specific and potent antiarrhythmic preparations as a means of improving clinical management of cardiac arrhythmias and quality of life of at-risk patients.

#### **Perspectives and conclusions**

The purpose of this review has been to highlight the (chrono) epidemiology and chronobiology of CVD and potential underlying pathogenic mechanisms. Clinically significant 24-h patterns in hypertensive BP levels, MI, AMI, SCD, and various atrial and ventricular arrhythmias are known, having been established by clinical case series and population-based studies plus meta-analysis of the data of individual reports. Clinical epidemiologists initially emphasized environmental factors as triggers of the morning-time excess in MI, AMI, and SCD, primarily the abrupt change from nocturnal rest and inactivity to extensive morning physical exertion and mental stress, in coincidence with less than optimal coverage by preventive therapy due to end of dosing-interval drug trough levels. Initial chronobiology studies identified several key circadian rhythms, e.g., in HR, BP, SNS, basal

vascular tone, vasoconstrictive hormones, prothrombotic tendency, platelet aggregability, plasma viscosity, and hematocrit, the phasing of which is hypothesized to result in elevated vulnerability to CVD events at the commencement of diurnal activity.<sup>198</sup> The physiology and neuroendocrinology of sleep also plays a role in the pathogenesis and triggering of hypertensive target-organ damage, MI and CVD events, and in certain patients, especially hypertensive ones, risk of MI and CVD events is increased also nocturnally. A recent area of research involves the circadian clock of cardiomyocytes and its role in CVD pathogenesis. Of interest are the findings on rodent models implicating the importance of the circadian clock of cardiomyocytes in the differential regulation during the 24 h of cardiac metabolism, contractile function, and ischemia/reperfusion tolerance, suggesting its role in the pathogenesis of ischemic heart disease and cardiac arrhythmias. Rodent models further suggest disruption (desynchronization) of this circadian clock gives rise to cardiovascular dysfunction, which seemingly helps explain why rotating and night shift workers, in comparison to day shift workers, are at greater risk of developing CVD.

The prominent time patterns in CVD events necessitate new approaches to the clinical management of at-risk patients. It is taken for granted that clinical cuff BP measures are sufficiently representative, apart from those prone to 'white coat' or 'pseudo' hypertension. However, this is not the case. The literature is convincing that the sleep-time BP mean and nature of the 24-h BP pattern i.e., dipping vs. non-dipping, better predict the risk of cardiac and other target injury – as well as immediate (5-year) CVD morbidity and mortality – than daytime cuff values. In the realm of diagnostic medicine, ambulatory 24-h ECG monitoring is clinically accepted as a standard method to properly assess MI and cardiac arrhythmias, because they cannot always be captured during daytime inpatient clinical ECG evaluation. Similarly, 24-h ABPM is justified, in spite of the fact it is not currently recommended or financially supported by health maintenance and preventive medicine programs, because representative diurnal and nocturnal SBP and DBP values and 24-h patterning cannot be captured during a single very brief patient–doctor daytime encounter in the clinic. In the realm of therapeutics, almost all prescribed preventive and curative MI, hypertension, and cardiac arrhythmia therapies are formulated and scheduled to achieve constant or near-constant 24-h blood and tissue concentrations, even though each one of these conditions, some of which are potentially life threatening, show rather high-amplitude, predictable-in-time 24-h patterning of risk. Indeed, it seems counter-intuitive to strive for constancy of anti-ischemic and antiarrhythmic medication levels when the data overwhelmingly support the notion therapeutic strategies ought to be systemically varied during the 24 h in synchrony with biological need, which in the case of MI and certain arrhythmias is greatest in the morning. The advantage of hypertension chronotherapy has recently been substantiated by the 5-year outcomes MAPEC trial; reestablishing the circadian BP pattern, i.e., normal dipping, by scheduling one or more hypertension medications at bedtime, rather than scheduling them all in the morning, was associated with significant reduction in CVD events. Nonetheless, extensive review of the medical literature reveals a disappointing paucity of CVD chronotherapeutic initiatives, despite the promise such an approach may result in better individualization of treatment, improved drug efficacy, better patient tolerance, and hence better therapeutic compliance. Indeed, only a few selected so-called CVD chronotherapies have been developed and approved for marketing, and their relative merit versus conventional therapies is unverified, since prospective outcome trials have yet to be conducted. Future chronobiologic developments hopefully will involve greater understanding of the role circadian phenomena play at all levels of

biological organization, especially at the cellular and molecular clock levels in CVD pathogenesis and event triggering and new circadian rhythm-based interventions to improve the clinical management and quality of life of at-risk patients.

### Practice points

- Most patients show significant variation in SBP and DBP during the 24 h, with different 24-h patterns exhibited by different patients.
- ABPM devices are needed to properly assess BP levels throughout the 24 h to make an accurate diagnosis of hypertension as well as to evaluate the attainment of treatment goals.
- The sleep-time mean BP level and the extent of the sleep-time mean BP decline relative to the daytime mean level are sensitive predictors of CVD risk and thus worthy therapeutic targets.
- The 24-h pattern in MI arises from circadian rhythms in underlying pathophysiologic mechanisms and coincident time-of-day cyclic variations in environmental stressors.
- In many patients, the manifestation of supraventricular (atrial arrhythmias, PSVT) and ventricular (VPBs, VT and VF) cardiac arrhythmias exhibits significant circadian rhythmicity.
- The ingestion time (with reference to the CTS) of BP and heart medications can be a significant determinant of the magnitude of their beneficial effects and also patient tolerance.

### Research agenda

- Explore the role of the cardiomyocyte circadian clock in cardiac tissue pathology.
- Knowledge of circadian rhythms in cardiac arrhythmias and MI should be used to better design improved drug-delivery formulations and to explore the advantage of a chronotherapeutic approach.
- Future research of 24-h patterns in supraventricular and ventricular cardiac arrhythmias must control for key factors extraneous to intrinsic arrhythmogenic activity, such as consumption of alcohol, caffeine and other sympathomimetic substances, time and dose of prescription medication, plus differences between subjects in sleep–wake routine which affect the phasing of involved endogenous circadian rhythm.

### References

1. Duguay D, Cermakian N. The crosstalk between physiology and circadian clock proteins. *Chronobiol Int* 2009;**26**:1479–513.
2. Balsalobre A, Damiola F, Schibler U. A serum shock induces circadian gene expression in mammalian tissue culture cells. *Cell* 1998;**93**:929–37.
3. Durgan DJ, Hotze MA, Tomlin TM, Egbejimi O, Graveleau C, Abel ED, et al. The intrinsic circadian clock within the cardiomyocyte. *Am J Physiol Heart Circ Physiol* 2005;**289**:H1530–41.
4. McNamara P, Seo SB, Rudic RD, Sehgal A, Chakravarti D, FitzGerald GA. Regulation of CLOCK and MOP4 by nuclear hormone receptors in the vasculature: a humoral mechanism to reset a peripheral clock. *Cell* 2001;**105**:877–89.
5. Nonaka H, Emoto N, Ikeda K, Fukuya H, Rohman MS, Raharjo SB, et al. Angiotensin II induces circadian gene expression of clock genes in cultured vascular smooth muscle cells. *Circulation* 2001;**104**:1746–8.

\* The most important references are denoted by an asterisk.

6. Takeda N, Maemura K, Horie S, Oishi K, Imai Y, Harada T, et al. Thrombomodulin is a clock-controlled gene in vascular endothelial cells. *J Biol Chem* 2007;**282**:32561–7.
7. Paschos GK, FitzGerald GA. Circadian clocks and vascular function. *Circ Res* 2010;**106**:833–41.
8. Rudic RD, Fulton DJ. Pressed for time: the circadian clock and hypertension. *J Appl Physiol* 2009;**107**:1328–38.
9. Young ME. The circadian clock within the heart: potential influence on myocardial gene expression, metabolism, and function. *Am J Physiol Heart Circ Physiol* 2006;**290**:H1–16.
10. Curtis AM, Cheng Y, Kapoor S, Reilly D, Price TS, Fitzgerald GA. Circadian variation of blood pressure and the vascular response to asynchronous stress. *Proc Natl Acad Sci USA* 2007;**104**:3450–5.
11. Viola AU, James LM, Archer SN, Dijk DJ. PER3 polymorphism and cardiac autonomic control: effects of sleep debt and circadian phase. *Am J Physiol Heart Circ Physiol* 2008;**295**:H2156–63.
12. Wang N, Yang G, Jia Z, Zhang H, Aoyagi T, Soodvilai S, et al. Vascular PPARgamma controls circadian variation in blood pressure and heart rate through Bmal1. *Cell Metab* 2008;**8**:482–91.
13. Bray MS, Shaw CA, Moore MW, Garcia RA, Zanquetta MM, Durgan DJ, et al. Disruption of the circadian clock within the cardiomyocyte influences myocardial contractile function, metabolism, and gene expression. *Am J Physiol Heart Circ Physiol* 2008;**294**:H1036–47.
14. Durgan DJ, Puliniilkunnil T, Villegas-Montoya C, Garvey ME, Frangogiannis NG, Michael LH, et al. Short communication: ischemia/reperfusion tolerance is time-of-day-dependent: mediation by the cardiomyocyte circadian clock. *Circ Res* 2010;**106**:546–50.
15. Tsai JY, Kienesberger PC, Puliniilkunnil T, Sailors MH, Durgan DJ, Villegas-Montoya C, et al. Direct regulation of myocardial triglyceride metabolism by the cardiomyocyte circadian clock. *J Biol Chem* 2010;**285**:2918–29.
16. Young ME. Anticipating anticipation: pursuing identification of cardiomyocyte circadian clock function. *J Appl Physiol* 2009;**107**:1339–47.
17. Young ME, Bray MS. Potential role for peripheral circadian clock dysynchrony in the pathogenesis of cardiovascular dysfunction. *Sleep Med* 2007;**8**:656–67.
18. Durgan DJ, Trexler NA, Egbejimi O, McElfresh TA, Suk HY, Petterson LE, et al. The circadian clock within the cardiomyocyte is essential for responsiveness of the heart to fatty acids. *J Biol Chem* 2006;**281**:24254–69.
19. Kohsaka A, Laposky AD, Ramsey KM, Estrada C, Joshi C, Kobayashi Y, et al. High-fat diet disrupts behavioral and molecular circadian rhythms in mice. *Cell Metab* 2007;**6**:414–21.
20. Kung TA, Egbejimi O, Cui J, Ha NP, Durgan DJ, Essop MF, et al. Rapid attenuation of circadian clock gene oscillations in the rat heart following ischemia-reperfusion. *J Mol Cell Cardiol* 2007;**43**:744–53.
21. Kunieda T, Minamino T, Katsuno T, Tateno K, Nishi J, Miyauchi H, et al. Cellular senescence impairs circadian expression of clock genes in vitro and in vivo. *Circ Res* 2006;**98**:532–9.
22. Mohri T, Emoto N, Nonaka H, Fukuya H, Yagita K, Okamura H, et al. Alterations of circadian expressions of clock genes in Dahl salt-sensitive rats fed a high-salt diet. *Hypertension* 2003;**42**:189–94.
23. Young ME, Razeghi P, Taegtmeyer H. Clock genes in the heart: characterization and attenuation with hypertrophy. *Circ Res* 2001;**88**:1142–50.
24. Young ME, Wilson CR, Razeghi P, Guthrie PH, Taegtmeyer H. Alterations of the circadian clock in the heart by streptozotocin-induced diabetes. *J Mol Cell Cardiol* 2002;**34**:223–31.
25. Knutsson A, Akerstedt T, Jonsson BG, Orth-Gomer K. Increased risk of ischaemic heart disease in shift workers. *Lancet* 1986;**2**:89–92.
26. Martino TA, Tata N, Belsham DD, Chalmers J, Straume M, Lee P, et al. Disturbed diurnal rhythm alters gene expression and exacerbates cardiovascular disease with rescue by resynchronization. *Hypertension* 2007;**49**:1104–13.
27. Martino TA, Oudit GY, Herzenberg AM, Tata N, Koletar MM, Kabir GM, et al. Circadian rhythm disorganization produces profound cardiovascular and renal disease in hamsters. *Am J Physiol Regul Integr Comp Physiol* 2008;**294**:R1675–83.
28. Sole MJ, Martino TA. Diurnal physiology: core principles with application to the pathogenesis, diagnosis, prevention, and treatment of myocardial hypertrophy and failure. *J Appl Physiol* 2009;**107**:1318–27.
29. Durgan DJ, Tsai JY, Grenett MH, Pat BM, Ratcliffe WF, Villegas-Montoya C, et al. Evidence suggesting that the cardiomyocyte circadian clock modulates responsiveness of the heart to hypertrophic stimuli in mice. *Chronobiol Int* 2011;**28**:187–203.
30. Portaluppi F, Smolensky MH. Circadian rhythm and environmental determinants of blood pressure regulation in normal and hypertensive conditions. In: White WB, editor. *Blood pressure monitoring in cardiovascular medicine and therapeutics*. Totowa (New Jersey): Humana Press; 2001. p. 79–138.
31. Portaluppi F, Vergnani L, Manfredini R, Degli Uberti EC, Fersini C. Time-dependent effect of isradipine on the nocturnal hypertension of chronic renal failure. *Am J Hypertens* 1995;**8**:719–26.
- \*32. Portaluppi F, Manfredini R, Fersini C. From a static to a dynamic concept of risk: the circadian epidemiology of cardiovascular events. *Chronobiol Int* 1999;**16**:33–49.
33. Snyder F, Hobson A, Morrison DF, Goldfrank F. Changes in respiration, heart rate and systolic blood pressure in human sleep. *J Appl Physiol* 1964;**19**:417–22.
34. Reinberg A, Ghata J, Halberg F, Gervais P, Abulker C, Dupont J, et al. Rythmes circadiens du pouls, de la pression arterielle, des excretions urinaires en 17-hydroxycorticosteroides catecholamines et potassium chez l'homme adulte sain, actif et au repos. *Ann Endocrinol (Paris)* 1970;**31**:277–87.
35. Tuck ML, Stern N, Sowers JR. Enhanced 24-hour norepinephrine and renin secretion in young patients with essential hypertension: relation with the circadian pattern of arterial blood pressure. *Am J Cardiol* 1985;**55**:112–5.
36. Davies AB, Gould BA, Cashman PM, Raftery EB. Circadian rhythm of blood pressure in patients dependent on ventricular demand pacemakers. *Br Heart J* 1984;**52**:93–8.
37. Broadhurst P, Brigden G, Dasgupta P, Lahiri A, Raftery EB. Ambulatory intra-arterial blood pressure in normal subjects. *Am Heart J* 1990;**120**:160–6.
38. Kario K, Pickering TG, Umeda Y, Hoshida S, Hoshida Y, Morinari M, et al. Morning surge in blood pressure as a predictor of silent and clinical cerebrovascular disease in elderly hypertensives: a prospective study. *Circulation* 2003;**107**:1401–6.
39. Mann S, Altman DG, Raftery EB, Bannister R. Circadian variation of blood pressure in autonomic failure. *Circulation* 1983;**68**:477–83.
40. Martinelli P, Coccagna G, Rizzuto N, Lugaresi E. Changes in systemic arterial pressure during sleep in Shy-Drager syndrome. *Sleep* 1981;**4**:139–46.
41. Tohgi H, Chiba K, Kimura M. Twenty-four-hour variation of blood pressure in vascular dementia of theBinswanger type. *Stroke* 1991;**22**:603–8.
42. Otsuka A, Mikami H, Katahira K, Nakamoto Y, Minamitani K, Imaoka M, et al. Absence of nocturnal fall in blood pressure in elderly persons with Alzheimer-type dementia. *J Am Geriatr Soc* 1990;**38**:973–8.
43. Tominaga M, Tsuchihashi T, Kinoshita H, Abe I, Fujishima M. Disparate circadian variations of blood pressure and body temperature in bedridden elderly patients with cerebral atrophy. *Am J Hypertens* 1995;**8**:773–81.
44. Shimada K, Kawamoto A, Matsubayashi K, Nishinaga M, Kimura S, Ozawa T. Diurnal blood pressure variations and silent cerebrovascular damage in elderly patients with hypertension. *J Hypertension* 1992;**10**:875–8.
45. Sander D, Klingelhöfer J. Changes of circadian blood pressure patterns after hemodynamic and thromboembolic brain infarction. *Stroke* 1994;**25**:1730–7.
46. Asmar RG, Julia PL, Mascarel VL, Fabiani JN, Benetos A, Safar ME. Ambulatory blood pressure profile after carotid endarterectomy in patients with ischaemic arterial disease. *J Hypertens* 1994;**12**:697–702.
47. Franklin SS, Sowers JR, Batzdorf U. Relationship between arterial blood pressure and plasma norepinephrine levels in a patient with neurogenic hypertension. *Am J Med* 1986;**81**:1105–7.
48. Portaluppi F, Cortelli P, Avoni P, Vergnani L, Contin E, Maltoni P, et al. Diurnal blood pressure variation and hormonal correlates in fatal familial insomnia. *Hypertension* 1994;**23**:569–76.
49. Hornung RS, Mahler RF, Raftery EB. Ambulatory blood pressure and heart rate in diabetic patients: an assessment of autonomic function. *Diabet Med* 1989;**6**:579–85.
50. Imai Y, Abe K, Miura Y, Nihei M, Sasaki S, Minami N, et al. Hypertensive episodes and circadian fluctuations of blood pressure in patients with pheochromocytoma: studies by long-term blood pressure monitoring based on a volume-oscillometric method. *J Hypertens* 1988;**6**:9–15.
51. Statius van Eps RG, van den Meiracker AH, Boomsma F, Man in't Veld AJ, Schalekamp MADH. Diurnal variation of blood pressure in patients with catecholamine-producing tumors. *Am J Hypertens* 1994;**7**:492–7.
52. Imai Y, Abe K, Sasaki S, Minami N, Nihei M, Munakata M, et al. Altered circadian blood pressure rhythm in patients with Cushing's syndrome. *Hypertension* 1988;**12**:11–9.
53. Imai Y, Abe K, Sasaki S, Minami N, Munakata M, Nihei M, et al. Exogenous glucocorticoid eliminates or reverses circadian blood pressure variations. *J Hypertens* 1989;**7**:113–20.
54. White WB, Malchoff C. Diurnal blood pressure variability in mineralocorticoid excess syndrome. *Am J Hypertens* 1992;**5**:414–8.
55. Fallo F, Fanelli G, Cipolla A, Betterle C, Boscaro M, Sonino N. 24-hour blood pressure profile in Addison's disease. *Am J Hypertens* 1994;**7**:1105–9.
56. Brickman AS, Stern N, Sowers JR. Circadian variations of catecholamines and blood pressure in patients with pseudohypoparathyroidism and hypertension. *Chronobiologia* 1990;**17**:37–44.
57. Tilkian AG, Guilleminault C, Schroeder JS, Lehrman KL, Simmons FB, Dement WC. Hemodynamics in sleep-induced apnea Studies during wakefulness and sleep. *Ann Intern Med* 1976;**85**:714–9.
58. Franz IW, Erb D, Tonnesmann U. Gestörte 24-Stunden-Blutdruckrhythmik bei normotensiven und hypertensiven Asthmatikern. *Z Kardiol* 1992;**81**:13–6.
59. Baumgart P, Walger P, Gerke M, Dorst KG, Vetter H, Rahn KH. Nocturnal hypertension in renal failure, haemodialysis and after renal transplantation. *J Hypertens* 1989;**7**(Suppl. 6):S70–1.
60. Portaluppi F, Montanari L, Massari M, Di Chiara V, Capanna M. Loss of nocturnal decline of blood pressure in hypertension due to chronic renal failure. *Am J Hypertens* 1991;**4**:20–6.
61. Shaw DB, Knapp MS, Davies DH. Variations in blood pressure in hypertensives during sleep. *Lancet* 1963;**1**:797–9.
62. de la Sierra A, del Mar Lluch M, Coca A, Aguilera MT, Sánchez M, Sierra C, et al. Assessment of salt sensitivity in essential hypertension by 24-h ambulatory blood pressure monitoring. *Am J Hypertens* 1995;**8**:970–7.
63. Olofsson P. Characteristics of a reversed circadian blood pressure rhythm in pregnant women with hypertension. *J Hum Hypertens* 1995;**9**:565–70.
64. Redman CW, Beilin LJ, Bonnar J. Reversed diurnal blood pressure rhythm in hypertensive pregnancies. *Clin Sci Mol Med* 1976;**3**:687s–9s.



65. Verdecchia P, Schillaci G, Guerrieri M, Gatteschi C, Benemio G, Boldrini F, et al. Circadian blood pressure changes and left ventricular hypertrophy in essential hypertension. *Circulation* 1990;**81**:528–36.
66. Soergel M, Maisin A, Azancot-Benisty A, Loirat C. Ambulante Blutdruckmessung bei nierentransplantierten Kindern und Jugendlichen. *Z Kardiol* 1992;**81**:67–70.
67. van de Borne P, Gelin M, Van de Stadt J, Degaute JP. Circadian rhythms of blood pressure after liver transplantation. *Hypertension* 1993;**21**:398–405.
68. Reeves RA, Shapiro AP, Thompson ME, Johnsen AM. Loss of nocturnal decline in blood pressure after cardiac transplantation. *Circulation* 1986;**73**:401–8.
69. Caruana MP, Lahiri A, Cashman PM, Altman DG, Raftery EB. Effects of chronic congestive heart failure secondary to coronary artery disease on the circadian rhythm of blood pressure and heart rate. *Am J Cardiol* 1988;**62**:755–9.
70. Portaluppi F, Montanari L, Ferlini M, Vergnani L, Bagni B, Degli Uberti EC. Differences in blood pressure regulation of congestive heart failure, before and after treatment, correlate with changes in the circulating pattern of atrial natriuretic peptide. *Eur Heart J* 1992;**13**:990–6.
71. van de Borne P, Tielemans C, Vanherweghem JL, Degaute JP. Effect of recombinant human erythropoietin therapy on ambulatory blood pressure and heart rate in chronic haemodialysis patients. *Nephrol Dial Transplant* 1992;**7**:45–9.
72. Sander D, Klingelhofer J. Circadian blood pressure patterns in four cases with hemodynamic brain infarction and prolonged blood–brain barrier disturbance. *Clin Neurol Neurosurg* 1993;**95**:221–9.
73. Degaute JP, Van Cauter E, van de Borne P, Linkowski P. Twenty-four-hour blood pressure and heart rate profiles in humans. A twin study. *Hypertension* 1994;**23**:244–53.
74. Lemmer B, Mattes A, Bohm M, Ganten D. Circadian blood pressure variation in transgenic hypertensive rats. *Hypertension* 1993;**22**:97–101.
75. Portaluppi F, Waterhouse J, Minors D. The rhythms of blood pressure in humans. Exogenous and endogenous components and implications for diagnosis and treatment. *Ann N Y Acad Sci* 1996;**783**:1–9.
76. Clark LA, Denby L, Pregibon D, Harshfield GA, Pickering TG, Blank S, et al. A quantitative analysis of the effects of activity and time of day on the diurnal variations of blood pressure. *J Chronic Dis* 1987;**40**:671–81.
77. Sundberg S, Kohvakka A, Gordin A. Rapid reversal of circadian blood pressure rhythm in shift workers. *J Hypertens* 1988;**6**:393–6.
78. Goto T, Yokoyama K, Araki T, Miura T, Saitoh H, Saitoh M, et al. Identical blood pressure levels and slower heart rates among nurses during night work and day work. *J Hum Hypertens* 1994;**8**:11–4.
79. Fava C, Burri P, Almgren P, Groop L, Hulthen UL, Melander O. Heritability of ambulatory and office blood pressure phenotypes in Swedish families. *J Hypertens* 2004;**22**:1717–21.
80. Fava C, Burri P, Almgren P, Arcaro G, Groop L, Lennart Hulthen U, et al. Dipping and variability of blood pressure and heart rate at night are heritable traits. *Am J Hypertens* 2005;**18**:1402–7.
81. Janssen BJ, Tyssen CM, Duindam H, Rietveld WJ. Suprachiasmatic lesions eliminate 24-h blood pressure variability in rats. *Physiol Behav* 1994;**55**:307–11.
82. Witte K, Schnecko A, Buijs R, Lemmer B. Circadian rhythms in blood pressure and heart rate in SCN-lesioned and unlesioned transgenic hypertensive rats [abstract]. *Biol Rhythm Res* 1995;**26**:458–9.
83. Smolensky MH, Hermida RC, Castriotta RJ, Portaluppi F. Role of sleep–wake cycle on blood pressure circadian rhythms and hypertension. *Sleep Med* 2007;**8**:668–80.
84. Coccagna G, Mantovani M, Brignani F, Manzini A, Lugaresi E. Laboratory note. Arterial pressure changes during spontaneous sleep in man. *Electroencephalogr Clin Neurophysiol* 1971;**31**:277–81.
85. Portaluppi F, Cortelli P, Provini F, Plazzi G, Manfredini R, Lugaresi E. Alterations of sleep and circadian blood pressure profile. *Blood Press Monit* 1997;**2**:301–13.
86. Portaluppi F, Vergnani L, Manfredini R, Fersini C. Endocrine mechanisms of blood pressure rhythms. *Ann N Y Acad Sci* 1996;**783**:113–31.
87. Suzuki Y, Kuwajima I, Mitani K, Miyao M, Uno A, Matsushita S, et al. The relation between blood pressure variation and daily physical activity in early morning surge in blood pressure. *Nippon Ronen Igakkai Zasshi* 1993;**30**:841–8.
88. Ferretti R, Charniak NS, Longobardo G, Levine OR, Morkin E, Singer DH, et al. Systemic and pulmonary vasomotor waves. *J Physiol (Lond)* 1965;**209**:37–50.
89. Charniak NS, Longobardo GS. Cheyne–Stokes breathing an instability in physiologic control. *N Engl J Med* 1973;**288**:952–7.
90. Lugaresi E, Coccagna G, Mantovani M, Lebrun R. Some periodic phenomena arising during drowsiness and sleep in man. *Electroencephalogr Clin Neurophysiol* 1972;**32**:701–5.
91. Phillipson EA. Control of breathing during sleep. *Am Rev Respir Dis* 1978;**118**:909–39.
92. Somers VK, Dyken ME, Mark AL, Abboud FM. Sympathetic-nerve activity during sleep in normal subjects. *N Engl J Med* 1993;**328**:303–7.
93. Conway J, Boon N, Davies C, Jones JV, Sleight P. Neural and humoral mechanisms involved in blood pressure variability. *J Hypertens* 1984;**2**:203–8.
94. Krum H, Louis WJ, Brown DJ, Jackman GP, Howes LG. Diurnal blood pressure variation in quadriplegic chronic spinal cord injury patients. *Clin Sci* 1991;**80**:271–6.
95. Coca A. Circadian rhythm and blood pressure control: physiological and pathophysiological factors. *J Hypertens* 1994;**12**(Suppl. 5):S13–21.
96. Dodt C, Breckling U, Derad I, Fehm HL, Born J. Plasma epinephrine and norepinephrine concentrations of healthy humans associated with nighttime sleep and morning arousal. *Hypertension* 1997;**30**:71–6.
97. Kirby DA, Verrier RL. Differential effects of sleep stage on coronary hemodynamic function during stenosis. *Physiol Behav* 1989;**45**:1017–20.
98. King MJ, Zir LM, Kaltman AJ, Fox AC. Variant angina associated with angiographically demonstrated coronary artery spasm and REM sleep. *Am J Med Sci* 1973;**265**:419–22.
99. Nowlin JB, Troyer WG, Collins WS. The association of nocturnal angina pectoris with dreaming. *Ann Intern Med* 1965;**63**:1040–6.
100. Tamura K, Karacan I, Williams RL, Meyer JS. Disturbances of the sleep–waking cycle in patients with vascular brain stem lesions. *Clin Electroencephalogr* 1983;**14**:35–46.
101. Ziegler B, Schneider E. Sleep EEG studies in ischemic brain stem lesions preliminary report. *Bull Soc Sci Med Grand Duché Luxemb* 1983;**120**:65–70.
102. Gadoth N, Sokol J, Lavie P. Sleep structure and nocturnal disordered breathing in familial dysautonomia. *J Neurol Sci* 1983;**60**:117–25.
103. McNicholas WT, Rutherford R, Grossman R, Moldofsky H, Zamel N, Phillipson EA. Abnormal respiratory pattern generation during sleep in patients with autonomic dysfunction. *Am Rev Respir Dis* 1983;**128**:429–33.
104. Sforza E, Zucconi M, Petronelli R, Lugaresi E, Cirignotta F. REM sleep behavioral disorders. *Eur Neurol* 1988;**28**:295–300.
105. Kavey NB, Whyte J, Blitzer A, Gidro-Frank S. Sleep-related laryngeal obstruction presenting as snoring or sleep apnea. *Laryngoscope* 1989;**99**:851–4.
106. Moldofsky H, Lue FA, Smythe HA. Alpha EEG sleep and morning symptoms in rheumatoid arthritis. *J Rheumatol* 1983;**10**:373–9.
107. Mahowald MW, Mahowald ML, Bundlie SR, Ytterberg SR. Sleep fragmentation in rheumatoid arthritis. *Arthritis Rheum* 1989;**32**:974–83.
108. Wooten V. Medical causes of insomnia. In: Kryger MH, Roth T, Dement WC, editors. *Principles and practice of sleep medicine*. 2nd ed. Philadelphia (Pennsylvania): W.B. Saunders; 1994. p. 509–22.
109. Passouant P, Passouant-Fontaine T, Cadilhac J. Influence on sleep of hyperthyroidism. *Electroencephalogr Clin Neurophysiol* 1967;**23**:283–4.
110. Milcu SM, Nicolescu-Catargi A. Deep sleep phase alterations in patients with loss of the adrenal secretory rhythm and in hypophysectomized patients (The effects of hypothalamo–hypophyseal stressing by metoppyron). *Electroencephalogr Clin Neurophysiol* 1967;**22**:574.
111. Krieger DT, Gewirtz GP. Recovery of hypothalamic–pituitary–adrenal function, growth hormone responsiveness and sleep EEG pattern in a patient following removal of an adrenal cortical adenoma. *J Clin Endocrinol Metab* 1974;**38**:1075–82.
112. Weitzman ED, Boyar RM, Kapen S, Hellman L. The relationship of sleep and sleep stages to neuroendocrine secretion and biological rhythms in man. *Recent Prog Horm Res* 1975;**31**:399–446.
113. Friess E, Wiedemann K, Steiger A, Holsboer F. The hypothalamic–pituitary–adrenocortical system and sleep in man. *Adv Neuroimmunol* 1995;**5**:111–25.
114. Lugaresi E, Medori R, Montagna P, Baruzzi A, Cortelli P, Lugaresi A, et al. Fatal familial insomnia and dysautonomia with selective degeneration of thalamic nuclei. *N Engl J Med* 1986;**315**:997–1003.
115. Manetto V, Medori R, Cortelli P, Montagna P, Tinuper P, Baruzzi A, et al. Fatal familial insomnia: clinical and pathologic study of five new cases. *Neurology* 1992;**42**:312–9.
116. Juniper EF. Rhinitis management: the patient's perspective. *Clin Exp Allergy* 1998;**28**(Suppl. 6):34–8.
117. Craig TJ, Hanks CD, Fisher LH. How do topical nasal corticosteroids improve sleep and daytime somnolence in allergic rhinitis? *J Allergy Clin Immunol* 2005;**116**:1264–6.
118. Juniper EF, Guyatt GH. Development and testing of a new measure of health status for clinical trials in rhinoconjunctivitis. *Clin Exp Allergy* 1991;**21**:77–83.
119. Bender BG, Leung SB, Leung DY. Actigraphy assessment of sleep disturbance in patients with atopic dermatitis: an objective life quality measure. *J Allergy Clin Immunol* 2003;**111**:598–602.
120. Turner-Warwick M. Epidemiology of nocturnal asthma. *Am J Med* 1988;**85**:6–8.
121. Sadeh A, Horowitz I, Wolach-Benodis L, Wolach B. Sleep and pulmonary function in children with well-controlled, stable asthma. *Sleep* 1998;**21**:379–84.
122. Bellamy N, Sothorn RB, Campbell J. Rhythmic variations in pain perception in osteoarthritis of the knee. *J Rheumatol* 1990;**17**:364–72.
123. Labrecque G, Vanier MC. Rhythms, pain and pain management. In: Redfern P, editor. *Chronotherapeutics*. London (U.K.): Pharmaceutical Press; 2003. p. 211–34.
124. Moore JG. Rhythms and therapeutics of diseases of the gastrointestinal tract. In: Redfern P, editor. *Chronotherapeutics*. London (U.K.): Pharmaceutical Press; 2003. p. 127–51.
125. Kroetz C. Ein biologischer 24-Stunden-Rhythmus des Blutkreislaufs bei Gesundheit und bei Herzschwäche zugleich ein Beitrag zur tageszeitlichen Häufung einiger akuter Kreislaufstörungen. *Munch Med Wochenschr* 1940;**87**:314–7.
126. McKeigue PM, Reynard JM. Relation of nocturnal polyuria of the elderly to essential hypertension. *Lancet* 2000;**355**:486–8.

127. Chasens ER, Umlauf MG. Nocturia: a problem that disrupts sleep and predicts obstructive sleep apnea. *Geriatr Nur (Lond)* 2003;**24**:76–81. 105.
128. Lose G, Mattiasson A, Walter S, Lalos O, van Kerrebroeck P, Abrams P, et al. Clinical experiences with desmopressin for long-term treatment of nocturia. *J Urol* 2004;**172**:1021–5.
129. Gangwisch JE, Heymsfield SB, Boden-Albala B, Buys RM, Kreier F, Pickering TG, et al. Short sleep duration as a risk factor for hypertension: analyses of the first National Health and Nutrition Examination Survey. *Hypertension* 2006;**47**:833–9.
130. Ringler J, Garpestad E, Basner RC, Weiss JW. Systemic blood pressure elevation after airway occlusion during NREM sleep. *Am J Respir Crit Care Med* 1994;**150**:1062–6.
131. Calverley PMA, Rees K. Systemic arterial blood pressure during obstructive sleep apnoea. *J Sleep Res* 1995;**4**:93–6.
132. Guilleminault C, Stoohs R. Arousal, increased respiratory efforts, blood pressure and obstructive sleep apnoea. *J Sleep Res* 1995;**4**:117–24.
133. Lugaesi E, Mondini S, Zucconi M, Montagna P, Cirignotta F. Staging of heavy snorers' disease: a proposal. *Bull Eur Physiopathol Respir* 1983;**19**:590–4.
- \*134. Cortelli P, Pierangeli G, Provini F, Plazzi G, Lugaesi E. Blood pressure rhythms in sleep disorders and dysautonomia. *Ann N Y Acad Sci* 1996;**783**:204–21.
135. Cortelli P, Parchi P, Sforza E, Contin M, Pierangeli G, Barletta G, et al. Cardiovascular autonomic dysfunction in normotensive awake subjects with obstructive sleep apnoea syndrome. *Clin Auton Res* 1994;**4**:57–62.
136. Guilleminault C, Tilkin A, Dement WC. The sleep apnea syndromes. *Annu Rev Med* 1976;**27**:465–84.
137. Kales A, Cadieux RJ, Bixler EO, Soldatos CR, Vela-Bueno A, Misoul CA, et al. Severe obstructive sleep apnea I: onset, clinical course, and characteristics. *J Chronic Dis* 1985;**38**:419–25.
138. Kales A, Bixler EO, Cadieux RJ, Schneck DW, Shaw LC, Locke TW, et al. Sleep apnoea in a hypertensive population. *Lancet* 1984;**2**:1005–8.
139. Williams AJ, Houston D, Finberg S, Lam C, Kinney JL, Santiago S. Sleep apnea syndrome and essential hypertension. *Am J Cardiol* 1985;**55**:1019–22.
140. Fletcher EC. An animal model of the relationship between systemic hypertension and repetitive episodic hypoxia as seen in sleep apnoea. *J Sleep Res* 1995;**4**:71–7.
141. Ehlenz K. Regulation of blood volume implications for cardiovascular pathophysiology in sleep apnoea. *J Sleep Res* 1995;**4**:30–3.
142. Wilcox I, Collins FL, Grunstein RR, Hedner J, Kelly DT, Sullivan CE. Relationship between chemosensitivity, obesity and blood pressure in obstructive sleep apnoea. *Blood Press* 1994;**3**:47–54.
143. Lugaesi E, Partinen M. Prevalence of snoring. In: Saunders NA, Sullivan CE, editors. *Sleep and breathing*. New York: Marcel Dekker; 1994. p. 337–61.
144. Portaluppi F, Provini F, Cortelli P, Plazzi G, Bertozzi N, Manfredini R, et al. Undiagnosed sleep-disordered breathing in male nondippers with essential hypertension. *J Hypertens* 1997;**15**:1227–33.
145. Bradbury S, Eggleston C. Postural hypotension a report of 3 cases. *Am Heart J* 1925;**1**:73.
146. Graham JG, Oppenheimer DR. Orthostatic hypotension and nicotine sensitivity in a case of multiple system atrophy. *J Neurol Neurosurg Psychiatry* 1969;**32**:28–34.
147. Delerive J, Thomas AA. L'atrophie olivo-ponto-cérébelleuse. *Nouv Iconog Salpetrière* 1900;**13**:330–70.
148. Shy GM, Drager GA. A neurological syndrome associated with orthostatic hypotension. *Arch Neurol* 1960;**2**:511–27.
149. Adams RD, Van Bogaert L, Van Der Eecken H. Dégénérescences nigro-striées et cérébello-nigro-striées. *Psychiatr Neurol (Basel)* 1961;**142**:219–59.
150. Wenning GK, Ben Shlomo Y, Magalhaes M, Daniel SE, Quinn NP. Clinical features and natural history of multiple system atrophy an analysis of 100 cases. *Brain* 1994;**117**:835–45.
151. Montagna P, Cortelli P, Avoni P, Tinuper P, Plazzi G, Gallassi R, et al. Clinical features of fatal familial insomnia: phenotypic variability in relation to a polymorphism at codon 129 of the prion protein gene. *Brain Pathol* 1998;**8**:515–20 [see comments].
152. Van Cauter E, Blackman JD, Roland D, Spire JP, Refetoff S, Polonsky KS. Modulation of glucose regulation and insulin secretion by circadian rhythmicity and sleep. *J Clin Invest* 1991;**88**:934–42.
153. Follenius M, Brandenberger G, Badesapt JJ, Libert JP, Ehrhart J. Nocturnal cortisol release in relation to sleep structure. *Sleep* 1992;**15**:21–7.
154. Späth-Schwalbe E, Uthgenannt D, Voget G, Kern W, Born J, Fehm H-L. Corticotropin-releasing hormone induced adrenocorticotropin and cortisol secretion depends on sleep and wakefulness. *J Clin Endocrinol Metab* 1993;**77**:1170–3.
155. Portaluppi F, Cortelli P, Avoni P, Vergnani L, Maltoni P, Pavani A, et al. Progressive disruption of the circadian rhythm of melatonin in fatal familial insomnia. *J Clin Endocrinol Metab* 1994;**78**:1075–8.
156. Portaluppi F, Cortelli P, Avoni P, Vergnani L, Maltoni P, Pavani A, et al. Dissociated 24-h patterns of somatotropin and prolactin in fatal familial insomnia. *Neuroendocrinology* 1995;**61**:731–7.
- \*157. Tobler I, Gaus SE, Deboer T, Achermann P, Fischer M, Rülicke T, et al. Altered circadian activity rhythms and sleep in mice devoid of prion protein. *Nature* 1996;**380**:639–42.
158. Portaluppi F, Bagni B, Degli Uberti E, Montanari L, Cavallini R, Trasforini G, et al. Circadian rhythms of atrial natriuretic peptide, renin, aldosterone, cortisol, blood pressure and heart rate in normal and hypertensive subjects. *J Hypertens* 1990;**8**:85–95.
159. Portaluppi F, Trasforini G, Margutti A, Vergnani L, Ambrosio MR, Rossi R, et al. Circadian rhythm of calcitonin gene-related peptide in uncomplicated essential hypertension. *J Hypertens* 1992;**10**:1227–34.
160. Portaluppi F, Vergnani L, degli Uberti EC. Atrial natriuretic peptide and circadian blood pressure regulation: clues from a chronobiological approach. *Chronobiol Int* 1993;**10**:176–89.
161. Portaluppi F, Montanari L, Ferlini M, Gilli P. Altered circadian rhythms of blood pressure and heart rate in non-hemodialysis chronic renal failure. *Chronobiol Int* 1990;**7**:321–7.
162. Portaluppi F, Montanari L, Ferlini M, Vergnani L, D'Ambrosi A, Cavallini AR, et al. Consistent changes in the circadian rhythms of blood pressure and atrial natriuretic peptide in congestive heart failure. *Chronobiol Int* 1991;**8**:432–9.
163. Portaluppi F. The time factor in the treatment of hypertension. *J Hypertens* 1997;**15**:327–8.
164. Verdecchia P, Porcellati C, Schillaci G, Borgioni C, Ciucci A, Battistelli M, et al. Ambulatory blood pressure: an independent predictor of prognosis in essential hypertension. *Hypertension* 1994;**24**:793–801.
165. Ohkubo T, Hozawa A, Yamaguchi J, Kikuya M, Ohmori K, Michimata M, et al. Prognostic significance of the nocturnal decline in blood pressure in individuals with and without high 24-h blood pressure: the Ohasama study. *J Hypertens* 2002;**20**:2183–9.
166. Dolan E, Stanton A, Thijs L, Hinedi K, Atkins N, McClory S, et al. Superiority of ambulatory over clinic blood pressure measurement in predicting mortality: the Dublin outcome study. *Hypertension* 2005;**46**:156–61.
- \*167. Boggia J, Li Y, Thijs L, Hansen TW, Kikuya M, Bjorklund-Bodegard K, et al. Prognostic accuracy of day versus night ambulatory blood pressure: a cohort study. *Lancet* 2007;**370**:1219–29.
168. Kikuya M, Ohkubo T, Asayama K, Metoki H, Obara T, Saito S, et al. Ambulatory blood pressure and 10-year risk of cardiovascular and noncardiovascular mortality: the Ohasama study. *Hypertension* 2005;**45**:240–5.
169. Ben-Dov IZ, Kark JD, Ben-Ishay D, Mekler J, Ben-Arie L, Bursztyn M. Predictors of all-cause mortality in clinical ambulatory monitoring: unique aspects of blood pressure during sleep. *Hypertension* 2007;**49**:1235–41.
- \*170. Fagard RH, Celis H, Thijs L, Staessen JA, Clement DL, De Buyzere ML, et al. Daytime and nighttime blood pressure as predictors of death and cause-specific cardiovascular events in hypertension. *Hypertension* 2008;**51**:55–61.
171. Fan HQ, Li Y, Thijs L, Hansen TW, Boggia J, Kikuya M, et al. Prognostic value of isolated nocturnal hypertension on ambulatory measurement in 8711 individuals from 10 populations. *J Hypertens* 2010;**28**:2036–45.
172. Hansen TW, Li Y, Boggia J, Thijs L, Richart T, Staessen JA. Predictive role of the night time blood pressure. *Hypertension* 2011;**57**:3–10.
- \*173. Hermida RC, Ayala DE, Mojon A, Fernandez JR. Influence of circadian time of hypertension treatment on cardiovascular risk: results of the MAPEC study. *Chronobiol Int* 2010;**27**:1629–51.
174. Lemmer B, Portaluppi F. Chronopharmacology of cardiovascular diseases. In: Redfern PH, Lemmer B, editors. *Physiology and Pharmacology of biological rhythms*. Heidelberg (Germany): Springer Verlag; 1997. p. 251–97.
175. Labrecque G, Beauchamp D. Rhythms and pharmacokinetics. In: Redfern P, editor. *Chronotherapeutics*. London (U.K.): Pharmaceutical Press; 2003. p. 75–110.
176. Hermida RC, Ayala DE, Calvo C, Portaluppi F, Smolensky MH. Chronotherapy of hypertension: administration-time-dependent effects of treatment on the circadian pattern of blood pressure. *Adv Drug Deliv Rev* 2007;**59**:923–39.
- \*177. Smolensky MH, Hermida RC, Ayala DE, Tiseo R, Portaluppi F. Administration-time-dependent effects of blood pressure-lowering medications: basis for the chronotherapy of hypertension. *Blood Press Monit* 2010;**15**:173–80.
178. Witte K, Lemmer B. Rhythms and pharmacodynamics. In: Redfern P, editor. *Chronotherapeutics*. London (U.K.): Pharmaceutical Press; 2003. p. 111–26.
179. Hermida RC, Ayala DE, Fernandez JR, Portaluppi F, Fabbian F, Smolensky MH. Circadian rhythms: role in blood pressure regulation and optimization of hypertension treatment with ACE inhibitor and ARB medications. *Am J Hypertens* 2011;**24**:383–91.
180. Hermida RC, Calvo C, Ayala DE, Dominguez MJ, Covelo M, Fernandez JR, et al. Administration time-dependent effects of valsartan on ambulatory blood pressure in hypertensive subjects. *Hypertension* 2003;**42**:283–90.
181. Hermida RC, Ayala DE, Fontao MJ, Mojon A, Alonso I, Fernandez JR. Administration-time-dependent effects of spirapril on ambulatory blood pressure in uncomplicated essential hypertension. *Chronobiol Int* 2010;**27**:560–74.
182. Bankir L, Bochud M, Maillard M, Bovet P, Gabriel A, Burnier M. Nighttime blood pressure and nocturnal dipping are associated with daytime urinary sodium excretion in African subjects. *Hypertension* 2008;**51**:891–8.
- \*183. Portaluppi F, Smolensky MH. Perspectives on the chronotherapy of hypertension based on the results of the MAPEC study. *Chronobiol Int* 2010;**27**:1652–67.
184. Waters DD, Miller DD, Bouchard A, Bosch X, Theroux P. Circadian variation in variant angina. *Am J Cardiol* 1984;**54**:61–4.
185. Rocco MB. Timing and triggers of transient myocardial ischemia. *Am J Cardiol* 1990;**66**:18G–21G.
186. Mulcahy D, Keegan J, Crean P, Quyyumi A, Shapiro L, Wright C, et al. Silent myocardial ischaemia in chronic stable angina: a study of its frequency and characteristics in 150 patients. *Br Heart J* 1988;**60**:417–23.

187. Araki H, Koiwaya Y, Nakagaki O, Nakamura M. Diurnal distribution of ST-segment elevation and related arrhythmias in patients with variant angina: a study by ambulatory ECG monitoring. *Circulation* 1983;**67**:995–1000.
188. Joy M, Pollard CM, Nunan TO. Diurnal variation in exercise responses in angina pectoris. *Br Heart J* 1982;**48**:156–60.
189. Ogawa H, Yasue H, Oshima S, Okumura K, Matsuyama K, Obata K. Circadian variation of plasma fibrinopeptide. A level in patients with variant angina. *Circulation* 1989;**80**:1617–26.
190. Yasue H, Omote S, Takizawa A, Nagao M, Miwa K, Tanaka S. Circadian variation of exercise capacity in patients with Prinzmetal's variant angina: role of exercise-induced coronary arterial spasm. *Circulation* 1979;**59**:938–48.
191. Pell S, D'Alonzo CA. Acute myocardial infarction in a large industrial population: report of a 6-year study of 1356 cases. *JAMA* 1963;**185**:831–8.
192. Cohen MC, Rohitla KM, Lavery CE, Muller JE, Middleman MA. Meta analysis of the morning excess of acute myocardial infarction and sudden cardiac death. *Am J Cardiol* 1997;**79**:1512–6.
193. Selvi Y, Smolensky MH, Boysan M, Aydin A, Bersiroglu L, Atli A, et al. Role of patient chronotype on circadian pattern of myocardial infarction: a pilot study. *Chronobiol Int* 2011;**28**:371–7.
194. Hjalmarson A, Gilpin EA, Nicod P, Dittrich H, Henning H, Engler R, et al. Differing circadian patterns of symptom onset in subgroups of patients with acute myocardial infarction. *Circulation* 1989;**80**:267–75.
195. Gilpin EA, Hjalmarson A, Ross J. Subgroups of patients with atypical circadian patterns of symptom onset in acute myocardial infarction. *Am J Cardiol* 1990;**66**:7G–11G.
196. Behar S, Halabi M, Reicher-Reiss H, Zion M, Kaplinsky E, Mandelzweig L, et al. Circadian variation and possible external triggers of onset of myocardial infarction SPRINT study group. *Am J Med* 1993;**94**:395–400.
197. Muller JE, Ludmer PL, Willich SN, Tofler GH, Aylmer G, Klangos I, et al. Circadian variation in the frequency of sudden cardiac death. *Circulation* 1987;**75**:131–8.
- \*198. Portaluppi F, Smolensky MH. Circadian rhythmic and environmental determinants of 24-hour blood pressure regulation in normal and hypertensive conditions. In: White WB, editor. *Blood pressure monitoring in cardiovascular medicine and therapeutics*. 2nd ed. Totowa (New Jersey): Humana Press; 2007. p. 135–58.
199. Benhorin J, Banai S, Moriel M, Gavish A, Keren A, Stern S, et al. Circadian variations in ischemic threshold and their relation to the occurrence of ischemic episodes. *Circulation* 1993;**87**:808–14.
200. Ino-Oka E, Sekino H, Ohtaki Y, Inooka H, Sagawa K, Hashimoto J, et al. Circadian variation of the myocardial ischemic threshold in patients with ischemic heart disease. *Clin Exp Hypertens* 2007;**29**:3–12.
201. Deedwania PC. *Circadian rhythms of cardiovascular disorders*. Armonk (New York): Futura Publishing; 1997.
202. Turtan MB, Deegan T. Circadian variations of plasma catecholamine, cortisol and immunoreactive insulin concentrations in supine subjects. *Clin Chim Acta* 1974;**55**:389–97.
203. Tofler GH, Brezinski D, Schafer AI, Czeisler CA, Rutherford JD, Willich SN, et al. Concurrent morning increase in platelet aggregability and the risk of myocardial infarction and sudden cardiac death. *N Engl J Med* 1987;**316**:1514–8.
204. Gordon RD, Wolfe LK, Island DP, Liddle GW. A diurnal rhythm in plasma renin activity in man. *J Clin Invest* 1966;**45**:1587–92.
205. Fujita M, Franklin D. Diurnal changes in coronary blood flow in conscious dogs. *Circulation* 1987;**76**:488–91.
206. Panza JA, Epstein SE, Quyyumi AA. Circadian variation in vascular tone and its relation to alpha-sympathetic vasoconstrictor activity. *N Engl J Med* 1991;**325**:986–90.
207. Quyyumi AA, Panza JA, Diodati JG, Lakatos E, Epstein SE. Circadian variation in ischemic threshold. A mechanism underlying the circadian variation in ischemic events. *Circulation* 1992;**86**:22–8.
208. Weitzman ED, Fukushima D, Nogeire C, Roffwarg H, Gallagher TF, Hellman L. Twenty-four hour pattern of the episodic secretion of cortisol in normal subjects. *J Clin Endocrinol Metab* 1971;**33**:14–22.
209. Rocco MB, Nabel EG, Selwyn AP. Circadian rhythms and coronary artery disease. *Am J Cardiol* 1987;**59**:13C–7C.
210. Barash D, Silverman RA, Gennis P, Budner N, Matos M, Gallagher EJ. Circadian variation in the frequency of myocardial infarction and death associated with acute pulmonary edema. *J Emerg Med* 1989;**7**:119–21.
211. Hansen O, Johansson BW, Gullberg B. The clinical outcome of acute myocardial infarction is related to the circadian rhythm of myocardial infarction onset. *Angiology* 1993;**44**:509–16.
212. Fujita M, Araie E, Yamanishi K, Miwa K, Kida M, Nakajima H. Circadian variation in the success rate of intracoronary thrombolysis for acute myocardial infarction. *Am J Cardiol* 1993;**71**:1369–71.
213. Baraldo M. The influence of circadian rhythms on the kinetics of drugs in humans. *Expert Opin Drug Metab Toxicol* 2008;**4**:175–92.
- \*214. Portaluppi F, Lemmer B. Chronobiology and chronotherapy of ischemic heart disease. *Adv Drug Deliv Rev* 2007;**59**:952–65.
215. Ezeugo U, Glasser SP. Clinical benefits versus shortcomings of diltiazem once-daily in the chronotherapy of cardiovascular diseases. *Expert Opin Pharmacother* 2009;**10**:485–91.
216. Irwin JM, McCarthy EA, Wilkinson WE, Pritchett EL. Circadian occurrence of symptomatic paroxysmal supraventricular tachycardia in untreated patients. *Circulation* 1988;**77**:298–300.
217. Manfredini R, Gallerani M, Portaluppi F, Salmi R, Chierici F, Fersini C. Circadian variation in the occurrence of paroxysmal supraventricular tachycardia in clinically healthy subjects. *Chronobiol Int* 1995;**12**:55–61.
218. Aronow WS, Ahn C. Circadian variation of primary cardiac arrest or sudden cardiac death in patients aged 62 to 100 years (mean 82). *Am J Cardiol* 1993;**71**:1455–6.
219. Hohnloser SH, Zabel M, Just H, Raeder EA. Relation of diurnal variation of ventricular repolarization to ventricular ectopic activity and modification by sotalol. *Am J Cardiol* 1993;**71**:475–8.
220. Behrens S, Galecka M, Bruggemann T, Ehlers C, Willich SN, Ziss W, et al. Circadian variation of sustained ventricular tachyarrhythmias terminated by appropriate shocks in patients with an implantable cardioverter defibrillator. *Am Heart J* 1995;**130**:79–84.
221. Peters RW, Mitchell LB, Brooks MM, Echt DS, Barker AH, Capone R, et al. Circadian pattern of arrhythmic death in patients receiving encainide, flecainide or moricizine in the cardiac arrhythmia suppression trial (CAST). *J Am Coll Cardiol* 1994;**23**:283–9.
222. Englund A, Behrens S, Wegscheider K, Rowland E. Circadian variation of malignant ventricular arrhythmias in patients with ischemic and non-ischemic heart disease after cardioverter defibrillator implantation. European 7219 Jewel Investigators. *J Am Coll Cardiol* 1999;**34**:1560–8.
223. Huikuri HV, Poutiainen AM, Makkilä TH, Koistinen MJ, Airaksinen KE, Mitrani RD, et al. Dynamic behavior and autonomic regulation of ectopic atrial pacemakers. *Circulation* 1999;**100**:1416–22.
224. Coumel P. Paroxysmal atrial fibrillation: a disorder of autonomic tone? *Eur Heart J* 1994;**15**(Suppl. A):9–16.
225. Mitchell AR, Spurrell PA, Sulke N. Circadian variation of arrhythmia onset patterns in patients with persistent atrial fibrillation. *Am Heart J* 2003;**146**:902–7.
226. Pritchett EL, Smith MS, McCarthy EA, Lee KL. The spontaneous occurrence of paroxysmal supraventricular tachycardia. *Circulation* 1984;**70**:1–6.
227. Kupari M, Koskinen P, Leinonen H. Double-peaking circadian variation in the occurrence of sustained supraventricular tachyarrhythmias. *Am Heart J* 1990;**120**:1364–9.
228. Rostagno C, Taddei T, Paladini B, Modesti PA, Utari P, Bertini G. The onset of symptomatic atrial fibrillation and paroxysmal supraventricular tachycardia is characterized by different circadian rhythms. *Am J Cardiol* 1993;**71**:453–5.
229. Raeder EA. Circadian fluctuations in ventricular response to atrial fibrillation. *Am J Cardiol* 1990;**66**:1013–6.
230. Manfredini R, Gallerani M, Boari B, Maraldi C, La Cecilia O, Rafael Ferreira AB, et al. Morning preference in onset of symptomatic third-degree atrio-ventricular heart block. *Chronobiol Int* 2002;**19**:785–91.
231. Soo LH, Gray D, Young T, Hampton JR. Circadian variation in witnessed out of hospital cardiac arrest. *Heart* 2000;**84**:370–6.
232. Lown B, Tykocinski M, Garfein A, Brooks P. Sleep and ventricular premature beats. *Circulation* 1973;**48**:691–701.
233. Raeder EA, Hohnloser SH, Graboyes TB, Podrid PJ, Lampert S, Lown B. Spontaneous variability and circadian distribution of ectopic activity in patients with malignant ventricular arrhythmia. *J Am Coll Cardiol* 1988;**12**:656–61.
234. Siegel D, Black DM, Seeley DG, Hulley SB. Circadian variation in ventricular arrhythmias in hypertensive men. *Am J Cardiol* 1992;**69**:344–7.
235. Goldstein S, Zoble RG, Akiyama T, Cohen JD, Lancaster S, Liebson PR, et al. Relation of circadian ventricular ectopic activity to cardiac mortality. CAST Investigators. *Am J Cardiol* 1996;**78**:881–5.
236. Lanza GA, Cortellessa MC, Rebuzzi AG, Scabbia EV, Costalunga A, Tamburi S, et al. Reproducibility in circadian rhythm of ventricular premature complexes. *Am J Cardiol* 1990;**66**:1099–106.
237. Morrison GW, Kumar EB, Portal RW, Aber CP. Cardiac arrhythmias 48 hours before, during, and 48 hours after discharge from hospital following acute myocardial infarction. *Br Heart J* 1981;**45**:500–11.
238. Smith R, Johnson L, Rothfeld D, Zir L, Tharp B. Sleep and cardiac arrhythmias. *Arch Intern Med* 1972;**130**:751–3.
239. Lichstein E, Morganroth J, Harrist R, Hubble E. Effect of propranolol on ventricular arrhythmia. The beta-blocker heart attack trial experience. *Circulation* 1983;**67**(6 Pt 2):15–10.
240. Goldstein S. Effect of beta-adrenergic blocking agents on the circadian occurrence of ischemic cardiovascular events. *Am J Cardiol* 1990;**66**:63G–5G.
241. Lucente M, Rebuzzi AG, Lanza GA, Tamburi S, Cortellessa MC, Coppola E, et al. Circadian variation of ventricular tachycardia in acute myocardial infarction. *Am J Cardiol* 1988;**62**:670–4.
242. Twidale N, Taylor S, Heddle WF, Ayres BF, Tonkin AM. Morning increase in the time of onset of sustained ventricular tachycardia. *Am J Cardiol* 1989;**64**:1204–6.
243. Kozak M, Krivan L, Semrad B. Circadian variations in the occurrence of ventricular tachyarrhythmias in patients with implantable cardioverter defibrillators. *Pacing Clin Electrophysiol* 2003;**26**:731–5.
244. Weisfeldt ML, Everson-Stewart S, Sitlani C, Rea T, Aufderheide TP, Atkins DL, et al. for the Resuscitation Outcomes Consortium (ROC) Investigators. Ventricular tachyarrhythmias after cardiac arrest in public versus at home. *N Engl J Med* 2011;**364**:313–21.
245. Weil ZM, Karelin K, Su AJ, Barker JM, Norman GJ, Zhang N, et al. Time-of-day determines neuronal damage and mortality after cardiac arrest. *Neurobiol Disease* 2009;**36**:352–60.

246. Fries R, Hein S, Kn J. Reversed circadian rhythms of heart rate variability and morning peak occurrence of sustained ventricular tachyarrhythmias in patients with implanted cardioverter defibrillator. *Med Sci Monit* 2002;**8**:CR751–6.
247. Smolensky MH, Tatar SE, Bergman SA, Losman JG, Barnard CN, Dacso CC, et al. Circadian rhythmic aspects of human cardiovascular function: a review by chronobiologic statistical methods. *Chronobiologia* 1976;**3**:337–71.
248. Oda E, Aizawa Y, Arai Y, Shibata A. Diurnal variation of QT interval in patients with VVI pacemaker. *Tohoku J Exp Med* 1985;**145**:419–26.
249. Molnar J, Zhang F, Weiss J, Ehler FA, Rosenthal JE. Diurnal pattern of QTc interval: how long is prolonged? Possible relation to circadian triggers of cardiovascular events. *J Am Coll Cardiol* 1996;**27**:76–83.
250. Dilaveris PE, Farbom P, Batchvarov V, Ghuran A, Malik M. Circadian behavior of P-wave duration, P-wave area, and PR interval in healthy subjects. *Ann Noninvasive Electrocardiol* 2001;**6**:92–7.
251. Christ JE, Hoff HE. An analysis of the circadian rhythmicity of atrial and ventricular rates in complete heart block. *J Electrocardiol* 1975;**8**:69–72.
252. Alboni P, Codeca L, Padovan G, Tomaini D, Destro A, Margutti A, et al. Variazioni circadiane della frequenza sinusale in soggetti con nodo del seno normale e patologico. *G Ital Cardiol* 1981;**11**:1211–8.
253. Cinca J, Moya A, Bardaji A, Rius J, Soler-Soler J. Circadian variations of electrical properties of the heart. *Ann N Y Acad Sci* 1990;**601**:222–33.
254. Mitsuoka T, Ueyama C, Matsumoto Y, Hashiba K. Influences of autonomic changes on the sinus node recovery time in patients with sick sinus syndrome. *Jpn Heart J* 1990;**31**:645–60.
255. Vinolas X, Moya A, Cinca J, Bayes de Luna A. Circadian variations in electrophysiological parameters and cardiac repolarization. In: Deedwania PC, editor. *Circadian rhythms of cardiovascular disorders*. Armonk (New York): Futura Publishing; 1997. p. 71–94.
256. Ishida S, Nakagawa M, Fujino T, Yonemochi H, Saikawa T, Ito M. Circadian variation of QT interval dispersion: correlation with heart rate variability. *J Electrocardiol* 1997;**30**:205–10.
257. Yetkin E, Senen K, Ileri M, Atak R, Topaloglu S, Ergun K, et al. Diurnal variation of QT dispersion in patients with and without coronary artery disease. *Angiology* 2001;**52**:311–6.
258. Manolis AG, Katsivas A, Koutsogeorgis D, Theodorakis A, Apostolopoulos G, Nikolaou P, et al. Congestive heart failure and VVI pacing mode: dynamic behavior of the dispersion of ventricular repolarization. *Pacing Clin Electrophysiol* 1996;**19**(1 Pt 2):1890–3.
259. Kong Jr TQ, Goldberger JJ, Parker M, Wang T, Kadish AH. Circadian variation in human ventricular refractoriness. *Circulation* 1995;**92**:1507–16.
260. Willich SN, Maclure M, Mittleman M, Arntz HR, Muller JE. Sudden cardiac death Support for a role of triggering in causation. *Circulation* 1993;**87**:1442–50.
261. Lown B, Calvert AF, Armington R, Ryan M. Monitoring for serious arrhythmias and high risk of sudden death. *Circulation* 1975;**52**:189–98.
262. Cinca J, Moya A, Bardaji A, Figueras J, Rius J. Daily variability of electrically induced reciprocating tachycardia in patients with atrioventricular accessory pathways. *Am Heart J* 1987;**114**:327–33.
263. Shimada R, Nakashima T, Nunoi K, Kohno Y, Takeshita A, Omae T. Arrhythmia during insulin-induced hypoglycemia in a diabetic patient. *Arch Intern Med* 1984;**144**:1068–9.
264. Gillis AM, Guilleminault C, Partinen M, Connolly SJ, Winkle RA. The diurnal variability of ventricular premature depolarizations: influence of heart rate, sleep, and wakefulness. *Sleep* 1989;**12**:391–9.
265. Gillis AM, Peters RW, Mitchell LB, Duff HJ, McDonald M, Wyse DG. Effects of left ventricular dysfunction on the circadian variation of ventricular premature complexes in healed myocardial infarction. *Am J Cardiol* 1992;**69**:1009–14.
266. Otsuka K, Sato T, Saito H, Kaba H, Otsuka K, Seto K, et al. Circadian rhythm of cardiac bradyarrhythmia episodes in rats. *Chronobiologia* 1985;**12**:11–28.
267. Cruz Filho FE, Maia IG, Fagundes ML, Barbosa RC, Alves PA, Sa RM, et al. Electrical behavior of T-wave polarity alternans in patients with congenital long QT syndrome. *J Am Coll Cardiol* 2000;**36**:167–73.
268. Novo S, Barbagallo M, Abrignani MG, Alaimo G, Nardi E, Corrao S, et al. Cardiac arrhythmias as correlated with the circadian rhythm of arterial pressure in hypertensive subjects with and without left ventricular hypertrophy. *Eur J Clin Pharmacol* 1990;**39**(Suppl. 1):S49–51.
269. Sideris DA, Tzoumanidis ST, Anastasiou-Nana M, Zakopoulos N, Kitsiou A, Tsagarakis K, et al. The circadian profile of extrasystolic arrhythmia: its relationship to heart rate and blood pressure. *Int J Cardiol* 1992;**34**:21–31.
270. McClelland J, Halperin B, Cutler J, Kudenchuk P, Kron J, McNulty J. Circadian variation in ventricular electrical instability associated with coronary artery disease. *Am J Cardiol* 1990;**65**:1351–7.
271. Shepard JW. Hypertension, circadian arrhythmias, myocardial infarction and stroke in relation to obstructive sleep apnea. *Clin Chest Med* 1992;**43**:437–58.
272. Guilleminault C, Connolly SJ, Winkle RA. Cardiac arrhythmia and conduction disturbances during sleep in 400 patients with sleep apnea syndrome. *Am J Cardiol* 1983;**52**:490–4.
273. Lanfranchi PA, Somers VK, Braghiroli A, Corra U, Eleuteri E, Giannuzzi P. Central sleep apnea in left ventricular dysfunction: prevalence and implications for arrhythmic risk. *Circulation* 2003;**107**:727–32.
274. Flick MR, Block AJ. Nocturnal vs diurnal cardiac arrhythmias in patients with chronic obstructive pulmonary disease. *Chest* 1979;**75**:8–11.
275. Svorc P, Benacka R, Petrasova D, Bracokova I, Kujanik S. Effect of systemic hypoxia and reoxygenation on electrical stability of the rat myocardium: chronophysiological study. *Physiol Res* 2005;**54**:319–25.
276. Pickering TG, Johnston J, Honour AJ. Comparison of the effects of sleep, exercise and autonomic drugs on ventricular extrasystoles, using ambulatory monitoring of electrocardiogram and electroencephalogram. *Am J Med* 1978;**65**:575–83.
277. Mulcahy D, Wright C, Sparrow J, Cunningham D, Curcher D, Purcell H, et al. Heart rate and blood pressure consequences of an afternoon SIESTA (Snooze-Induced Excitation of sympathetic triggered activity). *Am J Cardiol* 1993;**71**:611–4.
278. Frey B, Heinz G, Binder T, Wutte M, Schneider B, Schmidinger H, et al. Diurnal variation of ventricular response to atrial fibrillation in patients with advanced heart failure. *Am Heart J* 1995;**129**:58–65.
279. Hayano J, Sakata S, Okada A, Mukai S, Fujinami T. Circadian rhythms of atrioventricular conduction properties in chronic atrial fibrillation with and without heart failure. *J Am Coll Cardiol* 1998;**31**:158–66.
280. Simantirakis EN, Chrysostomakis SI, Marketou ME, Kochiadakis GE, Vardakis KE, Mavrikis HE, et al. Atrial and ventricular refractoriness in paced patients; circadian variation and its relationship to autonomic nervous system activity. *Eur Heart J* 2001;**22**:2192–200.
281. Culic V, Eterovic D, Miric D, Giunio L, Lukin A, Fabijanic D. Triggering of ventricular tachycardia by meteorologic and emotional stress: protective effect of beta-blockers and anxiolytics in men and elderly. *Am J Epidemiol* 2004;**160**:1047–58.
282. Behrens S, Ehlers C, Bruggemann T, Ziss W, Dischmann R, Galecka M, et al. Modification of the circadian pattern of ventricular tachyarrhythmias by beta-blocker therapy. *Clin Cardiol* 1997;**20**:253–7.
283. Peters RW, Muller JE, Goldstein S, Byington R, Friedman LM. Propranolol and the morning increase in the frequency of sudden cardiac death (BHAT study). *Am J Cardiol* 1989;**63**:1518–20.
284. Schuchert A, Behrens G, Meinertz T. Diurnal variations of the ventricular pacing threshold in patients with cardiac pacemakers are not related to changes in autonomic tone. *Am J Cardiol* 2000;**86**:226–9.
285. Svorc P, Wilk P, Murar J, Podlubny I, Kujanik S, Bracokova I, et al. Circadian rhythm of the ventricular fibrillation threshold in female Wistar rats. *Physiol Res* 1994;**43**:355–8.
286. Venditti Jr FJ, John RM, Hull M, Toffler GH, Shahian DM, Martin DT. Circadian variation in defibrillation energy requirements. *Circulation* 1996;**94**:1607–12.
287. Mitchell AR, Warman E, Schaaf K, Sulke N. The circadian variation of atrial defibrillation thresholds. *J Interv Card Electrophysiol* 2004;**10**:79–83.
288. Chakko S, Myerburg RJ. Cardiac arrhythmias and circadian changes. In: Deedwania PC, editor. *Circadian rhythms of cardiovascular disorders*. Armonk (New York): Futura Publishing; 1997. p. 129–45.
289. Linsell CR, Lightman SL, Mullen PE, Brown MJ, Causon RC. Circadian rhythms of epinephrine and norepinephrine in man. *J Clin Endocrinol Metab* 1985;**60**:1210–5.
290. Kawano Y, Tochikubo O, Minamisawa K, Miyajima E, Ishii M. Circadian variation of hemodynamics in patients with essential hypertension: comparison between early morning and evening. *J Hypertens* 1994;**12**:1405–12.
291. Sowers JR, Vlachakis N. Circadian variation in plasma dopamine levels in man. *J Endocrinol Invest* 1984;**7**:341–5.
292. Lown B, Verrier RL. Neural activity and ventricular fibrillation. *N Engl J Med* 1976;**294**:1165–70.
293. Huikuri HV, Valkama JO, Airaksinen KE, Seppanen T, Kessler KM, Takkunen JT, et al. Frequency domain measures of heart rate variability before the onset of nonsustained and sustained ventricular tachycardia in patients with coronary artery disease. *Circulation* 1993;**87**:1220–8.
294. Hayano J, Sakakibara Y, Yamada A, Yamada M, Mukai S, Fujinami T, et al. Accuracy of assessment of cardiac vagal tone by heart rate variability in normal subjects. *Am J Cardiol* 1991;**67**:199–204.
295. Malliani A, Pagani M, Lombardi F, Cerutti S. Cardiovascular neural regulation explored in the frequency domain. *Circulation* 1991;**84**:482–92.
296. Talan MI, Engel BT, Chew PH. Systematic nocturnal atrial demand pacing results in high-output heart failure. *J Appl Physiol* 1992;**72**:1803–9.
297. Andresen D, Bruggemann T. Heart rate variability preceding onset of atrial fibrillation. *J Cardiovasc Electrophysiol* 1998;**9**:S26–9.
298. Tomita T, Takei M, Saikawa Y, Hanaoka T, Uchikawa S, Tsutsui H, et al. Role of autonomic tone in the initiation and termination of paroxysmal atrial fibrillation in patients without structural heart disease. *J Cardiovasc Electrophysiol* 2003;**14**:559–64.
- \*299. Portaluppi F, Hermida RC. Circadian rhythms in cardiac arrhythmias and opportunities for their chronotherapy. *Adv Drug Deliv Rev* 2007;**59**:940–51.
300. Hermida RC, Ayala DE, Portaluppi F. Circadian variation of blood pressure: the basis for the chronotherapy of hypertension. *Adv Drug Deliv Rev* 2007;**59**:904–22.