CHEST

Original Research

PULMONARY VASCULAR DISEASE

CT Scan-Measured Pulmonary Artery to Aorta Ratio and Echocardiography for Detecting Pulmonary Hypertension in Severe COPD

Anand S. Iyer, MD; J. Michael Wells, MD; Sonia Vishin, MD, FCCP; Surya P. Bhatt, MD, FCCP; Keith M. Wille, MD, FCCP; and Mark T. Dransfield, MD

Background: COPD is associated with significant morbidity primarily driven by acute exacerbations. Relative pulmonary artery (PA) enlargement, defined as a PA to ascending aorta (A) diameter ratio greater than one (PA:A>1) identifies patients at increased risk for exacerbations. However, little is known about the correlation between PA:A, echocardiography, and invasive hemodynamics in COPD.

Methods: A retrospective observational study of patients with severe COPD being evaluated for lung transplantation at a single center between 2007 and 2011 was conducted. Clinical characteristics, CT scans, echocardiograms, and right-sided heart catheterizations were reviewed. The PA diameter at the bifurcation and A diameter from the same CT image were measured. Linear and logistic regression were used to examine the relationships between PA:A ratio by CT scan and PA systolic pressure (PASP) by echocardiogram with invasive hemodynamics. Receiver operating characteristic analysis assessed the usefulness of the PA:A ratio and PASP in predicting resting pulmonary hypertension (PH) (mean pulmonary artery pressure [mPAP] > 25 mm Hg). Results: Sixty patients with a mean predicted FEV₁ of 27% \pm 12% were evaluated. CT scan-measured PA:A correlated linearly with mPAP after adjustment for multiple covariates (r = 0.30, P = .03), a finding not observed with PASP. In a multivariate logistic model, mPAP was independently associated with PA:A > 1 (OR, 1.44; 95% CI, 1.02-2.04; P = .04). PA:A > 1 was 73% sensitive and 84% specific for identifying patients with resting PH (area under the curve, 0.83; 95% CI, 0.72-0.93; P < .001), whereas PASP was not useful.

Conclusions: A PA:A ratio > 1 on CT scan outperforms echocardiography for diagnosing resting PH in patients with severe COPD.

CHEST 2014; 145(4):824–832

Abbreviations: A = aorta; dPAP = diastolic pulmonary artery pressure; mPAP = mean pulmonary artery pressure; PA = pulmonary artery; PASP = pulmonary artery systolic pressure; PVR = pulmonary vascular resistance; RHC = right-sided heart catheterization; ROC = receiver operating characteristic; sPAP = systolic pulmonary artery pressure; UAB = University of Alabama at Birmingham

COPD is the third leading cause of death in the United States and is associated with significant morbidity and health-care-related expenditures mostly due to acute exacerbations. ^{1,2} Patients with COPD may develop subclinical or overt pulmonary vascular disease, and the presence of pulmonary hypertension (PH) is an important predictor of both COPD and exacerbation-related morbidity and mortality, particularly in patients with severe disease. ³⁻⁸ Although PH can occur in patients with mild to moderate COPD because of pulmonary inflammation and endothelial or cardiac dysfunction, it more commonly occurs and

becomes clinically apparent when airflow limitation is severe and is believed to be primarily due to hypoxic vasoconstriction of small pulmonary arteries that result in intimal hyperplasia as well as smooth muscle hypertrophy and hyperplasia.⁹⁻¹¹

Right-sided heart catheterization (RHC) remains the gold standard for diagnosis of pulmonary vascular disease. ¹² Although noninvasive strategies, such as Doppler echocardiography, allow the measurement of pulmonary artery systolic pressure (PASP), this modality is often problematic in patients with COPD because of lung hyperinflation and difficulty in obtaining clear

acoustic windows. 13 CT scanning has been used to assess the mean pulmonary artery (PA) diameter, the cross-sectional area of small pulmonary vasculature, and the PA to ascending aorta (A) ratio, and each has demonstrated varying associations with RHC data. However, most of these studies were conducted in heterogeneous populations with various underlying lung diseases, thus, limiting their application to patients with COPD.14-20

We showed that CT image-measured relative PA enlargement, defined as a PA:A ratio > 1, identifies patients at increased risk of COPD exacerbation and hospitalization²¹; however, the mechanism underlying relative PA enlargement has not been clearly elucidated.^{16,22} Although increases in the PA:A ratio could represent resting PH, other factors may contribute, including left ventricular dysfunction and hyperinflation due to emphysema. We hypothesized that the PA:A ratio would correlate with mean PA pressure (mPAP) by RHC and that the presence of PA:A > 1 could serve as a more useful screening test than echocardiography for identifying PH in patients with severe COPD.

MATERIALS AND METHODS

Study Design and Patient Selection

Records for patients who underwent lung transplant evaluation between 2007 and 2011 at the University of Alabama at Birmingham (UAB) were analyzed. Men and women were eligible if they were >40 years of age; had a history of COPD at the time of evaluation; lacked coexisting lung diseases such as cystic fibrosis, bronchiectasis, diffuse interstitial lung disease, or lung cancer; and had RHC, echocardiogram, and CT scan performed within 4 months of each other. The study was approved by the Institutional Review Board at the UAB (X120803011).

Clinical Characterization

Sociodemographic, clinical, and laboratory data were abstracted from medical records and the UAB Transplant Database at the time of transplant evaluation. RHC data collected included systolic PA pressure (sPAP), diastolic PA pressure (dPAP), mPAP, and mean

Manuscript received June 19, 2013; revision accepted September

Affiliations: From the Department of Internal Medicine (Drs Iyer, Wells, Vishin, Wille, and Dransfield), and the Division of Pulmonary, Allergy, and Critical Care (Drs Wells, Vishin, Bhatt, Wille, and Dransfield), Department of Medicine and University of Alabama at Birmingham Lung Health Center, University of Alabama at Birmingham; and the Birmingham Veterans Affairs Medical Center (Drs Wells and Dransfield), Birmingham, AL.

Drs Iyer and Wells contributed equally to the manuscript.

Funding/Support: Dr Wells is supported by the Walter B. Frommeyer Jr Fellowship in Investigational Medicine, University of Alabama at Birmingham.

Correspondence to: J. Michael Wells, MD, 1900 University Blvd, THT 422, Birmingham, AL 35294; e-mail: jmwells@uab.edu © **2014 American College of Chest Physicians.** Reproduction

of this article is prohibited without written permission from the American College of Chest Physicians. See online for more details.

DOI: 10.1378/chest.13-1422


pulmonary capillary wedge pressure, as well as mean right and left ventricular end-diastolic pressures, mean right atrial pressure, cardiac output by thermodilution, cardiac index, peripheral vascular resistance, and systemic vascular resistance. Echocardiography data collected included left ventricular ejection fraction, right ventricular ejection fraction, and PASP. Pulmonary hypertension was defined as a resting mPAP > 25 mm Hg by RHC. 12 Severe COPD was defined as airflow obstruction of < 50% defined by postbronchodilator FEV₁ in the setting of an FEV₁:FVC ratio of < 0.70.

Measuring the PA:A Ratio

One reviewer, blinded to hemodynamic information, analyzed CT scans using Philips iSite Enterprise Software (Koninklijke Philips N.V.). The PA diameter was measured at the level of the PA bifurcation, and the A diameter was averaged from two perpendicular measurements taken from the same CT image as previously reported (Fig 1).²¹ A second blinded reviewer measured the PA and A diameters to assess interobserver agreement. Cohen κ was 0.82 (95% CI, 0.68-0.97) for interobserver agreement in CT scan measurement of the PA:A ratio.

Statistical Analysis

Baseline data were expressed as means with SD for normally distributed values. Continuous variables were compared using twosided Student t tests, and categorical variables were examined by χ^2 testing as well as Fisher exact test as appropriate. Pearson correlation coefficients were determined for the relationships between the PA diameter, A diameter, and PA:A ratio, as well as echocardiography-measured PASP and hemodynamic parameters. Multiple linear regression, including demographic and clinical covariates, was used to examine the relationship between CT image metrics (PA, A, and PA:A ratio) as well as PASP and hemodynamic parameters. Univariate and multivariate logistic






FIGURE 1. Measurement of the PA and A diameters at the PA bifurcation. A = aorta; PA = pulmonary artery.

regression were used to determine the relationship between hemodynamic measurements and the presence of a PA:A > 1. Steiger Z-test was calculated to compare correlations between echocardiography-measured PASP and CT scan-measured PA:A with mPAP. Receiver operating characteristic (ROC) analysis assessed the usefulness of the PA:A ratio and PASP in predicting the presence of mPAP > 25 mm Hg. Cohen κ measured interobserver variability in CT scan diameter measurements for the cohort. SPSS for Windows version 20 (IBM) was used for all analyses. All statistical tests were two-sided, with significance assigned to tests with P values \leq .05.

RESULTS

Between 2007 and 2011, 134 of 690 patients referred for lung transplant evaluation had COPD, and 74 subjects were excluded on the basis of not having matched CT scan, echocardiography, or RHC data. The current analysis included 60 patients who were separated into groups based on a PA:A ratio > 1 (n = 22) and a PA:A ratio \leq 1 (n = 38). Participants had a mean (\pm SD) age of 55 (\pm 7) years; 50 were white (83%), and 34 were women (57%). Mean predicted FEV₁ was 27% (\pm 12%). The mean (\pm SD) number of days between RHC and CT scan was 18 (\pm 50) days and 1 (\pm 23) day between RHC and echocardiography. Of these patients, 22 (38%) had mPAP > 25 by RHC.

Differences in the PA:A ratio between both groups were driven by the PA diameter $(27.3 \pm 3.2 \text{ mm})$ in PA:A < 1 vs $34.3 \pm 4.1 \text{ mm}$ in PA:A > 1, P < .001), as no differences were detected between aortic diameters $(31.8 \pm 2.7 \text{ mm})$ vs $31.1 \pm 3.2 \text{ mm}$, respectively; P = .42; Table 1). There were no significant differences between the PA:A < 1 and PA:A > 1 groups regarding

Table 1—Baseline Characteristics of Participants

Characteristic	PA:A < 1 (n = 38)	PA:A > 1 (n = 22)	P Value	
Age, y	$55 (\pm 7.9)$	$57.5 (\pm 5.9)$.21	
Female sex	22 (58)	12 (55)	.80	
White	34 (90)	16 (73)	.09	
FEV ₁ % predicted	$26.3 (\pm 11.3)$	$29.1 \ (\pm 14.8)$.40	
FEV ₁ :FVC ratio	$37.3 (\pm 12.1)$	$38.4 \ (\pm 14.3)$.76	
6-min walk distance, ft	$915 \ (\pm 214)$	$776 (\pm 191)$.07	
Coronary artery disease	4 (10)	4 (18)	.41	
Hyperlipidemia	9 (24)	4 (18)	.63	
Congestive heart failure	2 (5)	5 (23)	.04	
Hypertension	18 (47)	11 (50)	.85	
Diabetes mellitus	1(3)	4 (18)	.04	
OSA	4 (10)	6 (27)	.10	
Currently smoking	1(3)	0 (0)	.45	
Smoking, pack-y	$45 \ (\pm 21)$	$54 \ (\pm 31)$.23	
Heart rate, beats/min	$91 \ (\pm 12)$	$94\ (\pm 19)$.46	
Percent oxygen saturation by pulse oximetry	$95 \ (\pm 3)$	$91 \ (\pm 6)$.004	
BMI, kg/m ²	$23.1 (\pm 3.9)$	$25.8 \ (\pm 5.5)$.04	
pH	$7.41 \ (\pm 0.03)$	$7.41 \ (\pm 0.03)$.94	
Pco_2	$45 \ (\pm 8)$	$50 \ (\pm 14)$.14	
Po_2	$65 (\pm 11)$	$54 \ (\pm 13)$.003	
$PHCO_3$	$27.9 (\pm 3.9)$	$29.9 \ (\pm 5.4)$.16	
Systolic PAP by RHC, mm Hg	$31 \ (\pm 6)$	$46 \ (\pm 14)$	< .001	
Diastolic PAP by RHC, mm Hg	$14\ (\pm5)$	$21 \ (\pm 8)$	< .001	
Mean PAP by RHC, mm Hg	$21 \ (\pm 4)$	$31 \ (\pm 10)$	< .001	
Mean PCWP by RHC, mm Hg	$11 \ (\pm 4)$	$13 (\pm 6)$.11	
Mean LVEDP by RHC, mm Hg	$11 \ (\pm 6)$	$15 \ (\pm 6)$.08	
Mean RVEDP by RHC, mm Hg	$8 (\pm 4)$	$11 \ (\pm 4)$	< .001	
Mean RAP by RHC, mm Hg	$6 \ (\pm 3)$	$9 \ (\pm 4)$.009	
Cardiae output, L/min	$5.0 \ (\pm 0.9)$	$5.3 (\pm 1.1)$.26	
Cardiac index, L/min/m ²	$2.9 \ (\pm 0.4)$	$2.9 \ (\pm 0.5)$.72	
Pulmonary vascular resistance, mm Hg×min/L	$2.1 (\pm 0.8)$	$4.5~(\pm 3.8)$	< .001	
Systemic vascular resistance, mm Hg×min/L	$18.4~(\pm 4.4)$	$17.1~(\pm 5.4)$.37	
PA diameter by CT scan, mm	$27.3 (\pm 3.2)$	$34.3 (\pm 4.1)$	< .001	
Ascending aortic diameter by CT scan, mm	$31.8 (\pm 2.7)$	$31.1 (\pm 3.2)$.42	
LVEF by echocardiography, %	$54 \ (\pm 5)$	$55 \ (\pm 1)$.51	
RVEF by echocardiography, %	$54 \ (\pm 2)$	$49 \ (\pm 12)$.009	
PASP by echocardiography, mm Hg	$34 (\pm 11, n = 25)$	$49 (\pm 21, n = 13)$	<.001	

Data are presented as No. (%) or mean (\pm SD). A = aorta; LVEDP = left ventricular end-diastolic pressure; LVEF = left ventricular ejection fraction; PA = pulmonary artery; PAP = pulmonary artery pressure; PASP = pulmonary artery systolic pressure; PCWP = pulmonary capillary wedge pressure; PHCO₃ = partial pressure of bicarbonate; RAP = right atrial pressure; RHC = right-sided heart catheterization; RVEDP = right ventricular end-diastolic pressure; RVEF = right ventricular ejection fraction.

age, sex, race, smoking status, lung function, or prevalence of coronary artery disease, hyperlipidemia, hypertension, and OSA. Those with PA:A>1 had higher BMIs and comorbid diabetes (18% vs 3%, P = .04), as well as lower percutaneous oxygen saturation (91% vs 95%, P = .004) and lower arterial oxygen tension $(PaO_{2}, 54 \text{ mm Hg vs } 65 \text{ mm Hg}; P = .003) \text{ (Table 1)}.$ The sPAP, dPAP, and mPAP were all higher in those with PA:A > 1 as compared with those with PA:A < 1. No difference in pulmonary capillary wedge pressure was observed despite the PA:A > 1 group having more patients with comorbid congestive heart failure (23% vs 5%, P = .04) (Table 1). Echocardiographymeasured PASP was higher in the group with PA:A>1 than in those with PA:A \leq 1 (49 \pm 21 mm Hg vs 34 \pm 11 mm Hg, P < .001), and the right ventricular ejection fraction was reduced $(49\% \pm 12\% \text{ vs } 54\% \pm 2\%)$, P = .009) as seen in Table 1.

CT scan-measured PA diameter correlated linearly with sPAP (r = 0.62, P < .001), dPAP (r = 0.53, P < .001),

and mPAP (r=0.60, P<.001) (Fig 2, Table 2). In contrast, the aortic diameter was not associated with any measure of PA pressure (Table 2). The PA:A ratio also demonstrated a significant linear correlation with sPAP (r=0.60, P<.001), mPAP (r=0.56, P<.001), and dPAP (r=0.47, P<.001) by RHC and remained significant after multiple linear regression analysis for sPAP (r=0.40, P=.005) and mPAP (r=0.30, P=.03) adjusting for age, race, sex, BMI, resting oxygen saturation, sleep apnea, congestive heart failure, and diabetes mellitus (Fig 2, Table 2). Measuring the PA diameter at a more proximal location yielded a similar correlation between the PA:A ratio and mPAP (e-Table 1).

Echocardiography-measured PASPs were available on 38 patients (63%) and in that subgroup correlated linearly with sPAP (r=0.42, P=.009), mPAP (r=0.33, P=.043), dPAP (r=0.35, P=.03), and pulmonary vascular resistance (PVR) (r=0.49, P=.002) (Fig 2, Table 3). In the 22 other patients, PASP was not measured because of the absence of a tricuspid regurgitant




FIGURE 2. Linear relationships between invasive hemodynamics, PA diameter and PA:A ratio from CT scan as well as PA systolic pressure by echocardiography. mPAP = mean PA pressure; mPCWP = mean pulmonary capillary wedge pressure; PASP = echocardiographic PA systolic pressure; sPAP = systolic PA pressure. See Figure 1 legend for expansion of other abbreviations.

Table 2—Univariate and Multiple Linear Regression Comparisons Between Invasive Hemodynamics and PA Diameter,
A Diameter, and PA:A

		Univariate Linear Regression							
Measure	PA	P Value	A	P Value	PA:A	P Value	PA:A	P Value	
sPAP	0.62	<.001	0.07	.59	0.60	<.001	0.40	.005	
mPAP	0.60	<.001	0.09	.50	0.56	<.001	0.30	.03	
dPAP	0.53	<.001	0.10	.46	0.47	<.001	0.33	.06	
LVEDP	0.23	.10	0.05	.97	0.27	.06	0.20	.35	
RVEDP	0.37	.004	-0.09	.48	0.42	.001	0.23	.22	
RAP	0.27	.03	-0.13	.32	0.36	.005	0.12	.56	
PVR	0.45	<.001	0.04	.79	0.45	<.001	0.49	.003	
PCWP	0.10	.46	-0.19	.15	0.20	.12	0.20	.37	

dPAP = diastolic PA pressure; mPAP = mean PA pressure; PVR = pulmonary vascular resistance; sPAP = systolic PA pressure. See Table 1 legend for expansion of other abbreviations.

^aVariables in the multiple linear regression model: PA:A value, age, race, sex, BMI, resting oxygen saturation, sleep apnea, congestive heart failure, and diabetes mellitus.

jet or an inadequate window. In a multiple linear regression model adjusting for age, race, sex, BMI, resting oxygen saturation, sleep apnea, congestive heart failure, and diabetes mellitus, the correlation between PASP and PVR remained significant (r = 0.34, P = .02) (Table 3), but the correlations between PASP and sPAP, dPAP, or mPAP were lost (Table 3). The correlation coefficients between PA:A ratio and echocardiographymeasured PASP with sPAP were not statistically different, but the correlation of the PA:A ratio with mPAP was significantly stronger than the correlation between echocardiography-derived PASP and mPAP (z = 2.01, P < .05). In a multivariate logistic model, mPAP was independently associated with the presence of PA:A > 1 (OR, 1.44; 95% CI, 1.02-2.04; P = .04) adjusting for age, sex, race, BMI, resting oxygen saturation, sleep apnea, congestive heart failure, and diabetes (Table 4).

ROC curves were performed to evaluate the sensitivity and specificity of both the PA:A ratio and echocardiography-measured PASP for diagnosing PH defined by an mPAP>25 mm Hg. The area under the curve for the PA:A ratio was 0.83 (95% CI, 0.72-0.93;

Table 3—Univariate and Multiple Linear Regression Comparisons Between Invasive Hemodynamics and Echocardiographic PA Systolic Pressure

Measure		ate Linear ression	Multiple Lines Regression ^a		
	PASP	P Value	PASP	P Value	
sPAP	0.42	.009	0.16	.23	
mPAP	0.33	.04	0.01	.91	
dPAP	0.35	.03	0.16	.34	
PVR	0.49	.002	0.34	.02	

See Table 1 and 2 legends for expansion of abbreviations.

^aVariables in the multiple linear regression model: PASP, age, race, sex, BMI, resting oxygen saturation, sleep apnea, congestive heart failure, and diabetes mellitus.

P < .001) with an mPAP > 25 mm Hg (Fig 3A). The sum of sensitivity and specificity between the PA:A ratio and an mPAP > 25 mm Hg was greatest at a PA:A ratio > 1 (73% sensitivity and 84% specificity) when compared in a range of PA:A ratio from > 0.70 to >1.20 (Table 5). The area under the curve between PASP and an mPAP > 25 mm Hg did not achieve statistical significance (0.60; 95% CI, 0.39-0.80; P = .33) (Fig 3B), corresponding to a significant difference from the area under the ROC curve for the PA:A ratio and an mPAP \geq 25 mm Hg (P = .045). The sum of sensitivity and specificity between the PASP and mPAP>25 mm Hg was greatest at a PASP>55 mm Hg (36% sensitivity and 96% specificity) (e-Table 2). We also evaluated the usefulness of the PA:A ratio and echocardiography-measured PASP for diagnosing severe PH, defined by an mPAP > 35 mm Hg.²³ The area under the ROC curve for the PA:A ratio was 0.85 (95% CI, 0.75 - 0.95; P = .003), whereas the area

Table 4—Relationship Between Invasive Hemodynamics and PA:A > 1 by Multivariate Logistic Regression

Measure	OR	95% CI	P Value
sPAP	1.50	1.02-2.20	.04
mPAP	1.44	1.02-2.04	.04
dPAP	1.38	0.94-2.05	.10
RAP	1.12	0.80-1.57	.52
LVEDP	1.09	0.92-1.29	.33
RVEDP	1.21	0.88-1.66	.25
PVR	1.04	1.004-1.09	.03
PCWP	1.08	0.86-1.37	.50
Cardiae output	0.33	0.05-2,28	.26
Cardiac index	0.58	0.02-16.8	.75

Variables in the multivariate logistic model: age, race, sex, BMI, resting oxygen saturation, sleep apnea, congestive heart failure, and diabetes mellitus, and the hemodynamic measurement of interest. A separate multivariate logistic regression model was used for each hemodynamic metric. See Table 1 and 2 legends for expansion of abbreviations.




FIGURE 3. A, Receiver operator characteristic curve with the PA:A ratio at identifying mPAP > 25 mm Hg. B, Receiver operating characteristic curve with the echocardiographic PA systolic pressure at identifying mPAP > 25 mm Hg. AUC = area under the curve. See Figure 1 and 2 legends for expansion of other abbreviations.

under the ROC curve for the PASP was again nonsignificant (0.70; 95% CI, 0.39-0.99; P = .13).

DISCUSSION

We have demonstrated that the CT scan-measured PA:A ratio correlates with pulmonary arterial pressures as determined by RHC in patients with severe COPD. This is the first study, to our knowledge, to specifically examine this relationship in a cohort consisting exclusively of well-characterized patients with severe COPD and provides a plausible mechanism for the development of relative pulmonary artery enlargement on CT scan, which has in turn been linked to the risk of exacerbation.²¹ Defining the relationship between the PA:A ratio and PH is important given the implications of comorbid COPD and PH on exacerbation risk, exacerbation-related mortality, and all-cause mortality.^{24,25} These findings also strengthen the pathophysiologic link between an elevated PA:A ratio and exacerbations.21 Measurement of the PA:A ratio is reproducible, requires minimal training, and performs better than echocardiography for the identification of PH in patients with severe COPD. 11,16,26,27

Estimates of the prevalence of PH in COPD vary widely from as low as 10% to as high as 85% depending on the severity of the underlying lung disease and the diagnostic criteria as well as approach used. 11,28-30 Accurate estimates have also been problematic, because RHC is not routinely used in patients with COPD, particularly in those with moderate disease.¹¹ In our population of patients with severe COPD, we found a 37% prevalence of PH using an mPAP≥25 mm Hg and a 37% prevalence using a PA:A>1. Overall, the presence of PA:A>1 was reproducible and 83% accurate in predicting resting PH as defined by RHC. We also explored the relationship between measurement of the PA diameter and invasive hemodynamics at a location more proximal to the bifurcation and found that the correlations with RHC data were similar. The interobserver agreement and anatomic consistency of the PA:A ratio support its clinical usefulness.

Table 5—Sensitivity and Specificity for PA:A Ratio in Identifying mPAP > 25

Measure	>0.70	>0.75	>0.80	> 0.85	>0.90	> 0.95	>1.00	>1.05	>1.10	>1.15	>1.20
Sensitivity	1.00	1.00	1.00	1.00	0.86	0.77	0.73	0.50	0.27	0.14	0.09
Specificity	0.05	0.13	0.21	0.39	0.53	0.66	0.84	0.92	0.92	0.95	1.00
PPV	0.38	0.40	0.42	0.49	0.51	0.57	0.73	0.79	0.67	0.60	1.00
NPV	1.00	1.00	1.00	1.00	0.87	0.83	0.84	0.76	0.69	0.65	0.66

NPV = negative predictive value; PPV = positive predictive value. See Table 1 and 2 legends for expansion of other abbreviations.

journal, publications, chestnet.org CHEST / 145 / 4 / APRIL 2014 **829**

Doppler echocardiography can provide an estimate of right ventricular systolic pressure (believed to reflect PASP¹³); however, this technique can be problematic in patients with COPD due in part to difficulties in obtaining adequate acoustic windows, a factor that does not impact the use of CT imaging. 13,31-33 PA pressures by echocardiography were only measured in 60% of the patients, a value not dissimilar from the 44% Arcasoy et al¹³ reported in a heterogeneous population with advanced lung disease undergoing lung transplant evaluation. Furthermore, our analysis showed that the correlation between CT scan-measured relative PA enlargement and invasive hemodynamics was more robust than echocardiography-measured PASP, thus, supporting CT scan as the preferred modality for the identification of PH in patients with severe COPD.

Although the correlation between relative PA enlargement and PA pressures is driven most by changes in PA diameter rather than changes in aortic diameter, relative PA enlargement is only partially explained by intrinsic PH itself, as indicated by the modest r value we report. Other mechanisms could contribute, including changes to pulmonary arterial distensibility,³⁴ redistribution of blood flow from peripheral capillary loss,³⁵ or left ventricular systolic or diastolic dysfunction,³⁶⁻³⁸ although we observed no clear correlation with capillary wedge pressures, cardiac output, or left ventricular ejection fraction in those with PA:A>1 and PA:A<1.

Prior studies have characterized the relationship between CT scan metrics and hemodynamics; however, these have been conducted in heterogeneous populations with various lung diseases and included a small number of patients with COPD. 17,39-41 In contrast, our study focuses on well-characterized patients with severe COPD and, thus, allows a better understanding of the relationship between the PA:A ratio, hemodynamics, and echocardiography in this population. This is of significant importance, as the correlation between PA diameter and the PA:A ratio varies based on the included patients' lung disease and the mechanisms leading to PA enlargement. 42

This study confirms our previous findings that an elevated PA:A ratio is associated with higher BMI, a greater requirement for oxygen, and lower arterial saturation, as well as a higher prevalence of congestive heart failure.²¹ These factors each contribute to or are a consequence of PH, thus, supporting the biologic plausibility that the PA:A ratio could serve as an integrated measure of intrinsic and secondary pulmonary vascular disease in COPD.

Our study is limited by its small size, single-center experience, and retrospective design, although we did observe robust and clinically relevant associations between the CT image-measured PA:A ratio and hemodynamics. The modest sample size may account

for the lack of association between echocardiographyderived PASP and hemodynamic metrics in multiple regression analyses. However, the relationships between the PA:A ratio and invasive metrics remained significant when controlled for other variables, highlighting the robust relationship between the PA:A ratio and hemodynamic measurements. Also, the association between mPAP and PA:A > 1 in patients with severe COPD may not be generalizable to those with more moderate disease in whom PH would be less prevalent. However, in a population of moderate to severe COPD (GOLD [Global Initiative for Chronic Obstructive Lung Disease] stage II-IV),²¹ the prevalence of PA:A > 1 was 24% compared with 37% in the current study, suggesting that this process also occurs in milder disease. Last, we cannot account for exacerbations or other events that may have occurred between CT scan and RHC; however, the potential for confounding was minimized by the short time between the two measurements.

In conclusion, the PA:A ratio and the presence of a PA:A > 1 correlate with invasive measurements of pulmonary arterial pressure and the presence of PH, respectively. These measurements are readily available, require minimal training, and show more robust linear correlation than echocardiography with invasive hemodynamics in patients with advanced COPD. CT scan-measured relative PA enlargement should be the preferred screening tool for PH in this patient population.

ACKNOWLEDGMENTS

Author contributions: Dr Dransfield is guarantor of the manuscript and is responsible for the work as a whole.

Dr *Îyer*: contributed as co-first author and to study design, vessel measurement, manuscript writing, and manuscript editing.

Dr Wells: contributed as co-first author and to study design, vessel measurement, data analysis, manuscript writing, and manuscript editing.

Dr Vishin: contributed to data collection, manuscript writing, and manuscript editing.

Dr Bhatt: contributed to manuscript writing and manuscript editing.

Dr Wille: contributed to study design, data collection, and manuscript editing.

Dr Dransfield: contributed as senior author and to study design, data analysis, manuscript writing, and manuscript editing.

Financial/nonfinancial disclosures: The authors have reported to CHEST the following conflicts of interest: Dr Dransfield has served as a consultant to Boehringer Ingelheim GmbH, GlaxoSmithKline, and Ikaria, Inc, and his institution has received research support from the National Institutes of Health; Aeris Therapeutics; AstraZeneca; Boehringer Ingelheim GmbH; Boston Scientific; Janssen Biotech, Inc; Forest Laboratories, Inc; GlaxoSmithKline; Ikaria, Inc; Medimmune, LLC; Olmsted Medical Center; Otsuka America Pharmaceutical, Inc; Pfizer, Inc; and Pulmonx. Drs Iyer, Wells, Vishin, Bhatt, and Wille have reported that no potential conflicts of interest exist with any companies/organizations whose products or services may be discussed in this article.

Role of sponsors: The sponsor had no role in the design of the study, the collection and analysis of the data, or the preparation of the manuscript.

Additional information: The e-Tables can be found in the "Supplemental Materials" area of the online article.

REFERENCES

- Petty TL. Scope of the COPD problem in North America: early studies of prevalence and NHANES III data: basis for early identification and intervention. *Chest.* 2000; 117(5_suppl_2):326S-331S.
- Centers for Disease Control and Prevention (CDC). Chronic obstructive pulmonary disease among adults—United States, 2011. MMWR Morb Mortal Wkly Rep. 2012;61(46):938-943.
- McGhan R, Radcliff T, Fish R, Sutherland ER, Welsh C, Make B. Predictors of rehospitalization and death after a severe exacerbation of COPD. Chest. 2007;132(6):1748-1755.
- Terzano C, Conti V, Di Stefano F, et al. Comorbidity, hospitalization, and mortality in COPD: results from a longitudinal study. *Lung*. 2010;188(4):321-329.
- Kessler R, Faller M, Fourgaut G, Mennecier B, Weitzenblum E. Predictive factors of hospitalization for acute exacerbation in a series of 64 patients with chronic obstructive pulmonary disease. Am J Respir Crit Care Med. 1999;159(1): 158-164.
- Barberà JA, Blanco I. Pulmonary hypertension in patients with chronic obstructive pulmonary disease: advances in pathophysiology and management. *Drugs*. 2009;69(9):1153-1171.
- Fayngersh V, Drakopanagiotakis F, Dennis McCool F, Klinger JR. Pulmonary hypertension in a stable community-based COPD population. *Lung*. 2011;189(5):377-382.
- Cuttica MJ, Kalhan R, Shlobin OA, et al. Categorization and impact of pulmonary hypertension in patients with advanced COPD. Respir Med. 2010;104(12):1877-1882.
- Ward JP, McMurtry IF. Mechanisms of hypoxic pulmonary vasoconstriction and their roles in pulmonary hypertension: new findings for an old problem. *Curr Opin Pharmacol*. 2009; 9(3):287-296.
- Peinado VI, Barbera JA, Ramirez J, et al. Endothelial dysfunction in pulmonary arteries of patients with mild COPD. Am J Physiol. 1998;274(6 pt 1):L908-L913.
- Minai OA, Chaouat A, Adnot S. Pulmonary hypertension in COPD: epidemiology, significance, and management: pulmonary vascular disease: the global perspective. *Chest*. 2010; 137(6_suppl):39S-51S.
- 12. McLaughlin VV, Archer SL, Badesch DB, et al; American College of Cardiology Foundation Task Force on Expert Consensus Documents; American Heart Association; American College of Chest Physicians; American Thoracic Society, Inc; Pulmonary Hypertension Association. ACCF/AHA 2009 expert consensus document on pulmonary hypertension a report of the American College of Cardiology Foundation Task Force on Expert Consensus Documents and the American Heart Association developed in collaboration with the American College of Chest Physicians; American Thoracic Society, Inc.; and the Pulmonary Hypertension Association. J Am Coll Cardiol. 2009;53(17):1573-1619.
- Arcasoy SM, Christie JD, Ferrari VA, et al. Echocardiographic assessment of pulmonary hypertension in patients with advanced lung disease. Am J Respir Crit Care Med. 2003; 167(5):735-740.
- Karazincir S, Balci A, Seyfeli E, et al. CT assessment of main pulmonary artery diameter. *Diagn Interv Radiol*. 2008;14(2): 72-74.
- Mahammedi A, Oshmyansky A, Hassoun PM, Thiemann DR, Siegelman SS. Pulmonary artery measurements in pulmonary hypertension: the role of computed tomography. *J Thorac Imag*ing. 2013;28(2):96-103.

- Matsuoka S, Washko GR, Yamashiro T, et al; National Emphysema Treatment Trial Research Group. Pulmonary hypertension and computed tomography measurement of small pulmonary vessels in severe emphysema. Am J Respir Crit Care Med. 2010;181(3):218-225.
- Ng CS, Wells AU, Padley SP. A CT sign of chronic pulmonary arterial hypertension: the ratio of main pulmonary artery to aortic diameter. *J Thorac Imaging*. 1999;14(4):270-278.
- Alhamad EH, Al-Boukai AA, Al-Kassimi FA, et al. Prediction of pulmonary hypertension in patients with or without interstitial lung disease: reliability of CT findings. *Radiology*. 2011; 260(3):875-883.
- Washko GR, Parraga G, Coxson HO. Quantitative pulmonary imaging using computed tomography and magnetic resonance imaging. *Respirology*. 2012;17(3):432-444.
- Devaraj A, Wells AU, Meister MG, Corte TJ, Wort SJ, Hansell DM. Detection of pulmonary hypertension with multidetector CT and echocardiography alone and in combination. *Radiology*. 2010;254(2):609-616.
- Wells JM, Washko GR, Han MK, et al; COPDGene Investigators; ECLIPSE Study Investigators. Pulmonary arterial enlargement and acute exacerbations of COPD. N Engl J Med. 2012;367(10):913-921.
- Hurdman J, Condliffe R, Elliot CA, et al. Pulmonary hypertension in COPD: results from the ASPIRE registry. Eur Respir J. 2013;41(6):1292-1301.
- Hoeper MM, Andreas S, Bastian A, et al. Pulmonary hypertension due to chronic lung disease: updated Recommendations of the Cologne Consensus Conference 2011. Int J Cardiol. 2011;154(suppl 1):S45-S53.
- 24. Kessler R, Faller M, Weitzenblum E, et al. "Natural history" of pulmonary hypertension in a series of 131 patients with chronic obstructive lung disease. Am J Respir Crit Care Med. 2001;164(2):219-224.
- 25. Weitzenblum E, Hirth C, Ducolone A, Mirhom R, Rasaholinjanahary J, Ehrhart M. Prognostic value of pulmonary artery pressure in chronic obstructive pulmonary disease. *Thorax*. 1981;36(10):752-758.
- Stevens GR, Fida N, Sanz J. Computed tomography and cardiac magnetic resonance imaging in pulmonary hypertension. *Prog Cardiovasc Dis.* 2012;55(2):161-171.
- Fisher MR, Criner GJ, Fishman AP, et al; NETT Research Group. Estimating pulmonary artery pressures by echocardiography in patients with emphysema. *Eur Respir J.* 2007; 30(5):914-921.
- Come CE, Divo MJ, San José Estépar R, et al; NETT Research Group. Lung deflation and oxygen pulse in COPD: results from the NETT randomized trial. Respir Med. 2012;106(1): 109-119.
- Weitzenblum E, Apprill M, Oswald M, Chaouat A, Imbs JL. Pulmonary hemodynamics in patients with chronic obstructive pulmonary disease before and during an episode of peripheral edema. *Chest.* 1994;105(5):1377-1382.
- Scharf SM, Iqbal M, Keller C, Criner G, Lee S, Fessler HE;
 National Emphysema Treatment Trial (NETT) Group. Hemodynamic characterization of patients with severe emphysema.
 Am J Respir Crit Care Med. 2002;166(3):314-322.
- Bach DS, Curtis JL, Christensen PJ, et al. Preoperative echocardiographic evaluation of patients referred for lung volume reduction surgery. *Chest.* 1998;114(4):972-980.
- 32. Burgess MI, Mogulkoc N, Bright-Thomas RJ, Bishop P, Egan JJ, Ray SG. Comparison of echocardiographic markers of right ventricular function in determining prognosis in chronic pulmonary disease. *J Am Soc Echocardiogr*. 2002; 15(6):633-639.
- 33. Fisher MR, Forfia PR, Chamera E, et al. Accuracy of Doppler echocardiography in the hemodynamic assessment of

journal.publications.chestnet.org CHEST / 145 / 4 / APRIL 2014 831

- pulmonary hypertension. Am J Respir Crit Care Med. 2009; 179(7):615-621.
- Jardim C, Rochitte CE, Humbert M, et al. Pulmonary artery distensibility in pulmonary arterial hypertension: an MRI pilot study. Eur Respir J. 2007;29(3):476-481.
- San Jose Estepar R, Kinney GL, Black-Shinn JL, et al. Computed tomographic measures of pulmonary vascular morphology in smokers and their clinical implications. Am J Respir Crit Care Med. 2013;188(2):231-239.
- Zakynthinos E, Daniil Z, Papanikolaou J, Makris D. Pulmonary hypertension in COPD: pathophysiology and therapeutic targets. Curr Drug Targets. 2011;12(4):501-513.
- Naeije R, Barberà JA. Pulmonary hypertension associated with COPD. Crit Care. 2001;5(6):286-289.
- Biernacki W, Flenley DC, Muir AL, MacNee W. Pulmonary hypertension and right ventricular function in patients with COPD. Chest. 1988;94(6):1169-1175.

- Chan AL, Juarez MM, Shelton DK, et al. Novel computed tomographic chest metrics to detect pulmonary hypertension. BMC Med Imaging. 2011;11:7.
- 40. Tan RT, Kuzo R, Goodman LR, Siegel R, Haasler GB, Presberg KW; Medical College of Wisconsin Lung Transplant Group. Utility of CT scan evaluation for predicting pulmonary hypertension in patients with parenchymal lung disease. Chest. 1998;113(5):1250-1256.
- 41. Haimovici JB, Trotman-Dickenson B, Halpern EF, et al; Massachusetts General Hospital Lung Transplantation Program. Relationship between pulmonary artery diameter at computed tomography and pulmonary artery pressures at right-sided heart catheterization. Acad Radiol. 1997;4(5):327-334.
- 42. Zisman DA, Karlamangla AS, Ross DJ, et al. High-resolution chest CT findings do not predict the presence of pulmonary hypertension in advanced idiopathic pulmonary fibrosis. *Chest.* 2007;132(3):773-779.