



Heart Failure and Cardiomyopathies

HEART RATE VARIABILITY DURING MENTAL STRESS PREDICTS HEART FAILURE OUTCOMES IN PATIENTS WITH STABLE CORONARY ARTERY DISEASE

Poster Contributions
Poster Hall_Hall F
Saturday, March 4, 2023, 11:45 a.m.-12:30 p.m.

Session Title: Heart Failure and Cardiomyopathies: Clinical Science 3
Abstract Category: 08. Heart Failure and Cardiomyopathies: Clinical Science
Presentation Number: 1279-165

Authors: *Maggie Wang, Chang Liu, Anish Shah, Yi-An Ko, Yan Sun, Rachel J. Lampert, Mariana Garcia, Kasra Moazzami, Zakaria Almuwaqqat, Laura Ward, Samaah Sullivan, Paolo Raggi, Doug Bremner, Arshed A. Quyyumi, Viola Vaccarino, Alanna A. Morris, Amit J. Shah, Emory University School of Medicine, Division of Cardiology, Atlanta, GA, USA*

Background: Heart failure (HF) is a common complication in patients with coronary artery disease (CAD) and leads to poor quality of life and increased mortality. Mental stress is recognized as a risk factor for HF, but the underlying mechanism is unclear. We predict that stress-induced autonomic dysfunction is associated with worse HF outcomes.

Methods: We examined 753 participants with stable CAD at an university-affiliated healthcare system from 2011 to 2016. Mental stress was evoked using a public speaking task and high-frequency heart rate variability (HFR HRV, signifying vagal withdrawal) was measured via Holter monitoring at rest and during mental stress. The primary outcome was a composite of acute (incident) and acute on chronic (worsening) HF in the inpatient or outpatient setting. Fine-Gray subdistribution hazard models were used.

Results: Participants had a mean age of 58 years, 35% were female, and 43% were Black. The median follow-up time was 5.66 years. In fully adjusted models for age, sex, race, and comorbidities, higher HFR HRV during mental stress was associated with a decreased risk of composite HF events (HR 0.80 [95% CI 0.73-0.87] per standard deviation increase, $P < .0001$). Stronger associations were observed with HF with reduced ejection fraction than HF with preserved ejection fraction, especially in fully adjusted models.

Conclusion: Stress-induced autonomic dysfunction predicts HF outcomes in patients with CAD and may be an independent mechanism in HF disease progression.

Table 1: Relationship between High-Frequency Heart Rate Variability and Heart Failure Events

Outcomes (n)	Stress High-Frequency HRV			
	Model 1		Model 2	
	SHR (95% CI)	P-value	SHR (95% CI)	P-value
HFrEF* (n=44)	0.76 (0.71-0.81)	$P < .0001$	0.77 (0.69-0.86)	$P < .0001$
HFpEF* (n=28)	0.82 (0.71-0.95)	$P = .01$	0.91 (0.75-1.12)	$P = .39$
Composite (n=76)	0.76 (0.70-0.82)	$P < .0001$	0.80 (0.73-0.87)	$P < .0001$
Outcomes (n)	Rest High-Frequency HRV			
	Model 1		Model 2	
	SHR (95% CI)	P-value	SHR (95% CI)	P-value
HFrEF* (n=44)	0.87 (0.77-0.99)	$P = .04$	0.94 (0.78-1.14)	$P = .54$
HFpEF* (n=28)	0.87 (0.77-0.99)	$P = .04$	0.94 (0.77-1.14)	$P = .52$
Composite (n=76)	0.87 (0.78-0.97)	$P = .01$	0.94 (0.82-1.09)	$P = .43$

*Four participants were excluded from sensitivity analyses due to unknown ejection fraction or moderately reduced ejection fraction.

Abbreviations: SHR, subdistribution hazard ratio; HRV, heart rate variability; HFrEF, heart failure with reduced ejection fraction; HFpEF, heart failure with preserved ejection fraction.

Model 1: Adjusted for age, sex, and race.

Model 2: Adjusted for age, sex, race, active smoking, myocardial infarction, coronary artery bypass graft, percutaneous transluminal coronary angioplasty, hypertension, hyperlipidemia, diabetes mellitus, and body mass index.