Question 1 of 8

A 45-year-old man has crushing chest pain for 30 minutes. ECG shows ST elevation in leads II, III, aVF. Which artery is most likely occluded?

A. Left anterior descending 8%
B. Right coronary artery 71%
C. Left circumflex 12%
D. Left main stem 6%
E. Posterior descending only 3%

Explanation

The correct answer is B.
Inferior ST elevation (II, III, aVF) points to the right coronary artery.
The left anterior descending artery supplies the anterior wall.

Question 2 of 8

Which electrolyte abnormality produces peaked T waves?

A. Hypokalaemia 5%
B. Hyperkalaemia 88%
C. Hypocalcaemia 4%
D. Hypernatraemia 2%
E. Hypomagnesaemia 1%

Explanation

The correct answer is B.
Hyperkalaemia causes peaked T waves, then wide QRS and a sine-wave pattern.
Hypokalaemia gives U waves.

Question 3 of 8

First-line drug for an acute asthma attack in a child with wheeze?

A. Oral prednisolone 20%
B. Inhaled salbutamol 70%
C. IV magnesium 3%
D. Nebulised ipratropium 6%
E. Oral montelukast 1%

Explanation

The correct answer is B.
Inhaled salbutamol is first line. Steroids follow if the response is poor.

Question 4 of 8

Which organism most commonly causes community-acquired pneumonia?

A. Staphylococcus aureus 4%
B. Streptococcus pneumoniae 80%
C. Klebsiella 3%
D. Legionella 5%
E. Mycoplasma 8%

Explanation

The correct answer is B.
Streptococcus pneumoniae is the commonest cause of community-acquired pneumonia.

Question 5 of 8

Best initial test for suspected hypothyroidism?

A. TSH 90%
B. Free T3 4%
C. Thyroid antibodies 3%
D. Ultrasound neck 2%
E. Radioiodine uptake 1%

Explanation

The correct answer is A.
TSH is the best initial test. Free T4 is added when TSH is abnormal.

Question 6 of 8

Which vitamin deficiency causes pellagra?

A. Thiamine 9%
B. Riboflavin 6%
C. Niacin 75%
D. Pyridoxine 6%
E. Cobalamin 4%

Explanation

The correct answer is C.
Niacin deficiency causes dermatitis, diarrhoea and dementia.

Question 7 of 8

Antidote for paracetamol overdose?

A. Naloxone 1%
B. Flumazenil 2%
C. N-acetylcysteine 95%
D. Atropine 1%
E. Pralidoxime 1%

Explanation

The correct answer is C.
N-acetylcysteine replenishes glutathione and prevents hepatotoxicity.

Question 8 of 8

Most common site of a carcinoid tumour in the gut?

A. Stomach 10%
B. Appendix 55%
C. Colon 20%
D. Oesophagus 5%
E. Duodenum 10%

Explanation

The correct answer is B.
The appendix and small bowel are the commonest sites of carcinoid tumours.