
Assumptions of Physics
Technical Brief
More technical briefs at https://assumptionsofphysics.org/resources/briefs



Quantum pure states are not enough

Gabriele Carcassi for the AoP collaboration

Category: Quantum mechanics - Tags: Quantum statistical mechanics, Convex spaces

Abstract

We show that the space of quantum pure states is not enough to uniquely recover the space of quantum ensembles, and therefore it is not enough to characterize quantum mechanics. The space of statistical ensembles, on the other hand, is enough to distinguish between the two.

1 Introduction

While most efforts within the foundations of quantum mechanics focus on pure states, we show they are not enough to characterize quantum mechanics. It is the space of statistical ensemble that marks the difference and, therefore, statistical mechanics plays a crucial role in the difference between the two theories.

2 Conditions

Since we want to compare the classical and quantum state space structures layer by layer, we break them up along similar lines: states and the convex structure of statistical ensembles.

The first two conditions just define the states, the full descriptions of the respective systems.

Condition CP (Classical state space only)**.** *The state space of a classical system is represented by a symplectic manifold* (X, ω)*.*

Condition QP-RAY (Quantum state space only)**.** *The state space of a quantum system is represented by the projective space* $\mathcal{P}(\mathcal{H})$ *of a complex Hilbert space* $\mathcal{H}$*.*

The next conditions specify the ensemble spaces and identify convex combinations with statistical mixtures.

Condition CE (Classical ensemble space)**.** *The statistical ensembles of a classical system are represented by probability measures over its state space. Statistical mixtures are represented by convex combinations of these measures.*

Condition QE-DENS (Quantum ensemble space)**.** *The statistical ensembles of a quantum system are represented by density operators* $\mathcal{D}(\mathcal{H})$*. Statistical mixtures are represented by convex combinations of density operators.*

Finally, we state the exclusivity condition whose relationship to these layers we want to investigate.

Condition CQE (Classical–quantum exclusivity)**.** *A non-trivial system, one that allows more than one possible state, cannot admit both a classical and a quantum representation.*

3 Theorems

The first theorem shows that, if we limit the mathematical representation to the state space definition, systems can be represented both by quantum Hilbert spaces and classical symplectic spaces.

Theorem 1. *Over CP+QP-RAY, CQE is not guaranteed.*

Proof. Recall that finite-dimensional complex projective spaces are symplectic spaces when equipped with the Fubini–Study symplectic form. Consider, then, the state space for a fixed-magnitude magnetic dipole, described by its orientation. This will be represented by a two-sphere, which is both a symplectic manifold and a complex projective line. Similarly, consider the state space of a qubit. This will be represented by a Bloch sphere, which is both a complex projective line and a symplectic manifold. Both these systems are non-trivial. Moreover, over CP+QP-RAY, both systems can be represented with both a classical and a quantum representation. This means that, over CP+QP-RAY, CQE is not guaranteed. □

The second theorem shows that, if we include ensembles, the mathematical representation of classical and quantum systems diverge.

Theorem 2. *Over CP+CE+QP-RAY+QE-DENS, CQE is guaranteed.*

Proof. For a system S to violate CQE over CP+CE+QP-RAY+QE-DENS its state space X must first allow to be represented as both a symplectic manifold and a complex projective space. Suppose S has such a property. Let $|\psi_0\rangle, |\psi_1\rangle \in \mathcal{H}$ be two normalized orthogonal state vectors in terms of the quantum representation, and $|\phi_\pm\rangle = \sqrt{\frac{1}{2}}(|\psi_0\rangle \pm |\psi_1\rangle)$. Consider the equal mixtures of the respective pairs in terms of density operators. We have $\frac{1}{2}\rho_{\psi_0} + \frac{1}{2}\rho_{\psi_1} = \frac{1}{2}\rho_{\phi_+} + \frac{1}{2}\rho_{\phi_-}$. Now consider the equal mixture of the respective pairs in terms of probability measures. We have $\frac{1}{2}\delta_{\psi_0} + \frac{1}{2}\delta_{\psi_1} \neq \frac{1}{2}\delta_{\phi_+} + \frac{1}{2}\delta_{\phi_-}$. That is, density operators allows multiple decompositions in terms of pure states while classical probability distributions do not. Given that, physically, a statistical mixture has a well-defined result, it cannot be represented in two inconsistent ways. Therefore, CP+CE+QP-RAY+QE-DENS, CQE is guaranteed. □

Insight 3. *The space of statistical ensembles is necessary to distinguish between classical and quantum spaces.*

Lastly, we can show that each quantum ensemble is an equivalence class of classical distributions over the pure states.

Theorem 4. *Over QE-DENS, let $M^1(P(\mathcal{H}))$ be the set of probability measures, non necessarily absolutely continuous, over the states. Then there exists an equivalence relationship $\sim \subseteq M^1(P(\mathcal{H})) \times M^1(P(\mathcal{H}))$ such that $D(\mathcal{H}) \equiv M^1(P(\mathcal{H}))_{/\sim}$.*

Proof. Note that $D(\mathcal{H})$ is a convex set with $P(\mathcal{H})$ as extreme points. Therefore, for every probability measure $\mu \in M^1(P(\mathcal{H}))$, there exists an element $\rho \in D(\mathcal{H})$ represented by μ in the sense of Choquet theory. That is, for every continuous linear functional $f : D(\mathcal{H}) \to \mathbb{R}$, $f(\rho) = \int f(\psi) d\mu$. Let $r : M^1(P(\mathcal{H})) \to D(\mathcal{H})$ be the map between the probability measure and the ensemble it represents. Define $\mu_1 \sim \mu_2$ if $r(\mu_1) = r(\mu_2)$. Then $D(\mathcal{H}) \equiv M^1(P(\mathcal{H}))_{/\sim}$. □

The equivalence class keeps track of all possible decompositions of mixed ensembles in terms of pure ensembles. Note that r defined about is affine, so any affine function of quantum ensembles is also an affine functional of the probability measures over quantum states. The converse is not true, and those are exactly the random variables that would distinguish between different possible preparations of the same ensemble.

4 Conclusion

Quantum mechanics cannot be understood by simply looking at pure states, their properties and their evolution. On the other hand, since the difference in the convex space of ensembles is enough to distinguish between the two theories, understanding the difference between classical and quantum mechanics requires understanding the difference between their statistical counterparts.