Assumptions of Physics
Technical Brief
More technical briefs at https://assumptionsofphysics.org/resources/briefs

Non-linear representations in quantum mechanics

Gabriele Carcassi for the AoP collaboration

Category: Quantum mechanics - Tags: Hilbert space, Projective space, collinear maps

Abstract

The linear representation of quantum theory in terms of vector spaces is not a physical requirement but a convenient mathematical representation. We show this by providing a non-linear map that preserves the Born rule, and therefore all the physics.

1 Introduction

It is often stated that the linearity of quantum mechanics is a critical part of the theory. That is, that physical concepts, like observables and time evolution, must be represented with linear structures. Here we show that this linearity is not actually a physical requirement, and one can, in fact, represent quantum theory with non-linear objects. This means that linearity is more a convenient choice of representation than a physically meaningful requirement.

2 Mathematical prerequisites

First, we construct a non-linear map on a two-dimensional complex Hilbert space that preserves the projective space and its geometry.

Proposition 1. *Let $\mathcal{H}$ be a two-dimensional complex Hilbert space. Let $v, w \in \mathcal{H}$ be two orthonormal vectors. Let $m : \mathcal{H} \to \mathcal{H}$ be the map defined by $m(0) = 0$ and, on non-zero vectors, by*

$$m(av + bw) = e^{\imath\theta \frac{|a|^2}{|a|^2+|b|^2}} (av + bw), \tag{2}$$

with $\theta \in (0, 2\pi)$. Then m is a bijective non-linear map that preserves rays and the Fubini–Study geodesic distance. That is, for all $x, y \in \mathcal{H} \setminus \{0\}$

$$m(v + w) \neq m(v) + m(w) \qquad [m(x)] = [x], \qquad \arccos \frac{|\langle m(x), m(y)\rangle|}{\|m(x)\|\|m(y)\|} = \arccos \frac{|\langle x, y\rangle|}{\|x\|\|y\|}.$$

The inverse is given by the same formula with θ replaced by $-\theta$.

Proof. First, let us show that the map preserves rays. For $x = av + bw \neq 0$, write

$$r(x) = \frac{|a|^2}{|a|^2 + |b|^2}, \qquad m(x) = e^{\imath\theta r(x)} x.$$

Since $m(x)$ differs from x only by a phase factor, they belong to the same ray. Therefore $[m(x)] = [x]$.

The Fubini–Study geodesic distance is also preserved. For non-zero $x, y \in \mathcal{H}$,

$$\begin{aligned} \langle m(x), m(y)\rangle &= e^{-\imath\theta r(x)} e^{\imath\theta r(y)} \langle x, y\rangle = e^{\imath\theta(r(y)-r(x))} \langle x, y\rangle, \\ \|m(x)\| &= |e^{\imath\theta r(x)}|\, \|x\| = \|x\|. \end{aligned}$$

The same norm identity holds for y. Therefore,

$$\frac{|\langle m(x), m(y)\rangle|}{\|m(x)\|\|m(y)\|} = \frac{|e^{\imath\theta(r(y)-r(x))}|\,|\langle x, y\rangle|}{\|x\|\|y\|} = \frac{|\langle x, y\rangle|}{\|x\|\|y\|}.$$

Taking the arccosine of both sides gives the equality of distances.

To establish bijectivity, note that for every $t \in \mathbb{R}$,

$$r(e^{\imath t}x) = \frac{|e^{\imath t}a|^2}{|e^{\imath t}a|^2 + |e^{\imath t}b|^2} = \frac{|e^{\imath t}|^2|a|^2}{|e^{\imath t}|^2(|a|^2 + |b|^2)} = r(x).$$

Define $n(0) = 0$ and $n(x) = e^{-\imath\theta r(x)}x$ for $x \neq 0$. Thus $r(n(x)) = r(m(x)) = r(x)$, and

$$\begin{aligned} m(n(x)) &= e^{\imath\theta r(n(x))} e^{-\imath\theta r(x)} x = e^{\imath\theta r(x)} e^{-\imath\theta r(x)} x = x, \\ n(m(x)) &= e^{-\imath\theta r(m(x))} e^{\imath\theta r(x)} x = e^{-\imath\theta r(x)} e^{\imath\theta r(x)} x = x. \end{aligned}$$

Both identities also hold at zero. Therefore m is bijective, with inverse n, given by replacing θ with $-\theta$.

Finally,

$$\begin{aligned} m(v) &= e^{\imath\theta} v, & m(w) &= w, \\ m(v + w) &= e^{\imath\theta/2} v + e^{\imath\theta/2} w, & m(v) + m(w) &= e^{\imath\theta} v + w. \end{aligned}$$

Since $0 < \theta/2 < \pi$, we have $e^{\imath\theta/2} \neq 1$. The coefficients of w therefore differ, so linear independence gives $m(v + w) \neq m(v) + m(w)$. Hence m is non-linear. □

Corollary 3. *Let m be the map in the preceding proposition. A linear operator $A : \mathcal{H} \to \mathcal{H}$ transported through m,*

$$\widetilde{A} = m \circ A \circ m^{-1},$$

need not be linear with respect to the ordinary vector-space structure on $\mathcal{H}$, even when A is both self-adjoint and unitary.

Proof. Define $A(av + bw) = bv + aw$. In the orthonormal basis (v, w),

$$A = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \qquad A^* = A, \qquad A^*A = A^2 = I.$$

Thus A is linear, self-adjoint and unitary. Using the inverse formula for m, we obtain

$$\begin{aligned}
\widetilde{A}(v) &= m\big(A(e^{-\imath\theta}v)\big) = m(e^{-\imath\theta}w) = e^{-\imath\theta}w,\\
\widetilde{A}(w) &= m(A(w)) = m(v) = e^{\imath\theta}v,\\
\widetilde{A}(v+w) &= m\big(A(e^{-\imath\theta/2}(v+w))\big)\\
&= m(e^{-\imath\theta/2}(v+w))\\
&= e^{\imath\theta/2}e^{-\imath\theta/2}(v+w) = v+w.
\end{aligned}$$

However,

$$\widetilde{A}(v)+\widetilde{A}(w) = e^{\imath\theta}v + e^{-\imath\theta}w.$$

Since $0<\theta<2\pi$, we have $e^{\imath\theta}\neq 1$. The coefficients of v therefore differ, so linear independence gives $\widetilde{A}(v+w)\neq\widetilde{A}(v)+\widetilde{A}(w)$. Hence $\widetilde{A}$ is non-linear. □

3 Conditions

We can sum up the definitions of a state space in quantum mechanics with the following condition.

Condition QST:H (Quantum states)**.** *The state space of a quantum system is represented by the projective space $\mathcal{P}(\mathcal{H})$ of a separable complex Hilbert space $\mathcal{H}$. The ensemble space is represented by the density operators $\mathcal{D}(\mathcal{H})$. The conditional probability is given by the Born rule $p(\psi|\phi)=\frac{\langle\phi|\psi\rangle\langle\psi|\phi\rangle}{\langle\psi|\psi\rangle\langle\phi|\phi\rangle}$. The entropy is given by the von Neumann entropy $S(\rho)=-\operatorname{tr}(\rho\log\rho)$.*

The following condition requires other physical objects to be represented by linear structures and operations on $\mathcal{H}$.

Condition QMLIN (Linear vector space representation)**.** *Quantum superpositions, observables, and time evolution are represented by linear structures over the Hilbert space associated to the state space. That is, quantum superpositions are represented by vector addition, observables by self-adjoint operators and system evolution by unitary operators.*

4 Theorem

Theorem 4 (Linearity is a choice)**.** *Over QST:H, QMLIN is neither guaranteed nor ruled out.*

Proof. To show that QMLIN is neither guaranteed nor ruled out over QST:H, it suffices to provide two valid representations for a single quantum system that satisfy QST:H, one satisfying QMLIN and one not. Consider a spin $\frac{1}{2}$ system represented in the usual way by a two-dimensional Hilbert space $\mathcal{H}$. That satisfies both QST:H and QMLIN, therefore QMLIN is not ruled out over QST:H.

Now consider m as defined by 1 with $v=|z^+\rangle$ and $w=|z^-\rangle$. Note that $\frac{|a|^2}{|a|^2+|b|^2}=p(z^+|\psi)$ with $|\psi\rangle = a|z^+\rangle+b|z^-\rangle$. This means that the phase shift introduced by the map is proportional to the probability of measuring z^+. Since m is a bijection, physical representatives on $\mathcal{H}$ transported through m will be physical representatives on $m(\mathcal{H})$. Since m preserves rays, the state representation is unchanged. Similarly, density operators will be left unchanged together with their entropy. The




Figure 1: Probability is vertical height on the Bloch sphere. The map m introduces a phase shift on the vectors that is proportional to the vertical height, while leaving each pure-state point unchanged on the Bloch sphere.

Born rule is left unchanged as it is the cosine square of the Fubini–Study distance. Therefore, QST:H remains valid on $m(\mathcal{H})$.

However, by 1 and 3, linear combinations and linear operators do not correspond, in general, to transported linear combinations and linear operators. Therefore, quantum superposition in one representation will not, in general, map to the same linear combinations. Similarly, observables will not be, in general, represented by self-adjoint operators and time evolution will not be, in general, represented by unitary maps.

We therefore have a representation where QST:H holds but superpositions, observables and time evolution are not represented by the ordinary linear structures. Therefore QMLIN is not guaranteed over QST:H. □

5 Conclusion

We have shown that non-linear maps that do not conserve the inner product but preserve the Born rule, and thus all the physics, exist and therefore the linear representation amounts to a choice of representation. We should also note that alternative representations exist that are not even tied to vector spaces: the Madelung equations (also known as the equations of quantum hydrodynamics) provide such a representation.